

## Washington University School of Medicine Digital Commons@Becker

---

### Open Access Publications

---

2018

# Genome sequence of *Acetomicrobium hydrogeniformans* OS1

Lauren E. Cook

*University of California, Los Angeles*

Spencer S. Gang

*University of California, Los Angeles*

Alicia Ihlan

*University of California, Los Angeles*

Matthew Maune

*University of Oklahoma*

Ralph S. Tanner

*University of Oklahoma*

*See next page for additional authors*

Follow this and additional works at: [https://digitalcommons.wustl.edu/open\\_access\\_pubs](https://digitalcommons.wustl.edu/open_access_pubs)

---

### Recommended Citation

Cook, Lauren E.; Gang, Spencer S.; Ihlan, Alicia; Maune, Matthew; Tanner, Ralph S.; McInerney, Michael J.; Weinstock, George; Lobos, Elizabeth A.; and Gunsalus, Robert P., "Genome sequence of *Acetomicrobium hydrogeniformans* OS1." *Genome Announcements*.6,26. e00581-18. (2018).

[https://digitalcommons.wustl.edu/open\\_access\\_pubs/7042](https://digitalcommons.wustl.edu/open_access_pubs/7042)

This Open Access Publication is brought to you for free and open access by Digital Commons@Becker. It has been accepted for inclusion in Open Access Publications by an authorized administrator of Digital Commons@Becker. For more information, please contact [engeszer@wustl.edu](mailto:engeszer@wustl.edu).

---

**Authors**

Lauren E. Cook, Spencer S. Gang, Alicia Ihlán, Matthew Maune, Ralph S. Tanner, Michael J. McInerney, George Weinstock, Elizabeth A. Lobos, and Robert P. Gunsalus



# Genome Sequence of *Acetomicrobium hydrogeniformans* OS1

Lauren E. Cook,<sup>a</sup> Spencer S. Gang,<sup>a</sup> Alicia Ihlan,<sup>a</sup> Matthew Maune,<sup>b</sup> Ralph S. Tanner,<sup>b</sup>  Michael J. McInerney,<sup>b</sup> George Weinstock,<sup>c</sup> Elizabeth A. Lobos,<sup>c</sup> Robert P. Gunsalus<sup>a,d</sup>

<sup>a</sup>Department of Microbiology, Immunology, and Molecular Genetics, University of California, Los Angeles, California, USA

<sup>b</sup>Department of Microbiology and Plant Biology, University of Oklahoma, Norman, Oklahoma, USA

<sup>c</sup>The Genome Institute, Washington University School of Medicine, St. Louis, Missouri, USA

<sup>d</sup>UCLA DOE Institute, University of California, Los Angeles, California, USA

**ABSTRACT** *Acetomicrobium hydrogeniformans*, an obligate anaerobe of the phylum *Synergistetes*, was isolated from oil production water. It has the unusual ability to produce almost 4 molecules H<sub>2</sub>/molecule glucose. The draft genome of *A. hydrogeniformans* OS1 (DSM 22491<sup>T</sup>) is 2,123,925 bp, with 2,068 coding sequences and 60 RNA genes.

*Acetomicrobium hydrogeniformans* strain OS1<sup>T</sup>, until recently known as *Anaerobaculum hydrogeniformans* (1), of the phylum *Synergistetes*, is capable of producing almost four molecules of hydrogen per glucose molecule, the theoretical maximum (2, 3). It ferments other substrates, including amino acids, dicarboxylic acids, and other sugars, and can respire using several sulfur compounds. NaCl is an absolute requirement for growth, which suggests a sodium-based energy strategy (2, 4, 5). Although it can reduce elemental sulfur, thiosulfate, and L-cysteine to sulfide, the pathways are unknown. The genome is currently divided among eight scaffolds, one of which contains 2,103,414 bp and seven of which are small scaffolds ranging from 507 bp to 7,345 bp. The genome lacks genes needed for respiration of sulfate and nitrate and genes for *b*- and *c*-type cytochromes. The metabolic strategy appears to be limited primarily to fermentation of simple sugars, amino acids, and certain dicarboxylic acids in addition to an as-yet undescribed respiratory mode for elemental sulfur or thiosulfate. Despite the ability to ferment sugars, the Embden-Meyerhof-Parnas, pentose phosphate, and Entner-Doudoroff pathways appear to be incomplete. It also lacks a complete set of trichloroacetic acid (TCA) genes. OS1<sup>T</sup> possesses two gene clusters with hydrogenase-related functions to form molecular hydrogen. It also possesses genes for two putative formate dehydrogenases and a formate hydrogenlyase complex. It has multiple genes for stress adaptation, including those for motility, pili, osmotic adaptation, heavy-metal resistance, multidrug export, and toxin/antitoxin synthesis. It has a 21-gene cassette for the synthesis of a carboxysome-like shell structure. Although motility was not previously reported (2), motility can be observed in wet mounts of cultures, and negative-stained cells exhibit polar flagella (R. S. Tanner, personal communication). The genome contains a complete inventory of genes needed to make flagella and multiple genes for chemoreception and chemotaxis. The genome also contains a complement of 10 *pil/pul* genes needed for synthesis and assembly of a type IV pilus apparatus. It has a relatively small number of primary transcription factors (43 genes) and six predicted two-component regulatory systems, including the aforementioned chemotaxis signaling systems. Thus, OS1 appears to have adopted a minimal regulatory strategy, although the substrate range is considerable with respect to the variety of sugars, amino acids, and dicarboxylates used (2). Little is known about translational or posttranslational control operating in this or other *Synergistetes* species. Strikingly, the genome

Received 30 May 2018 Accepted 30 May 2018 Published 28 June 2018

**Citation** Cook LE, Gang SS, Ihlan A, Maune M, Tanner RS, McInerney MJ, Weinstock G, Lobos EA, Gunsalus RP. 2018. Genome sequence of *Acetomicrobium hydrogeniformans* OS1. *Genome Announc* 6:e00581-18. <https://doi.org/10.1128/genomeA.00581-18>.

**Copyright** © 2018 Cook et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Robert P. Gunsalus, [rbg@microbio.ucla.edu](mailto:rbg@microbio.ucla.edu).

lacks genes for the prototypical bacterial F<sub>o</sub>F<sub>1</sub>-type ATP synthase but possesses instead three clusters of genes encoding either archaeal-like or vacuolar-like A/V ATP synthases. It also has other archaeal-like annotated genes, including those acting in glycolysis; of note, we found four putative candidates for glyceraldehyde-3-phosphate oxidoreductase, a ferredoxin-utilizing enzyme with archaeal properties. Other genes with archaeal properties include those related to formyl methanofuran dehydrogenase A, B, C, D, and E subunits, methanyltetrahydromethanopterin cyclohydrolase; and Formylmethanofuran-tetrahydromethanopterin *N*-formyltransferase. These genes may have been acquired by lateral gene transfer (6–8).

**Accession number(s).** This whole-genome shotgun project has been deposited in GenBank under the accession no. [ACJX00000000](https://www.ncbi.nlm.nih.gov/nuccore/ACJX00000000).

## ACKNOWLEDGMENTS

These sequence data were produced by the US Department of Energy Joint Genome Institute (<http://www.jgi.doe.gov>) in collaboration with the user community.

This work was supported by Department of Energy contract DE-FG02-08ER64689 (to R.P.G., R.S.T., and M.J.M.), UCLA-DOE Institute award DE-FC02-02ER63421 (to R.P.G.), and National Science Foundation grant NSF 1515843 (to R.P.G. and M.J.M.).

## REFERENCES

- Hania WB, Bouanane-Darenfed A, Cayol J-L, Ollivier B, Fardeau M-L. 2016. Reclassification of *Anaerobaculum mobile*, *Anaerobaculum thermoterrenum*, *Anaerobaculum hydrogeniformans* as *Acetomicrobium mobile* comb. nov., *Acetomicrobium thermoterrenum* comb. nov. and *Acetomicrobium hydrogeniformans* comb. nov., respectively, and emendation of the genus *Acetomicrobium*. *Int J Syst Evol Microbiol* 66:1506–1509. <https://doi.org/10.1099/ijsem.0.000910>.
- Maune MW, Tanner RS. 2012. Description of *Anaerobaculum hydrogeniformans* sp. nov., an anaerobe that produces hydrogen from glucose, and emended description of the genus *Anaerobaculum*. *Int J Syst Evol Microbiol* 62:832–838. <https://doi.org/10.1099/ijms.0.024349-0>.
- Thauer RK, Jungermann K, Decker K. 1977. Energy conservation in chemotrophic anaerobic bacteria. *Bacteriol Rev* 41:100–180.
- Menes RJ, Muxi L. 2002. *Anaerobaculum mobile* sp. nov., a novel anaerobic, moderately thermophilic, peptide-fermenting bacterium that uses crotonate as an electron acceptor, and emended description of the genus *Anaerobaculum*. *Int J Syst Evol Microbiol* 52:157–164. <https://doi.org/10.1099/00207713-52-1-157>.
- Buss KA, Sanders DA, Hasson MS, Ingram-Smith C, Ferry JG. 1997. Crystallization of acetate kinase from *Methanosarcina thermophila* and prediction of its fold. *Protein Sci* 6:2659–2662. <https://doi.org/10.1002/pro.5560061222>.
- Garcia-Vallvé S, Romeu A, Palau J. 2000. Horizontal gene transfer in bacterial and archaeal complete genomes. *Genome Res* 10:1719–1725. <https://doi.org/10.1101/gr.130000>.
- Nelson KE, Clayton RA, Gill SR, Gwinn ML, Dodson RJ, Haft DH, Hickey EK, Peterson JD, Nelson WC, Ketchum KA, McDonald L, Utterback TR, Malek JA, Linher KD, Garrett MM, Stewart AM, Cotton MD, Pratt MS, Phillips CA, Richardson D, Heidelberg J, Sutton GG, Fleischmann RD, Eisen JA, White O, Salzberg SL, Smith HO, Venter JC, Fraser CM. 1999. Evidence for lateral gene transfer between Archaea and Bacteria from genome sequence of *Thermotoga maritima*. *Nature* 399:323–329. <https://doi.org/10.1038/20601>.
- Deppenmeier U, Johann A, Hartsch T, Merkl R, Schmitz RA, Martinez-Arias R, Henne A, Wiezer A, Baumer S, Jacobi C, Bruggemann H, Lienard T, Christmann A, Bomeke M, Steckel S, Bhattacharyya A, Lykidis A, Overbeek R, Klenk HP, Gunsalus RP, Fritz HJ, Gottschalk G. 2002. The genome of *Methanosarcina mazei*: evidence for lateral gene transfer between bacteria and archaea. *J Mol Microbiol Biotechnol* 4:453–461.