



12-2017

Depression in stroke patients; a cross sectional study highlighting the association of stroke on age and gender basis.

Saira Abbas

Dow University of Health Sciences, Karachi, Pakistan

Naila Naeem Shahbaz

Civil Hospital Karachi

Muhammad Amir Umer

Civil Hospital Karachi

Sumera Rafat Umer

Civil Hospital Karachi

Asfia Irfan

National Medical Center Karachi

See next page for additional authors

Follow this and additional works at: <https://ecommons.aku.edu/pjns>

 Part of the [Neurology Commons](#)

Recommended Citation

Abbas, Saira; Shahbaz, Naila Naeem; Umer, Muhammad Amir; Umer, Sumera Rafat; Irfan, Asfia; and Anwar, Adnan (2017) "Depression in stroke patients; a cross sectional study highlighting the association of stroke on age and gender basis.," *Pakistan Journal of Neurological Sciences (PJNS)*: Vol. 12 : Iss. 4 , Article 7.
Available at: <https://ecommons.aku.edu/pjns/vol12/iss4/7>

Depression in stroke patients; a cross sectional study highlighting the association of stroke on age and gender basis.

Authors

Saira Abbas, Naila Naeem Shahbaz, Muhammad Amir Umer, Sumera Rafat Umer, Asfia Irfan, and Adnan Anwar

Depression in stroke patients; a cross sectional study highlighting the association of stroke on age and gender basis.

Dr. Saira Abbas¹, Dr. Naila Naeem Shahbaz², Dr. Muhammad Amir Umer³, Dr. Sumera Rafat Umer⁴,
Dr. Asfia Irfan⁵, Dr. Adnan Anwar⁶

¹Assistant Professor, Department of Neurology, Dow University Hospital/DIMC, DUHS

²Chairperson and Head of Department of Neurology, DUHS/Civil Hospital Karachi

³Assistant Professor, Department of Neurology, DUHS/Civil Hospital Karachi

⁴Senior Instructor, Department of Neurology, DUHS/Civil Hospital Karachi

⁵Consultant Neurologist, National Medical Center

⁶Senior lecturer, Department of Physiology, Altibri medical college

Corresponding to: Dr. Saira Abbas, M.B.B.S, F.C.P.S, Assistant Professor, Department of Neurology, Dow University Hospital/DIMC, DUHS, Email : dr.sairaabbas@gmail.com

Date of submission: April 26, 2017 **Date of revision:** June 29, 2017 **Date of acceptance:** July 19, 2017

ABSTRACT

INTRODUCTION

Stroke is the second leading cause of death. Depressive disorder is characterized by period of sad mood and anhedonia occurring for two consecutive weeks. Among stroke survivors, the consequence of physical and psychological changes can be devastating. One of those psychological changes is post-stroke depression (PSD). Stroke itself has debilitating morbidity and superimposed PSD further decreases the quality of life of patients and impairs recovery. This study leads us to know the magnitude of post stroke depression.

OBJECTIVE

To determine the frequency of post stroke depression in patients under treatment at tertiary care hospital Karachi.

METHODS

A cross-sectional study was conducted at the Neurology out-patient department, Civil Hospital, Karachi. A total of 340 patients of both male and female genders, aged between 20-70 years with post stroke duration of greater than three months were included in the study using no-probability purposive sampling. The patients were approached through neurology out-patient department. Informed consent was obtained from all the patients after explanation of the study protocol. Patients with cognitive impairment (modified 6-item Mini-Mental Status score < 3), with focal or diffuse organic brain disorder that may be associated with psychiatric manifestation like intellectual disability, A-V malformation, tumor, intracranial infections, systemic diseases that may predispose them to depression e.g. hypo/hyperthyroidism, SLE or those already diagnosed with depression or prior anti-depressant treatment were excluded from the study. Data was collected by using a predesigned questionnaire. SPSS version 19.0 was used for data entry and analysis. Chi square test was used to assess the association.

RESULTS

The average age of the patients was 38.45 ± 9.48 years. Post stroke depression was observed in 45% cases, out of which 13.2% had mild, 44.1% had moderate and 42.8% had severe depression. Prevalence of post stroke depression was significantly high in patients below 40 years of age and in patients who had ischemic type of stroke. The study results further revealed that patients younger than 40 years were more likely to have post stroke depression than those above 40 years of age ($p=0.009$) and patients who suffered from ischemic stroke were more likely to suffer from depression than those who suffered from hemorrhagic stroke (50% vs. 33.6%, $p=0.005$).

CONCLUSION

Prevalence of post stroke depression was significantly higher in patients with ischemic stroke than those with hemorrhagic stroke. We emphasize the importance of a thorough psychiatric evaluation of post stroke patients, particularly those who have a severe disability and a history of previous depressive episodes.

KEYWORDS

Depression, Ischemic Stroke, Hemorrhagic Stroke

INTRODUCTION

Stroke is the second leading cause of death after myocardial infarction and the most frequent cause of adult disability¹⁻³. In 2010, the people with first stroke were estimated to be 16.9 million, stroke survivors to be 33 million whereas stroke-related deaths to be 5.9 million⁴. Of all strokes, 88% are ischemic, 9% are intracerebral hemorrhagic and 3% are subarachnoid hemorrhagic⁵. Mortality and morbidity from stroke are on the rise. According to WHO, in 2002 the total number of deaths due to cerebrovascular accidents in Pakistan was 78,512. In Brazil, the absolute number of hospitalizations due to stroke ranged between 198,705 and 295,596 per annum between 1994 and 1997, and it was estimated that 25% were recurrent cases⁶. After suffering a stroke, 80% of patients present motor impairment⁷.

Depressive disorder is characterized by period of sad mood and anhedonia (inability to experience the pleasure from normally pleasurable life events such as eating, exercise, social or sexual interaction) occurring for two consecutive weeks as defined by DSM IV criteria for depression⁸. Among stroke survivors, the sequelae of physical and psychological changes can be devastating. One of those psychological changes is post-stroke depression (PSD) which constitutes an important complication of stroke, resulting in increased disability as well as mortality⁹. Morris et al., in 1993 reported that, over a 10-year period, depressed stroke patients were 3.4 more times likely to die than their non-depressed counterparts¹⁰. Depression is thought to have a detrimental effect on stroke recovery through a number of mechanisms. For instance, a depressed patient may be less motivated to participate in stroke rehabilitation because of persistent fatigue or lack of hope. Cognitive impairment may also impede the recovery process, causing non-adherence to treatment schedules, which may lead to increased mortality. PSD is common among both men and women. In 2005, a systematic review of 51 primary articles by Hackett et al., in 2005 estimated the overall frequency of PSD to be 33%¹¹. Stroke itself has debilitating morbidity and superimposed PSD further decreases the quality of life of patients and impairs the recovery.

This study enabled us to know the magnitude of post-stroke depression, so, that departmental protocol could be developed for early screening and referral for treatment of PSD. By early identification and treatment of PSD we could improve the quality of life of our patients, expedite their recovery and make them functional^{12,13}.

METHODS

After seeking ethical approval, a cross-sectional study was conducted at the Neurology out-patient

department, Civil Hospital, Karachi. As per the minimum required sample size, with 95% confidence level, anticipated frequency of PSD of 33%¹¹ and absolute precision of 5%, a total of 340 patients of either gender, aged between 20-70 years with post-stroke duration of greater than three months were included in the study using non-probability purposive sampling. The patients were approached through neurology out-patient department. Informed consent was obtained from all the patients after explanation of the study protocol. Patients with cognitive impairment (modified 6-item Mini-Mental Status score < 3), with focal or diffuse organic brain disorder that may be associated with psychiatric manifestation like intellectual disability A-V malformation, tumor, intracranial infections etc., with systemic diseases that may predispose them to depression e.g. hypo/hyperthyroidism, SLE or who were already diagnosed as having depression or prior antidepressant treatment were excluded from the study. The patients were interviewed by the principal investigator using a pre-designed questionnaire which had questions pertaining to age, gender, post-stroke duration, type of stroke (i.e. Ischemic or Hemorrhagic) and frequency of depression. All the stroke patients were evaluated by Beck Depression Inventory questionnaire for diagnosis of depression. The severity of depression was categorized as mild, with depression score 10-19, moderate, with depression score 20-29 and severe, with depression score ≥ 30 .

The data were analyzed on SPSS version 19. Frequency and percentage were calculated for qualitative variables like gender, type of stroke (ischemic and Hemorrhagic) and frequency of depression whereas mean \pm SD were computed for quantitative variables like age, and post-stroke duration. Variables considered as potential confounders and/or effect modifiers such as age and gender were stratified to find out the effect of these on the outcome. Post-stratification chi-square test was applied for inferential analysis and the significance level was set at 0.05. The duration of study was six months.

RESULTS

The study results showed that the mean age of the patients was 38.45 ± 9.48 years whereas 58% of them were male. The average duration of post-stroke cases was 4.25 ± 2.68 months whereas 68% of the stroke cases were of ischemic type. Prevalence of post-stroke depression was found to be 45% out of which 13.2% were mild, 44.1% were moderate and 42.8% were severe (table 1). The study results further revealed that patients younger than 40 years were more likely to have post-stroke depression (PSD) than those above 40 years of age ($p=0.009$). There was no significant gender difference among prevalence of PSD (43.1% vs. 46.9%, $p=0.49$). Patients who suffered from

ischemic stroke were more likely to suffer from depression than those who suffered from hemorrhagic stroke (50% vs. 33.6%, $p=0.005$) (table 2).

Table 1: Participants Profile

Variable (n=340)	Frequency(%) / Mean \pm S.D.
Age (Years)	38.45 \pm 9.48
Gender	
Male	197(58.0)
Female	143(42.0)
Type of Stroke	
Ischemic	230(68)
Hemorrhagic	110(32)
Post Stroke Depression	
Yes	152(45)
No	188(55)
Severity of Post Stroke Depression ¹	
Mild	20(13.2)
Moderate	67(44.1)
Severe	62(42.7)
Post Stroke Duration (Months)	4.25 \pm 2.68
¹ n=152	

Table 2: A univariate analysis of association between selected study variables and post stroke depression

Variable (n=340)	Post Stroke Yes (n=152) Frequency(%)	Depression No (n=188) Frequency(%)	p
Age Groups			
20-30 Years	45(57.0)	34(43.0)	0.009
31-40 Years	52(49.5)	53(50.5)	
41-60 Years	35(36.8)	60(63.2)	
61-70 Years	20(32.8)	41(67.2)	
Gender			
Male	85(43.1)	112(56.9)	0.49
Female	67(46.9)	76(53.1)	
Type of Stroke			
Ischemic	115(50.0)	115(50.0)	0.005
Hemorrhagic	37(33.6)	73(66.4)	

DISCUSSION

Stroke is a sudden non-convulsive focal neurological deficit produced by insufficiency of blood circulation to the brain¹⁴. In most western countries, 0.2% of the population suffers a stroke every year⁹. In addition to major impact of stroke on patients' physical health, many patients experience emotional disorders

following stroke. The frequent appearance of emotional disorders following stroke has been recognized for many years. Goldstein (1948) found that patients with left hemisphere injury developed depression, irritability and explosive outbursts which he termed the "catastrophic reaction"¹⁵.

Depression is a common neuropsychiatric consequence of stroke affecting approximately 40% of the patients. In addition to the psychosocial stress due to disability, loss of independence, and worsening of quality of life, neurobiological factors such as site of infarcts and brain atrophy have also been proposed to be related to depression after stroke¹⁶. In recent years, post-stroke depression (PSD) has attracted worldwide interest. Literature reports a prevalence rate of major depression of 19.3% among hospitalized patients and 23.3% among outpatient samples¹⁷. Longitudinal studies of stroke patients have shown that about 20% of these patients develop Major and another 20% develop Minor depression during the first year after stroke⁹.

In our study prevalence of post stroke depression was found to be 45% out of which 13.2% were mild, 44.1% were moderate and 42.8% were severe. The average duration of post stroke cases was found to be 4.25 \pm 2.68 months which is higher than those reported for outpatient samples by two reviews i.e. 23.3%¹⁷ and 33%¹⁸. This could be attributed to difference in the method of assessing depression or to population characteristics. The prevalence of post-stroke depression has been reported earlier to be about 30%^{19,20}. An earlier study also reported the prevalence of moderate to severe depression to be 15%²¹. The current study was done at 3 to 6 months post-stroke, whereas that study was done 4 to 8 weeks post stroke. Time lapsed after stroke has been mentioned as a variable in the prevalence of post stroke depression^{19,22}. The exclusion of a number of patients in this study as in other studies, because of severe cognitive or communicative deficits, may contribute to an underestimate of psychiatric morbidity. However, Starkstein and Robinson in 1989 suggested that aphasia does not cause depression, but the two may coexist. As yet, there remains no reliable method of assessing mood disorders for patients with severe comprehensive deficits¹⁵.

Demographic variables are important determinants of post stroke depression. Depressive symptoms were found statistically associated with young age group in our study. Prevalence of post stroke depression (PSD) was significantly high in patients below 40 years of age than those above 40 years of age ($p=0.009$). Our result has similarity with previous studies^{23,24}. However, most of the studies on association between age and post stroke depression show contradictory findings and reveals complex relationship between age and post stroke depression which could be dependent upon several factors.

A significant association was found between post stroke depression and Ischemic stroke type in our study. Prevalence of post stroke depression was significantly higher in patients suffering from ischemic stroke than those suffering from hemorrhagic stroke ($p=0.005$). This finding is in congruence with other studies^{16,25}. An earlier study reported high depression among patients having ischemic stroke in the left hemisphere as compared to those having it in the right hemisphere¹⁶. However, many studies report contradictory results as well^{26,27}. Just as functional depression may arise from several mechanisms, the cause of PSD is likely to represent a mixture of etiologic factors. Socio-demographic, psychological and biological risk factors mediate the relationship between stroke and depression²⁸. The association of depression with worse prognosis in stroke patients lends more support to the hypothesis that psychological rather than neurological factors are the main determinants of post stroke depression²⁹.

Furthermore, 68% of the patients in our study were found to have ischemic whereas 32% were found to have hemorrhagic stroke. Likewise, a number of earlier local studies have also consistently reported ischemic stroke to be more prevalent than hemorrhagic stroke. Marwat MA et al., in 2009 reported the prevalence of ischemic stroke to be 50% whereas that of hemorrhagic stroke to be 29.5%³⁰. Memon FA et al., in 2009 found the prevalence of ischemic stroke to be 62% and of hemorrhagic stroke to be 36% of stroke patients³¹. Lakhair MA et al., in 2008 reported the prevalence of ischemic stroke to be 75% whereas that of hemorrhagic stroke to be 25%³². Memon RA et al., in 2008 also reported the prevalence of ischemic stroke to be 77% whereas that of hemorrhagic stroke to be 23%³³. Likewise, Zafar A et al., in 2007 found the prevalence of ischemic stroke to be 88% whereas that of hemorrhagic stroke to be 12%³⁴. Similarly Khan J et al., in 2006 found ischemic stroke to be present in 71.4% whereas hemorrhagic stroke to be in 28.6% of the participants³⁵.

Limitation: It is acknowledged that the study may have suffered from recall bias, an inherent weakness of a cross-sectional study design. BDI is not designed to be used in stroke patients. How did patients who could not read and write complete this?

Recommendation: We emphasize the importance of a thorough psychiatric evaluation of post stroke patients, particularly those who have a severe disability and a history of previous depressive episodes.

CONCLUSION

The study findings revealed a high prevalence of post stroke depression among patients. The study

results further revealed that patients younger than 40 years were more likely to have post stroke depression than those above 40 years of age whereas patients who suffered from ischemic stroke were more likely to suffer from depression than those who suffered from hemorrhagic stroke.

REFERENCES:

1. Lopez AD, Mathers CD, Ezzati M, Jamison DT, Murray CJ. Global and regional burden of disease and risk factors, 2001: systematic analysis of population health data. *The Lancet*. 2006 Jun 2;367(9524):1747-57.
2. Kolominsky-Rabas PL, Heuschmann PU, Marschall D, Emmert M, Baltzer N, Neundörfer B, Schöffski O, Krobot KJ. Lifetime cost of ischemic stroke in Germany: Results and national projections from a population-based stroke registry. *Stroke*. 2006 May 1;37(5):1179-83.
3. Minnerup J, Seeger FH, Kuhnert K, Diederich K, Schilling M, Dimmeler S, Schäbitz WR. Intracarotid administration of human bone marrow mononuclear cells in rat photothrombotic ischemia. *Experimental & translational stroke medicine*. 2010 Feb 2;2(1):3.
4. Feigin VL, Forouzanfar MH, Krishnamurthi R, Mensah GA, Connor M, Bennett DA, Moran AE, Sacco RL, Anderson L, Truelsen T, O'Donnell M. Global and regional burden of stroke during 1990–2010: findings from the Global Burden of Disease Study 2010. *The Lancet*. 2014 Jan 24;383(9913):245-55.
5. Asfandiyarova N, Kolcheva N, Ryazantsev I, Ryazantsev V. Risk factors for stroke in type 2 diabetes mellitus. *Diabetes and Vascular Disease Research*. 2006 May;3(1):57-60.
6. Yamashita LF, Fukujima MM, Granitoff N, Prado GF. Paciente com acidente vascular cerebral isquêmico já é atendido com mais rapidez no Hospital São Paulo. *Arquivos de Neuro-Psiquiatria*. 2004 Mar 1.
7. Moura RD, Fukujima MM, Aguiar AS, Fontes SV, Dauar RF, Prado GF. Predictive factors for spasticity among ischemic stroke patients. *Arquivos de neuro-psiquiatria*. 2009 Dec;67(4):1029-36.
8. Poynter B, Hon MS, Diaz-Granados N, Kapral

- M, Grace SL, Stewart DE. Sex differences in the prevalence of post-stroke depression: a systematic review. *Psychosomatics*. 2009 Dec 31;50(6):563-9.
9. Robinson RG, Jorge RE. Post-stroke depression: a review. *American Journal of Psychiatry*. 2015 Dec 18;173(3):221-31.
 10. Morris PL, Robinson RG, Andrzejewski P, Samuels J, Price TR. Association of depression with 10-year poststroke mortality. *American Journal of Psychiatry*. 1993 Jan 1;150:124-.
 11. Hackett ML, Yapa C, Parag V, Anderson CS. Frequency of depression after stroke. *Stroke*. 2005 Jun 1;36(6):1330-40.
 12. Stroke WH. Recommendations on stroke prevention, diagnosis, and therapy. Report of the WHO Task Force on Stroke and other Cerebrovascular Disorders. *Stroke*. 1989 Oct;20(10):1407-31.
 13. Williams LS, Brizendine EJ, Plue L, Bakas T, Tu W, Hendrie H, Kroenke K. Performance of the PHQ-9 as a screening tool for depression after stroke. *stroke*. 2005 Mar 1;36(3):635-8.
 14. Adams RD, Victor M, Ropper AH, Daroff RB. Principles of neurology.
 15. Starkstein SE, Robinson RG. Post-Stroke Mood Disorders. In *Neurology and Psychiatry* 1989 (pp. 192-209). Karger Publishers.
 16. Vataja R, Pohjasvaara T, Leppävuori A, Mäntylä R, Aronen HJ, Salonen O, Kaste M, Erkinjuntti T. Magnetic resonance imaging correlates of depression after ischemic stroke. *Archives of general psychiatry*. 2001 Oct 1;58(10):925-31.
 17. Robinson RG. Poststroke depression: prevalence, diagnosis, treatment, and disease progression. *Biological psychiatry*. 2003 Aug 1;54(3):376-87.
 18. Hackett ML, Yapa C, Parag V, Anderson CS. Frequency of depression after stroke. *Stroke*. 2005 Jun 1;36(6):1330-40.
 19. Aström M, Adolfsson R, Asplund K. Major depression in stroke patients. A 3-year longitudinal study. *Stroke*. 1993 Jul 1;24(7):976-82.
 20. Dam H, Pedersen HE, Ahlgren P. Depression among patients with stroke. *Acta Psychiatrica Scandinavica*. 1989 Aug 1;80(2):118-24.
 21. Glamcevski MT, Chong-Tin T. Prevalence of post-stroke depression, a Malaysian study. 2000.
 22. Robinson RG, Starr LB, Price TR. A two year longitudinal study of mood disorders following stroke. Prevalence and duration at six months follow-up. *The British Journal of Psychiatry*. 1984 Mar 1;144(3):256-62.
 23. Verdelho A, Henon H, Lebert F, Pasquier F, Leys D. Depressive symptoms after stroke and relationship with dementia A three-year follow-up study. *Neurology*. 2004 Mar 23;62(6):905-11.
 24. Eriksson M, Asplund K, Glader EL, Norrving B, Stegmayr B, Terént A, Åsberg KH, Wester PO. Self-reported depression and use of antidepressants after stroke: a national survey. *Stroke*. 2004 Apr 1;35(4):936-41.
 25. Raju RS, Sarma PS, Pandian JD. Psychosocial problems, quality of life, and functional independence among Indian stroke survivors. *Stroke*. 2010 Dec 1;41(12):2932-7.
 26. House A, Dennis M, Mogridge L, Warlow C, Hawton KE, Jones LE. Mood disorders in the year after first stroke. *The British Journal of Psychiatry*. 1991 Jan 1;158(1):83-92.
 27. Kotila M, Numminen H, Waltimo O. Depression after Stroke: Results of the FINNSTROKE Study. *Year Book of Psychiatry and Applied Mental Health*. 2000 Jan 1;2000(1):251-2.
 28. Truelsen T, Heuschmann PU, Bonita R, Arjundas G, Dalal P, Damasceno A, Nagaraja D, Ogunniyi A, Oveisgharan S, Radhakrishnan K, Skvortsova VI. Standard method for developing stroke registers in low-income and middle-income countries: experiences from a feasibility study of a stepwise approach to stroke surveillance (STEPS Stroke). *The Lancet Neurology*. 2007 Feb 28;6(2):134-9.
 29. Cramer SC. Repairing the human brain after stroke: I. Mechanisms of spontaneous recovery. *Annals of neurology*. 2008 Mar 1;63(3):272-87.
 30. Marwat MA, Usman M, Hussain M. Stroke and its relationship to risk factors. *Gomal Journal of Medical Sciences*. 2004 Jun 1;7(1).

31. Memon FA, Khooharo Y, Ali S, Sajjad. Clinical audit of stroke patients presenting at a teaching hospital. *PJMS*. 2009 Oct 1;25(6):968-71.
32. Lakhair MA, Memon RA, Rahu QA et al. Clinical spectrum of stroke in young adults. *Medical Channel*. 2008; 14: 181-84.
33. Memon RA, Lakhair MA, Rahu QA et al. Risk factors for stroke: An experience at Nawabshah Medical College Hospital Nawabshah. *Medical Channel*. 2008; 14: 175-77.
34. Zafar A, Shahid SK, Siddiqui M, Khan FS. Pattern of stroke in type 2 diabetic subjects versus non diabetic subjects. *J Ayub Med Coll Abbottabad*. 2007;19(4):64-7.
35. Khan J, Rehman A, Shah AA, Jielani A. Frequency of hypertension in stroke patients presenting at Ayub Teaching Hospital. *J Ayub Med Coll Abbottabad*. 2006;18(1):36-9.

Conflict of interest: Author declares no conflict of interest.

Funding disclosure: Nil

Author's contribution:

Saira Abbas; concept, data collection, data analysis, manuscript writing, manuscript review

Naila Naeem Shahbaz; data collection, data analysis, manuscript writing, manuscript review

Muhammad Amir Umer; data analysis, manuscript writing, manuscript review

Sumera Rafat Umer; data analysis, manuscript writing, manuscript review

Asfia Irfan; data analysis, manuscript writing, manuscript review

Adnan Anwer; data analysis, manuscript writing, manuscript review