



THE AGA KHAN UNIVERSITY

eCommons@AKU

Department of Biological & Biomedical Sciences

Medical College, Pakistan

April 2018

Stress of infertility: Can the couple cope?

Faiza Alam

Karachi University

Taseer Ahmed Khan

Karachi University

Rehana Rehman

Aga Khan University, rehana.rehman@aku.edu

Follow this and additional works at: https://ecommons.aku.edu/pakistan_fhs_mc_bbs

 Part of the [Obstetrics and Gynecology Commons](#), [Psychiatric and Mental Health Commons](#), [Psychiatry and Psychology Commons](#), and the [Women's Health Commons](#)

Recommended Citation

Alam, F., Khan, T. A., Rehman, R. (2018). Stress of infertility: Can the couple cope?. *JPMA. The Journal of the Pakistan Medical Association*, 68(4), 679-680.

Available at: https://ecommons.aku.edu/pakistan_fhs_mc_bbs/363

Stress of infertility: Can the couple cope?

Faiza Alam,¹ Taseer Ahmed Khan,² Rehana Rehman³

Madam, health impact of infertility has resulted in an aggravation of medical, social, psychological and economic burden of developing countries. Female partners in particular, suffer a lot with respect to their health, self-esteem self-assurance, personal control, genetic legacy and optimism in future. Infertile females

specially living in joint family system tend to stay within themselves under such pressures and drift to a world of depression and stress. Stress has an impact on all systems of the body (Figure), however the effect of stress on pathogenesis of infertility and the precise casual mechanism has yet to be explored.




Figure: Response of different systems of the body towards stress.

.....
^{1,2}Karachi University, ³Biological & Biomedical Sciences, Aga Khan University, Karachi.

Correspondence: Rehana Rehman. Email: drrehana7@gmail.com

Fertility predominantly depends on the maintenance of the quantity and quality of the ovarian reserve which is determined by the extrinsic and intrinsic factors.¹ The reserve declines with increase in age of a female, owing to

the consumption of the follicle, and majorly due to the imbalance in the redox activity. Oxidative stress (OS) produced by mitochondrial dysfunction within the ovaries is considered as the main cause for chromosomal segregation disorders, maturation and fertilization failures, or oocyte/embryo fragmentation. Cellular redox activity is a normal mechanism of the male and female reproductive system but its inequilibrium affects the fertility by hindering attainment and maintenance of oocyte developmental potential during in vivo processes.² OS is a multifactorial process that implicates several different outcomes especially the oocyte maturation with its impact on fertility irrespective of female previous health, genetic predisposition as well as environmental condition.³

Sirtuins (silent information regulator, SIRT), a family of Nicotinamide Adenine Dinucleotide (NAD) dependent enzymes is evolving as main regulators of OS. SIRT1 signaling initiates a positive ovarian cells response by stimulating the expression of antioxidants and thus preventing dysfunction of the ovarian cells.⁴ Moreover, significant change of SIRT1 mRNA has also been observed with increase in reactive oxidative species.⁵ The detection of its role in OS with reference to oocyte maturity can pave new pathways for reduction of cost and trauma of procedures in assisted fertilization techniques such as in vitro fertilization and intracytoplasmic sperm injection by increasing oocyte maturity, fertilization and hence

embryo implantation.

Although there are extensive facts available in literature indicating relationship between increased OS, infertility and decreased level of antioxidants, yet human studies to explain role of SIRT1 for expression of antioxidants, repair of cells, damaged by oxidative stress in order to prevent dysfunction of the ovarian cells have yet to be accomplished.

Disclaimer: None to declare.

Conflict of Interest: None to declare.

Funding Disclosure: None to declare.

Reference

1. Altaee H, Al-Madfai ZAM, Alkhafaji ZH. Effect of obesity on ovarian reserve parameters in mid-reproductive age women. *F1000Research* 2012, 1:43 (doi: 10.12688/f1000research.1-43.v1)
2. Ishii T, Miyazawa M, Takanashi Y, Tanigawa M, Yasuda K, Onouchi H, et al. Genetically induced oxidative stress in mice causes thrombocytosis, splenomegaly and placental angiodysplasia that leads to recurrent abortion. *Redox Biol.* 2014; 2: 679-85.
3. Zhang X, Li X-H, Ma X, Wang Z-H, Lu S, Guo Y-L. Redox-induced apoptosis of human oocytes in resting follicles in vitro. *J Soc Gynecol Investig.* 2006; 13: 451-8.
4. Matos L, Stevenson D, Gomes F, Silva-Carvalho JL, Almeida H. Superoxide dismutase expression in human cumulus oophorus cells. *Mol hum reprod.* 2009; 15: 411-9.
5. Ruder EH, Hartman TJ, Reindollar RH, Goldman MB. Female dietary antioxidant intake and time to pregnancy among couples treated for unexplained infertility. *Fertil Steril.* 2014; 101: 759-66.