



THE AGA KHAN UNIVERSITY

eCommons@AKU

Community Health Sciences

Department of Community Health Sciences

January 2016

Frequency and Etiology of Community-Acquired Bloodstream Infection in Hospitalized Febrile Children

Tufail Soomro

Ghulam Mohammad Mahar Medial College, Sukkur, Karachi, Pakistan

Shiyam Sunder Tikmani

Aga Khan University, shiyam.sunder@aku.edu

Sumera Aziz Ali

Aga Khan University, sumera.ali@aku.edu

Follow this and additional works at: https://ecommons.aku.edu/pakistan_fhs_mc_chs_chs

 Part of the [Bacteria Commons](#), [Bacteriology Commons](#), [Fungi Commons](#), [Pathogenic Microbiology Commons](#), and the [Public Health Commons](#)

Recommended Citation

Soomro, T., Tikmani, S. S., Ali, S. A. (2016). Frequency and Etiology of Community-Acquired Bloodstream Infection in Hospitalized Febrile Children. *Journal of Medical Diagnostic Methods*, 5(3).

Available at: https://ecommons.aku.edu/pakistan_fhs_mc_chs_chs/365

Frequency and Etiology of Community-Acquired Bloodstream Infection in Hospitalized Febrile Children

Tufail Soomro¹, Shiyam Sunder Tikmani² and Sumera Aziz Ali^{2*}

¹Department of Pediatrics, Ghulam Mohammad Mahar Medial College, Sukkur, Karachi, Pakistan

²Department of Community health Sciences, Aga Khan University, Karachi, Pakistan

*Corresponding author: Sumera Aziz Ali, Department of Community health Sciences, Aga Khan University, Karachi, Pakistan, Tel: +9234864837; E-mail: sumera.ali@aku.edu

Received date: June 29, 2016; Accepted date: July 15, 2016; Published date: July 22, 2016

Copyright: © 2016 Soomro T, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Background: Blood stream infection is a serious problem that needs immediate attention and treatment. We aim to identify the frequency of common organisms in blood culture of febrile pediatric patients so that empirical antibiotic therapy can be started timely.

Methods: This cross-sectional study was conducted in pediatric ward Civil Hospital Sukkur from 1st May 2013 to 31st October 2014. Children of both genders who are between the age of one month to 15 years admitted to the pediatric ward with a fever of > 38.0 c and with a history of fever > two days and whose blood culture has been sent were included in this study. Informed consent was taken prior to enrollment. The patient's blood culture was taken under aseptic technique. Samples were sent to microbiology lab civil hospital Sukkur. 1st culture on admission to be taken to avoid confounding with hospital-acquired the infection. The frequency of organisms grown in the culture was documented.

Results: The median age of enrolled participants was 48.5 (46.6) months, 153 (38.5%) were male and 158 (61.5%) were the female and median weight of enrolled participants was 15.9 (9.7) kg. Overall, 178 (69.3%) cases had a history of prior antibiotic use; blood cultures sent within 24 hours were 117 (45.5%) cases. Twenty-two (8.4%) cases were positive. Six were positive for *Salmonella typhi* cases followed by *E. coli*.

Conclusions: It was concluded from this study that the frequency of community acquired blood stream infection was 7.6%. *Salmonella typhi*, *E. coli* were found to be common pathogens.

Keywords: Infection; Community acquired; Blood culture; Pathogens

Introduction

Bloodstream infection is a severe problem that needs immediate attention and treatment as it can lead to morbidity and mortality. Fever is often the most common reason for presentation to the hospital in developing countries. Most patients with febrile illness have self-limiting viral infections, but some have a serious bacterial infection that needs urgent and appropriate antimicrobial therapy [1] advance in blood culture technique have resulted in efficient and reliable techniques for detection of bacterial and fungal pathogens [1]. Blood stream infection is a cause of high mortality especially if caused by multidrug-resistant bacteria. Bacteriologic culture to isolate the offending pathogen and knowledge about sensitivity pattern of isolates remains the mainstay of definitive diagnosis and management of bloodstream infection. However, blood culture is time-consuming and takes at least 2 to 5 days until the identification of organism [2] one key determinant in the ultimate outcome of a patient with sepsis is an institution of early empirical and appropriate antimicrobial therapy. This is only possible with knowledge of community-acquired bacteria and their likely susceptibility to antibiotics in a given set up. There are limited reports on community-acquired bacteremia in children,

particularly from developing countries. This could be due to either poor lab infrastructure or poor health seeking behavior and treatment of febrile illnesses mostly by primary care physicians without a bacteriologic diagnosis. A study done in developing countries shows 8% children had blood stream infection [3] several studies have shown that rate of isolation of pathogens from blood culture increases with the quantity of blood submitted [4]. Small study of community-acquired infections from Pakistan revealed five hundred and eight blood culture samples were positive for bacterial growth. Of these 465 (91.5%) having monomicrobial growth and 43 (8.5%) having polymicrobial growth. Among them, Klebsiella spp (24.1%) were most common amongst gram negative and staph aureus (24.1%) amongst gram positive [5]. An interesting finding in the subcontinent is the isolation of more gram negative bacteria than gram positive there is a high prevalence of enteric fever caused by *Salmonella typhi* and *Salmonella paratyphi* in Pakistan. More than 51% pediatric blood culture isolates contained *S. typhi* or *S. paratyphi A*, *S. typhi* alone was detected in 43% of positive blood culture. The other organisms identified were coagulase-negative *Staphylococci* (20%), *Acinetobacter* species (5.1%), *E. coli* (4.5%), and *Pseudomonas* species (11.4%) [6,7].

There are limited number of studies from Pakistan, despite the high incidence of community-acquired bacteremia and case fatality among hospitalized bacteremic children [8,9]. We aim to conduct this study to identify the frequency of common organisms in blood culture of

febrile pediatric patients so that empirical antibiotic therapy can be started with bacteriology of community-acquired bacteremia. The objective of this study was to determine the frequency of community acquired blood stream infection in febrile children and to identify the commonly isolated organisms.

Material and Methods

This cross-sectional study was conducted at pediatric ward off civil hospital Sukkur from 1st May 2013 to 31st October 2014. Children of both genders who are between the age of one month to 15 years admitted to the pediatric ward with recorded temperature of > 38.0 c and with a history of fever more than two days and whose blood culture has been sent. Neonates, post-surgical patients, patients on chemotherapy and on antibiotic therapy, children with congenital anomaly, those children who have central line for antibiotic administration over long periods, children who are referred from another hospital or health care center for further management.

Cases diagnosed with HIV and tuberculosis, and children of prolonged course of prophylactic antibiotics for any associated comorbid condition like repeated UTI were excluded. Community-acquired bacteremia was defined as bloodstream infection with the growth of pathogenic bacteria in a blood culture obtained within first 48 hours after admission in febrile children between the age of 01 months to 15 years. Children who fulfilled study criteria were enrolled in the study after taking informed consent was taken from the parents. The demographic information of patient was taken on structured proforma by the investigator.

The patient's blood cultures were performed in microbiology lab according to standard guidelines at civil hospital Sukkur. 1st culture on admission to be taken to avoid confounding with hospital-acquired the infection. The culture was followed by the primary researcher. The frequency of organisms grown in the culture was documented.

Non-probability consecutive sampling was done. The frequency of bloodstream infection in febrile pediatrics patients is 8%, considering the bond on the error of estimation of 3% and 95% confidence interval the estimated sample size was 257. Data was analyzed using SPSS version 20 (SPSS Inc. Chicago, IL, USA). All quantitative variables like age, weight, were expressed as median (interquartile range).

Qualitative variables like gender, commonly isolated community-acquired bloodstream organisms, and history of prior antibiotics were expressed in percentages. Chi-square or Fisher exact test was used to compare categorical variables and Man-Whitney U test was used to compare median age and weight. Effect modifiers like age, gender, weight and history of prior antibiotics were stratified to see their effect on the final outcome. The study is approved by ethical review of the institute (Figures 1 and 2).




Figure 1: Independent samples Mann Whitney U Test.




Figure 2: Frequency of isolated cultures from blood culture.

Results

A total of 257 patients were enrolled in this study during the study period. The median age of enrolled participants was 48.5 (46.6) months, 153 (38.5%) were male and 158 (61.5%) were the female and median weight of enrolled participants was 15.9 (9.7) kg. Overall, 178 (69.3%) cases had a history of prior antibiotic use; blood cultures sent within 24 hours were 117 (45.5%) cases. Twenty-two (8.4%) cases were positive. Six were positive for *Salmonella typhi* cases followed by *E. coli*. Comparative analysis is shown in Table 1 and stratified analysis summarized in Table 2. Stratified analysis showed, of children <30 months of age, 10 (6.5%) had positive blood culture compared to 12 (11.5%) cases among children of age \geq 30 months ($p>0.120$); of female children, 10 (10.1%) had positive blood culture compared to 12 (7.6%) cases among male children ($p>0.079$); of children weigh <13 kg, 10 (6.4%) had positive blood culture compared to 12 (11.9%) cases among children weigh \geq 13 kg ($p>0.138$) and of children with no history taking antibiotics prior to culture, 17 (21.5%) has positive blood culture compare to 5 (2.8%) cases children who had history of taking antibiotics prior to blood culture.

	Negative blood culture (n=235)		Positive blood culture (n=22)		Total
	n	%	n	%	
Age Months (Mean)	42.6 ± 4.7		61 ± 5.5		
less than 30 months	143	60.9%	10	45.5%	153
More than 30 months	92	39.1%	12	54.5%	104
Gender					
Male	146	62.1%	12	54.5%	99
Female	89	37.9%	10	45.5%	158
Weight kg (Mean)	10.2 ± 10.2		15.1 ± 9		
Less than 13 kg	146	62.1%	10	45.5%	156
More than 13 Kg	89	37.9%	12	54.5%	101
Blood sample taken					
Within 24 hr	97	41.3%	20	90.9%	117
24-48 hr	138	58.7%	2	9.1%	140
History of antibiotic use	173	73.6%	5	22.7%	178

Table 1: Comparison of baseline characteristics between children with positive or negative blood culture.

Blood culture	Age		Gender		Weight		Prior use of antibiotics	
	less than 30 months	More than 30 months	Female	Male	Less than 13 kg	More than 13 Kg	No	Yes
Negative	143	92	89	146	146	89	62	173
	93.5%	88.5%	89.9%	92.4%	93.6%	88.1%	78.5%	97.2%
Positive	10	12	10	12	10	12	17	5
	6.5%	11.5%	10.1%	7.6%	6.4%	11.9%	21.5%	2.8%
Total	153	104	99	158	156	101	79	178
	p>0.120		p>0.316		p>0.079			p>0.001

Table 2: Stratified analysis by age, sex, weight and prior use of antibiotics.

Discussion

Febrile illness is a leading reason for admission to hospital in Pakistan [10-12]. Despite the major contribution of infectious diseases to hospital admission, the availability of diagnostic microbiology services for bloodstream infections other than malaria is often limited by cost, infrastructure, and personnel constraints [11] consequently, health-care workers must often rely on syndrome-oriented empirical approaches to treatment and might underestimate or overestimate the likelihood of certain diseases, risking poor clinical outcomes and the promotion of antimicrobial resistance [12,13] understanding the causes and prevalence of community-acquired bloodstream infection, which is associated with high risk of death, can inform efforts to improve health outcomes in Pakistan and promote the meeting of millennium development goals for the reduction of child mortality, malaria, and tuberculosis.

Blood cultures are vital for recognizing pathogens causing severe infections and in leading appropriate antibiotic therapy. Furthermore, they remain the standard method for identifying bacteraemia in the assessment of sick patients [14] inappropriately; blood culture contamination is a common event and may lead to confusion regarding the significance of a positive blood culture. The most common contaminants are coagulase-negative staphylococcus species which are also becoming more prevalent as a primary pathogen in immunocompromised patients and in patients with indwelling intravascular devices [9,15] the uncertain clinical significance of potential contaminants leads to longer hospital stays, unnecessary antibiotic therapy, and additional laboratory testing; as a result, the cost incurred by a hospital is many times that incurred by the laboratory.

In this study 184 patients were enrolled during the study period with a median age of 48.52±46.6 months, the majority was females' 59.8% and median weight of 15.9±9.7 kg. Overall, 118 (64.1%) cases had a history of prior antibiotic use blood cultures sent within 24 hours were 175 (95.1%) cases. Fourteen (7.6%) cases were positive. In our study *Salmonella*, enteric serotype typhi was the most common bacterial pathogen accounting for 4.9% of cases followed by *E.coli* and *Pseudomonas* spp. Similar findings had been reported in the systemic review where salmonella is most common pathogen isolated.

In another study from Cambodia, it was reported that salmonella is most common infection followed by *E.coli* [15].

We also did a stratified analysis of positive blood cultures based on age and weight. We did not find statistically significant difference across strata. Children <30 months of age, 5.4% had positive blood culture compared to 9.9% cases among children of age ≥ 30 months (p>0.191) and children weigh <13 kg, 5.7% had positive blood culture compared to 10.5% cases among children weigh ≥ 13 kg (p>0.138).

There are some important limitations to this study. We examined hospital-based cohorts to define community-acquired bloodstream infections, and thereby might have included patients with more severe illness and included only those who have access to health-care facilities.

This study presents a strong case for the importance of considering bloodstream infections in patients admitted to civil hospital Sukkur, particularly those with fever. Our findings underscore the importance of continuous re-evaluation of who's integrated management of childhood illness and other empirical treatment algorithms for febrile illness in view of changing patterns of disease. When used, empirical treatment with antimicrobial drugs should include coverage for gram-negative, especially against *Salmonella*, *E.coli* and *Pseudomonas*.

It was concluded from this study that the frequency of bloodstream infections was 7.6%. *Salmonella typhi*, *E.coli*, and *Pseudomonas* were found to be common pathogens. No relationship of positive blood culture or pathogen was found with age, weight, and gender.

References

1. Deen J, Seidlein LV, Anderson F, Elle N, White JW, et al. (2012) Community-acquired bacterial blood stream infections in developing countries in south and southeast Asia: a systematic review. *Lancet Inf Dis* 12: 480-487.
2. Latif S, Anwar MS, Ahmad I (2009) Bacterial pathogens responsible for blood stream infections (bsi) and pattern of drug resistance in a tertiary care hospital of Lahore. *Biomedica* 25: 101-105.
3. Gonsalves WI, Cornish N, Moore M, Chen A, Varman M (2009) Effects of volume and site of blood drawn on blood cultures results. *J Clin Microbiol* 47: 3482-3485.
4. Podda A, Saul A, Arora R, Bhutta Z, Sinha A, et al. (2010) Conjugate vaccines for enteric fever: proceedings of a meeting organized in new Delhi, India in 2009. *J Infect Dev Ctries* 4: 404-411.
5. Sigauque B, Roca A, Mandomando I, Morais L, Quinto L, et al. (2009) Community-acquired bacteremia among children admitted to a rural hospital in Mozambique. *Pediatr Infect Dis J* 28: 108-113.
6. Blomberg B, Manji KP, Urassa WK, Tamim BS, Mwakagile DS, et al. (2007) Antimicrobial resistance predicts death in Tanzanian children with bloodstream infections: a prospective cohort study. *BMJ Infect Dis* 7: 43.
7. Claudias I, Barray LJ (2010) Pediatric emergencies associated with fever. *Emerg Med Clin North Am* 28: 67-84.
8. Wilson M, Mitchell M, Morris AJ, Murray PR, Reimer LG, et al. (2007) Principles and procedures for blood cultures: approved guideline. *Clinical and laboratory standards institute* 27.
9. Zhou L, Pollard AJ (2010) A fast and highly sensitive blood culture PCR method for clinical detection of salmonella enteric serovar typhi. *Ann Clin Microbiol Antimicrob* 9: 14.
10. Mintegi S, Benito J, Sanchez J, Azkunaga B, Iturralde I, et al. (2009) Predictors of occult bacteremia in young febrile children in the era of the heptavalent pneumococcal conjugate vaccine. *Eur J Emerg Med* 16: 199-205.
11. Wilkinson M, Bulloch B, Smith M (2009) Prevalence of occult bacteremia in children aged 3 to 36 months presenting to the emergency department with fever in the post-pneumococcal conjugate vaccine era. *Acad Emerg Med* 16: 220-225.
12. Brauner M, Goldman M, Kozer E (2010) Extreme leukocytosis and the risk of serious bacterial infections in febrile children. *Arch Dis Child* 95: 209-212.
13. Huppler AR, Eickhoff JC, Wald ER (2010) Performance of low-risk criteria in the evaluation of young infants with fever: a review of the literature. *Pediatrics* 125: 228-233.
14. Herz AM, Greenhow TL, Alcantara J, Hansen J, Baxter RP, et al. (2006) Changing epidemiology of outpatient bacteremia in 3- to 36-month-old children after the introduction of the heptavalent conjugated pneumococcal vaccine. *Pediatr Infect Dis J* 25: 293-300.
15. Whitney CG, Farley M, Hadler J, Harrison LH, Bennett NM, et al. (2003) Decline in invasive pneumococcal disease after the introduction of protein-polysaccharide conjugate vaccine. *N Engl J Med* 348: 1737-1746.