



THE AGA KHAN UNIVERSITY

eCommons@AKU

Section of Gastroenterology

Department of Medicine

February 2018

Impact of direct acting antivirals on occurrence and recurrence of hepatocellular carcinoma: Biologically plausible or an epiphenomenon?

Amna Subhan Butt

Aga Khan University, amna.subhan@aku.edu

Fatima Sharif

Aga Khan University

Shahab Abid

Aga Khan University, shahab.abid@aku.edu

Follow this and additional works at: https://ecommons.aku.edu/pakistan_fhs_mc_med_gastroenterol

 Part of the [Gastroenterology Commons](#)

Recommended Citation

Butt, A., Sharif, F., Abid, S. (2018). Impact of direct acting antivirals on occurrence and recurrence of hepatocellular carcinoma: Biologically plausible or an epiphenomenon?. *World journal of hepatology*, 10(2), 267-276.

Available at: https://ecommons.aku.edu/pakistan_fhs_mc_med_gastroenterol/153

World Journal of *Hepatology*

World J Hepatol 2018 February 27; 10(2): 172-351



REVIEW

- 172 Glycogenic hepatopathy: A narrative review
Sherigar JM, Castro JD, Yin YM, Guss D, Mohanty SR
- 186 Hepatitis C virus: Morphogenesis, infection and therapy
Morozov VA, Lagaye S
- 213 Cell fusion in the liver, revisited
Lizier M, Castelli A, Montagna C, Lucchini F, Vezzoni P, Faggioli F
- 222 Management of bacterial infection in the liver transplant candidate
Ferrarese A, Zanetto A, Becchetti C, Sciarrone SS, Shalaby S, Germani G, Gambato M, Russo FP, Burra P, Senzolo M
- 231 Digital liver biopsy: Bio-imaging of fatty liver for translational and clinical research
Mancini M, Summers P, Faita F, Brunetto MR, Callea F, De Nicola A, Di Lascio N, Farinati F, Gastaldelli A, Gridelli B, Mirabelli P, Neri E, Salvadori PA, Rebelos E, Tiribelli C, Valenti L, Salvatore M, Bonino F
- 246 Alkaline sphingomyelinase (NPP7) in hepatobiliary diseases: A field that needs to be closely studied
Duan RD

MINIREVIEWS

- 254 Spontaneous bacterial and fungal peritonitis in patients with liver cirrhosis: A literature review
Shizuma T
- 267 Impact of direct acting antivirals on occurrence and recurrence of hepatocellular carcinoma: Biologically plausible or an epiphenomenon?
Butt AS, Sharif F, Abid S

ORIGINAL ARTICLE

Basic Study

- 277 Homologous recombination mediates stable *Fah* gene integration and phenotypic correction in tyrosinaemia mouse-model
Junge N, Yuan Q, Huong Vu T, Krooss S, Bednarski C, Balakrishnan A, Cathomen T, Mams MP, Baumann U, Sharma AD, Ott M
- 287 Multipotent stromal cells stimulate liver regeneration by influencing the macrophage polarization in rat
Elchaninov A, Fatkhudinov T, Usman N, Arutyunyan I, Makarov A, Lokhonina A, Eremina I, Surovtsev V, Goldshtein D, Bolshakova G, Glinkina V, Sukhikh G

- 297 Morphological and biochemical effects of weekend alcohol consumption in rats: Role of concentration and gender

Morales-González JA, Sernas-Morales ML, Morales-González Á, González-López LL, Madrigal-Santillán EO, Vargas-Mendoza N, Fregoso-Aguilar TA, Anguiano-Robledo L, Madrigal-Bujaidar E, Álvarez-González I, Chamorro-Cevallos G

Retrospective Cohort Study

- 308 Survival outcomes of liver transplantation for hepatocellular carcinoma in patients with normal, high and very high preoperative alpha-fetoprotein levels

She WH, Chan ACY, Cheung TT, Lo CM, Chok KSH

Clinical Practice Study

- 319 Hepatitis C virus knowledge improves hepatitis C virus screening practices among primary care physicians

Samuel ST, Martinez AD, Chen Y, Markatou M, Talal AH

Observational Study

- 329 Outcomes assessment of hepatitis C virus-positive psoriatic patients treated using pegylated interferon in combination with ribavirin compared to new Direct-Acting Antiviral agents

Damiani G, Franchi C, Pigatto P, Altomare A, Pacifico A, Petrou S, Leone S, Pace MC, Fiore M

META-ANALYSIS

- 337 Outcomes of kidney transplantation in patients with hepatitis B virus infection: A systematic review and meta-analysis

Thongprayoon C, Kaewput W, Sharma K, Wijarnpreecha K, Leeaphorn N, Ungprasert P, Sakhuja A, Cabeza Rivera FH, Cheungpasitporn W

CASE REPORT

- 347 Primary hepatic peripheral T-cell lymphoma associated with Epstein-Barr viral infection

Ramai D, Ofori E, Nigar S, Reddy M

ABOUT COVER

Editorial Board Member of *World Journal of Hepatology*, Ji Won Kim, MD, PhD, Associate Professor, Department of Internal Medicine, Seoul National University College of Medicine, SMG-SNU Medical Center, Seoul 156-707, South Korea

AIM AND SCOPE

World Journal of Hepatology (*World J Hepatol*, *WJH*, online ISSN 1948-5182, DOI: 10.4254), is a peer-reviewed open access academic journal that aims to guide clinical practice and improve diagnostic and therapeutic skills of clinicians.

WJH covers topics concerning liver biology/pathology, cirrhosis and its complications, liver fibrosis, liver failure, portal hypertension, hepatitis B and C and inflammatory disorders, steatohepatitis and metabolic liver disease, hepatocellular carcinoma, biliary tract disease, autoimmune disease, cholestatic and biliary disease, transplantation, genetics, epidemiology, microbiology, molecular and cell biology, nutrition, geriatric and pediatric hepatology, diagnosis and screening, endoscopy, imaging, and advanced technology. Priority publication will be given to articles concerning diagnosis and treatment of hepatology diseases. The following aspects are covered: Clinical diagnosis, laboratory diagnosis, differential diagnosis, imaging tests, pathological diagnosis, molecular biological diagnosis, immunological diagnosis, genetic diagnosis, functional diagnostics, and physical diagnosis; and comprehensive therapy, drug therapy, surgical therapy, interventional treatment, minimally invasive therapy, and robot-assisted therapy.

We encourage authors to submit their manuscripts to *WJH*. We will give priority to manuscripts that are supported by major national and international foundations and those that are of great basic and clinical significance.

INDEXING/ABSTRACTING

World Journal of Hepatology is now indexed in Emerging Sources Citation Index (Web of Science), PubMed, PubMed Central, and Scopus.

EDITORS FOR THIS ISSUE

Responsible Assistant Editor: *Xiang Li*
Responsible Electronic Editor: *Rui-Fang Li*
Proofing Editor-in-Chief: *Lian-Sheng Ma*

Responsible Science Editor: *Li-Jun Cui*
Proofing Editorial Office Director: *Xiu-Xia Song*

NAME OF JOURNAL
World Journal of Hepatology

ISSN
 ISSN 1948-5182 (online)

LAUNCH DATE
 October 31, 2009

FREQUENCY
 Monthly

EDITOR-IN-CHIEF
Wan-Long Chuang, MD, PhD, Doctor, Professor, Hepatobiliary Division, Department of Internal Medicine, Kaohsiung Medical University Hospital, Kaohsiung Medical University, Kaohsiung 807, Taiwan

EDITORIAL BOARD MEMBERS
 All editorial board members resources online at <http://www.wjgnet.com/1948-5182/editorialboard.htm>

EDITORIAL OFFICE
 Xiu-Xia Song, Director

World Journal of Hepatology
 Baishideng Publishing Group Inc
 7901 Stoneridge Drive, Suite 501,
 Pleasanton, CA 94588, USA
 Telephone: +1-925-2238242
 Fax: +1-925-2238243
 E-mail: editorialoffice@wjgnet.com
 Help Desk: <http://www.f6publishing.com/helpdesk>
<http://www.wjgnet.com>

PUBLISHER
 Baishideng Publishing Group Inc
 7901 Stoneridge Drive, Suite 501,
 Pleasanton, CA 94588, USA
 Telephone: +1-925-2238242
 Fax: +1-925-2238243
 E-mail: bpgoffice@wjgnet.com
 Help Desk: <http://www.f6publishing.com/helpdesk>
<http://www.wjgnet.com>

PUBLICATION DATE
 February 27, 2018

COPYRIGHT

© 2018 Baishideng Publishing Group Inc. Articles published by this Open Access journal are distributed under the terms of the Creative Commons Attribution Non-commercial License, which permits use, distribution, and reproduction in any medium, provided the original work is properly cited, the use is non commercial and is otherwise in compliance with the license.

SPECIAL STATEMENT

All articles published in journals owned by the Baishideng Publishing Group (BPG) represent the views and opinions of their authors, and not the views, opinions or policies of the BPG, except where otherwise explicitly indicated.

INSTRUCTIONS TO AUTHORS

<http://www.wjgnet.com/bpg/gerinfo/204>

ONLINE SUBMISSION

<http://www.f6publishing.com>

Impact of direct acting antivirals on occurrence and recurrence of hepatocellular carcinoma: Biologically plausible or an epiphenomenon?

Amna Subhan Butt, Fatima Sharif, Shahab Abid

Amna Subhan Butt, Fatima Sharif, Shahab Abid, Section of Gastroenterology, Department of Medicine, Aga Khan University Hospital, Karachi 74800, Pakistan

ORCID number: Amna Subhan Butt (0000-0002-7311-4055); Fatima Sharif (0000-0002-3366-9830); Shahab Abid (0000-0003-2530-0378).

Author contributions: All authors have contributed to the study concept, manuscript writing, editing and approval of the final manuscript for publication.

Conflict-of-interest statement: No conflicts of interest to declare.

Open-Access: This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Invited manuscript

Correspondence to: Shahab Abid, FACG, FCPS, MD, PhD, Professor, Section of Gastroenterology at Department of Medicine, Aga Khan University, Stadium Road, Karachi 74800, Pakistan. shahab.abid@aku.edu
Telephone: +92-21-34930051
Fax: +92-21-34932095

Received: November 28, 2017

Peer-review started: December 5, 2017

First decision: January 15, 2018

Revised: February 6, 2018

Accepted: February 9, 2018

Article in press: February 9, 2018

Published online: February 27, 2018

Abstract

Hepatocellular carcinoma (HCC) is a major cause of morbidity and mortality worldwide. Chronic hepatitis C virus infection (HCV) is the most common cause of HCC in many European countries, Japan and Pakistan. Introduction of the new direct acting antivirals (DAAs) has revolutionized the management of HCV worldwide, with high rates of sustained virologic response in patients who could not have tolerated the previous interferon based treatments. However, recently there have been reports raising caution about the long term effects of DAAs, particularly a possible increased risk of HCC. Therefore this review explores the current molecular studies as well as clinical data that investigate the impact of DAAs on occurrence and recurrence of HCC.

Key words: Hepatocellular carcinoma; Direct acting antivirals; Hepatitis C

© **The Author(s) 2018.** Published by Baishideng Publishing Group Inc. All rights reserved.

Core tip: Our aim is to consolidate the existing literature as well as to identify whether there is a particular subset of the population in which this phenomenon was witnessed. The ground-breaking discovery of the new group of direct acting antiviral agents (DAAs) had led to a paradigm shift in the management of chronic hepatitis C (CHC). Wide variations have been observed in the studies assessing the long-term role of DAA based therapy on occurrence and recurrence of HCC. There is a need to differentiate whether the reported higher occurrence and recurrence rates are due to DAA or host and disease related factors and to identify subset of individuals particularly at risk. Also, future investigations should be directed towards

assessing the long-term effects of DAAs on group of patients that have not been studied thus far. Some important Centers in Europe and United States have been delaying antiviral treatment for 6 mo or more after the recent treatment for HCC. Hence, until more robust data is available, clinical practices should continue as per current guidelines in those patient groups who can benefit from DAA therapy with close surveillance of patients with advance fibrosis.

Butt AS, Sharif F, Abid S. Impact of direct acting antivirals on occurrence and recurrence of hepatocellular carcinoma: Biologically plausible or an epiphenomenon? *World J Hepatol* 2018; 10(2): 267-276 Available from: URL: <http://www.wjgnet.com/1948-5182/full/v10/i2/267.htm> DOI: <http://dx.doi.org/10.4254/wjh.v10.i2.267>

INTRODUCTION

The ground breaking discovery of the new group of direct acting antiviral agents (DAAs) had led to a paradigm shift in the management of chronic hepatitis C (CHC) which is the most common cause of hepatocellular carcinoma (HCC) in Japan, Pakistan, United States and many European countries^[1,2]. With annual incidence of HCC ranging from 1% to 7% in patients with HCV related cirrhosis, (HCC) is a leading cause of morbidity and the second most common cause of cancer related deaths worldwide^[3,4]. Besides contribution of several host and viral factors in the pathogenesis of disease progression, achieving sustained virologic response (SVR) has been found as the single most important factor in reducing HCV associated HCC incidence^[5].

The novel DAAs not only provided a potent, oral alternative to injectable interferons, but also had a shorter duration of treatment, better efficacy with over 90% achievement of SVR and a more favorable side effect profile^[6]. However, since 2016, concerns were raised regarding the effect of DAAs on progression to HCC^[7]. In addition, their long-term benefits including impact on HCC have been questioned in the context of specific populations and subgroups which were not included in the landmark trials investigating DAA based therapy^[3].

Therefore, this review aims to explore existing molecular studies as well as clinical observations in order to determine whether there is an association between the use of DAAs and the occurrence or recurrence of HCC among patients with HCV related liver disease. We also aim to evaluate whether there is a subset of the population in which this phenomenon has been observed.

DAA AND CARCINOGENESIS: BIOLOGICALLY PLAUSIBLE OR NOT?

The carcinogenesis of HCC is a chronic process with

several steps that may serve as potential targets for drug therapy. Unlike the hepatitis B virus, HCV is an RNA virus which is unable to integrate into the host genome and thus it is unlikely to have direct carcinogenic activity^[8]. While the mechanism of carcinogenesis due to HCV is not completely understood, observations from transgenic mice models suggest that liver cancer occurs because of rapid hepatocyte turnover, dysregulation of apoptosis and generation of reactive oxygen species, arising in the setting of a chronic inflammatory state induced by HCV^[9,10]. This indirect mechanism of cirrhosis driven carcinogenesis is supported by clinical data which shows a greater risk of HCC with chronic HCV infection and worsening liver fibrosis^[8].

One of the mechanisms proposed regarding the risk of DAAs towards development of HCC is that DAAs downregulate interferon genes, disrupting the innate immunosurveillance of the body^[8]. In the chronic phase of HCV, an estimated 10¹² virions are produced per day by infected hepatocytes^[11]. These trigger an immune response mediated primarily by natural killer (NK) cells which release cytokines such as interferon (IFN)- γ . IFNs upregulate interferon stimulated genes which have an anti-proliferative response by prolonging all phases of the cell cycle and decreasing viral replication. When HCV infection becomes persistent, NK cells become dysfunctional due to continuous antigenic stimulation by the high load of virions, resulting in impaired production of IFNs^[12]. In mice models, it was observed that decreased levels of IFN- γ independently control tumorigenesis^[13]. Thus the downregulation of these genes in IFN free therapy could contribute to development of HCC.

Analysis of the peripheral blood mononuclear cells (PBMCs) of CHC patients treated with DAAs has shown that in comparison to healthy controls, these patients have attenuated activity of both NK cells and monocytes, reflected by a decreased level of inflammatory cytokines in patients who had achieved rapid virological response (RVR), *i.e.*, undetectable HCV RNA at the end of 4 wk of DAA treatment^[14]. Natural Killer cell group 2D (NKG2D) is an activating receptor of immune responses which has been studied in the context of HCV associated HCC. It has been found that in HCV patients treated with IFN free DAA therapy, an on treatment decrease in the expression of NKG2D correlated to the early occurrence and recurrence of clinically evident HCC within the 6-mo surveillance period following treatment^[15].

With DAA based therapy, HCV RNA becomes undetectable in days to weeks, a much more rapid response than that observed with IFN based regimens. It is hypothesized that rapid eradication of HCV and subsequent abrupt resolution of the chronic inflammatory state disrupts the natural immune response of the body, possibly favoring the proliferation of neoplastic cells^[16]. Since clinical studies have shown no difference among different DAA regimens and development of HCC, it is hypothesized that if this effect does exist then it would have to be a class effect of the DAAs^[17,18].

Table 1 Possible factors contributing to hepatocellular carcinoma after hepatitis C virus eradication by direct acting antivirals

| |
|---|
| Down regulation of IFN genes |
| Presence of fibrosis |
| Sudden disruption of chronic inflammatory state |
| Impaired immune response by NK cells |
| T cell dysfunction |
| Decreased microRNA-122 |

IFN: Interferon; NK: Natural killer.

An experimental study analyzing the soluble inflammatory milieu from plasma cells of cirrhotic HCV patients found that HCV specific CD8+ T cells failed to recover from the baseline after DAA therapy. These cells play a role in HCC surveillance; therefore, their reduced activity in IFN free therapy could potentially affect development of HCC^[19]. Furthermore, serum levels of microRNA (miRNA) 122 were found to be reduced in DAA treated HCV patients after achieving SVR^[20]. Previous evidence shows that miRNA-122, which is the most abundant miRNA in the liver, functions as a tumor suppressor against HCC^[21]. Thus it is hypothesized that decreased miRNA-122 in DAA treated patients could contribute to an increased risk of HCC recurrence^[22]. Table 1 summarizes the possible factors that could lead to HCC in DAA treated HCV patients.

DAA AND CARCINOGENESIS: REVIEW OF EXISTING EVIDENCE/DATA

Studies that observed increased incidence of HCC

While DAAs represented a major breakthrough in the treatment of CHC, one of the first reports questioning their long-term role in development of HCC came from Reig *et al.*^[8] in 2016. The authors retrospectively assessed a cohort of 58 patients who had a history of HCC secondary to HCV and had received different regimens of DAA based therapy. After a median follow up of 5.7 mo (range 0.4-14.6 mo), 16 out of their 58 patients (27.6%) showed radiographic evidence of HCC. This study alerted the scientific community regarding the potential risks associated with use of DAAs and the authors called for a large-scale assessment to confirm their findings^[8]. However, their method of statistical analysis was questioned by Camma and colleagues who felt that reporting the crude rate of recurrence was a weakness of the study because of the variation in time elapsed from treatment of HCC and starting DAAs (median 11.2 mo, range 1.2-87.7 mo)^[23]. Cammà *et al.*^[23] used the data presented by Reig *et al.*^[24] to calculate the actuarial probability of HCC recurrence by plotting a Kaplan Meier curve. For their analysis, Cammà *et al.*^[23] used the time of HCC treatment as the initiation point, not the time of initiating DAA treatment as used by the original authors. With this method they found a much lower recurrence rate than that reported in the original study (7% and 13% at 6 and 12 mo

respectively).

Reig *et al.*^[24] later went on to present a follow up of their original cohort. In 2017, they reported that not only did they observe a higher recurrence rate of HCC among the DAA treated patients in their study; they also found a more aggressive pattern of recurrence in terms of tumor staging and subsequent treatment options. Renzulli *et al.*^[25] have also reported a rapid development of HCC following DAA treatment with a more aggressive pattern of microvascular invasion. The median duration between completion of DAA treatment and diagnosis of HCC in their patients was 82 d (range 0-318).

A retrospective study from Italy by Conti *et al.*^[18] reported HCC occurrence and recurrence rates of 3.16% (95%CI: 1.45-5.90) and 28.81% (95%CI: 17.76-42.07) respectively in a cohort of 344 CHC patients treated with different DAA regimens over a follow up of 24 wk. Approximately 69% of this patient population had HCV genotype 1 and 91% had achieved SVR. However, the lack of a control group makes the interpretation of these findings difficult. The authors attempted to account for this limitation by comparing their findings to those of a historic cohort of untreated cirrhotic patients at their center. They found an HCC occurrence rate of 3.2%, which was similar to their current study. Conti *et al.*^[18] have interpreted their results with caution. The authors claim that while DAA treatment of HCV does not seem to reduce the occurrence or recurrence of HCC, antiviral treatment should be started as early as possible to prevent the development of cirrhosis and recommend active surveillance of all cirrhotic patients, during and after DAA therapy.

In Portugal, Cardoso *et al.*^[26] found HCC incidence to be 7.4% within a one year follow up of cirrhotic patients that had achieved SVR after being treated with sofosbuvir and ledipasvir. These *de novo* HCC patients had been asymptomatic, and were detected on radiological screening. This emphasizes the recommendation of Conti *et al.*^[18] that there should be close monitoring of CHC patients for development of HCC, despite achieving SVR^[25]. In a larger study from Belgium, it was found that there was no difference in the early occurrence of HCC among patients treated with DAAs with or without Peg-IFN. However, this study reported HCC recurrence of 15% in patients treated with DAAs alone as compared to 0% in those who received a combination of Peg-IFN and DAAs. This study had a predominantly HCV Genotype 1 population. In these patients the authors also noted that those who developed HCC had a higher baseline risk of HCC which is a potential confounder^[27].

The most recent study on this research question is by Ida *et al.*^[28], published in October 2017. The study population comprised 100 patients from Japan with HCV genotype 1, treated with Daclastavir and Asunaprevir, who were followed for 15 mo. In this group, there were 5 new cases and 12 recurrences of HCC. The authors

have hypothesized that the high rate of HCC seen in this study could be related to a history of HCC, as these patients already had advanced fibrosis which is known to be implicated in the process of hepatocarcinogenesis. Table 2 summarizes the studies that report an increased incidence of HCC in DAA treated HCV patients.

Studies that did not note any significant effect

In 2016 and 2017 there have been several other similar reports, fueling the debate regarding the role of DAAs in HCC. Two large retrospective cohort studies have been conducted in the United States to investigate this matter; one by Ioannou *et al.*^[33] with a sample size of 62354 and the other by Kanwal *et al.*^[34] with 22500 study participants. Both studies concluded that DAAs are not associated with a significant risk of HCC as compared to IFN based treatment. Ioannou *et al.*^[33] found that DAA induced SVR reduced the risk of HCC by 71%. This effect was similar in groups who had received DAAs alone, DAAs in combination with IFN or IFN alone regimens, thus suggesting that achieving SVR could be the crucial factor for risk reduction of HCC, regardless of the therapeutic agents used^[33]. Kanwal *et al.*^[34] found that while there was a relative risk reduction in HCC, the absolute risk of HCC still persisted in those patients who had DAA induced SVR. In both studies the risk of HCC was greater in cirrhotic patients. Additionally, Kanwal *et al.*^[34] found that diabetes mellitus, alcohol use and a higher Fib-4 Index for assessment of fibrosis were risk factors for occurrence of HCC in patients who had achieved SVR. However, both study populations were restricted to United States veterans and were mostly patients with HCV genotype 1. The specific population included in these studies might limit the generalization of their findings.

The ANRS Collaborative Study Group assessed HCC recurrence rates among 3 French multicenter prospective cohorts who had received DAA based therapy after HCC curative treatment^[35]. Their study included a diverse patient population with cirrhotics, non-cirrhotics and liver transplant recipients. They did not find any evidence that DAAs increases the risk of HCC recurrence. The strength of this study was that analysis of data from 3 distinct patient cohorts yielded fairly consistent results. Secondly, they included patients who had received curative therapy for HCC as opposed to non-curative therapies such as chemoembolization, leading to speculation that some of the earlier studies that had reported a higher recurrence rate might have included patients in whom the initial tumor staging was incorrect or was incompletely treated^[35]. Table 3 summarizes the studies that do not show an increased risk of HCC in DAA treated patients.

Geographical variation of incidence of HCC in DAA treated patients

Many of the studies investigating the link between DAAs and HCC have been from Japan or European countries. Racial differences are known to be implicated in the

progression to HCC among HCV infected patients^[52]. In a large cohort of United States veterans with HCV, it was found that Hispanics were at greater risk of developing cirrhosis and HCC. Black race and Hispanic ethnicity have also been identified as independent predictors of treatment failure^[53]. Indeed the first report raising caution about the possible association of DAAs with recurrence of HCC in an aggressive form did come from the Spanish cohort followed by Reig *et al.*^[8].

A systematic review and meta-analysis by Waziry *et al.*^[54] found that there was no difference in HCC occurrence and recurrence in CHC patients who received IFN based or DAA based regimens. However, the authors acknowledge that in their analysis they were unable to account for geographical variations. Due to the heterogeneity in the studies included in this meta-analysis, the results should be interpreted with caution. In this meta-analysis most of the IFN based studies were from Japan whereas DAA based studies were from Europe. Hence, due to baseline difference in between two populations it's hard to draw an accurate conclusion regarding occurrence or recurrence of HCC among DAA and IFN based therapy. Furthermore, they had to exclude several studies which had incomplete data regarding BCLC staging of HCC.

In our literature review, the populations that we have found to be under-represented in terms of the current research question include Indians, Arabs and Africans. Additionally, while the 2 studies with the largest sample size are from the United States, these were limited to veterans only^[33,34].

Genotype based variation of HCC in DAA treated patients

HCV genotype is an important consideration when considering progression to HCC. HCV genotype 3 is generally more aggressive and is associated with a higher risk of progression to cirrhosis and HCC^[55]. To the best of our knowledge, the studies published so far suggesting a greater risk of HCC with DAA treatment have not identified a particular genotype of HCV that is significantly associated with this disease progression. However, as most of these reports are from Japan or European countries such as France and Italy, therefore the major disease burden studied has been of HCV genotype 1^[35,37]. Table 4 lists the factors predisposing to development of HCC in DAA treated HCV patients.

WAY FORWARD

There is wide variation in the studies assessing the long-term role of DAA based therapy on occurrence and recurrence of HCC, both in terms of the baseline characteristics of study population and the different DAA regimens used. At present, most of the studies have been reported from regions where HCV genotype 1 is the most prevalent one^[34,35]. Data is very limited from Asian populations where HCV Genotype 3 is the most common genotype. This is an important consideration as the different genotypes of HCV have a unique

Table 2 Studies showing an increased incidence of hepatocellular carcinoma in direct acting antivirals treated patients

| Ref. | Treatment groups | Country/sample size | Male gender n (%) | Age (yr) | Genotype n (%) | Duration of follow up | Cirrhosis n (%) | History of HCC n (%) | HCC occurrence | HCC recurrence |
|--|--|-------------------------|---|---|--|--------------------------------|--|--|--|--|
| Reig <i>et al</i> ^[8] | All DAA treated | Spain n = 58 | 40 (69) | Median 66.3 (45-83) | G1a = 8 (13.8%), G1b = 45 (77.6%), G3 = 2 (3.4%), G4 = 3 (5.2%) | median 5.7 mo | 55 (94.8) | 58 (100) | Not applicable | 16 (27.6%) |
| Conti <i>et al</i> ^[68] | All DAA treated | Italy n = 344 | 207 (60.2) | Median = 63 (29-85) | G1 = 237 (69), G2 = 40 (11.6), G3 = 38 (11), G4 = 29 (8.4) | 24 wk | All cirrhotic | 59 (17) | 9 of 285 patients (3.16%, 95% CI: 1.45-5.90) | 17 of 59 patients (28.81%, 95% CI: 17.76-42.07) |
| Cardoso <i>et al</i> ^[26] | All DAA treated | Portugal n = 54 | > 70% | No HCC: 59 yr (41-81). Patients with HCC = 58 yr (55-72) | No HCC: G1 = 78%, G3 = 18%. Patients who developed HCC: G1 = 75%, G3 = 25% | Median = 12.0 mo (9.4-12.5 mo) | All cirrhotic | 0 (0) | 4 (7.4%) | Not applicable |
| Ida <i>et al</i> ^[28] | All DAA treated | Japan n = 100 | 46 (46) | Median 72.5 (26-87) | G1 = 100 (100) | 15 mo | | 26 (26) | 5 (6) | 12 (12) |
| Kozbial <i>et al</i> ^[29] | All DAA treated | Austria | - | 56-74 yr | G1a = 3 (15.8), G1b = 13 (68.4), G3a = 2 (10.5), G4 = 1 (5.2) | | | 3 (15.8) | 6.60% | 3 patients |
| Issachar <i>et al</i> ^[30] | All DAA treated | Israel n = 273 | - | Median 55 (49-60) yr | - | 15 mo | All cirrhotic | - | 6 (2.1) | 3 (1.05) |
| Kwong <i>et al</i> ^[31] | patients divided into 3 eras: IFN era (2003-2010), protease inhibitor era (2011-2013), and DAA era (2014-2015) | United States n = 48158 | 29858 (62) | Median 55 (49-60) yr | - | Median 493 d | All cirrhotic | - | Incidence Rate of HCC was 49% higher in the DAA era (IR 6.6/100 person-years [py], 95%CI: 5.6-7.9) vs the IFN era (4.5/100 py, 95%CI: 4.2-4.7; IRR 1.49, 95%CI: 1.24-1.79, P < 0.001). | Not applicable |
| Strazulla <i>et al</i> ^[32] | daclatasvir and simeprevir for 24 weeks to reach SVR | Italy n = 1 case report | 1 male | 53 | G1 | 24 wk | Decompe-nated cirrhosis | Treated | Not applicable | 1 recurrence |
| Bielen <i>et al</i> ^[27] | PEG-IFN+ DAA = 77 (13.5), DAAs only = 490 (86.4) | Belgium n = 567 | PEG IFN + DAA group: 55 (71.4) DAA only group: 307 (62.7) | PEG IFN + DAA group age : 52 ± 9, DAA only group: 59 ± 12 | PEG IFN + DAA group: all genotype 1. DAA only: G1 = 69%, G4 = 14.7% | 6 mo | Metavir fibrosis score F3/F4 included only | PEG IFN + DAA = 1/77 (1.3%) DAA alone: 41/490 (8.4%) | Early occurrence rate of HCC = 1.7% and 1.1% in patients treated with DAA and without PEG-IFN, respectively | Early recurrence rate was 0% in patients treated with PEG-IFN+DAA and 15.0% in patients treated with DAA without PEG-IFN |

DAAs: Direct acting antivirals; IFN: Interferon; HCC: Hepatocellular carcinoma.

Table 3 Studies that do not report an increased risk of hepatocellular carcinoma with direct acting antivirals

| Ref. | Treatment N /sample size | Country | Male gender n (%) | Age (yr) | Genotype n (%) | Duration of follow up | Cirrhosis n (%) | History of HCC n (%) | HCC occurrence | HCC recurrence |
|---|---|---------------|---|---|---|---|---|--|--|--|
| Ioannou <i>et al</i> ^[63] | IFN only = 35871 (58%), DAA + IFN = 4535 (7.2%), DAA only = 21948 (35%) n = 62354 | United States | 96.60% | Mean 55.8 ± 7.6 | G1 = 77.4 %, G2 = 13.5%, G3 = 8.3%, G4 = 0 | Mean follow-up DAA only group = 1.53 years, DAA+IFN group = 3.6 yr, IFN only group = 9.1 yr | Cirrhosis: 16.8%, decompensated cirrhosis: 4.7% | None | Total 3271 incident cases. IFN group = 0.81/100 person years, DAA + IFN = 1.06 /100 py, DAA only = 1.32/100 py | Not applicable |
| Kanwal <i>et al</i> ^[64] | All DAA treated n = 22500 | United States | 21761 (96.7%) | Mean 61.6 ± 6.1 | G1 = 19531 (86.8%), G2 = 1422 (6.3%), G3 = 940 (4.2%), G4 = 217 (1%) | Median: 20.2 mo after DAA initiation and 26.1 mo for untreated patients | Cirrhosis: DAA group = 152 (80%), no DAAs = 55 (72 %) | All treated | Not Applicable | DAA group = 24 (12.7%), no DAAs = 16 (20.5%) |
| ANRS CO22 HEP ATH-ER <i>et al</i> ^[65] | DAA group = 189, no DAA = 78 n = 267 | France | DAA group = 78% (78%) | DAA group = 62 ± 9 yr, no DAAs = 66 ± 10 yr | 65 % genotype 1 | | | | | |
| ANRS CO12 CIRVIR <i>et al</i> ^[66] | DAA group = 13, no DAA = 66 n = 79 | France | DAA group = 11 (85%), no DAA = 39 (59%) | DAA group = 61 ± 10 yr, no DAA group = 65 ± 9 yr | Genotype 1: DAA group = 53/63 (84 %) | | | | | |
| ANRS CO23 CUPILT <i>et al</i> ^[65] | All DAA treated n = 314 | France | 257 (82%) | 61 ± 8 yr | 212 (67.5%) genotype 1 | | 49 (15.6%) | Treated | Not Applicable | 7 (2.2%) |
| Cabibbo <i>et al</i> ^[64] | All DAA treated n = 143 | Italy | 80 (60.1) | Mean 70.4 ± 8.9 | G1a: 9 (6.3), 1b: 114 (79.7), G2: 9 (6.3), G3: 7 (4.9), G4: 4 (2.8) | 6, 12 and 18 mo | All cirrhotic | All treated | Not applicable | 6-, 12- and 18-mo HCC recurrence rates were 12%, 26.6% and 29.1%, respectively |
| Nagata <i>et al</i> ^[71] | IFN-based: 1145, IFN-free DAA group: 752 n = 1897 | Tokyo | IFN group: 621 (54), IFN free: 340 (45) | Median: IFN group: 59 (19-79); IFN free: 69 (24-87) | IFN group: G1a = 8 (7), G1b = 833 (73), G2a = 182 (16), G2b = 105 (9), G3 = 1 (0) | Median for IFN group: 6.8 (0.2-22.0); IFN free: 1.8 (0.1-7.7) | | 5% of IFN group, 11% of IFN free group | IFN group: 18 (2.5%), IFN-free group: 7 (1.1%) | IFN group: 18 (53%), IFN-free group: 22 (29%) |
| Ikedo <i>et al</i> ^[88] | All DAA treated n = 177 | Japan | M:F = 52: 37 in each group | DAA group: 71 (39-85) | | Median 20.7 mo | | All treated | Not applicable | HCC recurrence rates at 1 st and 2 nd year were 18.1 and 25.0% in pts with DAA therapy and 21.8 and 46.5% in those without DAAs, (P = 0.003) |
| Zanetto <i>et al</i> ^[69] | DAA treated = 23, control = 23 n = 46 | Italy | | DAA group = 59 (49-69), controls = 58 (46-70) | DAA group: G1a = 5 (22), Gb = 9 (39), G2 = 1 (4), G3 = 5 (22), G4 = 3 (13) | Median = DAA group = 10 mo, Control group = 7 mo | All cirrhotic | All treated | Not applicable | 12.5% of DAA-treated patients and 8.3% of control group had HCC recurrence (P = 0.60) |

| Author | All DAA treated n = | Italy | 20 (64.5) | Mean 65 ± 8 | G1a = 4 (13), G1b = 23(74), G2 = 2 (6.5), G4 = 2 (6.5) | Median 8 mo | All cirrhotic | All treated | Not applicable | 1 (3.2) |
|--|--|----------------|--|---|---|---|-----------------------------|-------------|--|--|
| Zavaglia <i>et al</i> ^[40] | 31 | | | | | | | | | |
| Ogata <i>et al</i> ^[41] | All DAA treated n = 1170 | Japan | 493 (42) | Median = 67 (21-88) | All genotype 1 | Time from the end of DAA therapy until last visit: 1.3 yr | | None | 22 cases (1.8%) | Not applicable |
| Minami <i>et al</i> ^[42] | DAA group = 27, IFN group = 38, Controls = 861 n = 926 | Japan | DAA group: 18 (67), IFN group: 27 (71), Controls: 489 (57) | Median age: DAA group = 71 (48-82) IFN group = 66 (49-79), Control = 71 (44-91) | Genotype 1: DAA = 21 (78), IFN = 29 (76), Controls = 633 (74). Genotype 2: DAA = 6 (22), IFN = 9 (24), Control = 147 (17) G1 = 743 (76.2) | 1 and 2 yr | All had advanced cirrhosis | All treated | Not applicable | Cumulative recurrence rates at 1 and 2 yr were 21.1% and 29.8%, respectively, in the DAA group, 26.3% and 52.9%, respectively, in the IFN group, and 30.5% and 61.0%, respectively, in the control group |
| Deterding <i>et al</i> ^[43] | n = 974 | Germany | | | | | All had advanced cirrhosis | | 12 (1.2) | |
| Degasperi <i>et al</i> ^[44] | n = 565 | Italy | 60% | Median age = 65 (30-87) yr | G1a = 15%, G1b = 49%, G2 = 13%, G3 = 11%, G4 = 12%, G5 = 1% | Median 42 wk for occurrence, 39 wk for recurrence | All cirrhotic | 48 (8%) | 20 (4%) estimated annual incidence of 1.6% | 9 (19%), annual incidence of 7.7% |
| Bouliere <i>et al</i> ^[45] | DAA + RBV ± PEG IFN = 21%, IFN free DAA therapy = 79% n = 1393 | | | 47 (19-79) yr | | 2.5 (0.6-4.3) yr | | | 0 (0) | 0 (0) |
| Nagaoki <i>et al</i> ^[46] | PEG-IFN/ RBV = 244, DCV/ ASV = 154 n = 398 | Japan | | | All genotype 1 | Median for PEG-IFN/ RBV = 96 (10-196) and DCV/ ASV group = 23 (4-78) mo | | | PEG-IFN/ RBV = 13 (5.3%), DCV/ ASV group = 7 (4.5%) | |
| Cheung <i>et al</i> ^[47] | All DAA | United Kingdom | | | | 15 mo | All decompensated cirrhosis | 29 (71.4%) | 17 (5%) | 2 |
| Mettke <i>et al</i> ^[48] | 158 DAA treated, 184 controls | Germany | | | | Median = 440 (91-908) and 592 (90-1000) d | All cirrhotic | | 6 and 14 patients during follow-up, resulting in an HCC incidence of 2.9 (AVT) and 4.48 (Con) per 100 py, respectively | |
| Ji <i>et al</i> ^[49] | | China | 51% | Mean age 51 | 82.2% genotype 1b | Median 14 (3-35) mo | 48% cirrhotic | None | | Not applicable |
| Innes <i>et al</i> ^[50] | | Scotland | | | | 1.8 yr | | None | 44 (5.1%) | Not applicable |
| Torres <i>et al</i> ^[51] | All DAA treated | United States | 7 (87.5%) | Median 64 (57-87) years | G1 = 6 (75%), mixed genotype = 2 (25%) | 12 mo | 7 (87.5%) | All treated | Not applicable | 0 (0) |

DAA: Direct acting antivirals; IFN: Interferon; HCC: Hepatocellular carcinoma.

Table 4 Factors predisposing to hepatocellular carcinoma in hepatitis C virus patients treated with direct acting antivirals after sustained virologic response

| |
|--|
| Past history of hepatocellular carcinoma ^[18,28,36] |
| Male gender ^[28] |
| Cirrhosis ^[34] |
| Hypoalbuminemia ^[41] |
| Thrombocytopenia ^[41] |
| Raised AFP levels ^[41] |

response to DAA based therapy and are associated with a unique burden of HCC^[56]. Beside male patients, group of individuals with prior history of HCC, cirrhosis and elevated AFP at baseline were found with greater risk of HCC if treated with DAAs. Currently, the phenomenon appears to be a class effect rather than an individual drug effect. Hence, focusing on these HCV patients and measuring the impact of DAAs on progression or development of HCC will help to estimate the more accurate risk. In certain studies higher rates of HCC recurrence was found. There is a need to differentiate whether the reported higher recurrence rate and more aggressive pattern of recurrence are due to DAA or host or disease related factors including presence of fibrosis, gaps in initial tumor staging and receiving non-curative therapies such as chemoembolization. Also, future investigations should be directed towards assessing the long-term effects of DAAs on these populations that have not been studied thus far^[3]. In the meantime, a consensus recommendation seen in most of the studies at present is that even after achieving SVR, there should be close surveillance of patients with CHC especially with advance fibrosis and those who received a recent treatment for HCC in order to detect HCC at an early stage^[18,19,26,32]. Some important Centers in Europe and United States have been delaying antiviral treatment for 6 mo or more after recent treatment for HCC in these patients. Moreover, until more robust data is available to investigate the role of DAAs in HCV related HCC cases, clinical practice should continue as per current guidelines in those patient groups who can benefit from DAA therapy^[51].

REFERENCES

- 1 **Hsu CS**, Chao YC, Lin HH, Chen DS, Kao JH. Systematic Review: Impact of Interferon-based Therapy on HCV-related Hepatocellular Carcinoma. *Sci Rep* 2015; **5**: 9954 [PMID: 25963067 DOI: 10.1038/srep09954]
- 2 **El-Serag HB**. Epidemiology of viral hepatitis and hepatocellular carcinoma. *Gastroenterology* 2012; **142**: 1264-1273 [PMID: 22537432 DOI: 10.1053/j.gastro.2011.12.061]
- 3 **Goossens N**, Hoshida Y. Hepatitis C virus-induced hepatocellular carcinoma. *Clin Mol Hepatol* 2015; **21**: 105-114 [PMID: 26157746 DOI: 10.3350/cmh.2015.21.2.105]
- 4 **Siegel R**, Ma J, Zou Z, Jemal A. Cancer statistics, 2014. *CA Cancer J Clin* 2014; **64**: 9-29 [PMID: 24399786 DOI: 10.3322/caac.21208]
- 5 **Moon C**, Jung KS, Kim DY, Baatarkhuu O, Park JY, Kim BK, Kim SU, Ahn SH, Han KH. Lower incidence of hepatocellular carcinoma and cirrhosis in hepatitis C patients with sustained virological response by pegylated interferon and ribavirin. *Dig Dis Sci* 2015; **60**: 573-581 [PMID: 25236421 DOI: 10.1007/s10620-014-3361-6]
- 6 **Cortez KJ**, Kottlilil S. Beyond interferon: rationale and prospects for newer treatment paradigms for chronic hepatitis C. *Ther Adv Chronic Dis* 2015; **6**: 4-14 [PMID: 25553238 DOI: 10.1177/2040622314551934]
- 7 **Grandhe S**, Frenette CT. Occurrence and Recurrence of Hepatocellular Carcinoma After Successful Direct-Acting Antiviral Therapy for Patients With Chronic Hepatitis C Virus Infection. *Gastroenterol Hepatol (NY)* 2017; **13**: 421-425 [PMID: 28867970]
- 8 **Reig M**, Mariño Z, Perelló C, Iñarrairaegui M, Ribeiro A, Lens S, Diaz A, Vilana R, Darnell A, Varela M, Sangro B, Calleja JL, Forns X, Bruix J. Unexpected high rate of early tumor recurrence in patients with HCV-related HCC undergoing interferon-free therapy. *J Hepatol* 2016; **65**: 719-726 [PMID: 27084592 DOI: 10.1016/j.jhep.2016.04.008]
- 9 **Moriya K**, Fujie H, Shintani Y, Yotsuyanagi H, Tsutsumi T, Ishibashi K, Matsuura Y, Kimura S, Miyamura T, Koike K. The core protein of hepatitis C virus induces hepatocellular carcinoma in transgenic mice. *Nat Med* 1998; **4**: 1065-1067 [PMID: 9734402 DOI: 10.1038/2053]
- 10 **Kamegaya Y**, Hiasa Y, Zukerberg L, Fowler N, Blackard JT, Lin W, Choe WH, Schmidt EV, Chung RT. Hepatitis C virus acts as a tumor accelerator by blocking apoptosis in a mouse model of hepatocarcinogenesis. *Hepatology* 2005; **41**: 660-667 [PMID: 15723444 DOI: 10.1002/hep.20621]
- 11 **Neumann AU**, Lam NP, Dahari H, Gretch DR, Wiley TE, Layden TJ, Perelson AS. Hepatitis C viral dynamics in vivo and the antiviral efficacy of interferon-alpha therapy. *Science* 1998; **282**: 103-107 [PMID: 9756471 DOI: 10.1126/science.282.5386.103]
- 12 **Mondelli MU**, Varchetta S, Oliviero B. Natural killer cells in viral hepatitis: facts and controversies. *Eur J Clin Invest* 2010; **40**: 851-863 [PMID: 20597961 DOI: 10.1111/j.1365-2362.2010.02332.x]
- 13 **Street SE**, Cretney E, Smyth MJ. Perforin and interferon-gamma activities independently control tumor initiation, growth, and metastasis. *Blood* 2001; **97**: 192-197 [PMID: 11133760 DOI: 10.1182/blood.V97.1.192]
- 14 **Ning G**, Li YT, Chen YM, Zhang Y, Zeng YF, Lin CS. Dynamic Changes of the Frequency of Classic and Inflammatory Monocytes Subsets and Natural Killer Cells in Chronic Hepatitis C Patients Treated by Direct-Acting Antiviral Agents. *Can J Gastroenterol Hepatol* 2017; **2017**: 3612403 [PMID: 28567369 DOI: 10.1155/2017/3612403]
- 15 **Chu PS**, Nakamoto N, Taniki N, Ojio K, Amiya T, Makita Y, Murata H, Yamaguchi A, Shiba S, Miyake R, Katayama T, Ugamura A, Ikura A, Takeda K, Ebinuma H, Saito H, Kanai T. On-treatment decrease of NKG2D correlates to early emergence of clinically evident hepatocellular carcinoma after interferon-free therapy for chronic hepatitis C. *PLoS One* 2017; **12**: e0179096 [PMID: 28617830 DOI: 10.1371/journal.pone.0179096]
- 16 **Llovet JM**, Villanueva A. Liver cancer: Effect of HCV clearance with direct-acting antiviral agents on HCC. *Nat Rev Gastroenterol Hepatol* 2016; **13**: 561-562 [PMID: 27580683 DOI: 10.1038/nrgastro.2016.140]
- 17 **Blanco JR**, Rivero-Juárez A. The risk of hepatocellular carcinoma after sustained virological response in patients treated with the new direct-acting antiviral drugs: should we be worry about it? *Expert Rev Anti Infect Ther* 2016; **14**: 993-996 [PMID: 27686837 DOI: 10.1080/14787210.2016.1243466]
- 18 **Conti F**, Buonfiglioli F, Scuteri A, Crespi C, Bolondi L, Caraceni P, Foschi FG, Lenzi M, Mazzella G, Verucchi G, Andreone P, Brillanti S. Early occurrence and recurrence of hepatocellular carcinoma in HCV-related cirrhosis treated with direct-acting antivirals. *J Hepatol* 2016; **65**: 727-733 [PMID: 27349488 DOI: 10.1016/j.jhep.2016.06.015]
- 19 **Sekyere SO**, Falk C, Aregay A, Schlevogt B, Deterding K, Manns MP, Vogel A, Wirth T, Cornberg M, Wedemeyer H. GS-003-Interferon-free cure of hepatitis C virus infection

- does alter the soluble inflammatory milieu which could affect hepatocellular carcinoma surveillance by CD8+ T cells in patients with liver cirrhosis. *J Hepatol* 2017; **66**: S2 [DOI: 10.1016/S0168-8278(17)30266-0]
- 20 **Waring JF**, Dumas EO, Abel S, Coakley E, Cohen DE, Davis JW, Podsadecki T, Dutta S. Serum miR-122 may serve as a biomarker for response to direct acting antivirals: effect of paritaprevir/R with dasabuvir or ombitasvir on miR-122 in HCV-infected subjects. *J Viral Hepat* 2016; **23**: 96-104 [PMID: 26436610 DOI: 10.1111/jvh.12470]
 - 21 **Nakao K**, Miyaaki H, Ichikawa T. Antitumor function of microRNA-122 against hepatocellular carcinoma. *J Gastroenterol* 2014; **49**: 589-593 [PMID: 24531873 DOI: 10.1007/s00535-014-0932-4]
 - 22 **Liu CH**, Kao JH. Hepatocellular carcinoma recurrence after interferon-free direct acting antiviral treatment for chronic hepatitis C virus infection: fact or fiction? *Transl Cancer Res* 2016; **5**: S192-195 [DOI: 10.21037/8773]
 - 23 **Cammà C**, Cabibbo G, Craxi A. Direct antiviral agents and risk for HCC early recurrence: Much ado about nothing. *J Hepatol* 2016; **65**: 861-862 [PMID: 27255578 DOI: 10.1016/j.jhep.2016.04.033]
 - 24 **Reig M**, Mariño Z, Perelló C, Iñarrairaegui M, Lens S, Díaz A, Vilana R, Darnell A, Varela M, Sangro B, Calleja JL. PS-031-Tumour recurrence after Interferon-free treatment for hepatitis C in patients with previously treated hepatocellular carcinoma discloses a more aggressive pattern and faster tumour growth. *J Hepatol* 2017; **66**: S20 [DOI: 10.1016/S0168-8278(17)30302-1]
 - 25 **Renzulli M**, Buonfiglioli F, Conti F, Brocchi S, Serio I, Foschi FG, Caraceni P, Mazzella G, Verucchi G, Golfieri R, Andreone P, Brillanti S. Imaging features of microvascular invasion in hepatocellular carcinoma developed after direct-acting antiviral therapy in HCV-related cirrhosis. *Eur Radiol* 2018; **28**: 506-513 [PMID: 28894901 DOI: 10.1007/s00330-017-5033-3]
 - 26 **Cardoso H**, Vale AM, Rodrigues S, Gonçalves R, Albuquerque A, Pereira P, Lopes S, Silva M, Andrade P, Morais R, Coelho R, Macedo G. High incidence of hepatocellular carcinoma following successful interferon-free antiviral therapy for hepatitis C associated cirrhosis. *J Hepatol* 2016; **65**: 1070-1071 [PMID: 27476768 DOI: 10.1016/j.jhep.2016.07.027]
 - 27 **Bielen R**, Moreno C, Van Vlierberghe H, Bourgeois S, Mulkay JP, Vanwolleghem T, Verlinden W, Brixco C, Decaestecker J, de Galocsy C, Janssens F, Van Overbeke L, Van Steenkiste C, D'Heygere F, Cool M, Wuyckens K, Nevens F, Robaey G. The risk of early occurrence and recurrence of hepatocellular carcinoma in hepatitis C-infected patients treated with direct-acting antivirals with and without pegylated interferon: A Belgian experience. *J Viral Hepat* 2017; **24**: 976-981 [PMID: 28504854 DOI: 10.1111/jvh.12726]
 - 28 **Ida H**, Hagiwara S, Kono M, Minami T, Chishina H, Arizumi T, Takita M, Yada N, Minami Y, Ueshima K, Nishida N, Kudo M. Hepatocellular Carcinoma after Achievement of Sustained Viral Response with Daclatasvir and Asunaprevir in Patients with Chronic Hepatitis C Virus Infection. *Dig Dis Sci* 2017; **35**: 565-573 [PMID: 29040989 DOI: 10.1159/000480183]
 - 29 **Kozbial K**, Moser S, Schwarzer R, Laferl H, Al-Zoairy R, Stauber R, Stättermayer AF, Beinhardt S, Graziadei I, Freissmuth C, Maieron A, Gschwantler M, Strasser M, Peck-Radosaljevic M, Trauner M, Hofer H, Ferenci P. Unexpected high incidence of hepatocellular carcinoma in cirrhotic patients with sustained virologic response following interferon-free direct-acting antiviral treatment. *J Hepatol* 2016; **65**: 856-858 [PMID: 27318327 DOI: 10.1016/j.jhep.2016.06.009]
 - 30 **Issachar A**, Sneh-Arbib O, Braun M, Shlomai A, Oxtrud E, Harif Y, Karavani C, Tur-Kaspa R, Cohen-Naftaly M. LBP-509-Occurrence and recurrence of malignancies post DAA Treatment in 5.1% of patients-single center experience. *J Hepatol* 2017; **66**: S97 [DOI: 10.1016/S0168-8278(17)30454-3]
 - 31 **Kwong AJ**, Kim W, Flemming JA. Continued Increase in Incidence of De Novo Hepatocellular Carcinoma Among Liver Transplant Registrants with Hepatitis C Virus Infection. *Hepatology* 2017; **66**: 71A [DOI: 10.1002/hep.29500]
 - 32 **Strazzulla A**, Iemmolo RMR, Carbone E, Postorino MC, Mazzitelli M, De Santis M, Di Benedetto F, Cristiani CM, Costa C, Pisani V, Torti C. The Risk of Hepatocellular Carcinoma After Directly Acting Antivirals for Hepatitis C Virus Treatment in Liver Transplanted Patients: Is It Real? *Hepat Mon* 2016; **16**: e41933 [PMID: 28070200 DOI: 10.5812/hepatmon.41933]
 - 33 **Ioannou GN**, Green PK, Berry K. HCV eradication induced by direct-acting antiviral agents reduces the risk of hepatocellular carcinoma. *J Hepatol* 2017; pii: S0168-8278 [PMID: 28887168 DOI: 10.1016/j.jhep.2017.08.030]
 - 34 **Kanwal F**, Kramer J, Asch SM, Chayanupatkul M, Cao Y, El-Serag HB. Risk of Hepatocellular Cancer in HCV Patients Treated With Direct-Acting Antiviral Agents. *Gastroenterology* 2017; **153**: 996-1005.e1 [PMID: 28642197 DOI: 10.1053/j.gastro.2017.06.012]
 - 35 **ANRS collaborative study group on hepatocellular carcinoma (ANRS CO22 HEPATHER, CO12 CirVir and CO23 CUPILT cohorts)**. Lack of evidence of an effect of direct-acting antivirals on the recurrence of hepatocellular carcinoma: Data from three ANRS cohorts. *J Hepatol* 2016; **65**: 734-740 [PMID: 27288051 DOI: 10.1016/j.jhep.2016.05.045]
 - 36 **Cabibbo G**, Petta S, Calvaruso V, Cacciola I, Cannavò MR, Madonia S, Distefano M, Larocca L, Prestileo T, Tinè F, Bertino G, Giannitrapani L, Benanti F, Licata A, Scalisi I, Mazzola G, Cartabellotta F, Alessi N, Barbàra M, Russello M, Scifo G, Squadrito G, Raimondo G, Craxi A, Di Marco V, Cammà C; Rete Sicilia Selezione Terapia - HCV (RESIST-HCV). Is early recurrence of hepatocellular carcinoma in HCV cirrhotic patients affected by treatment with direct-acting antivirals? A prospective multicentre study. *Aliment Pharmacol Ther* 2017; **46**: 688-695 [PMID: 28791711 DOI: 10.1016/j.dld.2017.01.119]
 - 37 **Nagata H**, Nakagawa M, Asahina Y, Sato A, Asano Y, Tsunoda T, Miyoshi M, Kaneko S, Otani S, Kawai-Kitahata F, Murakawa M, Nitta S, Itsui Y, Azuma S, Kakinuma S, Nouchi T, Sakai H, Tomita M, Watanabe M; Ochanomizu Liver Conference Study Group. Effect of interferon-based and -free therapy on early occurrence and recurrence of hepatocellular carcinoma in chronic hepatitis C. *J Hepatol* 2017; **67**: 933-939 [PMID: 28627363 DOI: 10.1016/j.jhep.2017.05.028]
 - 38 **Ikeda K**, Kawamura Y, Kobayashi M, Kominami Y, Fujiyama S, Sezaki H, Hosaka T, Akuta N, Saitoh S, Suzuki F, Suzuki Y, Arase Y, Kumada H. Direct-Acting Antivirals Decreased Tumor Recurrence After Initial Treatment of Hepatitis C Virus-Related Hepatocellular Carcinoma. *Dig Dis Sci* 2017; **62**: 2932-2942 [PMID: 28884320 DOI: 10.1007/s10620-017-4739-z]
 - 39 **Zanetto A**, Shalaby S, Vitale A, Mescoli C, Ferrarese A, Gambato M, Franceschet E, Germani G, Senzolo M, Romano A, Angeli P, Ruge M, Farinati F, Forton DM, Cillo U, Burra P, Russo FP. Dropout rate from the liver transplant waiting list because of hepatocellular carcinoma progression in hepatitis C virus-infected patients treated with direct-acting antivirals. *Liver Transpl* 2017; **23**: 1103-1112 [PMID: 28544587 DOI: 10.1002/lt.24790]
 - 40 **Zavaglia C**, Okolicsanyi S, Cesarini L, Mazzarelli C, Pontecorvi V, Ciaccio A, Strazzabosco M, Belli LS. Is the risk of neoplastic recurrence increased after prescribing direct-acting antivirals for HCV patients whose HCC was previously cured? *J Hepatol* 2017; **66**: 236-237 [PMID: 27592303 DOI: 10.1016/j.jhep.2016.08.016]
 - 41 **Ogata F**, Kobayashi M, Akuta N, Osawa M, Fujiyama S, Kawamura Y, Sezaki H, Hosaka T, Kobayashi M, Saitoh S, Suzuki Y, Suzuki F, Arase Y, Ikeda K, Kumada H. Outcome of All-Oral Direct-Acting Antiviral Regimens on the Rate of Development of Hepatocellular Carcinoma in Patients with Hepatitis C Virus Genotype 1-Related Chronic Liver Disease. *Oncology* 2017; **93**: 92-98 [PMID: 28448999 DOI: 10.1159/000470910]
 - 42 **Minami T**, Tateishi R, Nakagomi R, Fujiwara N, Sato M, Enoku K, Nakagawa H, Asaoka Y, Kondo Y, Shiina S, Koike K. The impact of direct-acting antivirals on early tumor recurrence after radiofrequency ablation in hepatitis C-related hepatocellular carcinoma. *J Hepatol* 2016; **65**: 1272-1273 [PMID: 27524465 DOI: 10.1016/j.jhep.2016.07.0]

- 43 **Deterding K**, Mauss S, Pathil A, Buggisch P, Schott E, Cornberg M, Zimmermann T, Simon KG, Klinker H, Günther R, Pfeiffer-Vornkahl H. PS-096-Long-term follow-up after IFN-free therapy of advanced HCV-associated liver cirrhosis: continued improvement of liver function parameters—Results from the German Hepatitis C-Registry (DHC-R). *J Hepatol* 2017; **66**: S55 [DOI: 10.1016/S0168-8278(17)30372-0]
- 44 **Degasperi E**, D'Ambrosio R, Sangiovanni A, Aghemo A, Soffredini R, De Nicola S, Lunghi G, Lampertico P. Low rates of de novo or recurrent hepatocellular carcinoma in HCV cirrhotic patients treated with direct-acting antivirals (DAAs): a single-center experience. *Hepatology* 2017; **66**: 46A
- 45 **Bourliere M**, Gane EJ, Jacobson I, Gordon SC, Sulkowski MS, McNabb BL, Chen F, Dvory-Sobol H, McNally J, Brainard DM, Subramanian M. Long-Term Follow Up of Patients with Chronic HCV and No or Minimal Fibrosis Shows Low Risk for Liver-Related Morbidity and Mortality After Achieving SVR with DAA-Based Therapy: Results from the Gilead SVR Registry. *Hepatology* 2017; **66**: 518A-519A
- 46 **Nagaoki Y**, Imamura M, Aikata H, Daijo K, Teraoka Y, Honda F, Nakamura Y, Hatooka M, Morio R, Morio K, Kan H, Fujino H, Kobayashi T, Masaki K, Ono A, Nakahara T, Kawaoka T, Tsuge M, Hiramatsu A, Kawakami Y, Hayes CN, Miki D, Ochi H, Chayama K. The risks of hepatocellular carcinoma development after HCV eradication are similar between patients treated with peg-interferon plus ribavirin and direct-acting antiviral therapy. *PLoS One* 2017; **12**: e0182710 [PMID: 28797106 DOI: 10.1371/journal.pone.0182710]
- 47 **Cheung MCM**, Walker AJ, Hudson BE, Verma S, McLaughlan J, Mutimer DJ, Brown A, Gelson WTH, MacDonald DC, Agarwal K, Foster GR, Irving WL; HCV Research UK. Outcomes after successful direct-acting antiviral therapy for patients with chronic hepatitis C and decompensated cirrhosis. *J Hepatol* 2016; **65**: 741-747 [PMID: 27388925 DOI: 10.1016/j.jhep.2016.06.019]
- 48 **Mettke F**, Schlevogt B, Deterding K, Wranke A, Port K, Manns MP, Vogel A, Cornberg M, Wedemeyer H. THU-081-Interferon-free therapy of chronic hepatitis C does not change the short-term risk for de novo hepatocellular carcinomas in patients with liver cirrhosis. *J Hepatol* 2017; **66**: S210-211
- 49 **Ji D**, Wang C, Shao Q, Li F, Li B, Wu V, Wong A, Wang YD, Chen J, Chen GF, Lau G. No increase in the occurrence rate of hepatocellular carcinoma in Chinese treated by direct-acting antivirals compared to Interferon after eradication of hepatitis c virus: a long-term follow-up. *J Hepatol* 2017; **66**: S23 [DOI: 10.1016/S0168-8278(17)30308-2]
- 50 **Innes H**, Barclay ST, Hayes PC, Fraser A, Dillon JF, Stanley A, Bathgate A, McDonald S, Goldberg D, Valerio H, Fox R. PS-035-Among Cirrhotic Patients with a Hepatitis C Sustained Viral Response, the Risk of De-novo Hepatocellular Carcinoma Relates to Baseline Factors and Not the Use of Direct Acting Antivirals: Results from a Nationwide Cohort. *J Hepatol* 2017; **66**: S22-23 [DOI: 10.1016/S0168-8278(17)30306-9]
- 51 **Torres HA**, Vauthey JN, Economides MP, Mahale P, Kaseb A. Hepatocellular carcinoma recurrence after treatment with direct-acting antivirals: First, do no harm by withdrawing treatment. *J Hepatol* 2016; **65**: 862-864 [PMID: 27255582 DOI: 10.1016/j.jhep.2016.05.034]
- 52 **El-Serag HB**, Kramer J, Duan Z, Kanwal F. Racial differences in the progression to cirrhosis and hepatocellular carcinoma in HCV-infected veterans. *Am J Gastroenterol* 2014; **109**: 1427-1435 [PMID: 25070058 DOI: 10.1038/ajg.2014.214]
- 53 **Su F**, Green PK, Berry K, Ioannou GN. The association between race/ethnicity and the effectiveness of direct antiviral agents for hepatitis C virus infection. *Hepatology* 2017; **65**: 426-438 [PMID: 27775854 DOI: 10.1002/hep.28901]
- 54 **Waziry R**, Hajarizadeh B, Grebely J, Amin J, Law M, Danta M, George J, Dore GJ. Hepatocellular carcinoma risk following direct-acting antiviral HCV therapy: A systematic review, meta-analyses, and meta-regression. *J Hepatol* 2017; **67**: 1204-1212 [PMID: 28802876 DOI: 10.1016/j.jhep.2017.07.025]
- 55 **Kanwal F**, Kramer JR, Ilyas J, Duan Z, El-Serag HB. HCV genotype 3 is associated with an increased risk of cirrhosis and hepatocellular cancer in a national sample of U.S. Veterans with HCV. *Hepatology* 2014; **60**: 98-105 [PMID: 24615981 DOI: 10.1002/hep.27095]
- 56 **Ampuero J**, Romero-Gómez M, Reddy KR. Review article: HCV genotype 3 – the new treatment challenge. *Aliment Pharmacol Ther* 2014; **39**: 686-698 [PMID: 24612116 DOI: 10.1111/apt.12646]

P- Reviewer: Brillanti S S- Editor: Song XX L- Editor: A
E- Editor: Li RF





Published by **Baishideng Publishing Group Inc**
7901 Stoneridge Drive, Suite 501, Pleasanton, CA 94588, USA
Telephone: +1-925-223-8242
Fax: +1-925-223-8243
E-mail: bpgoffice@wjgnet.com
Help Desk: <http://www.f6publishing.com/helpdesk>
<http://www.wjgnet.com>

