



THE AGA KHAN UNIVERSITY

eCommons@AKU

Department of Radiology

Medical College, Pakistan

September 2003

Extrarenal Wilms' tumor

M Yunus

R Hashmi

S H. Hasan

H M. Brohi

Follow this and additional works at: https://ecommons.aku.edu/pakistan_fhs_mc_radiol



Part of the [Radiology Commons](#)

Recommended Citation

Yunus, M., Hashmi, R., Hasan, S. H., Brohi, H. M. (2003). Extrarenal Wilms' tumor. *Journal of Pakistan Medical Association*, 53(9), 436-139.

Available at: https://ecommons.aku.edu/pakistan_fhs_mc_radiol/193

Case Reports

Extrarenal Wilms' Tumor

M. Yunus, R. Hashmi, S. H. Hasan*, H. M. Y. Brohi**

Departments of Radiology, Pathology*, Neurology**, Aga Khan University and Hospital, Karachi.

Introduction

Extrarenal Wilms' tumor, which by definition excludes primary tumor in the kidney, is extremely rare. It occurs predominantly in children. Most of the cases that have been reported involved the retroperitoneum.

We are reporting a case of retroperitoneal (extrarenal) Wilms' tumor extending into the left psoas muscle and through neural foramina into the spinal canal with meningeal metastasis in a neonate. To the best of our knowledge, such an extensive extrarenal Wilms' tumor at this age with intraspinal extension and meningeal metastasis has not been reported previously

Case Report

An 8-day-old full term baby boy presented with jaundice, developed on the 3rd day of life, and vomiting with hematemesis since the 5th day of life. No antenatal ultrasound examination had been performed. The baby was full term and had a normal vaginal delivery.

Clinical examination revealed a firm and non-tender mass in the right iliac fossa. X-ray of abdomen revealed a mass of soft tissue density, occupying lower abdomen and pelvis, displacing bowel loops. The ultrasound examination showed a large, solid, predominantly hyperechoic mass, measuring 85 mm in longest dimension, with internal vascularity suggestive of neoplastic lesion. Posteriorly the mass was immediately adjacent to the spine. There was no connection of the mass to the kidneys. Mild hydronephrosis of the right kidney was identified. As the lesion was suggestive of malignancy, CT of abdomen was done which showed a heterogeneously enhancing, solid, retroperitoneal lobulated mass about 100x80mm in dimensions (Figure 1). The mass was extending from the level of lower pole of both kidneys, displacing them and pushing the aorta and iliac vessels anterolaterally. In addition, at the level of L2/3, the mass was encroaching into the spinal canal through the left sided neural foramen expanding it without bony destruction. Hydronephrosis of the right kidney was also noted. The rest of the abdominal viscera were normal. There

was no evidence of lymphadenopathy on CT examination. Differential diagnosis included neuroblastoma and teratodermoid tumor.

The child underwent exploratory laparotomy. The surgeons extracted a retroperitoneal mass of 10x8cm from the lumbosacral region which was extending upward with the aorta and iliac vessels stretched over the tumor. The tumor was extending into the spinal canal at the L2/3 level. Tumor site, extent and appearance resembled that of neuroblastoma and the intraspinal component was not removed. The operation was successful and recovery was uneventful.

On gross examination, the specimen consisted of a large nodular tissue measuring 9x7.5x5 cm. The cut surface was solid, lobulated and grayish white in color. Histopathological examination showed features of Wilms' tumor, exhibiting both epithelial and mesenchymal differentiation. The features of anaplasia were not appreciated (Figure 2). Resected lymph nodes showed benign lymphoid hyperplasia.

A magnetic resonance imaging (MRI) was not carried out preoperatively. Follow up MRI examination showed a 3 cm mass in the left psoas muscle extending through a left neural foramina into the extradural space from L4 to S1 (Figures 3a, b). Pial enhancement over the conus medullaris of the spinal cord was also identified suggesting meningeal metastasis (Figure 4). L4/5 laminotomy and removal of extradural tumor was carried out. The patient received chemotherapy (Vincristine and Actinomycin D). Follow-up to date has been over 1 yr and 10 months and the patient is currently well.

Discussion

Extrarenal Wilms' tumor is extremely rare. According to our review of the literature 72 cases have been reported¹⁻⁴⁰ (including the present case report as shown in Table). These tumors can occur in isolation or in association with other tumors usually teratomas.

The diagnostic criteria necessary to establish the diagnosis include absence of primary kidney tumor and supernumerary kidney (radiologically and surgically).

The origin of Wilms' tumor is controversial. An extrarenal location supports more frequent occurrence of ectopic metanephric blastema.¹ But, the presence of tumor cephalad to a horseshoe kidney favors origin from primitive mesodermal tissue, probably mesonephric rests.² Presence of persistent mesonephric duct remnants in the walls of cervix, vagina and inguinal canal and occurrence of extrarenal Wilms' tumor in these locations supports the theory of mesonephros as the source of origin.³ Another possible mechanism involves Connheim's cell rest theory in which cells with persistent embryonal potential undergo malignant transformation.³ These tumors can also arise from other neoplasms, generally teratoma.²

A palpable mass is the most common presentation. Patients with uterine extrarenal Wilms' tumor may present with irregular menstrual bleeding. Two patients with thoracic lesions were discovered on chest radiograph for nonspecific pulmonary symptoms.⁴⁻⁶

Age of presentation is reported to be between 2 months to 10 years. The oldest recorded patient suffering from uterine extrarenal Wilms' tumor was a 77 years old female.⁷ It is more common in females, 56.6% of reported cases occurred in females and 43.4% in males.¹⁻⁴⁰

Extrarenal Wilms' tumors are mostly solid large masses. Cystic cavities associated with a solid tumor can occur.^{13,14} One unusual manifestation presented from a large cystic component mimicked ascites clinically.⁴ Extrarenal Wilms' tumor has the potential for local recurrence¹⁵⁻¹⁷ and distant metastasis. Metastasis have been reported in lungs¹⁵, liver, pancreas¹⁸ and brain.¹⁹ Venous invasion of a right suprarenal tumor into the IVC and right atrium has also been reported.²

Six patients of extrarenal Wilms' tumor were associated with horseshoe kidney, i.e., 7.6% of patients having extrarenal Wilms' tumor had horseshoe kidneys in association.²⁰ This tumor has also been reported in patients with spinal dysraphism.^{19,21}

Regarding staging and management of extrarenal Wilms' tumor, the behavior of this tumor appears to parallel that of a similarly staged intrarenal Wilms' tumor. Hence, the staging and management protocols of intrarenal Wilm's tumor can be applied to an extrarenal location.²² An adaptation of the National Wilms' Tumor Study (NWTS) protocol can be carried out.³ Currently TNM classification is used for staging purpose.²²

References

1. Broecker BH, Caldames AA, McWilliams NB, et al. Primary Extrarenal Wilms' tumor in children. *J Paediatr Surg* 1989; 24:1283-8.
2. Fernandes ET, Kumar M, Doughlas EC, et al. Extrarenal Wilms' tumor. *Rev. Paediatr Surg* 1989;24:1283-8.
3. Andrews PE, Kelalis PP, Haase GM. Extrarenal Wilms' tumor: results of National Wilms' Tumor Study. *J Paediatr Surg* 1992; 27:1181-4.
4. Narasimharao KL, Marwaha RK, Kaushik S, et al. Extrarenal Wilms' tumor: a case report. *Paediatr Surg* 1989; 24:212-14.
5. Sahin A, Benda JA. Primary ovarian Wilm's tumor. *Cancer* 1988; 61:1460-3.
6. Madanat F, Osborne B, Cangir A, et al. Extrarenal Wilms' tumor. *J Paediatr* 1978; 93:439-43.
7. Jiscoot P, Aertsens W, Degels MA, et al. Extrarenal Wilm's tumor of the uterus. *Eur J Gynaecol Oncol* 1999; 20:195-7.
8. Johnson F, Lutertton C, Limbert D. Extrarenal and urothelial Wilm's tumor. *Urology* 1980;15:370-3.
9. Ho J, Ma L, Wong KC. An extrarenal Wilm's tumor arising from an undescended testis. *Pathology* 1981;13:619.
10. Gillis AJM, Oosterhuis JW, Schipper MEI, et al. Origin and histology of testicular Wilm's tumor. *Genes Chromosomes Cancer* 1994;11:126-35.
11. Heyns CF, Van Niekerk DJT. Nephroblastoma in an ovotestis of a true hermaphrodite. A case report. *J Urol* 1987; 137:1003.
12. Fahner JB, Switzer R, Freyer DR, et al. Extrarenal Wilm's tumor, unusual presentation in the lumbosacral region. *Am J Pediatr Hematol Oncol* 1993;15:117-19.
13. Edelstein G, Webb RS, Jr, Romsdahl MM et al. Extrarenal Wilm's tumor *Am J Surg* 1965;109:509-12.
14. Orłowski JP, Levin HS, Dyment PG. Intrascrotal Wilms tumor developing in heterotopic anlage of probable mesonephric origin. *J Paediatr Surg* 1980;15:679-82.
15. Thompson MR, Emmanuel IG, Campbell MS, et al. Extrarenal Wilm's tumor. *J Paediatr Surg* 1973;8:37-41.
16. Wu JP, Garcia J. Supernumerary kidney with Wilm's tumor. *Wis Med J* 1971;70:211-16.
17. Tamaro P, Fonda E, Valente M, et al. A new case report of extrarenal Wilm's tumor. *Pediatr Med Chir* 1982; 4:143-8.
18. Mantke R, Manger T, Bedwelski K, et al. Hepatic and retroperitoneal tumor resection for late metastasis of a Wilm's tumor in an adult patient. A case report. *Hepatogastroenterol* 1999;46:2289-92.
19. Mirkin LD, Azzarelli B, Seo IS. Extrarenal Wilm's tumor with cerebellar metastasis in a four-year-old girl with spina bifida. *Am J Clin Path* 1990;93:805-9.
20. Vinod K, Kapur, RP, Sakalkale KV, et al. Association of Extrarenal Wilm's tumor with a horseshoe kidney *J. Paediatr Surg* 1998;6:935-7.
21. Abraham JM, Pawel BB, Duhaime AC, et al. Extrarenal nephroblastic proliferation in spinal dysraphism. A case report. *Paediatr Neurosurg* 1999;31:40-4.
22. Coppes MJ, Wilson PC, Weetzman S. Extrarenal Wilm's tumor: staging, treatment and prognosis. Retrospective multicase study. *J Clin Oncol* 1991;9:167-74.
23. Suzuki K, Miyake H, Tashiro M, et al. Extrarenal Wilm's tumor. A case report. *Paediatr Radiol* 1993;23:149-50.
24. Lopez CP, Nortes CL, Martinez BE. Extrarenal Wilm's tumor, diagnosis and review. *Arch Esp Urol* 1997;50:999-1001.
25. Koretz MJ, Wang S, Klein FA, et al. Extrarenal Wilm's tumor. *Cancer* 1987;60:2484-8.
26. Fukutomi Y, Shibuya C, Yamamoto S, et al. Extrarenal Wilm's tumor in the adult patient. A case report and review of world literature. *Am J Clin Pathol* 1988;90:618-22.
27. Wakely PE Jr, Sprangue RI, Kornstein MJ. Extrarenal Wilm's tumor: an analysis of 4 cases. *Hum Pathol* 1989;20: 691-5.
28. Kim YW, Park YK, Oh SM, et al. Retroperitoneal teratoma with predominance of nephroblastic elements. A case report. *J Korean Med Sci* 1990;5:237-42.
29. Tebbi K, Ragab AH, Ternberg JL, et al. An extrarenal Wilm's tumor arising from sacrococcygeal teratoma. *Clin Pediatr* 1974;13:1019-21.
30. Arkovitz MS, Ginsburg HB, Eidelman J, et al. Primary extrarenal Wilm's tumor in the inguinal canal. A case report and review of literature. *J Paediatr Surg* 1996;13:957-9.
31. Strand WR, Chou P, Pero JE, et al. Extrarenal Wilm's tumor occurring in inguinal canal. *J Urol* 1990;143:783-5.
32. Simha MR, Doctor VM. Extrarenal Wilm's tumor. A case report and review of literature. *Indian J Cancer* 1991;28:16-21.
33. Benatar B, Wright C, Freinkel AL, et al. Primary extrarenal Wilm's tumor of the uterus presenting as a cervical polyp. *Int J Gynecol Pathol* 1998;17:277-80.
34. Iraniha S, Shen V, Kruppe CN, et al. Uterine cervical extrarenal Wilm's tumor managed without hysterectomy. *J Paediatr Hematol Oncol* 1999;21:548-50.
35. Babin EA, Davis JR, Hatch KD, et al. Wilm's tumor of cervix. *Gynecol Oncol* 2000;76:107-11.
36. Muc RS, Grayson W, Grobelaar JJ. Adult extrarenal Wilm's tumor occurring in uterus. *Arch Pathol Lab Med* 2001;125:1081-3.
37. Bittencourt AL, Britto JF, Fonseca LF. Wilm's tumor of the uterus. First tumor of the literature. *Cancer* 1981;47:2496-99.
38. Bell DA, Shimm DS, Gang DL. Wilm's tumor of the endocervix. *Arch Path Lab Med* 1985;109:371.
39. .Pereira F, Carrascal E, Canas C, et al. Extrarenal Wilm's tumor of left ovary. *J Paediatr Oncol* 2000;22:88-9.
40. Isaac MA, Vijayalakshmi S, Madhu CS, et al. Pure cystic nephroblastoma of the ovary with a review of extrarenal Wilm's tumor. *Hum Pathol* 2000;31:761-4.