



THE AGA KHAN UNIVERSITY

eCommons@AKU

Community Health Sciences

Department of Community Health Sciences

February 1998

An epidemiologic study of cystitis among Muslim women attending an out-patients clinic in Karachi

N Atiullah
Aga Khan University

F F. Fikree
Aga Khan University

I Husain

Follow this and additional works at: https://ecommons.aku.edu/pakistan_fhs_mc_chs_chs

Recommended Citation

Atiullah, N., Fikree, F. F., Husain, I. (1998). An epidemiologic study of cystitis among Muslim women attending an out-patients clinic in Karachi. *Journal of Pakistan Medical Association*, 48(2), 35-37.

Available at: https://ecommons.aku.edu/pakistan_fhs_mc_chs_chs/396

An Epidemiologic Study of Cystitis among Muslim Women Attending An Out-Patients Clinic in Karachi

Pages with reference to book, From 35 To 37

N. Atiullah, F.E Fikree (Department of Community Health Sciences, The Aga Khan University, Karachi.)

I. Husain (Department of Urological Surgery, Bedford Hospital, Bedford, England.)

Abstract

Cystitis is considered to be one of the most common infections afflicting women of all age groups. It has been suggested that personal hygiene customs may influence the prevalence of cystitis. In this study, 130 women of child bearing age attending the out-patients clinic were interviewed in order to determine the frequency of cystitis and its associated risk factors including personal hygiene practices. Twenty seven percent of the women reported suffering from cystitis atleast once in the past, comparable to reported studies from the United States but higher than reported from other Muslim countries. Sixty-three percent of women reported their first episode during pregnancy and a significant relationship was observed between parity and cystitis (p value <0.000). However, personal hygiene practices following micturition or sexual intercourse bore no significant association with cystitis. Cystitis, commonly reported among Pakistani married women, occurs primarily during pregnancy. Preventive measures during pregnancy are suggested (JPMA 48:35.1998).

Introduction

Cystitis, an inflammation of the urinary bladder characterised by symptoms of frequency and burning dysuria, is most commonly due to infection. Cystitis is considered by some physicians to be the most common bacterial infection in women¹ and is second to common cold as a cause for out-patient visits. This is reflected in the high rates of cystitis: four out of five women in Britain will suffer the symptoms of cystitis at some time in their lives². The vaginal reservoir and the short urethra in the female predispose to bacterial advancement resulting in much higher rates of cystitis among women than men. Predisposing risk factors include sexual activity^{3,4}, use of diaphragm for contraception, catheterization^{5,6} and a past history of cystitis⁷. Personal hygiene customs have been shown to affect the prevalence of cystitis. It has been observed that careful washing with soap and water before and after sexual intercourse has proved most effective in reducing risk of re-infection in susceptible women⁸. On the other hand, a study to demonstrate a linkage between methods of wiping with toilet paper and the genesis of urinary tract infections was inconclusive due to difficulty in interpretation of responses from patients⁹. The use of toilet paper is virtually unknown in Muslim communities, where personal cleansing is effected exclusively with water¹⁰. It is feasible that this fact alone could simply account for the smaller number of women experiencing non-specific cystitis in these populations¹¹. This paper reports on the frequency of cystitis among women from a facility-based population survey of an indigenous Muslim population in Karachi and examines the role played by Islamic personal hygiene customs in the frequency of cystitis. This study is part of a multi-centered study which aims to relate differences in the frequency of cystitis to variation in socio-cultural and personal hygiene practices among Islamic and non-Islamic communities in Norway, England, Saudi Arabia and Pakistan.

Materials and Methods

The study was carried out among women presenting to the consulting clinics of the Aga Khan

University Hospital (AKUH), Karachi, during the period November 27 -December 1, 1994. Though AKUH is a private, tertiary care center, it caters to all socioeconomic strata of the city. Therefore, it may be assumed that a random selection of women attending the consulting clinics would be a true representation of the female population of Karachi. Pre-coded, close-ended, standard questionnaires were administered to 130 women in the child bearing age range (15-45 years) by a single interviewer, a physician. She randomly selected respondents from the pool of women in the waiting room area and based on our selection criteria, proceeded to conduct the interviews. To eliminate selection bias, the eligibility criteria for selecting subjects was restricted to Muslim females who were accompanying patients to the clinics and were not patients themselves. The information obtained included past history of ever suffering from cystitis [frequency, burning and dysuria], age, marital status, sexual activity, parity and personal hygiene habits related to micturition and sexual activity. To assess socio-economic status, women were asked to report the gross monthly family income and the area of their residence. Different areas of the city were classified, according to standard norms, as high, upper middle, middle, lower middle and lower income group areas. There was a statistically significant correlation between reported household income and area of residence ($p=0.000$). The range in reported gross monthly family income and area of residence supported our assumption that AKUH caters to all socioeconomic strata of the city (Table I). The frequency of cystitis was defined as the percentage of women who ever reported suffering from cystitis while demographic and socio-cultural practices are those reported currently.

Results

The mean age of the 130 women interviewed was 31 ± 7.2 years. Of these, 28 (21.5%) were single and the remainder were currently either married (73.1%) or widowed/divorced (5.4%). The mean monthly income reported was Rs.8,900 \pm 7,003 (Table 1). Regarding socio-cultural practices among currently married women, 82% (78/95) reported frequency of sexual intercourse at least once per week, 62% (59/95) washed both before and after sexual intercourse while only 24% (23/95) reported washing only after sexual intercourse (Table I).

Table I. Demographic and socio-cultural practices among 130 Muslim women attending an out-patient clinic, Karachi, Pakistan.

Characteristics	Women (#)	Percentage (%)
Age:		
20	4	3.1
20-29	52	40.0
30-39	53	40.8
≥40	21	16.6
Marital status:		
Single	28	21.5
Married	95	73.1
Widowed/Divorced	7	5.4
Parity:		
Nulliparous	19	14.6
1 - 3	47	36.2
>3	36	27.7
Single	28	21.5
SES status¹		
High	38	29.2
Middle		
upper	23	17.7
middle	44	33.8
low	19	14.6
Lower	6	4.6
Family income:		
<5000	61	46.9
5000-9999	12	9.2
10000-14999	29	22.3
≥15000	28	21.5
Occupation:		
Housewife	107	82.3
Fulltime employee	12	9.2
Part time employee	11	8.5
Intercourse frequency³		
Atleast once	78	82.1
More than once	17	17.9
Personal hygiene:		
<i>After micturition:</i>		
Wash ⁴ only	115	88.5
Wash ⁴ - Toilet paper	15	11.5
<i>Pre-sex washing⁵:</i>		
Yes	59	62.1
No	36	37.9
<i>Post-sex washing⁶</i>		
Yes	82	86.3
No	13	13.7

1 = Socioeconomic (SES) status as described by residential area.

2 = Mean family income in Rs/month

3 = Per week; n=95 - single women excluded

4 = Usage of water for personal cleansing

5 = Usage of water for personal cleansing prior to sexual intercourse; n=95, single women excluded

6 = Usage of water for personal cleansing prior after sexual intercourse; n=95 - single women excluded

Thirty-five (26.9%) women reported suffering from cystitis atleast once in the past. Repeated episodes of cystitis were reported by 40%(14/35) of cystitis sufferers. Twenty-two (63%) women had their first episode during pregnancy while the remaining reported the first episode either during childhood (2/35=5.7%) or during their honeymoon (6/35=17%). Six (17%) of these 35 cystitis sufferers reported

diabetes as a concurrent illness (Table II).

Table II. Characteristics of women ever reporting cystitis, Karachi, Pakistan.

Characteristics	Women (#)	Percentage (%)
Ever suffered:		
Yes	35	26.9
No	95	73.1
First episode:		
Childhood	2	5.7
Honeymoon	6	17.1
Pregnancy	22	62.9
Any later time	5	14.3
Number of episodes:		
One	21	60.0
More than once	14	40.0

Ever-married women were at a significantly higher risk of developing cystitis as compared to single women. In addition, a significant linear trend was observed for parity with women reporting three or more live births at greatest risk of cystitis as compared to single women. Women reporting an average monthly income of Rs.5,000 and over were at a significantly higher risk of cystitis as compared to lower income women. However, there was no significant association between cystitis and personal hygiene habits (Table III).

Table III. Relationship of selected demographic and socio-cultural practices to cystitis and non-cystitis women.

Characteristics	Women reporting cystitis				P value
	Yes (n=35)		No,(n=95)		
	n	%	n	%	
Age:					
<20	0	0.0	4	4.2	0.09
20-29	10	28.6	42	44.2	
30-39	17	48.6	36	37.9	
≥40	8	22.8	13	13.7	
Marital status:					
Single	1	2.8	27	28.4	0.004
Ever-married	34	97.2	68	71.6	
Parity:					
Single	1	2.8	27	28.4	0.000
0-3	19	54.3	47	49.5	
>3	15	42.9	21	22.1	
Family income ¹ :					
<5000	6	17.1	55	57.9	0.000
≥5000	29	82.9	40	42.1	
Personal hygiene:					
After bowel movement:					
Wash ² only	29	82.9	86	90.5	0.36
Wash ² + Toilet paper	6	17.1	9	9.5	
Pre-sex washing ³					
Yes	21	65.6	38	60.3	0.78
No	11	34.4	25	39.7	
Post-sex washing ⁴					
Yes	27	84.4	55	87.3	0.94
No	5	15.6	8	12.7	

1 = Mean family income in Rs/month.

2 = Usage of water for personal cleansing

3 = Usage of water for personal cleansing prior to sexual intercourse

4 = Usage of water for personal cleansing prior after sexual intercourse.

3 and 4 = Cystitis n=32. Non-cystitis n=63; single and divorced/widowed women excluded.

Discussion

A frequency of ever suffering from cystitis of 27% is reported here which suggests that almost three out

of every ten women suffer from cystitis sometime during their life. Remis et al reports that an estimated 25-35% of women aged 20-40 years in the United States suffer atleast one episode of urinary tract infection while a higher prevalence of four out of every five women is reported from Britain³. A frequency of 27% is comparable to reports from the United States but much lower than those from Great Britain³. However, reports from other muslim communities suggest that the frequency in this study is much higher¹¹. One possible explanation of the higher frequency of cystitis may be the high fertility rate of Pakistani. Not only was a significant relationship between parity and cystitis observed but approximately two-thirds of the cystitis was reported as occurring during pregnancy. It is therefore suggested that preventive measures, such as careful washing with soap and water before and after sexual intercourse and after defecation, will reduce the risk of re-infection in susceptible pregnant women. In this study no association was found between personal cleansing hygiene following defecation or sexual activity and cystitis, though ever-married women are observed to be at significantly higher risk of suffering from cystitis as compared to single women (Table III). It appears that the lack of an association is primarily due to the majority of women following the Islamic tradition of washing before and after sexual intercourse as shown in Table I. The larger multi-centered study aimed to relate differences in the frequency of cystitis to variation in socio-cultural and personal hygiene practices among Islamic and non-Islamic communities in Norway, England, Saudi Arabia and Pakistan will demonstrate the association of the Islamic tradition of cleansing following defecation and prior and after sexual activity with cystitis.

References

1. Tolckoff, N.E. and Rubin, R.H. New approaches to the treatment of UTI. *Am. J. Med.*, 1987;82:270-277,
2. Williams, J. The burning issue. Cystitis update. *Fam. Health*, 1994;10:17-19.
3. Kunin, C.M and McCormack, R.C. An epidemiologic study of bacteriuria and blood pressure among nuns and working women. *N. Engl. J. Med.*, 1968;278 :635-642.
4. Remia, R.S., Gurwith, M.J., Gurwith, D. et al. Risk factors for U.T.I. *Am.J. Epidemiol.*, 1987;126:685-689.
5. Kaye, M., de-Vriea, J. and Mac-Farlane, K.T. The initiation of U.T.I following a single bladder catheterization. *Can. Med. Assoc. J.*, 1962;86 :9-12.
6. Theil, O., Spuhler, O. U.T.I. by catheter and the so called infectious (episomal) resistance. *Schweiz. Med. Wochenachr.*, 1965;95: 1155-1163.
7. Kraft, J.K. and Stamey, T.A. The natural history of symptomatic recurrent bacteriuria in women. *Medicine*, 1977;56:55-60.
8. Kilmartin, A. *Understanding cystitis*. London, Arrow, 1992.
9. Marsh, P.P., Murray, A., Panchamia, P. The relationship between bacterial cultures of the vaginal introitus and urinary infection. *Br. J. Urol.*, 1972;44:368-472.
10. Husain, I. and Kazim, E. Urological evaluation of the tropical patient. In: Husain, I. (ed): *Tropical Urology*. Edinburgh and London, Churchill Living stone, 1984, pp. 11-12.
11. Kazim, E., Ansari, E.M. and Husain, I. The spectrum of genitourinary disease seen in Abu Dhabi. *Emirates Med. J.*, 1981;2:76-83.
12. *Pakistan Demographic and Health Survey 1990/91*. National Institute of Population Studies, Islamabad. Pakistan and IRD. Calverton, MD, USA, Marco International Inc., 1992.