



7-2012

Diffusion Tractography in Spinal Cord Injury: Preliminary Experience

Vikas Bhattia
IGMC, Shimla

R.G Sood
IGMC, Shimla

Sanjiv Sharma
IGMC, Shimla

Prashant Panda
IGMC, Shimla

Follow this and additional works at: <https://ecommons.aku.edu/pjns>

 Part of the [Neurology Commons](#)

Recommended Citation

Bhattia, Vikas; Sood, R.G; Sharma, Sanjiv; and Panda, Prashant (2012) "Diffusion Tractography in Spinal Cord Injury: Preliminary Experience," *Pakistan Journal of Neurological Sciences (PJNS)*: Vol. 7 : Iss. 2 , Article 6.
Available at: <https://ecommons.aku.edu/pjns/vol7/iss2/6>

DIFFUSION TRACTOGRAPHY IN SPINAL CORD INJURY: PRELIMINARY EXPERIENCE

Vikas Bhattia, R. G. Sood, Sanjiv Sharma, Prashant Panda
Department of radiodiagnosis, IGMC, Shimla

Correspondence to: Dr. Vikas Bhattia, M.D., Department of radiodiagnosis, IGMC, Shimla. Email: drvikasbhattia@gmail.com

ABSTRACT

Diffusion tractography is a new imaging technique to display the traumatic region for valuable preoperative information with prognostic significance. We present a case of 54 year male with quadriplegia showing reduced FA values and increased ADC in injured segment with depiction of disrupted tracts on tractography. In the near future, DTI will become a part of routine imaging in trauma patients as experience with this novel method gains acceptance.

Key words: DTI, Spinal trauma, MRI

INTRODUCTION

Spinal cord injuries result in damage to the myelinated fibers of the spinal cord and/or nerve roots, MRI can detect these changes as increased signal intensity on T2W and STIR images. However diffusion tensor imaging (DTI) can be used to detect abnormalities in the spinal cord, even in cases where routine MRI may be normal. Role of diffusion tensor imaging has been established in brain imaging however spinal cord DTI has remained largely as a research tool due to issues like small size of the spinal cord, its compactness and susceptibility to motion artifacts.

CASE REPORT

We present a case of 54 year old male who sustained trauma at a road side accident, following trauma he developed quadriplegia. There was evidence of bladder and bowel incontinence. On CNS examination there was evidence of power grade 0 in all the limbs with areflexia. Cervical and dorsal spine MRI was requested to rule out cord injury. MRI was done using a 1.5-T machine (Siemens, somatom) 1.5 T image. STIR and T2W images in the sagittal and axial planes were obtained for cervical and dorsal spine. DTI was performed in the axial and sagittal planes, using a phased array spine coil with the following parameters: 20 directions EPI tensor imaging TR: 8000; TE: 97.6; b value: 1000; frequency: 128; phase: 128; FOV (field of view): 26 X 20.8; slice thickness: 5 mm with no interslice gap. On MRI there was altered SI involving the C7 vertebra which was hypointense on T1w and hyperintense on T2w and STIR sequences s/o edema. There was altered SI and expansion of cord involving the C7-T1 level with similar SI characteristics as C7

vertebra suggestive of cord oedema. On DTI imaging the FA value was $.25 + .18$ which was significantly reduced than control value $.56 + .14$ at proximal to the injury. ADC value of $937 + 112$ was present at the site of injury which was increased as compared to $432 + 110$ at proximal level. Tractographic image clearly demonstrated disruption of the fibres bilaterally. Patient was managed conservatively and later underwent screw fixation to stabilize the vertebrae. There was only minimal recovery of the paresis in all the limbs.



Figure 1: sagittal STIR image showing hyperintensity involving the cord (white arrow) and C7 vertebra (black arrow)



Figure 2: sagittal T2w image showing hyperintensity involving the cord with oedema(white arrow)and C7 vertebra(black arrow)

DISCUSSION

Demonstration of human spinal cord injury by DTI tractography techniques has been described by only few studies previously. Role of diffusion tensor imaging has been established in brain imaging however spinal cord DTI has remained largely as a research tool due to issues like small size of the spinal cord, its compactness and susceptibility to motion artifacts.¹ Some studies have used 9 Tesla DTI to clearly demonstrate the discontinuous cortico-spinal tracts after a hemisection injury in common marmosets.² But with current setup high strength MRI is not clinically applicable for imaging humans in routine clinical practice. Studies documenting the role of DTI in acute spinal cord

injuries have focused mainly on DTI anisotropy indices rather than tractography, which is more representative than the former.³ The clinical application of tensor imaging in spinal cord lesions due to trauma, tumors, and inflammation has shown the usefulness of this technique. DTI has been successfully utilised to demonstrate displaced white matter tracts or their involvement by lesions in the cord. This has revolutionised the treatment planning and follow-up of cases.⁴ Tensor imaging can show changes in white matter tracts even in cases where routine imaging is normal. Even when routine CT scan and MRI normal, the tractography has been proven to be of utility as there was reduction in diffusion anisotropy after suggesting axonal injury.⁵ Reduced FA in the cervical cord has been demonstrated in patients as compared to controls, in demyelinating disease such as multiple sclerosis, even when routine MRI imaging was normal.⁶ It has been seen that DTI offers better clinical correlation with neurological deficit whereas the routine MRI may not show the abnormality and shows lesser correlation with the motor deficits.⁷ Only experimental and high strength use of DTI in spinal trauma patients has misled the investigators regarding the practical use of DTI in evaluating spinal cord diseases. But results from newer studies suggest that DTI of spinal cord is technically and practically possible with a 1.5 Tesla MRI.⁸ DTI may be beneficial in quantification of spinal injury prior to surgical treatment and help in prognostic postsurgical outcome. It can also determine the extent of injury particular below the level of diagnosed injury.⁸ Studies have shown that injured segments show evidence of reduced FA and increased ADC values as compared to normal areas with depiction of disruption on tractography images.⁹




Figure 3: tractography images showing the disrupted fibres (thin white arrow)




Figure 4: tractography images showing the disrupted fibres (thin black arrow)

REFERENCES

1. Diffusion Tensor Tractography Demonstration of Partially Injured Spinal Cord Tracts in a Patient with Posttraumatic Brown Sequard Syndrome Shanmughanathan Rajasekaran, PhD,* Rishi Mugesh Kanna, MS, Rajamanickam Karunanithi, PhD, and Ajoy Prasad Shetty, MS *JOURNAL OF MAGNETIC RESONANCE IMAGING* 32:978–981 (2010)
2. Fujiyoshi K, Yamada M, Nakamura M, et al. In vivo tracing of neural tracts in the intact and injured spinal cord of marmosets by diffusion tensor tractography. *J Neurosci* 2007;27:11991–11998
3. Shanmuganathan K, Gullapalli RP, Zhuo J, Mirvis SE. Diffusion tensor imaging in cervical spine trauma. *AJNR Am J Neuroradiol* 2008;29:655–659
4. Vargas MI, Delavelle J, Jlassi H, Rilliet B, Viallon M, Becker CD, et al. Clinical applications of diffusion tensor tractography of the spinal cord. *Neuroradiology* 2008;50:25-9
5. Inglese M, Makani S, Johnson G, Cohen BA, Silver JA, Gonen O, et al. Diffuse axonal injury in mild traumatic brain injury: A diffusion tensor imaging study. *J Neurosurg* 2005;103:298-303.
6. Hesseltine SM, Law M, Babba J, Rada M, Lopeza S, Gea Y, et al. Diffusion tensor imaging in multiple sclerosis: Assessment of regional differences in the axial plane within normal-appearing cervical spinal cord. *AJNR Am J Neuroradiol* 2006;27:1189-93.
7. Chang Y, Jung TD, Yoo DS, Hyun JK. Diffusion tensor imaging and fiber tractography of patients with cervical spinal cord injury. *J Neurotrauma* 2010;27:2033-40.
8. Ellingson BM, Schmit BD, Ulmer JL, Kurpad SN. Diffusion tensor magnetic resonance imaging in spinal cord injury. *Concepts Magn Reson Part A* 2008;32A:219–237.
9. Kamble RB, Venkataramana NK, Naik AL, Rao SV. Diffusion tensor imaging in spinal cord injury. *Indian J Radiol Imaging* 2011;21:221-4