

12-10-2018

# A Proficient Two-Stage Stratified Randomized Response Strategy


Tanveer A. Tarray

*Islamic University of Science and Technology, Awantipora, India, tanveerstat@gmail.com*

Housila P. Singh

*Vikram University, Ujjain, India, hpsujn@gmail.com*

Follow this and additional works at: <https://digitalcommons.wayne.edu/jmasm>

 Part of the [Applied Statistics Commons](#), [Social and Behavioral Sciences Commons](#), and the [Statistical Theory Commons](#)

## Recommended Citation

Tarray, Tanveer A. and Singh, Housila P. (2018) "A Proficient Two-Stage Stratified Randomized Response Strategy," *Journal of Modern Applied Statistical Methods*: Vol. 17 : Iss. 1 , Article 29.

DOI: 10.22237/jmasm/1544453468

Available at: <https://digitalcommons.wayne.edu/jmasm/vol17/iss1/29>

This Regular Article is brought to you for free and open access by the Open Access Journals at DigitalCommons@WayneState. It has been accepted for inclusion in Journal of Modern Applied Statistical Methods by an authorized editor of DigitalCommons@WayneState.

# A Proficient Two-Stage Stratified Randomized Response Strategy

**Tanveer A. Tarray**

Islamic University of Science and Technology  
Awantipora, India

**Housila P. Singh**

Vikram University  
Ujjain, India

---

A stratified randomized response model based on R. Singh, Singh, Mangat, and Tracy (1995) improved two-stage randomized response strategy is proposed. It has an optimal allocation and large gain in precision. Conditions are obtained under which the proposed model is more efficient than R. Singh et al. (1995) and H. P. Singh and Tarray (2015) models. Numerical illustrations are also given in support of the present study.

*Keywords:* Randomized response technique, stratified random sampling, dichotomous population, Sensitive attribute, estimation of proportion, optimum allocation

---

## Introduction

Warner (1965) introduced a randomized response (RR) model to estimate proportion for sensitive attributes including sexual orientation, criminal activity, child abuse, suicidal tendency in teenagers, all cases of AIDS, abortion, or drug addiction. Greenberg, Abul-Ela, Simmons, and Horvitz (1969) envisaged an unrelated question randomized response model using Warner's sensitive question and unrelated question. This technique has generated much interest in the statistical literature since the publication of Warner's RR model. The RR model has been studied by many authors, such as Chaudhuri and Muherjee (1988), Ryu, Hong, and Lee (1993), Tracy and Mangat (1996), S. Singh (2003), and Kim and Elam (2007).

Hong, Yum, and Lee (1994) suggested a stratified RR technique using a proportional allocation. Kim and Warde (2004) presented a stratified RR technique based on Warner's (1965) model that has an optimal allocation. Kim and Elam (2005) have applied Kim and Warde's (2004) stratified Warner's RR model to the two-stage model of Mangat and Singh (1990). Kim, Tebbs, and An (2006) proposed a Bayesian version of the Mangat (1994) model. Kim and Warde (2005) have

---

suggested a mixed randomized response model using simple random sampling which rectifies the privacy problem and extended the proposed model to stratified sampling. Lee, Uhm, and Kim (2011) have extended the work of Land, Singh, and Sedory (2011) in stratified sampling using a Poisson distribution.

Recently, H. P. Singh and Tarray (2015) have applied a stratified RR model to the Tracy and Osahan (1999) two-stage model and derived the probability  $\theta_i$  of “Yes” answer in stratum  $i$  for this procedure as:

$$\theta_i = T_i \pi_{Si} + (1 - T_i) \left[ P_i \pi_{Si} + \frac{(1 - P_i)}{2} \right], \quad \text{for } i = 1, 2, \dots, k \quad (1)$$

where  $\pi_{Si}$  is the proportion of people with the sensitive trait in a stratum  $i$ . The unbiased estimator  $\hat{\pi}_{ST}$  of a sensitive proportion estimate of  $\pi_S = \sum_{i=1}^k w_i \pi_{Si}$  is given by

$$\hat{\pi}_{ST} = \sum_{i=1}^k w_i \hat{\pi}_{Si} = \sum_{i=1}^k w_i \left[ \frac{\hat{\theta}_i - (1 - T_i)(1 - P_i)/2}{T_i + P_i(1 - T_i)} \right] \quad (2)$$

where  $w_i = (N_i / N)$  for  $i = 1, 2, \dots, k$ , so that  $w = \sum_{i=1}^k w_i = 1$ ;  $N$  is the number of units in the entire population;  $N_i$  is the total number of units in the stratum  $i$ ; and  $\hat{\theta}_i$ , the proportion of “Yes” answers obtained from the  $n_i$  sampled respondents using simple random sampling with replacement (SRSWR) of stratum  $i$ , is a point estimate of  $\theta_i$  in (1).

The minimal variance of the estimator  $\hat{\pi}_{ST}$  under optimal allocation is given by

$$V(\hat{\pi}_{ST}) = \frac{1}{n} \left[ \sum_{i=1}^k w_i \left\{ \pi_{Si} (1 - \pi_{Si}) + \frac{(1 - T_i)(1 - P_i) \{ 2 - (1 - T_i)(1 - P_i) \}}{4 \{ T_i + P_i(1 - T_i) \}^2} \right\}^{1/2} \right]^2 \quad (3)$$

In this paper, we have suggested a new stratified randomized response model based on the R. Singh et al. (1995) two-stage RR model. It is demonstrated that the estimator resulting from the proposed model is more efficient than those of its

## A PROFICIENT TWO-STAGE STRATIFIED RRS

competent models under the condition presented in the case of completely truthful reporting.

### Suggested Model

The population is divided in to  $k$  strata and a sample is selected by SRSWR in each stratum. To get the full benefit from stratification, we assume that the number of units in each stratum is known. In the first stage of the survey interview, an individual respondent in the sample of stratum  $i$  is instructed to use the randomization device  $R_{1i}$ , which consists of a sensitive question ( $S$ ) card with probability  $T_i$  and a “Go to randomization device  $R_{2i}$  in the second stage” direction card with probability  $(1 - T_i)$ . The respondents in the second stage of the stratum  $i$  are instructed to use the randomization device  $R_{2i}$  which uses three statements: (i) “I possess the sensitive attribute  $A$ ,” (ii) “Yes;” and (iii) “No;” with probabilities  $P_i$ ,  $(1 - P_i)\alpha_i$ , and  $(1 - P_i)(1 - \alpha_i)$ , respectively, where  $\alpha_i \in [0, 1]$ . Let  $n_i$  denote the number of units in the sample from stratum  $i$  and  $n$  denote the total number of units in sample from all stratum so that  $n = \sum_{i=1}^k n_i$ . Thus, when respondents report truthfully, the probability of a "Yes" answer in stratum  $i$  for this procedure is given by

$$\theta_{ai} = T_i\pi_{Si} + (1 - T_i)[P_i\pi_{Si} + \alpha_i(1 - P_i)], \quad \text{for } i = 1, 2, \dots, k \quad (4)$$

where  $\theta_{ai}$  is the proportion of “Yes” responses and  $\pi_{Si}$  is the proportion of respondents with the sensitive trait in the population from stratum  $i$ .

If  $\hat{\theta}_{ai}$  denotes the estimate of the proportion of “Yes” answers in a stratum  $i$ , the relation (4) yields

$$\hat{\pi}_{aSi} = \frac{\hat{\theta}_{ai} - (1 - T_i)(1 - P_i)\alpha_i}{T_i + P_i(1 - T_i)}, \quad \text{for } i = 1, 2, \dots, k \quad (5)$$

as an unbiased estimator of the proportion  $\pi_{Si}$  of respondents with the sensitive trait in the population from stratum  $i$ .

The variance of the estimator  $\hat{\pi}_{aST}$  is given by

$$V(\hat{\pi}_{\alpha Si}) = \left[ \frac{\pi_{Si}(1-\pi_{Si})}{n_i} + \frac{(1-T_i)(1-P_i) \left[ \pi_{Si} \{T_i + P_i(1-T_i)\} (1-2\alpha_i) + \alpha_i \{1-(1-T_i)(1-P_i)\alpha_i\} \right]}{n_i [T_i + (1-T_i)P_i]^2} \right] \quad (6)$$

Because the selection in different strata are made independently, the estimator for individual strata can be added together to obtain an estimator for the whole population. The maximum likelihood estimator of  $\pi_s$ , the proportion of respondents with the sensitive trait, is

$$\hat{\pi}_s = \sum_{i=1}^k w_i \hat{\pi}_{\alpha Si} = \sum_{i=1}^k w_i \left[ \frac{\hat{\theta}_{\alpha i} - (1-T_i)(1-P_i)\alpha_i}{T_i + P_i(1-T_i)} \right] \quad (7)$$

where  $N$  denotes the number of units in the entire population,  $N_i$  is the total number of units in the stratum  $i$ , and  $w_i = (N_i / N)$  for  $i = 1, 2, \dots, k$ , so  $w = \sum_{i=1}^k w_i = 1$ . The proposed estimator  $\hat{\pi}_s$  is an unbiased estimator for the population proportion  $\pi_s$ .

**Theorem 1.** The variance of the estimator  $\hat{\pi}_s$  is

$$V(\hat{\pi}_s) = \sum_{i=1}^k \frac{w_i^2}{n_i} \left[ \frac{\pi_{Si}(1-\pi_{Si})}{n_i} + \frac{(1-T_i)(1-P_i) \left[ \pi_{Si} \{T_i + P_i(1-T_i)\} (1-2\alpha_i) + \alpha_i \{1-(1-T_i)(1-P_i)\alpha_i\} \right]}{n_i [T_i + (1-T_i)P_i]^2} \right] \quad (8)$$

**Proof.** This follows from taking the variance of (7) and corollary 1 in section 5.5 of Cochran (1977).

In practice, information on  $\pi_{Si}$  is usually not known. But if prior information on  $\pi_{Si}$  is available from past experience then it helps to establish the following optimal allocation formula:

**Theorem 2.** The optimal allocation of  $n$  to  $n_1, n_2, \dots, n_{k-1}$ , and  $n_k$  to derive the minimum variance of  $\hat{\pi}_s$  subject to  $n = \sum_{i=1}^k n_i$  is approximately given by

## A PROFICIENT TWO-STAGE STRATIFIED RRS

$$\begin{aligned}
 \frac{n_i}{n} = & \\
 & w_i \left[ \pi_{Si} (1 - \pi_{Si}) \right. \\
 & \left. + \frac{(1 - T_i)(1 - P_i) \left[ \pi_{Si} \{T_i + P_i(1 - T_i)\} (1 - 2\alpha_i) + \alpha_i \{1 - (1 - T_i)(1 - P_i)\alpha_i\} \right]}{n_i [T_i + (1 - T_i)P_i]^2} \right]^{\frac{1}{2}} \\
 & \frac{\sum_{i=1}^k w_i \left[ \pi_{Si} (1 - \pi_{Si}) \right. \\
 & \left. + \frac{(1 - T_i)(1 - P_i) \left[ \pi_{Si} \{T_i + P_i(1 - T_i)\} (1 - 2\alpha_i) + \alpha_i \{1 - (1 - T_i)(1 - P_i)\alpha_i\} \right]}{n_i [T_i + (1 - T_i)P_i]^2} \right]^{\frac{1}{2}}}{\sum_{i=1}^k} \quad (9)
 \end{aligned}$$

**Proof.** Follows from section 5.5 of Cochran (1977).  
The minimal variance of the estimator  $\hat{\pi}_s$  is given by

$$\begin{aligned}
 V(\hat{\pi}_s) = & \frac{1}{n} \left[ \sum_{i=1}^k w_i \left\{ \frac{\pi_{Si} (1 - \pi_{Si})}{n_i} \right. \right. \\
 & \left. + \frac{\pi_{Si} \{T_i + P_i(1 - T_i)\} (1 - 2\alpha_i) (1 - T_i)(1 - P_i)}{n_i [T_i + (1 - T_i)P_i]^2} \right. \\
 & \left. \left. + \frac{\alpha_i \{1 - (1 - T_i)(1 - P_i)\alpha_i\} (1 - T_i)(1 - P_i)}{n_i [T_i + (1 - T_i)P_i]^2} \right\}^{\frac{1}{2}} \right]^2 \quad (10)
 \end{aligned}$$

By substituting  $(n_i - 1)$  for  $n_i$  in (8), the unbiased minimal variance of the estimator  $\hat{\pi}_s$  can be obtained.

**Remark 1.** The proposed procedure is equivalent to the H. P. Singh and Tarray (2015) procedure for  $\alpha = 1/2$ .

**Remark 2.** For any given pair  $(P_i, T_i)$ , the variance  $V(\hat{\pi}_{\alpha Si})$  at  $(\pi_{Si}, \alpha_i)$  and  $[(1 - \pi_{Si}), (1 - \alpha_i)]$  has equal value. This means that  $V(\hat{\pi}_{\alpha Si})$  is symmetric about the

point  $(1/2, 1/2)$  along any line in the  $(\pi_{Si}, \alpha_i)$  plane passing through this point; see Corollary 2.3 in R. Singh et al. (1995, p. 268).

### Relative Efficiency

An efficiency comparison of the proposed stratified randomized response technique and the two-stage randomized response techniques that were presented by R. Singh et al. (1995) will be conducted by comparing variances.

**Theorem 3.** Suppose there are two strata in the population,  $n = n_1 + n_2$ ,  $P = P_1 = P_2$  ( $P$  is the probability of selecting the sensitive question in the second stage),  $T = T_1 = T_2$  ( $T$  is the probability of selecting the sensitive question in the first stage), and  $\hat{\pi}_s = w_1 \hat{\pi}_{\alpha s_1} + w_2 \hat{\pi}_{\alpha s_2}$ . The proposed estimator  $\hat{\pi}_s$  is more efficient than the R. Singh et al. (1995) estimator  $\hat{\pi}_{SSMT}$  (say), where  $\pi_{s1} \neq \pi_{s2}$ , under the following conditions:

$$\left[ (\pi_{s1} - \pi_{s2})^2 + (\sqrt{A_1} - \sqrt{A_2})^2 \right] > \left\{ 1 + (w_1 w_2)^{-1} \right\} B \quad (11)$$

where

$$\begin{aligned} A_1 &= [\pi_{s1}(1 - \pi_{s1}) + A\pi_{s1} + B], A_2 = [\pi_{s2}(1 - \pi_{s2}) + A\pi_{s2} + B] \\ A &= \frac{(1-T)(1-P)\{T + P(1-T)\}(1-2\alpha)}{\{T + P(1-T)\}^2} \\ B &= \frac{(1-T)(1-P)\alpha[1 - (1-T)(1-P)\alpha]}{\{T + P(1-T)\}^2} \end{aligned}$$

The following theorem exhibits that the proposed estimator  $\hat{\pi}_s$  is more efficient than the H. P. Singh and Tarray (2015) estimator  $\hat{\pi}_{ST}$ :

**Theorem 4.** Assume that there are two strata in the population,  $n = n_1 + n_2$ ,  $P = P_1 = P_2$ , and  $T = T_1 = T_2$ . The proposed estimator  $\hat{\pi}_s$  will be always more efficient than the H. P. Singh and Tarray (2015) estimator  $\hat{\pi}_{ST}$  (where  $\pi_{s1} \neq \pi_{s2}$ ) under the following condition:

## A PROFICIENT TWO-STAGE STRATIFIED RRS

$$\text{Either } \alpha < \left[ \frac{1}{(1-T)(1-P)} - \frac{1}{2} \right] \text{ or } \alpha > \left[ \frac{1}{(1-T)(1-P)} - \frac{1}{2} \right] \quad (12)$$

### Empirical Study

To see the tangible idea about the performance of the proposed estimator  $\hat{\pi}_s$  over the H. P. Singh and Tarray (2015) estimator  $\hat{\pi}_{ST}$ , compute the percent relative efficiency (PRE) of  $\hat{\pi}_s$  with respect to  $\hat{\pi}_{ST}$  for two strata, i.e.  $k = 2$ ,  $P = P_1 = P_2$ , and  $T = T_1 = T_2$ , by using the formula

$$\begin{aligned} \text{PRE}(\hat{\pi}_s, \hat{\pi}_{ST}) &= \frac{V(\hat{\pi}_{ST})}{V(\hat{\pi}_s)} \times 100 \\ &= \frac{\left[ w_1 \sqrt{(G_1 + H)} + w_2 \sqrt{(G_2 + H)} \right]^2}{\left[ w_1 \sqrt{(G_1 + H_1)} + w_2 \sqrt{(G_2 + H_2)} \right]^2} \times 100 \end{aligned} \quad (13)$$

where

$$\begin{aligned} G_1 &= \pi_{s1}(1 - \pi_{s1}), G_2 = \pi_{s2}(1 - \pi_{s2}) \\ H &= \frac{(1-T_i)(1-P_i)\{2 - (1-T)(1-P)\}}{4\{T + P(1-T)\}^2} \\ H_1 &= \frac{(1-T)(1-P)\left[\pi_{s1}\{T + P(1-T)\}(1-2\alpha) + \alpha\{1 - (1-T)(1-P)\alpha}\right]}{\left[T + (1-T)P\right]^2} \\ H_2 &= \frac{(1-T)(1-P)\left[\pi_{s2}\{T + P(1-T)\}(1-2\alpha) + \alpha\{1 - (1-T)(1-P)\alpha}\right]}{\left[T + (1-T)P\right]^2} \end{aligned}$$

The computed  $\text{PRE}(\hat{\pi}_s, \hat{\pi}_{ST})$  for  $n = 1000$ ,  $\alpha = 0 (0.1) 0.4$ ,  $T = 0.1 (0.4) 0.9$ ,  $P = 0.6, 0.8$ ; and for  $\pi_{s1} = 0.28$ ,  $\pi_{s2} = 0.33$  and different values of  $w_1$  and  $w_2$  such that  $\pi_s \leq 0.5$ . Findings are shown in Table 1.



TARRAY & SINGH

**Table 1.** The relative efficiency of the proposed estimator  $\hat{\pi}_s$  with respect to the H. P. Singh and Tarray (2015) estimator  $\hat{\pi}_{ST}$ , i.e.  $RE(\hat{\pi}_s, \hat{\pi}_{ST}) = V(\hat{\pi}_{ST} / \hat{\pi}_s)$ , when  $n = 1000$ ,  $T = 0.1, 0.4, 0.9$ ,  $\pi_{S1} = 0.28$ ,  $\pi_{S2} = 0.33$ , and different values of  $w_1$  and  $w_2$

$\pi_{S1}$	$\pi_{S2}$	$w_1$	$w_2$	$\pi_s$	$\alpha$	$T$	$P$				
							0.30	0.33	0.36	0.39	0.42
0.28	0.33	0.9	0.1	0.285	0.0	0.6	145.05	141.75	138.58	135.60	132.80
0.28	0.33	0.9	0.1	0.285	0.0	0.7	128.99	127.10	125.29	123.55	121.88
0.28	0.33	0.9	0.1	0.285	0.0	0.8	118.54	115.59	114.66	113.76	112.89
0.28	0.33	0.8	0.2	0.290	0.1	0.6	140.19	137.56	134.56	131.56	129.53
0.28	0.33	0.8	0.2	0.290	0.1	0.7	125.53	123.89	122.32	120.82	119.37
0.28	0.33	0.8	0.2	0.290	0.1	0.8	114.28	113.47	112.67	111.90	111.50
0.28	0.33	0.7	0.3	0.295	0.4	0.6	129.61	127.61	125.70	123.90	122.18
0.28	0.33	0.7	0.3	0.295	0.4	0.7	117.43	116.34	115.29	114.29	113.32
0.28	0.33	0.7	0.3	0.295	0.4	0.8	108.64	108.14	107.65	107.18	106.72
0.28	0.33	0.6	0.4	0.300	0.0	0.6	144.20	140.91	137.82	134.90	132.15
0.28	0.33	0.6	0.4	0.300	0.0	0.7	128.39	126.54	124.76	123.05	121.41
0.28	0.33	0.6	0.4	0.300	0.0	0.8	116.16	115.23	114.33	113.44	112.59
0.28	0.33	0.5	0.5	0.305	0.1	0.6	139.46	136.61	133.91	131.37	128.96
0.28	0.33	0.5	0.5	0.305	0.1	0.7	125.02	123.41	121.87	120.38	118.96
0.28	0.33	0.5	0.5	0.305	0.1	0.8	113.97	113.17	112.39	111.63	110.90
0.28	0.33	0.4	0.6	0.310	0.4	0.6	129.11	127.13	125.25	123.47	121.78
0.28	0.33	0.4	0.6	0.310	0.4	0.7	117.10	116.03	115.00	114.01	113.05
0.28	0.33	0.4	0.6	0.310	0.4	0.8	108.46	107.96	107.49	107.02	106.13
0.28	0.33	0.3	0.7	0.315	0.0	0.6	143.37	140.13	137.09	134.22	131.51
0.28	0.33	0.3	0.7	0.315	0.0	0.7	127.81	125.99	124.24	122.56	120.95
0.28	0.33	0.3	0.7	0.315	0.0	0.8	115.80	114.89	114.00	113.14	112.30
0.28	0.33	0.2	0.8	0.320	0.1	0.6	138.74	135.93	133.28	130.77	128.40
0.28	0.33	0.2	0.8	0.320	0.1	0.7	124.52	122.94	121.42	119.96	118.56
0.28	0.33	0.2	0.8	0.320	0.1	0.8	113.66	112.87	112.11	111.37	110.65
0.28	0.33	0.1	0.9	0.325	0.4	0.6	128.62	126.66	124.81	123.05	121.38
0.28	0.33	0.1	0.9	0.325	0.4	0.7	116.78	115.72	114.71	113.73	112.79
0.28	0.33	0.1	0.9	0.325	0.4	0.8	108.28	107.80	107.32	106.87	106.43

Seven values of  $P$  and three different values of  $\alpha$  were set to verify the relative efficiency of the suggested estimator  $\hat{\pi}_s$  with respect to the H. P. Singh and Tarray (2015) estimator  $\hat{\pi}_{ST}$ . Shown in Table 1, the value of percent relative efficiency  $PRE(\hat{\pi}_s, \hat{\pi}_{ST})$  decreases as the values of  $P$  and  $\alpha$  increase.

The values of  $PRE(\hat{\pi}_s, \hat{\pi}_{ST})$  are greater than 100 for all values of  $\pi_{S1}$ ,  $\pi_{S2}$ ,  $w_1$ ,  $w_2$ ,  $P$ ,  $\alpha$ , and  $T$  considered here. The proposed estimator  $\hat{\pi}_s$  is more efficient than the H. P. Singh and Tarray (2015) estimator  $\hat{\pi}_{ST}$  in the case of the two strata in the population. It is further noted that, depending on the values of  $\pi_{S1}$ ,  $\pi_{S2}$ ,  $w_1$ , and  $w_2$ ,

## A PROFICIENT TWO-STAGE STRATIFIED RRS

and keeping the cooperation from the respondents in view, the value of  $\alpha$  as near to “Zero” or “Unity” as possible should be taken as an indication of a more efficient strategy.

### Acknowledgements

The authors are thankful to the Editor-in-Chief and to the anonymous learned referee for their valuable suggestions regarding improvement of the paper.

### References

- Chaudhuri, A., & Mukerjee, R. (1988). *Randomized response: Theory and techniques*. New York, NY: Marcel-Dekker.
- Cochran, W. G. (1977). *Sampling technique* (3rd ed.). New York: John Wiley and Sons.
- Greenberg, B., Abul-Ela, A., Simmons, W. R., & Horvitz, D. G. (1969). The unrelated question randomized response: Theoretical framework. *Journal of the American Statistical Association*, 64(326), 529-539. doi: 10.1080/01621459.1969.10500991
- Hong, K. H., Yum, J. K., & Lee, H. Y. (1994). A stratified randomized response technique. *Korean Journal of Applied Statistics*, 7(1), 141-147. Retrieved from [http://koreascience.or.kr/article/ArticleFullRecord.jsp?cn=GCGHDE\\_1994\\_v7n1\\_141](http://koreascience.or.kr/article/ArticleFullRecord.jsp?cn=GCGHDE_1994_v7n1_141)
- Kim, J.-M., & Elam, M. E. (2005). A two-stage stratified Warner's randomized response model using optimal allocation. *Metrika*, 61(1), 1-7. doi: 10.1007/s001840400319
- Kim, J.-M., & Elam, M. E. (2007). A stratified unrelated randomized response model. *Statistical Papers*, 48(2), 215-233. doi: 10.1007/s00362-006-0327-6
- Kim, J.-M., Tebbs, J. M., & An, S.-W. (2006). Extensions of Mangat's randomized-response model. *Journal of Statistical Planning and Inference*, 136(4), 1554-1567. doi: 10.1016/j.jspi.2004.10.005
- Kim, J.-M., & Warde, W. D. (2004). A stratified Warner's randomized response model. *Journal of Statistical Planning and Inference*, 120(1-2), 155-165. doi: 10.1016/s0378-3758(02)00500-1

- Kim, J.-M., & Warde, W. D. (2005). A mixed randomized response model. *Journal of Statistical Planning and Inference*, 133(1), 211-221. doi: 10.1016/j.jspi.2004.03.011
- Land, M., Singh, S., & Sedory, S. A. (2012). Estimation of a rare attribute using Poisson distribution. *Statistics*, 46(3), 351-360. doi: 10.1080/02331888.2010.524300
- Lee, G.-S., Uhm, D., & Kim, J.-M. (2011). Estimation of a rare sensitive attribute in a stratified sample using Poisson distribution. *Statistics*, 47(3), 575-589. doi: 10.1080/02331888.2011.625503
- Mangat, N. S. (1994). An improved randomized response strategy. *Journal of the Royal Statistical Society. Series B (Methodological)*, 56(1), 93-95. Retrieved from <https://www.jstor.org/stable/2346030>
- Mangat, N. S., & Singh, R. (1990). An alternative randomized procedure. *Biometrika*, 77(2), 439-442. doi: 10.1093/biomet/77.2.439
- Ryu, J. B., Hong, K. H., & Lee, G. S. (1993). *Randomized response model* [Unpublished manuscript]. Seoul, Korea: Freedom Academy.
- Singh, H. P., & Tarray, T. A. (2015). A stratified Tracy and Osahan's two-stage randomized response model. *Communications in Statistics – Theory and Methods*, 45(11), 3126-3137. doi: 10.1080/03610926.2014.895839
- Singh, R., Singh, S., Mangat, N. S., & Tracy, D. S. (1995). An improved two stage randomized response strategy. *Statistical Papers*, 36(1), 265-271. doi: 10.1007/bf02926038
- Singh, S. (2003). *Advanced sampling theory with applications*. Dordrecht, Netherlands: Kluwer Academic Publishers.
- Tracy, D. S., & Mangat, N. S. (1996). Some development in randomized response sampling during the last decade – A follow up of review by Chaudhuri and Mukherjee. *Journal of Applied Statistical Science*, 4(2/3), 147-158.
- Tracy, D. S., & Osahan, S. S. (1999). An improved randomized response technique. *Pakistan Journal of Statistics*, 15(1), 1-6.
- Warner, S. L. (1965). Randomized response: A survey technique for eliminating evasive answer bias. *Journal of the American Statistical Association*, 60(309), 63-69. doi: 10.1080/01621459.1965.10480775