

THE  
UNIVERSITY  
OF RHODE ISLAND

University of Rhode Island  
DigitalCommons@URI

---

Mathematics Faculty Publications

Mathematics

---

2005

# Augmented Implicitly Restarted Lanczos Bidiagonalization Methods

James Baglama

*University of Rhode Island*, [jbaglama@uri.edu](mailto:jbaglama@uri.edu)

Lothar Reichel

Follow this and additional works at: [https://digitalcommons.uri.edu/math\\_facpubs](https://digitalcommons.uri.edu/math_facpubs)

Terms of Use

All rights reserved under copyright.

---

## Citation/Publisher Attribution

Baglama, J., & Reichel, L. (2005). Augmented Implicitly Restarted Lanczos Bidiagonalization Methods. *SIAM J. Sci. Comput.*, 27(1), 19-42. doi: 10.1137/04060593X

Available at: <https://doi.org/10.1137/04060593X>

This Article is brought to you for free and open access by the Mathematics at DigitalCommons@URI. It has been accepted for inclusion in Mathematics Faculty Publications by an authorized administrator of DigitalCommons@URI. For more information, please contact [digitalcommons@etal.uri.edu](mailto:digitalcommons@etal.uri.edu).

## AUGMENTED IMPLICITLY RESTARTED LANCZOS BIDIAGONALIZATION METHODS\*

JAMES BAGLAMA<sup>†</sup> AND LOTHAR REICHEL<sup>‡</sup>

*Dedicated to Henk van der Vorst on the occasion of his 60th birthday*

**Abstract.** New restarted Lanczos bidiagonalization methods for the computation of a few of the largest or smallest singular values of a large matrix are presented. Restarting is carried out by augmentation of Krylov subspaces that arise naturally in the standard Lanczos bidiagonalization method. The augmenting vectors are associated with certain Ritz or harmonic Ritz vectors. Computed examples show the new methods to be competitive with available schemes.

**Key words.** singular value computation, partial singular value decomposition, iterative method, large-scale computation

**AMS subject classifications.** 65F15, 65F50, 15A18

**DOI.** 10.1137/04060593X

**1. Introduction.** Many problems in scientific computation require knowledge of a few of the largest or smallest singular values of a matrix and associated left and right singular vectors. These problems include the approximation of a large matrix by a matrix of low rank, the computation of the null space of a matrix, total least-squares problems (see, e.g., Björck [4, sections 4.6 and 7.6.5]), as well as the tracking of signals; see, e.g., Comon and Golub [8] for a discussion of the latter.

Let  $A \in \mathbb{R}^{\ell \times n}$  be a large sparse matrix. We may assume that  $\ell \geq n$ , because otherwise we replace the matrix by its transpose. Let

$$(1.1) \quad \sigma_1^{(A)} \geq \sigma_2^{(A)} \geq \dots \geq \sigma_n^{(A)} \geq 0$$

denote the singular values of  $A$ , and let  $u_j^{(A)} \in \mathbb{R}^\ell$  and  $v_j^{(A)} \in \mathbb{R}^n$ ,  $1 \leq j \leq n$ , be associated left and right singular vectors, respectively. Hence,

$$(1.2) \quad Av_j^{(A)} = \sigma_j^{(A)} u_j^{(A)}, \quad A^T u_j^{(A)} = \sigma_j^{(A)} v_j^{(A)}, \quad 1 \leq j \leq n,$$

and

$$A = \sum_{j=1}^n \sigma_j^{(A)} u_j^{(A)} (v_j^{(A)})^T.$$

The matrices  $U_n^{(A)} = [u_1^{(A)}, u_2^{(A)}, \dots, u_n^{(A)}]$  and  $V_n^{(A)} = [v_1^{(A)}, v_2^{(A)}, \dots, v_n^{(A)}]$  have orthonormal columns. We refer to  $\{\sigma_j^{(A)}, u_j^{(A)}, v_j^{(A)}\}$  as a singular triplet of  $A$ . Singular

---

\*Received by the editors March 29, 2004; accepted for publication (in revised form) October 26, 2004; published electronically August 17, 2005.

<http://www.siam.org/journals/sisc/27-1/60593.html>

<sup>†</sup>Department of Mathematics, University of Rhode Island, Kingston, RI 02881 (jbaglama@math.uri.edu, <http://hypatia.math.uri.edu/~jbaglama>). This author's research was supported in part by NSF grant DMS-0311786.

<sup>‡</sup>Department of Mathematical Sciences, Kent State University, Kent, OH 44242 (reichel@math.kent.edu, <http://www.math.kent.edu/~reichel>). This author's research was supported in part by NSF grant DMS-0107858.



restarts and in their implementation of the restarting procedure. Ideally, we would like  $p_1$  to be a linear combination of the right singular vectors of  $A$  associated with the desired singular values.

Sorensen [29] proposed efficient approaches for restarting the Arnoldi and the Lanczos tridiagonalization procedures. These approaches can be thought of as curtailed QR-algorithms, and, similarly to the QR-algorithms, their performance depends critically on the selection of shifts; see also [1, 7, 21, 30] for discussions and extensions. Björck, Grimme, and Van Dooren [5] derived analogous recursion formulas for a restarted Lanczos bidiagonalization method and presented an application to the solution of ill-posed problems. Recently, Kokiopoulou, Bekas, and Gallopoulos [17] applied these recursion formulas to compute a few desired singular triplets of a large sparse matrix. The shifts are applied by “chasing the bulge” in a curtailed QR-algorithm. Other implementations of restarted Lanczos bidiagonalization are discussed in [13, 14, 15, 16, 28]. The present paper describes mathematically equivalent, but numerically more robust, implementations of the methods discussed by Kokiopoulou, Bekas, and Gallopoulos [17].

This paper is organized as follows. Section 2 reviews Lanczos bidiagonalization and introduces notation used in the remainder of the paper. Section 3 describes our implementations of restarted Lanczos bidiagonalization. Restarting is carried out by augmenting the Krylov subspaces that arise naturally in the standard Lanczos bidiagonalization method by Ritz vectors or harmonic Ritz vectors associated with desired singular triplets. A few computed examples, which compare the performance of several methods and implementations, are presented in section 4.

**2. Lanczos bidiagonalization.** The following algorithm determines the partial Lanczos bidiagonalization (1.3)–(1.4) of the large sparse matrix  $A \in \mathbb{R}^{\ell \times n}$ . The number of Lanczos bidiagonalization steps,  $m$ , is typically much smaller than either one of the matrix dimensions  $\ell$  and  $n$ . Throughout this paper  $\|\cdot\|$  denotes the Euclidean vector norm or the associated induced matrix norm. We remark that the methods described can easily be modified to apply to matrices  $A$  with complex-valued entries; the main modification required is that the matrices  $P_m$ ,  $Q_m$ , and  $B_m$  in (1.3)–(1.4) may have complex-valued entries, and transposition has to be replaced by transposition and complex conjugation. The MATLAB code used for the computed examples of section 4 can be applied to matrices  $A$  with complex-valued entries.

ALGORITHM 2.1. LANCZOS BIDIAGONALIZATION.

*Input:*  $A \in \mathbb{R}^{\ell \times n}$  or functions for evaluating matrix-vector products  
with the matrices  $A$  and  $A^T$ ,  
 $p_1 \in \mathbb{R}^n$  : initial vector of unit length,  
 $m$  : number of bidiagonalization steps.

*Output:*  $P_m = [p_1, p_2, \dots, p_m] \in \mathbb{R}^{n \times m}$  : matrix with orthonormal columns,  
 $Q_m = [q_1, q_2, \dots, q_m] \in \mathbb{R}^{\ell \times m}$  : matrix with orthonormal columns,  
 $B_m \in \mathbb{R}^{m \times m}$  : upper bidiagonal matrix (1.6) with entries  $\alpha_j$  and  $\beta_j$ ,  
 $r_m \in \mathbb{R}^n$  : residual vector.

1.  $P_1 := p_1$ ;  $q_1 := Ap_1$ ;
2.  $\alpha_1 := \|q_1\|$ ;  $q_1 := q_1/\alpha_1$ ;  $Q_1 := q_1$ ;
3. for  $j = 1 : m$ 
  4.  $r_j := A^T q_j - \alpha_j p_j$ ;

5. *Reorthogonalization*:  $r_j := r_j - P_j(P_j^T r_j)$ ;
6. *if*  $j < m$ , *then*
  7.  $\beta_j := \|r_j\|$ ;  $p_{j+1} := r_j/\beta_j$ ;  $P_{j+1} := [P_j, p_{j+1}]$ ;
  8.  $q_{j+1} := Ap_{j+1} - \beta_j q_j$ ;
  9. *Reorthogonalization*:  $q_{j+1} := q_{j+1} - Q_j(Q_j^T q_{j+1})$ ;
  10.  $\alpha_{j+1} := \|q_{j+1}\|$ ;  $q_{j+1} := q_{j+1}/\alpha_{j+1}$ ;  $Q_{j+1} := [Q_j, q_{j+1}]$ ;
11. *endif*
12. *endfor*

When the computations with Algorithm 2.1 are carried out in finite precision arithmetic and the columns of  $P_m$  and  $Q_m$  are not reorthogonalized, the computed columns might be far from orthogonal. We therefore reorthogonalize the columns of these matrices in lines 5 and 9 of the algorithm.

Several reorthogonalization strategies for the columns of the matrices  $P_m$  and  $Q_m$  are discussed in the literature. Larsen [18] found that when  $m$  is fairly large and one is interested in computing a few of the largest singular triplets of  $A$ , partial reorthogonalization gives comparable accuracy and requires less computational work than full reorthogonalization, but when a few of the smallest singular triplets are desired, often (essentially) full reorthogonalization is required to achieve high accuracy. Wu and Simon [33] report that when  $m$  is not large, full reorthogonalization should be carried out, because due to the overhead associated with partial reorthogonalization, the latter is not competitive. Moreover, Simon and Zha [28] show that when the matrix  $A$  is not very ill-conditioned, only the columns of one of the matrices  $P_m$  or  $Q_m$  need to be reorthogonalized. Reorthogonalization of the columns of  $P_m$  only can reduce the computational effort required to compute the partial Lanczos bidiagonalization (1.3)–(1.4) considerably when  $\ell \gg n$ . Algorithm 2.1 can easily be modified to implement the latter approach.

The Lanczos bidiagonalization algorithm (Algorithm 2.1) requires the diagonal entries  $\alpha_j > 0$ ,  $1 \leq j \leq m$ , as well as the superdiagonal entries  $\beta_j$ ,  $1 \leq j < m$ , of  $B_m$  to be nonvanishing. Assume for the moment that  $\alpha_j > 0$  for  $1 \leq j < m$ , and  $\alpha_m = 0$ . Then the vector  $q_m$  determined in line 8 of the algorithm vanishes, and the decompositions (1.3)–(1.4) can be expressed as

$$AP_m = Q_{m-1}B_{m-1,m}, \quad A^T Q_{m-1} = P_m B_{m-1,m}^T,$$

where  $B_{m-1,m}$  denotes the leading  $(m-1) \times m$  submatrix of  $B_m$ . It follows that  $A$  is singular and that the singular values of  $B_{m-1,m}$  are singular values of  $A$ . The associated singular triplets of  $A$  can be determined from  $P_m$ ,  $Q_{m-1}$ , and the singular triplets of  $B_{m-1,m}$ . The singular values of  $B_{m-1,m}$  are nonvanishing. Moreover, the right singular vector of  $A$  associated with the zero singular value can be expressed as a linear combination of the columns of  $P_m$ , e.g., by using the singular value decomposition of  $B_m$ . However, the corresponding left singular vector of  $A$  is not readily available. For simplicity, we will in the remainder of this paper assume that all  $\alpha_j$  are positive.

It follows from Algorithm 2.1 that  $\beta_m$ , the last superdiagonal element of the upper bidiagonal matrix  $B_{m+1} \in \mathbb{R}^{(m+1) \times (m+1)}$  obtained by  $m+1$  steps of partial Lanczos bidiagonalization, is given by

$$(2.1) \quad \beta_m := \|r_m\|$$

and therefore can be computed when the decompositions (1.3)–(1.4) are available. We may assume that  $\beta_m > 0$ , because otherwise the singular values of  $B_m$  are singular

values of  $A$ , and the associated singular triplets of  $A$  can be determined from the singular value decomposition of  $B_m$  and the matrices  $P_m$  and  $Q_m$ . If  $\beta_m = 0$  and the determined singular triplets of  $A$  include all the desired ones, then we are done; otherwise we proceed to compute additional singular triplets by the methods described in this paper.

When  $\beta_m > 0$ , the last column of  $P_{m+1} := [P_m, p_{m+1}]$  is given by

$$(2.2) \quad p_{m+1} := r_m / \beta_m,$$

and the relation (1.4) can be expressed as

$$(2.3) \quad A^T Q_m = P_{m+1} B_{m,m+1}^T,$$

where the matrix  $B_{m,m+1} \in \mathbb{R}^{m \times (m+1)}$  is obtained by appending the column  $\beta_m e_m$  to  $B_m$ . In particular, the matrices in the decomposition (2.3) are available after  $m$  Lanczos bidiagonalization steps. We also note that  $B_{m,m+1}$  is the leading  $m \times (m+1)$  submatrix of the matrix  $B_{m+1}$  obtained after  $m+1$  steps of Lanczos bidiagonalization.

We will use the connection between partial Lanczos bidiagonalization (1.3)–(1.4) of  $A$  and partial Lanczos tridiagonalization of the matrix  $A^T A$ . Multiplying (1.3) by  $A^T$  from the left-hand side yields

$$(2.4) \quad A^T A P_m = P_m B_m^T B_m + r_m e_m^T B_m = P_m B_m^T B_m + \alpha_m r_m e_m^T,$$

where the last equality follows from the fact that  $B_m$  is upper bidiagonal. The matrix

$$(2.5) \quad T_m := B_m^T B_m \in \mathbb{R}^{m \times m}$$

is symmetric and tridiagonal, and the expression (2.4) is a partial Lanczos tridiagonalization of  $A^T A$  with initial vector  $p_1 = P_m e_1$ ; see, e.g., [11, section 9.1.2] for a discussion on Lanczos tridiagonalization. Since  $T_m$  is tridiagonal, (2.4) shows that the columns of  $P_m$  satisfy a three-term recurrence relation; the columns form an orthonormal basis of the Krylov subspace

$$(2.6) \quad \mathbb{K}_m(A^T A, p_1) = \text{span}\{p_1, A^T A p_1, (A^T A)^2 p_1, \dots, (A^T A)^{m-1} p_1\}.$$

Similarly, multiplying (1.4) by  $A$  from the left-hand side yields

$$(2.7) \quad A A^T Q_m = Q_m B_m B_m^T + A r_m e_m^T.$$

The columns of  $Q_m$  form an orthonormal basis of the Krylov subspace

$$(2.8) \quad \mathbb{K}_m(A A^T, q_1) = \text{span}\{q_1, A A^T q_1, (A A^T)^2 q_1, \dots, (A A^T)^{m-1} q_1\}$$

with  $q_1 := A p_1$ . We remark that since the columns of  $Q_m$  are not, in general, orthogonal to  $A r_m$ , the decomposition (2.7) typically is not a Lanczos tridiagonalization of  $A A^T$ .

Let  $\{\sigma_j^{(B_m)}, u_j^{(B_m)}, v_j^{(B_m)}\}$ ,  $1 \leq j \leq m$ , be the singular triplets of  $B_m$  enumerated so that

$$(2.9) \quad \sigma_1^{(B_m)} \geq \sigma_2^{(B_m)} \geq \dots \geq \sigma_m^{(B_m)} \geq 0.$$

Then, analogously to (1.2),

$$(2.10) \quad B_m v_j^{(B_m)} = \sigma_j^{(B_m)} u_j^{(B_m)}, \quad B_m^T u_j^{(B_m)} = \sigma_j^{(B_m)} v_j^{(B_m)}, \quad 1 \leq j \leq m,$$

and the  $m \times m$  matrices of left and right singular vectors

$$U_m^{(B_m)} := [u_1^{(B_m)}, u_2^{(B_m)}, \dots, u_m^{(B_m)}], \quad V_m^{(B_m)} := [v_1^{(B_m)}, v_2^{(B_m)}, \dots, v_m^{(B_m)}]$$

are orthogonal.

We determine approximate singular triplets  $\{\tilde{\sigma}_j^{(A)}, \tilde{u}_j^{(A)}, \tilde{v}_j^{(A)}\}$ ,  $1 \leq j \leq m$ , of  $A$  from the singular triplets of  $B_m$  by

$$(2.11) \quad \tilde{\sigma}_j^{(A)} := \sigma_j^{(B_m)}, \quad \tilde{u}_j^{(A)} := Q_m u_j^{(B_m)}, \quad \tilde{v}_j^{(A)} := P_m v_j^{(B_m)}.$$

Combining (2.10) with (1.3)–(1.4) shows that

$$(2.12) \quad A\tilde{v}_j^{(A)} = \tilde{\sigma}_j^{(A)}\tilde{u}_j^{(A)}, \quad A^T\tilde{u}_j^{(A)} = \tilde{\sigma}_j^{(A)}\tilde{v}_j^{(A)} + r_m e_m^T u_j^{(B_m)}, \quad 1 \leq j \leq m.$$

The equations (2.12) suggest that an approximate singular triplet  $\{\tilde{\sigma}_j^{(A)}, \tilde{u}_j^{(A)}, \tilde{v}_j^{(A)}\}$  be accepted as a singular triplet of  $A$  if  $r_m e_m^T u_j^{(B_m)}$  is sufficiently small. Specifically, our numerical method accepts  $\{\tilde{\sigma}_j^{(A)}, \tilde{u}_j^{(A)}, \tilde{v}_j^{(A)}\}$  as a singular triplet of  $A$  if

$$(2.13) \quad \beta_m |e_m^T u_j^{(B_m)}| \leq \delta \|A\|$$

for a user-specified value of  $\delta$ , where we have used (2.1). The quantity  $\|A\|$  in (2.13) is easily approximated by the singular value  $\sigma_1^{(B_m)}$  of largest magnitude of the bidiagonal matrix  $B_m$ . The computation of  $\sigma_1^{(B_m)}$  is inexpensive because the matrix  $B_m$  is small. During the computations of the desired singular triplets of  $A$ , typically, several matrices  $B_m$  and their singular value decompositions are computed. We approximate  $\|A\|$  by the largest of the singular values of all the matrices  $B_m$  generated. This generally gives a good estimate of  $\|A\|$ .

**3. Augmented Lanczos bidiagonalization methods.** It is well known that the implicitly restarted Arnoldi and Lanczos tridiagonalization methods described by Sorensen [29] can suffer from numerical instability due to propagated round-off errors. The instability can delay or prevent convergence of desired eigenvalues and eigenvectors; see Lehoucq and Sorensen [21] for a discussion and remedies. Morgan [25] showed that the implicitly restarted Arnoldi method by Sorensen [29] can be implemented by augmenting the available Krylov subspace basis by certain Ritz vectors. Such an implementation can be less sensitive to propagated round-off errors than the implementation in [29]. Recently, Wu and Simon [33] described a so-called thick-restarted Lanczos tridiagonalization method for the symmetric eigenvalue problem. The method is based on augmenting Krylov subspaces by certain Ritz vectors; it is simple to implement and is mathematically equivalent to the implicitly restarted Lanczos tridiagonalization method of Sorensen [29]. This paper presents thick-restarted Lanczos bidiagonalization methods. We remark that thick-restarting techniques have also been used in the context of the Jacobi–Davidson and Arnoldi methods for eigenvalue computations; see Stathopoulos and Saad [31] and Stathopoulos, Saad, and Wu [32] for discussions and analyses.

**3.1. Augmentation by Ritz vectors.** Let the partial Lanczos bidiagonalization (1.3)–(1.4) be available, and assume that we are interested in determining the  $k$  largest singular triplets of  $A$ , where  $k < m$ . Note that the approximate right singular

vector  $\tilde{v}_j^{(A)}$  of  $A$ , defined by (2.11), is a Ritz vector of  $A^T A$  associated with the Ritz value  $(\tilde{\sigma}_j^{(A)})^2$ . We have

$$(3.1) \quad A^T A \tilde{v}_j^{(A)} - (\tilde{\sigma}_j^{(A)})^2 \tilde{v}_j^{(A)} = \alpha_m r_m e_m^T v_j^{(B_m)}, \quad 1 \leq j \leq m.$$

The observation that the right-hand sides are parallel to  $r_m$  for all  $j$  forms the basis of the restarted Lanczos tridiagonalization method by Wu and Simon [33], as well as of the restarted Lanczos bidiagonalization method of this subsection.

We derive decompositions of the form (1.3)–(1.4) which allow us to choose the first  $k$  columns of the matrix  $P_m$  as Ritz vectors. The accuracy of these approximate left singular vectors and available approximate right singular vectors is then improved by restarting the computations. It follows from the orthonormality of the columns of the matrices  $P_m$  and  $V_m^{(B_m)}$  that the Ritz vectors  $\tilde{v}_j^{(A)}$  defined by (2.11) are orthonormal. Moreover, (1.5) shows that the  $\tilde{v}_j^{(A)}$  are orthogonal to  $r_m$ .

Let the Ritz vectors  $\tilde{v}_j^{(A)}$ ,  $1 \leq j \leq k$ , associated with the  $k$  largest Ritz values be available, assume that  $r_m \neq 0$ , and introduce the matrix

$$(3.2) \quad \tilde{P}_{k+1} := [\tilde{v}_1^{(A)}, \tilde{v}_2^{(A)}, \dots, \tilde{v}_k^{(A)}, p_{m+1}].$$

In view of (2.2), the last column of  $\tilde{P}_{k+1}$  is parallel to the residual error (3.1). It follows from (2.11) that

$$(3.3) \quad A \tilde{P}_{k+1} = [\tilde{\sigma}_1^{(A)} \tilde{u}_1^{(A)}, \tilde{\sigma}_2^{(A)} \tilde{u}_2^{(A)}, \dots, \tilde{\sigma}_k^{(A)} \tilde{u}_k^{(A)}, Ap_{m+1}].$$

Orthogonalization of  $Ap_{m+1}$  against the vectors  $\tilde{u}_j^{(A)}$  yields

$$(3.4) \quad Ap_{m+1} = \sum_{j=1}^k \tilde{\rho}_j \tilde{u}_j^{(A)} + \tilde{r}_k,$$

where the remainder  $\tilde{r}_k$  is orthogonal to the vectors  $\tilde{u}_j^{(A)}$ ,  $1 \leq j \leq k$ . The coefficients  $\tilde{\rho}_j := (\tilde{u}_j^{(A)})^T Ap_{m+1}$  can be evaluated inexpensively by using the right-hand side of

$$(\tilde{u}_j^{(A)})^T Ap_{m+1} = p_{m+1}^T A^T \tilde{u}_j^{(A)} = p_{m+1}^T (\tilde{\sigma}_j^{(A)} \tilde{v}_j^{(A)} + r_m e_m^T u_j^{(B_m)}) = \beta_m e_m^T \tilde{u}_j^{(B_m)}.$$

We may assume that the vector  $\tilde{r}_k$  is nonvanishing, because otherwise we can terminate the iterations; see below. Introduce the matrices

$$(3.5) \quad \tilde{Q}_{k+1} := \left[ \tilde{u}_1^{(A)}, \tilde{u}_2^{(A)}, \dots, \tilde{u}_k^{(A)}, \frac{\tilde{r}_k}{\|\tilde{r}_k\|} \right]$$

and

$$(3.6) \quad \tilde{B}_{k+1} := \begin{bmatrix} \tilde{\sigma}_1^{(A)} & & 0 & \tilde{\rho}_1 \\ & \ddots & & \vdots \\ & & \tilde{\sigma}_k^{(A)} & \tilde{\rho}_k \\ 0 & & & \tilde{\alpha}_{k+1} \end{bmatrix} \in \mathbb{R}^{(k+1) \times (k+1)}.$$

Thus,  $\tilde{B}_{k+1}$  may have nonvanishing entries only on the diagonal and in the last column. Substituting (3.4) into (3.3) now yields the decomposition

$$(3.7) \quad A \tilde{P}_{k+1} = \tilde{Q}_{k+1} \tilde{B}_{k+1},$$



which is analogous to (1.3).

We turn to the matrix

$$(3.8) \quad A^T \tilde{Q}_{k+1} = \left[ A^T \tilde{u}_1^{(A)}, A^T \tilde{u}_1^{(A)}, \dots, A^T \tilde{u}_k^{(A)}, A^T \frac{\tilde{r}_k}{\|\tilde{r}_k\|} \right],$$

which we would like to express in terms of  $\tilde{P}_{k+1}$  and  $\tilde{B}_{k+1}^T$ . This will give an analogue of the decomposition (1.4). The first  $k$  columns of (3.8) are linear combinations of the vectors  $\tilde{v}_j^{(A)}$  and  $p_{m+1}$ ; specifically,

$$(3.9) \quad A^T \tilde{u}_j^{(A)} = \tilde{\sigma}_j^{(A)} \tilde{v}_j^{(A)} + r_m e_m^T u_j^{(B_m)} = \tilde{\sigma}_j^{(A)} \tilde{v}_j^{(A)} + p_{m+1} \tilde{\rho}_j, \quad 1 \leq j \leq k,$$

where we have used (2.2). The last column of (3.8) is orthogonal to the Ritz vectors  $\tilde{v}_j^{(A)}$ ,

$$(\tilde{v}_j^{(A)})^T A^T \frac{\tilde{r}_k}{\|\tilde{r}_k\|} = \tilde{\sigma}_j^{(A)} \frac{(\tilde{u}_j^{(A)})^T \tilde{r}_k}{\|\tilde{r}_k\|} = 0, \quad 1 \leq j \leq k,$$

and therefore it can be expressed as

$$(3.10) \quad A^T \frac{\tilde{r}_k}{\|\tilde{r}_k\|} = \tilde{\gamma}_1 p_{m+1} + \tilde{f}_{k+1},$$

where  $\tilde{f}_{k+1} \in \mathbb{R}^n$  is orthogonal to the vectors  $\tilde{v}_j^{(A)}$ ,  $1 \leq j \leq k$ , as well as to  $p_{m+1}$ . Since

$$p_{m+1}^T A^T \frac{\tilde{r}_k}{\|\tilde{r}_k\|} = \frac{\tilde{r}_k^T}{\|\tilde{r}_k\|} A p_{m+1} = \frac{\tilde{r}_k^T}{\|\tilde{r}_k\|} \left( \sum_{j=1}^k \tilde{\rho}_j \tilde{u}_j^{(A)} + \tilde{r}_k \right) = \|\tilde{r}_k\|,$$

it follows from (3.10) that  $\tilde{\gamma}_1 = \|\tilde{r}_k\|$ . This observation, together with (3.9) and (3.10), gives the expression

$$(3.11) \quad A^T \tilde{Q}_{k+1} = \tilde{P}_{k+1} \tilde{B}_{k+1}^T + \tilde{f}_{k+1} e_{k+1}^T,$$

which is the desired analogue of the decomposition (1.4). We remark that  $\tilde{f}_{k+1}$  can be computed from (3.10).

The similarity of the decompositions (3.7) and (3.11), and (1.3)–(1.4), suggests that it may be possible to append new columns to the matrices  $\tilde{P}_{k+1}$  and  $\tilde{Q}_{k+1}$  in a way similar to Lanczos bidiagonalization. We will show that this is indeed the case. For notational simplicity, denote the columns of  $\tilde{P}_{k+1}$  and  $\tilde{Q}_{k+1}$  by  $\tilde{p}_j$  and  $\tilde{q}_j$ , respectively, i.e.,

$$\tilde{P}_{k+1} = [\tilde{p}_1, \tilde{p}_2, \dots, \tilde{p}_{k+1}] \in \mathbb{R}^{n \times (k+1)}, \quad \tilde{Q}_{k+1} = [\tilde{q}_1, \tilde{q}_2, \dots, \tilde{q}_{k+1}] \in \mathbb{R}^{\ell \times (k+1)}.$$

We may assume that  $\tilde{f}_{k+1} \neq 0$ , because otherwise it follows from (3.7) and (3.11) that the singular values of  $\tilde{B}_{k+1}$  are singular values of  $A$ , and we are done. Thus, let  $\tilde{\beta}_{k+1} := \|\tilde{f}_{k+1}\|$  and  $\tilde{p}_{k+2} := \tilde{f}_{k+1}/\tilde{\beta}_{k+1}$ . Then the matrix  $\tilde{P}_{k+2} := [\tilde{P}_{k+1}, \tilde{p}_{k+2}]$  has orthonormal columns.

Let

$$(3.12) \quad \tilde{\alpha}_{k+2} \tilde{q}_{k+2} := (I - \tilde{Q}_{k+1} \tilde{Q}_{k+1}^T) A \tilde{p}_{k+2},$$

where  $\tilde{\alpha}_{k+2} > 0$  is a scaling factor, such that  $\tilde{q}_{k+2}$  is of unit length. We comment on the possibility of  $\tilde{\alpha}_{k+2}$  vanishing below. Equation (3.11) yields

$$(3.13) \quad A^T \tilde{Q}_{k+1} = \tilde{P}_{k+1} \tilde{B}_{k+1}^T + \tilde{\beta}_{k+1} \tilde{p}_{k+2} e_{k+1}^T,$$

and substituting (3.13) into (3.12) shows that

$$(3.14) \quad \begin{aligned} \tilde{\alpha}_{k+2} \tilde{q}_{k+2} &= A \tilde{p}_{k+2} - \tilde{Q}_{k+1} (A^T \tilde{Q}_{k+1})^T \tilde{p}_{k+2} \\ &= A \tilde{p}_{k+2} - \tilde{Q}_{k+1} (\tilde{B}_{k+1} \tilde{P}_{k+1}^T + \tilde{\beta}_{k+1} e_{k+1} \tilde{p}_{k+2}^T) \tilde{p}_{k+2} \\ &= A \tilde{p}_{k+2} - \tilde{\beta}_{k+1} \tilde{Q}_{k+1} e_{k+1} = A \tilde{p}_{k+2} - \tilde{\beta}_{k+1} \tilde{q}_{k+1}. \end{aligned}$$

Let  $\tilde{Q}_{k+2} := [\tilde{Q}_{k+1}, \tilde{q}_{k+2}] \in \mathbb{R}^{\ell \times (k+2)}$ , and define  $\tilde{B}_{k+2} \in \mathbb{R}^{(k+2) \times (k+2)}$  by first appending the column  $\tilde{\beta}_{k+1} e_{k+1}$  and then the row  $\tilde{\alpha}_{k+2} e_{k+2}^T$  to  $\tilde{B}_{k+1}$ , i.e.,

$$\tilde{B}_{k+2} := \begin{bmatrix} \tilde{\sigma}_1^{(A)} & & 0 & \tilde{\rho}_1 & 0 \\ & \ddots & & \vdots & \vdots \\ & & \tilde{\sigma}_k^{(A)} & \tilde{\rho}_k & 0 \\ & & & \tilde{\alpha}_{k+1} & \tilde{\beta}_{k+1} \\ 0 & & & & \tilde{\alpha}_{k+2} \end{bmatrix}.$$

It now follows from (3.7) and (3.14) that

$$(3.15) \quad A \tilde{P}_{k+2} = \tilde{Q}_{k+2} \tilde{B}_{k+2}.$$

We turn to the derivation of a decomposition of the form (3.13) with  $k+1$  replaced by  $k+2$ . Let

$$(3.16) \quad \tilde{\beta}_{k+2} \tilde{p}_{k+3} := (I - \tilde{P}_{k+2} \tilde{P}_{k+2}^T) A^T \tilde{q}_{k+2},$$

where  $\tilde{\beta}_{k+2} > 0$  is a scaling factor, such that  $\tilde{p}_{k+3}$  is of unit length. We may assume that a positive coefficient  $\tilde{\beta}_{k+2}$  exists; otherwise we are done; see below. Substituting (3.15) into (3.16) yields

$$\tilde{\beta}_{k+2} \tilde{p}_{k+3} = A^T \tilde{q}_{k+2} - \tilde{P}_{k+2} \tilde{B}_{k+2}^T e_{k+2} = A^T \tilde{q}_{k+2} - \tilde{\alpha}_{k+2} \tilde{p}_{k+2},$$

which, together with (3.13), shows that

$$(3.17) \quad A^T \tilde{Q}_{k+2} = \tilde{P}_{k+2} \tilde{B}_{k+2}^T + \tilde{\beta}_{k+2} \tilde{p}_{k+3} e_{k+2}^T.$$

The decompositions (3.15) and (3.17) are analogous to (3.7) and (3.13). We therefore can continue in the same fashion by appending new columns to the matrices  $\tilde{P}_j$  and  $\tilde{Q}_j$ , and new rows and columns to the matrices  $\tilde{B}_j$ , for  $j = k+2, k+3, \dots$ . After a total of  $m-k$  steps, we obtain the decompositions

$$(3.18) \quad A \tilde{P}_m = \tilde{Q}_m \tilde{B}_m, \quad A^T \tilde{Q}_m = \tilde{P}_m \tilde{B}_m^T + \tilde{\beta}_m \tilde{p}_{m+1} e_m^T,$$

where the matrices  $\tilde{P}_m$  and  $\tilde{Q}_m$  have orthonormal columns and

$$\tilde{B}_m = \begin{bmatrix} \tilde{\sigma}_1^{(A)} & & 0 & \tilde{\rho}_1 & & 0 \\ & \ddots & & \vdots & & \\ & & \tilde{\sigma}_k^{(A)} & \tilde{\rho}_k & & \\ & & & \tilde{\alpha}_{k+1} & \tilde{\beta}_{k+1} & \\ & & & & \ddots & \\ & 0 & & & & \tilde{\beta}_{m-1} \\ & & & & & \tilde{\alpha}_m \end{bmatrix}.$$

The method now proceeds by first computing the singular value decomposition of  $\tilde{B}_m$  and then determining the  $k$  largest approximate singular triplets of  $A$  from the  $k$  largest singular triplets of  $\tilde{B}_m$ ; cf. (2.11). These triplets define new decompositions of the form (3.7) and (3.11), from which we compute new decompositions of the form (3.18). The computations proceed in this manner until sufficiently accurate approximations of the  $k$  largest singular triplets of  $A$  have been determined. An algorithm is presented in subsection 3.3 below.

We remark that the computations are analogous for determining the  $k$  smallest singular triplets of  $A$ . The vectors  $\tilde{v}_j^{(A)}$ ,  $1 \leq j \leq k$ , in (3.2) should then be replaced by the right approximate singular vectors of the  $k$  smallest available approximate singular triplets of  $A$ . These approximate singular triplets are used to define decompositions (3.7) and (3.11), from which we compute decompositions (3.18) in the same manner as described above. We are interested in the  $k$  smallest singular triplets of the matrix  $\tilde{B}_m$  in the decompositions (3.18). These triplets yield approximations of the  $k$  smallest triplets of  $A$ . The computations are continued until sufficiently accurate approximations of the  $k$  smallest singular triplets of  $A$  have been found. However, we note that when the  $k$  smallest singular triplets of  $A$  are desired, it can be advantageous to augment by harmonic Ritz vectors instead. This is discussed in subsection 3.2.

Finally, we comment on the cases when  $\tilde{r}_k$  vanishes in (3.4) and when the left-hand sides of (3.12) and (3.16) vanish. In all these cases, one can show that the singular values of the matrix  $\tilde{B}_j$  also are singular values of  $A$  and that the singular triplets of  $\tilde{B}_j$  yield singular triplets of  $A$ .

**3.2. Augmentation by harmonic Ritz vectors.** Augmenting by Ritz vectors, as described in the previous subsection, or, equivalently, shifting by Ritz values, often gives good approximations to the largest singular triplets of  $A$ . However, Kokiopoulou, Bekas, and Gallopoulos [17] observed that when seeking to compute the smallest singular triplets of  $A$ , shifting by harmonic Ritz values can give faster convergence than shifting by Ritz values. This section describes how shifting by harmonic Ritz values can be implemented via augmentation by harmonic Ritz vectors. Harmonic Ritz vectors are approximate eigenvectors of  $A^T A$  associated with harmonic Ritz values of  $A^T A$ .

Let the partial Lanczos bidiagonalization (1.3)–(1.4) of  $A$  be available, and assume that all the diagonal and superdiagonal entries of  $B_m$ , as well as  $\beta_m$  given by (2.1), are nonvanishing. Then, in particular,  $B_m$  is nonsingular. The harmonic Ritz values  $\hat{\theta}_j$  of  $A^T A$  associated with the partial Lanczos tridiagonalization (2.4) are the eigenvalues

of the generalized eigenvalue problem

$$(3.19) \quad ((B_m^T B_m)^2 + \alpha_m^2 \beta_m^2 e_m e_m^T) \hat{w}_j = \hat{\theta}_j B_m^T B_m \hat{w}_j, \quad 1 \leq j \leq m,$$

with  $\hat{w}_j \in \mathbb{R}^m \setminus \{0\}$ ; see, e.g., Morgan [24] or Paige, Parlett, and van der Vorst [27] for properties of harmonic Ritz values.

The eigenpairs  $\{\hat{\theta}_j, \hat{w}_j\}$  of (3.19) can be computed without forming the matrix  $B_m^T B_m$  as follows. Let

$$(3.20) \quad w_j := B_m \hat{w}_j.$$

Then (3.19) can be expressed as

$$(3.21) \quad (B_m B_m^T + \beta_m^2 e_m e_m^T) w_j = \hat{\theta}_j w_j,$$

where we may choose the eigenvectors  $w_j$  to be orthonormal. Let  $B_{m,m+1}$  be the matrix in (2.3), and note that

$$(3.22) \quad B_{m,m+1} B_{m,m+1}^T = B_m B_m^T + \beta_m^2 e_m e_m^T.$$

Introduce the singular triplets  $\{\sigma_j^{(B_{m,m+1})}, u_j^{(B_{m,m+1})}, v_j^{(B_{m,m+1})}\}$ ,  $1 \leq j \leq m$ , of the matrix  $B_{m,m+1}$ , and let them be enumerated so that

$$(3.23) \quad 0 < \sigma_1^{(B_{m,m+1})} \leq \sigma_2^{(B_{m,m+1})} \leq \dots \leq \sigma_m^{(B_{m,m+1})}.$$

This enumeration differs from the one for the singular values of  $B_m$  (cf. (2.9)), because in the present subsection we are concerned with the computation of the  $k < m$  smallest singular triplets of  $A$ . Throughout this subsection, we use the following simplified notation:

$$\sigma'_j = \sigma_j^{(B_{m,m+1})}, \quad u'_j = u_j^{(B_{m,m+1})}, \quad v'_j = v_j^{(B_{m,m+1})}.$$

The  $k$  smallest singular triplets of  $B_{m,m+1}$  determine the matrices

$$(3.24) \quad \begin{aligned} U'_k &:= [u'_1, u'_2, \dots, u'_k] \in \mathbb{R}^{m \times k}, \\ V'_k &:= [v'_1, v'_2, \dots, v'_k] \in \mathbb{R}^{(m+1) \times k}, \\ \Sigma'_k &:= \text{diag}[\sigma'_1, \sigma'_2, \dots, \sigma'_k] \in \mathbb{R}^{k \times k}, \end{aligned}$$

where  $U'_k$  and  $V'_k$  have orthonormal columns, and

$$(3.25) \quad B_{m,m+1} V'_k = U'_k \Sigma'_k, \quad B_{m,m+1}^T U'_k = V'_k \Sigma'_k.$$

We refer to (3.25) as a partial singular value decomposition of  $B_{m,m+1}$ . It follows from (3.22) and (3.25) that  $\{(\sigma'_j)^2, u'_j\}$  is an eigenpair of (3.21), and (3.20) shows that  $\{(\sigma'_j)^2, B_m^{-1} u'_j\}$  is an eigenpair of (3.19). Thus, the eigenpairs of (3.19) and (3.21) associated with the  $k$  smallest eigenvalues can be determined from the partial singular value decomposition (3.25). Gu and Eisenstat [12] describe how the singular value decomposition of  $B_{m,m+1}$  can be computed by updating the singular value decomposition of  $B_m$ ; see also Bunch and Nielsen [6]. However, when  $A$  is large and  $m$  is small, the computational effort required for determining the singular value decompositions of  $B_m$  and  $B_{m,m+1}$  is negligible. We will therefore not dwell on the computation of (3.25).

The harmonic Ritz vector of  $A^T A$  associated with the harmonic Ritz value  $\hat{\theta}_j$  is given by

$$(3.26) \quad \hat{v}_j := P_m \hat{w}_j;$$

see, e.g., [24, 27]. Morgan and Zeng [26] recently pointed out that the residual errors associated with different harmonic Ritz pairs  $\{\hat{\theta}_j, \hat{v}_j\}$  are parallel. We show this result for the problem at hand, because this property is central for our augmentation method. Thus, using (2.4), (3.20), (3.21), and (3.26), we obtain

$$\begin{aligned} A^T A \hat{v}_j - \hat{\theta}_j \hat{v}_j &= A^T A P_m \hat{w}_j - \hat{\theta}_j P_m \hat{w}_j \\ &= (P_m B_m^T B_m + \alpha_m r_m e_m^T) \hat{w}_j - \hat{\theta}_j P_m \hat{w}_j \\ &= P_m (B_m^T B_m - \hat{\theta}_j I_m) \hat{w}_j + \alpha_m r_m e_m^T \hat{w}_j \\ &= P_m B_m^{-1} (B_m B_m^T - \hat{\theta}_j I_m) w_j + r_m e_m^T w_j \\ &= P_m B_m^{-1} (-\beta_m^2 e_m e_m^T w_j) + r_m e_m^T w_j \\ &= e_m^T w_j (r_m - \beta_m^2 P_m B_m^{-1} e_m). \end{aligned}$$

It is convenient to define the scaled residual vector

$$(3.27) \quad \hat{r}_m := p_{m+1} - \beta_m P_m B_m^{-1} e_m,$$

where we have used (2.2).

We are in a position to derive relations analogous to (3.7) and (3.11) for harmonic Ritz vectors. Equations (3.20), (3.26), and (3.27) yield

$$[\hat{v}_1 \sigma'_1, \hat{v}_2 \sigma'_2, \dots, \hat{v}_k \sigma'_k, \hat{r}_m] = P_{m+1} \begin{bmatrix} B_m^{-1} U'_k \Sigma'_k & -\beta_m B_m^{-1} e_m \\ 0 & 1 \end{bmatrix},$$

where the matrix  $P_{m+1}$  is the same as in (2.3). Introduce the QR-factorization

$$(3.28) \quad \begin{bmatrix} B_m^{-1} U'_k \Sigma'_k & -\beta_m B_m^{-1} e_m \\ 0 & 1 \end{bmatrix} = Q'_{k+1} R'_{k+1},$$

where  $Q'_{k+1} \in \mathbb{R}^{(m+1) \times (k+1)}$  has orthonormal columns and  $R'_{k+1} \in \mathbb{R}^{(k+1) \times (k+1)}$  is upper triangular.

Consider the matrix

$$(3.29) \quad \hat{P}_{k+1} = [\hat{p}_1, \hat{p}_2, \dots, \hat{p}_{k+1}] := P_{m+1} Q'_{k+1}.$$

Its columns  $\hat{p}_j$  are orthonormal since both  $P_{m+1}$  and  $Q'_{k+1}$  have orthonormal columns. Equations (1.3) and (3.28) yield

$$\begin{aligned} A \hat{P}_{k+1} &= [A P_m, A p_{m+1}] Q'_{k+1} \\ &= [Q_m B_m, A p_{m+1}] \begin{bmatrix} B_m^{-1} U'_k \Sigma'_k & -\beta_m B_m^{-1} e_m \\ 0 & 1 \end{bmatrix} (R'_{k+1})^{-1} \\ &= [Q_m U'_k \Sigma'_k, -\beta_m q_m + A p_{m+1}] (R'_{k+1})^{-1}. \end{aligned}$$

In the derivation of our method, we assume only that the matrix  $B_m$  is upper triangular, because  $B_m$  has this property after restarts. Below we comment on possible simplifications when  $B_m$  is upper bidiagonal.

Let

$$(3.30) \quad \hat{Q}_k := Q_m U'_k.$$

The columns of this matrix are orthonormal, and we define the Fourier coefficients

$$\hat{c}_k = [\hat{\gamma}_1, \hat{\gamma}_2, \dots, \hat{\gamma}_k]^T := \hat{Q}_k^T (-\beta_m q_m + Ap_{m+1}).$$

The vector

$$\hat{\alpha}_{k+1} \hat{q}_{k+1} := -\beta_m q_m + Ap_{m+1} - \hat{Q}_k \hat{c}_k$$

is orthogonal to the columns of  $\hat{Q}_k$ , and the scaling factor  $\hat{\alpha}_{k+1} > 0$  is chosen so that  $\hat{q}_{k+1}$  is of unit length. It follows that

$$(3.31) \quad A\hat{P}_{k+1} = \hat{Q}_{k+1}\hat{B}_{k+1},$$

where

$$(3.32) \quad \hat{Q}_{k+1} := [\hat{Q}_k, \hat{q}_{k+1}] \in \mathbb{R}^{\ell \times (k+1)},$$

$$(3.33) \quad \hat{B}_{k+1} := \begin{bmatrix} \sigma'_1 & & & 0 & \hat{\gamma}_1 \\ & \sigma'_2 & & & \hat{\gamma}_2 \\ & & \ddots & & \vdots \\ & & & \sigma'_k & \hat{\gamma}_k \\ 0 & & & & \hat{\alpha}_{k+1} \end{bmatrix} (R'_{k+1})^{-1} \in \mathbb{R}^{(k+1) \times (k+1)}.$$

Thus,  $\hat{Q}_{k+1}$  has orthonormal columns, and  $\hat{B}_{k+1}$  is the product of two upper triangular matrices, one of which has nonzero entries only on the diagonal and in the last column. In particular,  $\hat{B}_{k+1}$  is upper triangular. The decomposition (3.31) is the desired analogue of (3.7). When  $B_m$  is given by (1.6), one can show that  $\hat{\alpha}_{k+1} = \alpha_{m+1}$ ,  $\hat{q}_{k+1} = q_{m+1}$ , and  $\hat{c}_k = 0$ .

We now derive an analogue of the decomposition (3.11). Let  $\hat{Q}_k$  be given by (3.30). Then (3.25) yields

$$(3.34) \quad A^T \hat{Q}_k = A^T Q_m U'_k = P_{m+1} B_{m,m+1}^T U'_k = P_{m+1} V'_k \Sigma'_k.$$

It follows from the decomposition on the left-hand side of (3.25) that

$$[I_m, \beta_m B_m^{-1} e_m] V'_k = B_m^{-1} U'_k \Sigma'_k,$$

and therefore

$$(3.35) \quad V'_k = \begin{bmatrix} B_m^{-1} U'_k \Sigma'_k & -\beta_m B_m^{-1} e_m \\ 0 & 1 \end{bmatrix} \begin{bmatrix} I_k \\ e_{m+1}^T V'_k \end{bmatrix}.$$

Substituting (3.35) into (3.34), using (3.28) and (3.29), gives

$$(3.36) \quad A^T \hat{Q}_k = P_{m+1} Q'_{k+1} R'_{k+1} \begin{bmatrix} I_k \\ e_{m+1}^T V'_k \end{bmatrix} \Sigma'_k = \hat{P}_{k+1} R'_{k+1} \begin{bmatrix} I_k \\ e_{m+1}^T V'_k \end{bmatrix} \Sigma'_k.$$

Let  $\hat{B}_{k,k+1}$  be the leading  $k \times (k+1)$  submatrix of the upper triangular matrix  $\hat{B}_{k+1}$  in (3.31). Then (3.31) yields

$$(3.37) \quad \hat{Q}_k^T A \hat{P}_{k+1} = \hat{B}_{k,k+1}.$$



has the  $(k+1) \times (k+1)$  leading principal submatrix  $\hat{B}_{k+1}$  and an  $(m-k-1) \times (m-k-1)$  upper bidiagonal trailing principal submatrix. The residual vector  $\check{r}_m$  is orthogonal to the columns of  $\hat{P}_m$ . The method of this subsection can be restarted by letting  $B_m := \hat{B}_m$ ,  $P_m := \hat{P}_m$ ,  $Q_m := \hat{Q}_m$ ,  $r_m := \check{r}_m$ , and  $\beta_m := \|\check{r}_m\|$ .

We remark that the accurate computation of the vector  $B_m^{-1}e_m$ , used in (3.28), can be difficult when the matrix  $B_m$  has a large condition number  $\kappa(B_m) := \sigma_m^{(B_m)}/\sigma_1^{(B_m)}$ , where the singular values are enumerated as in (3.23). In this case, we switch from augmentation by harmonic Ritz vectors to augmentation by Ritz vectors. Details are provided in the following subsection.

**3.3. An augmented Lanczos bidiagonalization algorithm.** We describe an algorithm for the computation of a few of the largest or smallest singular triplets of a large matrix  $A$ . The algorithm is based on augmentation by Ritz vectors or harmonic Ritz vectors as described in the previous subsections. The Boolean variable *harmonic* suggests the type of augmentation used. If *harmonic* is true and  $B_m$  is not too ill-conditioned, then the augmentation scheme of subsection 3.2 is applied; otherwise augmentation is carried out according to subsection 3.1.

ALGORITHM 3.1. AUGMENTED LANCZOS BIDIAGONALIZATION.

*Input:*  $A \in \mathbb{R}^{\ell \times n}$  or functions for evaluating matrix-vector products with the matrices  $A$  and  $A^T$ ,  
 $p_1 \in \mathbb{R}^n$  : initial vector of unit length,  
 $m$  : number of bidiagonalization steps,  
 $k$  : number of desired singular triplets,  
 $\delta$  : tolerance for accepting computed approximate singular triplet; cf. (2.13),  
 $\epsilon$  : machine epsilon,  
*harmonic* : Boolean variable that suggests type of augmentation; see above.

*Output:* Computed set of approximate singular triplets  $\{\sigma_j, u_j, v_j\}_{j=1}^k$  of  $A$ .

1. Compute the partial Lanczos bidiagonalization (1.3)–(1.4) using Algorithm 2.1.
2. Compute the singular value decomposition (2.10) of  $B_m$ .
3. Check convergence: If all  $k$  desired singular triplets satisfy (2.13), then exit.
4. Compute the augmenting vectors:
  - 4a. if not harmonic or  $\kappa(B_m) > \epsilon^{-1/2}$ , then  
 Determine the matrices  $P := \tilde{P}_{k+1}$ ,  $Q := \tilde{Q}_{k+1}$ ,  $B := \tilde{B}_{k+1}$ , and the vector  $r := \tilde{f}_{k+1}$  by (3.2), (3.5), (3.6), and (3.10), respectively.
  - 4b. if harmonic and  $\kappa(B_m) \leq \epsilon^{-1/2}$ , then  
 Compute the partial singular value decomposition (3.24) of  $B_{m,m+1}$  and the QR-factorization (3.28).  
 Determine the matrices  $P := \hat{P}_{k+1}$ ,  $Q := \hat{Q}_{k+1}$ ,  $B := \hat{B}_{k+1}$ , and the vector  $r := \check{r}_{k+1}$  by (3.29), (3.32), (3.33), and (3.39), respectively.
5. The available matrices  $P$ ,  $Q$ ,  $B$ , and the vector  $r$  satisfy

$$AP = QB, \quad A^T Q = PB^T + r e_k^T.$$

Append  $m - k$  columns to the matrices  $P$  and  $Q$ , and  $m - k$  rows and columns to the matrix  $B$ . Denote the matrices so obtained by  $P_m$ ,  $Q_m$ , and  $B_m$ , respectively. Determine a new residual vector and denote it by  $r_m$ .

6. Goto 2.



The above algorithm is a simplification of the actual computations carried out. For instance, the algorithm exits when a matrix  $B_m$  that is numerically singular has been detected, but the available decompositions of  $A$  can be used to determine singular triplets of  $A$ ; see the discussion in section 1. Moreover, the number of augmented vectors used at each restart is typically larger than the number of desired singular triplets. Assume that  $k'$  of the desired  $k$  singular triplets have been found. We then augment by  $k + k''$  (instead of  $k$ ) singular triplets, where  $k''$  is chosen as large as possible, such that  $k'' \leq k'$  and  $k + k'' \leq m - 3$ . The term  $-3$  secures that at least three orthogonalization steps can be carried out between restarts. This approach has been advocated by Lehoucq [20] in the context of the implicitly restarted Arnoldi method. It often yields faster convergence without increasing the memory requirement. Finally, our implementation enforces two-sided reorthogonalization when  $\kappa(B_m) > \epsilon^{-1/2}$ . MATLAB code is available at the authors' home pages.

**4. Numerical examples.** All computations were carried out using MATLAB version 6.5 R13 on a Dell 530 workstation with two 2.4 GHz (512k cache) Xeon processors and 2 GB (400 MHz) of memory running under the Windows XP operating system. Machine epsilon is  $\epsilon = 2.2 \cdot 10^{-16}$ .

We compare our methods, outlined by Algorithm 3.1, with methods recently proposed by Hochstenbach [14, 13] and Kokiopoulou, Bekas, and Gallopoulos [17], as well as with the MATLAB internal function `svds` and the scheme proposed in [2]. Hochstenbach [14, 13] presents a Jacobi–Davidson method. This is a powerful scheme when a good preconditioner for the linear system of equations that has to be solved is available. In our computed examples, we assume that no good preconditioner is known, and we apply the method without preconditioner. The linear system of equations is solved by the GMRES iterative method. The MATLAB implementation<sup>1</sup> `jdsvd` offers several extraction choices, such as standard, u-harmonic, v-harmonic, double-harmonic, and refined. In our numerical examples, refined extraction often gave best accuracy. This is consistent with results reported by Hochstenbach [14]. We refer to the Jacobi–Davidson method with refined extraction as `jdsvd(Ref)`. The code `jdsvd` is still under development, and we used the version available to us at the time of the numerical experiments.

Our methods are mathematically, but not numerically, equivalent to the methods proposed by Kokiopoulou, Bekas, and Gallopoulos [17]. We used the MATLAB implementation `irlanb` by Kokiopoulou, Bekas, and Gallopoulos [17] in our comparison. The code `irlanb` calls Larsen's MATLAB code `lanbpro` [18] to compute partial Lanczos bidiagonalizations with partial reorthogonalization. `irlanb` is designed for computing a few of the smallest singular triplets but not for computing a few of the largest ones. The code therefore is not used in Examples 4 and 5 below. Ritz values and harmonic Ritz values can be used as shifts. We refer to `irlanb` with these shift selections as `irlanb(R)` and `irlanb(H)`, respectively. The code `irlanb` is still under development, and we report results for the version available to us at the time of the numerical experiments.

The internal MATLAB function `svds` uses FORTRAN codes of ARPACK [22]. It calls an eigenvalue routine to compute eigenvalue-eigenvector pairs associated with positive eigenvalues of the symmetric matrix

$$(4.1) \quad Z := \begin{bmatrix} 0 & A \\ A^T & 0 \end{bmatrix} \in \mathbb{R}^{(\ell+n) \times (\ell+n)}.$$

<sup>1</sup>Code is available at <http://www.case.edu/artsci/math/hochstenbach/software/jdsvd.html>

The matrix  $Z$  has the eigenvalues

$$\pm\sigma_1^{(A)}, \pm\sigma_2^{(A)}, \dots, \pm\sigma_n^{(A)},$$

as well as  $\ell - n$  zero eigenvalues, where we as usual assume that  $\ell \geq n$ . The eigenvectors of  $Z$  yield both the right and left singular vectors of  $A$ . The method requires the computation of eigenpairs associated with eigenvalues near zero when determining the smallest singular triplets of  $A$ . This can be difficult when  $Z$  has many positive and negative eigenvalues of large magnitude. For this reason, the `svds` function uses a shift-and-invert approach for computing the smallest singular triplets. This requires factorization of the matrix  $Z$ , and therefore the function `svds` typically demands much more storage than the other methods in our comparison. Since `svds` does not yield the number of linear systems of equations that have to be solved, we report only the CPU time required.

The MATLAB code `irblsvds` implements an implicitly restarted block-Lanczos method applied to the matrix (4.1) for the computation of a few singular triplets of  $A$ . The method and code are described in [2, 3].<sup>2</sup> The matrix  $Z$  is used only for evaluation of matrix-vector products; in particular, the matrix  $Z$  is not factored. The code `irblsvds` is not, in general, well suited for computing the smallest singular triplets of  $A$  when  $\ell - n$  is large, because then  $Z$  has  $\ell - n$  zero eigenvalues, and the code may determine these eigenvalues and associated eigenvectors instead of eigenpairs associated with tiny singular triplets of  $A$ . Unless indicated otherwise, we use the default value 3 for the block-size in our experiments with `irblsvds`. The largest number of consecutive block-Lanczos steps is chosen so that `irblsvds` has about the same storage requirement as the other codes. We note that of the methods used in the examples of this section, only the ones of the present paper are based on augmented matrix formulations.

The schemes of subsections 3.1 and 3.2 are implemented by the MATLAB code `irlba`.<sup>2</sup> The execution of `irlba` is determined by certain user-specified parameters; see Table 4.1. We refer to the augmentation method of subsection 3.1, based on Ritz vectors, as `irlba(R)` and to the augmentation method of subsection 3.2, based on harmonic Ritz vectors, as `irlba(H)`. The scheme `irlba(R)` with one- and two-sided full reorthogonalization is referred to as `irlba(R1)` and `irlba(R2)`, respectively. The analogous implementations of `irlba(H)` are denoted by `irlba(H1)` and `irlba(H2)`. One-sided full reorthogonalization reorthogonalizes the columns of the smaller of the matrices  $P_m$  and  $Q_m$ . Thus, when  $\ell \geq n$ , the columns of  $P_m$  are reorthogonalized.

The codes `irblsvds`, `irlanb`, `jdsvd`, and `svds` allow a user to choose numerous parameters that affect their performance. Unless stated otherwise, we use the default values for the parameters. Except for the function `svds`, we choose the parameters that affect storage so that all codes require about the same maximum computer storage.

It is impossible to use the same starting vector for all the methods, since some routines work with  $A$  and  $A^T$  and others with the matrix  $Z$ . To make our comparison less dependent on the choice of starting vector, we record the best results for each method over five runs using default random starting vector(s) generated by each code. We use the same starting vector when the same method is applied with different parameter values. The reported number of matrix-vector products is the total number of matrix-vector product evaluations with  $A$  and  $A^T$  for `irlba`, `irlanb`, and `jdsvd` and with  $Z$  for `irblsvds`.

---

<sup>2</sup>Code is available at the authors' home pages.

TABLE 4.1  
Parameters for `irlba`.

<i>adjust</i>	Initial number of vectors added to the $k$ restart vectors to speed up convergence. Default value: <i>adjust</i> = 3.
<i>aug</i>	A 4-letter string. The value RITZ yields the augmentation described in subsection 3.1; the value HARM gives augmentation according to subsection 3.2. Default value: <i>aug</i> = HARM if <i>sigma</i> = SS, and <i>aug</i> = RITZ if <i>sigma</i> = LS.
<i>disps</i>	When <i>disps</i> > 0, available approximations of the $k$ desired singular values and norms of associated residual errors are displayed each iteration; <i>disps</i> = 0 inhibits display of these quantities. Default value: <i>disps</i> = 0.
<i>k</i>	Number of desired singular triplets. Default value: <i>k</i> = 6.
<i>maxit</i>	Maximum number of restarts. Default value: <i>maxit</i> = 100.
<i>steps</i>	Maximum number of Lanczos bidiagonalization steps. The parameter specifies the largest value of steps $m$ in (1.3)–(1.4) and determines the storage requirement of the method. Default value: <i>steps</i> = 20.
<i>reorth</i>	A 3-letter string. The value ONE yields one-sided full reorthogonalization of the “shorter” vectors; the value TWO gives two-sided full reorthogonalization. When our available estimate of $\kappa(A)$ (see the discussion following (2.13)) is larger than $\epsilon^{-1/2}$ , two-sided full reorthogonalization is used. Default value: <i>reorth</i> = ONE.
<i>sigma</i>	A 2-letter string (SS for smallest and LS for largest) which specifies which extreme singular triplets are to be computed. Default value: <i>sigma</i> = LS.
$\delta$	Tolerance used for convergence check; see (2.13). Default value: $\delta = 10^{-6}$ .
$v_0$	Initial vector for Lanczos bidiagonalization. When $\ell \geq n$ , $p_1 := v_0$ ; cf. Algorithm 2.1. Default value: $v_0$ is a random vector with normally distributed entries.

*Example 1* (smallest singular value). We would like to compute the smallest singular triplet of the diagonal matrices

$$A := \text{diag}[1, 2, \dots, n], \quad n = 100, 200, 300, 400.$$

Each code was instructed to determine only one singular triplet, the smallest one. This corresponds to the parameter values *sigma* = SS and *k* = 1 for `irlba`. We allowed each of the codes `irlba`, `irlanb`, `jdsvd`, and `irblsvds` about the same amount of storage as required for carrying out 20 Lanczos bidiagonalization steps. In particular, the largest number of consecutive block-Lanczos steps allowed by `irblsvds` was limited to 7 (with block-size 3). This corresponds to about the same storage requirement as 21 Lanczos bidiagonalization steps. We let *adjust* := 4 for `irlba`; see Table 4.1. This forces both the `irlba` and `irlanb` codes to apply the same number of bidiagonalization steps in the first restart. We chose the tolerance  $10^{-6}$  for all codes, i.e.,  $\delta := 10^{-6}$  in (2.13). For the `jdsvd` code we report only refined extraction, because this extraction method was the fastest. Figure 4.1 displays the CPU times required for the different methods. The MATLAB `svds` function is seen to be fastest. This may depend on the fact that `svds` is not coded in MATLAB and that it is based on a shift-and-invert approach. For large problems shift-and-invert may require unacceptable amounts of memory and execution time; however, when computation and storage of factors of matrices of the form  $Z - \tau I$ , where  $\tau$  is a scalar, is feasible, then this approach is attractive. Among the methods that use only the matrices  $A$  and  $A^T$  for evaluating matrix-vector products, `irlba` is seen to be competitive.

*Example 2* (smallest singular values). We are interested in determining the smallest singular triplet of four matrices generated in MATLAB with the commands

$$A = \text{randn}(n), \quad A(:, 1) = A(:, 10),$$

for  $n = 200, 400, 600, 800$ . Here  $A = \text{randn}(n)$  determines an  $n \times n$  matrix with




FIG. 4.1. Example 1: CPU times for computing the smallest singular triplet.

normally distributed entries with mean zero and variance one. The command  $A(:, 1) = A(:, 10)$  overwrites column 1 by column 10 and secures that the matrix obtained has a zero singular value. This example illustrates that augmentation by harmonic Ritz vectors can give substantially faster convergence than augmentation by Ritz vectors.

The code `irlba` was used with  $k = 1$ ,  $\delta = 10^{-16}$ ,  $reorth = TWO$ , and  $steps = 30$ . Figure 4.2 shows the convergence to zero of the smallest computed singular value. The vertical axis displays the absolute error in the smallest computed singular value, i.e., the smallest computed singular value, and the horizontal axis shows the number of restarts. The cross-over label indicates when `irlba` switched from augmenting with harmonic Ritz vectors to augmenting with Ritz vectors. As mentioned in subsection 3.2, augmentation by harmonic Ritz vectors requires the solution of a linear system of equations with the matrix  $B_m$  (while augmentation by Ritz vectors does not). We switch from augmentation by harmonic Ritz vectors to augmentation by Ritz vectors when the condition number of the matrix  $B_m$  is larger than the square root of the reciprocal of machine epsilon; cf. Algorithm 3.1.

We remark that computing a singular triplet with a numerically vanishing singular value is not always possible since the approximated left singular vector  $\tilde{u}_j^{(A)}$  lives in the Krylov subspace (2.8), which is restricted to the range of  $A$ . However, in the presence of round-off errors, `irlba` is often able to successfully compute singular triplets associated with zero singular values.




FIG. 4.2. Example 2: Absolute error in the smallest singular value computed by `irlba` versus number of restarts using augmentation with Ritz vectors (solid curve) and augmentation with harmonic Ritz vectors (dotted curve). After the cross-over point augmentation is with Ritz vectors.

*Example 3* (smallest singular values). We consider the  $1033 \times 320$  matrix WELL1033 and the  $1850 \times 712$  matrix WELL1850 from the set LSQ in the Harwell–Boeing sparse matrix collection [9]. These matrices arise from surveying problems. We would like to determine the six smallest singular triplets of these matrices and select the parameters for the different codes accordingly. For instance, for `irlba` we let  $\sigma := \text{SS}$  and  $k := 6$ . All codes are allowed the amount of storage required for 40 Lanczos bidiagonalization steps. The number of implicit QR-steps in `irlanb` is chosen to be 31; this choice is consistent with the default value of the parameter `adjust` of `irlba`. It forces `irlba` and `irlanb` to apply the same number of Lanczos bidiagonalization steps in the first restart. The tolerance for all methods is set to  $\delta = 10^{-6}$ . `irlanb` with shifts chosen to be harmonic Ritz values gave better results than when Ritz values were used as shifts. We therefore report only results for the former. The restarted block-Lanczos method `irblsvds` used block-size 4 and was allowed to carry out at most 10 consecutive block-Lanczos steps between restarts. `jdsvd` gave the best results for refined extraction. Therefore we do not report results for other extraction methods. The MATLAB `svds` function was unable to find any of the desired singular triplets to requested accuracy for either of the matrices. Table 4.2 summarizes the computed results. The table shows that `irlba` was able to compute

TABLE 4.2

Example 3: Computation of the six smallest singular triplets of the matrices WELL1033 and WELL1850.

irlba(H1)

Matrix	# Matrix-vector products	CPU time	Magnitude of largest error
WELL1033	638	0.36s	$2.41 \cdot 10^{-15}$
WELL1850	1442	1.39s	$1.72 \cdot 10^{-13}$

irlba(H2)

Matrix	# Matrix-vector products	CPU time	Magnitude of largest error
WELL1033	638	0.59s	$2.14 \cdot 10^{-15}$
WELL1850	1442	2.41s	$1.72 \cdot 10^{-13}$

irblsvds

Matrix	# Matrix-vector products	CPU time	Magnitude of largest error
WELL1033	7520	18.67s	$8.53 \cdot 10^{-16}$
WELL1850	16080	77.53s	$1.11 \cdot 10^{-15}$

irlanb(H)

Matrix	# Matrix-vector products	CPU time	Magnitude of largest error
WELL1033*	—	—	—
WELL1850	1578	8.31s	$3.03 \cdot 10^{-10}$

\* Method failed to convergence.

jdsvd(Ref)

Matrix	# Matrix-vector products	CPU time	Magnitude of largest error
WELL1033	1978	2.66s	$4.15 \cdot 10^{-10}$
WELL1850	4244	7.69s	$6.84 \cdot 10^{-12}$

all six singular values with the fewest matrix-vector product evaluations and the least CPU time for both matrices.

Example 4 (largest singular values). The matrices MEDLINE, CRANFIELD, and CISI are standard term-by-document test matrices and can be obtained from the Cornell SMART FTP server<sup>3</sup> or the TMG web page.<sup>4</sup> They are of size  $5735 \times 1033$ ,  $4563 \times 1398$ , and  $5544 \times 1460$ , respectively. We also consider the term-by-document matrix HYPATIA<sup>5</sup> of size  $11390 \times 1265$  with 109056 nonzero terms from the web server at the Department of Mathematics, University of Rhode Island. HYPATIA was created in the same manner as standard term-by-document test matrices. All test matrices use the local term frequency (i.e., number of times a word occurs on a website) and have no global weighting or normalization. The parameter values chosen for the different methods are consistent with our desire to determine the 10 largest

<sup>3</sup><ftp://ftp.cs.cornell.edu/pub/smart>.

<sup>4</sup><http://scgroup.hpclab.ceid.upatras.gr/scgroup/Projects/TMG/>.

<sup>5</sup>The matrix is available at <http://math.uri.edu/~jbaglama>.

TABLE 4.3

Example 4: Computation of the 10 largest singular values of the matrices MEDLINE, CRANFIELD, CISI, and HYPATIA.

irlba(R1)					
Matrix	# Mat.-vec. products	CPU time			Magn. largest error
		mat.-vec. products	one-sided reorthog.	total	
MEDLINE	74	0.23s	0.05s	0.33s	$1.11 \cdot 10^{-10}$
CRANFIELD	78	0.35s	0.03s	0.48s	$1.29 \cdot 10^{-12}$
CISI	76	0.26s	0.08s	0.48s	$2.77 \cdot 10^{-12}$
HYPATIA	78	0.41s	0.17s	0.76s	$1.54 \cdot 10^{-11}$

irlba(R2)					
Matrix	# Mat.-vec. products	CPU time			Magn. largest error
		mat.-vec. products	two-sided reorthog.	total	
MEDLINE	74	0.22s	0.11s	0.41s	$1.11 \cdot 10^{-10}$
CRANFIELD	78	0.35s	0.12s	0.53s	$1.39 \cdot 10^{-12}$
CISI	76	0.26s	0.14s	0.53s	$2.79 \cdot 10^{-12}$
HYPATIA	78	0.40s	0.28s	0.89s	$1.55 \cdot 10^{-11}$

irblsvds			
Matrix	# Mat.-vec. products	CPU time	Magn. largest error
MEDLINE	436	4.55s	$5.09 \cdot 10^{-10}$
CRANFIELD	508	5.72s	$3.28 \cdot 10^{-10}$
CISI	564	6.39s	$2.41 \cdot 10^{-10}$
HYPATIA	432	9.36s	$2.15 \cdot 10^{-09}$

jdsvd(Ref)			
Matrix	# Mat.-vec. products	CPU time	Magn. largest error
MEDLINE	218	2.25s	$7.32 \cdot 10^{-10}$
CRANFIELD	240	2.58s	$7.16 \cdot 10^{-11}$
CISI	218	2.39s	$2.93 \cdot 10^{-10}$
HYPATIA	240	4.70s	$4.43 \cdot 10^{-10}$

svds		
Matrix	CPU time	Magn. largest error
MEDLINE	1.06s	$4.94 \cdot 10^{-12}$
CRANFIELD	1.48s	$2.64 \cdot 10^{-11}$
CISI	1.58s	$2.73 \cdot 10^{-12}$
HYPATIA	2.53s	$1.02 \cdot 10^{-11}$

singular triplets for each of these matrices; for `irlba` we let  $\sigma := LS$  and  $k := 10$ . The available storage is assumed to be large enough to simultaneously store all vectors generated during 20 consecutive steps of Lanczos bidiagonalization. The tolerance  $\delta$  in the stopping criterion is  $10^{-6}$ . Since we seek to determine the largest singular triplets, `irbla` uses the augmentation method of section 3.1. We report the results for one-sided and two-sided full reorthogonalization. The code `irblsvds` was used with block-size 4 and was allowed to carry out at most five consecutive block-Lanczos steps between restarts. The fastest extraction method for `jdsvd` was refined. Table 4.3 displays the performance of the methods and illustrates the competitiveness of

`irlba`. The code `irlanb` is not part of our comparison, since it is designed for the computation of only a few of the smallest singular triplets.

*Example 5* (condition number). We would like to determine the condition number  $\kappa(A) := \sigma_1^{(A)}/\sigma_n^{(A)}$ , where the singular values are enumerated according to (1.1), of the Läuchli matrix  $A := L(n, \mu) \in \mathbb{R}^{(n+1) \times n}$  for  $n = 20000$  and the default value of  $\mu$ .<sup>6</sup> Thus,  $A$  has ones across the top row and  $\mu$  on the subdiagonal; the remaining matrix entries are zero. This matrix often is used to illustrate the drawback of forming  $A^T A$  in least-squares computations; see [19]. The Läuchli matrix  $A$  is nonsingular; it has the simple singular value  $\sqrt{n + \mu^2}$  and the singular value  $\mu$  of multiplicity  $n - 1$  giving the condition number  $\kappa(A) = 9.490724975767860 \cdot 10^9$ , and therefore  $A^T A$  is numerically singular. `irlba` only implicitly works with the matrix  $A^T A$  and is able to compute  $\kappa(A)$ . Specifically, we let  $\text{sigma} := \text{LS}$  and  $\text{SS}$ ,  $k := 1$ , and we allow the storage required for all vectors generated by 20 consecutive steps of Lanczos bidiagonalization. The tolerance  $\delta$  in the stopping criterion is set to machine epsilon  $\epsilon$ . Two-sided full reorthogonalization was employed. We augmented by Ritz vectors when computing the largest singular value and, until the matrices  $B_m$  became ill-conditioned, by harmonic Ritz vectors when determining the smallest singular value. `irlba` required 0.703 and 0.796 seconds of CPU time to compute the largest and smallest singular values of  $A$ , respectively, and gave the condition number  $9.490724975767925 \cdot 10^9$  with a relative error of  $6.83 \cdot 10^{-15}$ . Thus, the fact that  $A^T A$  is numerically singular does not cause `irlba` problems.

**Acknowledgments.** We would like thank Stratis Gallopoulos for a copy of the paper [17], Michiel Hochstenbach and Efi Kokiopoulou for providing copies of their codes, and Costas Bekas, Michiel Hochstenbach, and Efi Kokiopoulou for valuable comments and discussions.

## REFERENCES

- [1] J. BAGLAMA, D. CALVETTI, AND L. REICHEL, *Iterative methods for the computation of a few eigenvalues of a large symmetric matrix*, BIT, 36 (1996), pp. 400–421.
- [2] J. BAGLAMA, D. CALVETTI, AND L. REICHEL, *IRBL: An implicitly restarted block-Lanczos method for large-scale Hermitian eigenproblems*, SIAM J. Sci. Comput., 24 (2003), pp. 1650–1677.
- [3] J. BAGLAMA, D. CALVETTI, AND L. REICHEL, *Algorithm 827: irbleigs: A MATLAB program for computing a few eigenpairs of a large sparse Hermitian matrix*, ACM Trans. Math. Software, 29 (2003), pp. 337–348.
- [4] Å. BJÖRCK, *Numerical Methods for Least Squares Problems*, SIAM, Philadelphia, 1996.
- [5] Å. BJÖRCK, E. GRIMME, AND P. VAN DOOREN, *An implicit shift bidiagonalization algorithm for ill-posed systems*, BIT, 34 (1994), pp. 510–534.
- [6] J. R. BUNCH AND C. P. NIELSEN, *Updating the singular decomposition*, Numer. Math., 31 (1978), pp. 111–129.
- [7] D. CALVETTI, L. REICHEL, AND D. C. SORENSEN, *An implicitly restarted Lanczos method for large symmetric eigenvalue problems*, Electron. Trans. Numer. Anal., 2 (1994), pp. 1–21.
- [8] P. COMON AND G. H. GOLUB, *Tracking a few extreme singular values and vectors in signal processing*, Proc. IEEE, 78 (1990), pp. 1327–1343.
- [9] I. S. DUFF, R. G. GRIMES, AND J. G. LEWIS, *User’s Guide for the Harwell-Boeing Sparse Matrix Collection (Release I)*, Technical report TR/PA/92/86, CERFACS, Toulouse, France, 1992. The matrices are available at <http://math.nist.gov/MatrixMarket/>.
- [10] G. GOLUB AND W. KAHAN, *Calculating the singular values and pseudo-inverse of a matrix*, J. Soc. Indust. Appl. Math. Ser. B Numer. Anal., 2 (1965), pp. 205–224.

---

<sup>6</sup>The Läuchli matrix can be obtained from the Matrix Market website, <http://math.nist.gov/MatrixMarket/deli/Lauchli/>. We used  $\mu = 1.4901006677403 \cdot 10^{-8}$ .



- [11] G. H. GOLUB AND C. F. VAN LOAN, *Matrix Computations*, 3rd ed., The Johns Hopkins University Press, Baltimore, MD, 1996.
- [12] M. GU AND S. C. EISENSTAT, *A Stable and Fast Algorithm for Updating the Singular Value Decomposition*, Technical report YALEU/DCS/RR-966, Department of Computer Science, Yale University, New Haven, CT, 1993.
- [13] M. E. HOCHSTENBACH, *A Jacobi–Davidson type SVD method*, SIAM J. Sci. Comput., 23 (2001), pp. 606–628.
- [14] M. E. HOCHSTENBACH, *Harmonic and refined extraction methods for the singular value problem, with applications to least-squares problems*, BIT, to appear.
- [15] M. E. HOCHSTENBACH, *Subspace Methods for Eigenvalue Problems*, Ph.D. thesis, Department of Mathematics, University of Utrecht, Utrecht, The Netherlands, 2003; also available online from <http://www.case.edu/artsci/math/hochstenbach/>.
- [16] Z. JIA AND D. NIU, *An implicitly restarted refined bidiagonalization Lanczos method for computing a partial singular value decomposition*, SIAM J. Matrix Anal. Appl., 25 (2003), pp. 246–265.
- [17] E. KOKIOPOULOU, C. BEKAS, AND E. GALLOPOULOS, *Computing smallest singular triplets with implicitly restarted Lanczos bidiagonalization*, Appl. Numer. Math., 49 (2004), pp. 39–61.
- [18] R. M. LARSEN, *Lanczos bidiagonalization with partial reorthogonalization*, Chapter of Ph.D. thesis, Department of Computer Science, University of Aarhus, Aarhus, Denmark, 1998; also available online from <http://soi.stanford.edu/~rmunk/>.
- [19] P. LÄUCHLI, *Jordan-Elimination und Ausgleichung nach kleinsten Quadraten*, Numer. Math., 3 (1961), pp. 226–240.
- [20] R. B. LEHOUCQ, *Implicitly restarted Arnoldi methods and subspace iteration*, SIAM J. Matrix Anal. Appl., 23 (2001), pp. 551–562.
- [21] R. B. LEHOUCQ AND D. C. SORENSEN, *Deflation techniques for an implicitly restarted Arnoldi iteration*, SIAM J. Matrix Anal. Appl., 17 (1996), pp. 789–821.
- [22] R. B. LEHOUCQ, D. C. SORENSEN, AND C. YANG, *ARPACK Users’ Guide: Solution of Large-Scale Eigenvalue Problems with Implicitly Restarted Arnoldi Methods*, Software Environ. Tools 6, SIAM, Philadelphia, 1998.
- [23] THE MATHWORKS, INC., *MATLAB, Version 6.5 R13*, Natick, MA, 2002.
- [24] R. B. MORGAN, *Computing interior eigenvalues of large matrices*, Linear Algebra Appl., 154/156 (1991), pp. 289–309.
- [25] R. B. MORGAN, *On restarting the Arnoldi method for large nonsymmetric eigenvalue problems*, Math. Comp., 65 (1996), pp. 1213–1230.
- [26] R. B. MORGAN AND M. ZENG, *Harmonic Restarted Arnoldi Algorithm for Calculating Eigenvalues and Determining Multiplicity*, preprint, 2003.
- [27] C. C. PAIGE, B. N. PARLETT, AND H. A. VAN DER VORST, *Approximate solutions and eigenvalue bounds from Krylov subspaces*, Numer. Linear Algebra Appl., 2 (1995), pp. 115–134.
- [28] H. D. SIMON AND H. ZHA, *Low-rank matrix approximation using the Lanczos bidiagonalization process with applications*, SIAM J. Sci. Comput., 21 (2000), pp. 2257–2274.
- [29] D. C. SORENSEN, *Implicit application of polynomial filters in a k-step Arnoldi method*, SIAM J. Matrix Anal. Appl., 13 (1992), pp. 357–385.
- [30] D. C. SORENSEN, *Numerical methods for large eigenvalue problems*, Acta Numer., 11 (2002), pp. 519–584.
- [31] A. STATHOPOULOS AND Y. SAAD, *Restarting techniques for the (Jacobi-)Davidson symmetric eigenvalue methods*, Electron. Trans. Numer. Anal., 7 (1998), pp. 163–181.
- [32] A. STATHOPOULOS, Y. SAAD, AND K. WU, *Dynamic thick restarting of the Davidson, and the implicitly restarted Arnoldi methods*, SIAM J. Sci. Comput., 19 (1998), pp. 227–245.
- [33] K. WU AND H. SIMON, *Thick-restart Lanczos method for large symmetric eigenvalue problems*, SIAM J. Matrix Anal. Appl., 22 (2000), pp. 602–616.