

2016

# Spatial distribution of the persistent organic pollutants across the Tibetan Plateau and its linkage with the climate systems: a 5-year air monitoring study

Xiaoping Wang

Jiao Ren

*See next page for additional authors*

Creative Commons License



This work is licensed under a [Creative Commons Attribution 3.0 License](https://creativecommons.org/licenses/by/3.0/).

Follow this and additional works at: <https://digitalcommons.uri.edu/gsofacpubs>

## Citation/Publisher Attribution

Wang, X., Ren, J., Gong, P., Wang, C., Xue, Y., Yao, T., and Lohmann, R.: Spatial distribution of the persistent organic pollutants across the Tibetan Plateau and its linkage with the climate systems: a 5-year air monitoring study, *Atmos. Chem. Phys.*, 16, 6901-6911, <https://doi.org/10.5194/acp-16-6901-2016>, 2016.

Available at: <https://doi.org/10.5194/acp-16-6901-2016>

This Article is brought to you for free and open access by the Graduate School of Oceanography at DigitalCommons@URI. It has been accepted for inclusion in Graduate School of Oceanography Faculty Publications by an authorized administrator of DigitalCommons@URI. For more information, please contact [digitalcommons@etal.uri.edu](mailto:digitalcommons@etal.uri.edu).

---

**Authors**

Xiaoping Wang, Jiao Ren, Ping Gong, Chuanfei Wang, Yonggang Xue, Tandong Yao, and Rainer Lohmann



# Spatial distribution of the persistent organic pollutants across the Tibetan Plateau and its linkage with the climate systems: a 5-year air monitoring study

Xiaoping Wang<sup>1,2</sup>, Jiao Ren<sup>1</sup>, Ping Gong<sup>1,2</sup>, Chuanfei Wang<sup>1</sup>, Yonggang Xue<sup>1</sup>, Tandong Yao<sup>1,2</sup>, and Rainer Lohmann<sup>3</sup>

<sup>1</sup>Key Laboratory of Tibetan Environment Changes and Land Surface Processes, Institute of Tibetan Plateau Research, Chinese Academy of Sciences (CAS), Beijing, 100101, China

<sup>2</sup>CAS Center for Excellence in Tibetan Plateau Earth Sciences, Beijing, 100101, China

<sup>3</sup>Graduate School of Oceanography, University of Rhode Island, Narragansett, Rhode Island 02882-1197, USA

Correspondence to: Xiaoping Wang (wangxp@itpcas.ac.cn)

Received: 28 January 2016 – Published in Atmos. Chem. Phys. Discuss.: 23 February 2016

Revised: 17 May 2016 – Accepted: 18 May 2016 – Published: 6 June 2016

**Abstract.** The Tibetan Plateau (TP) has been contaminated by persistent organic pollutants (POPs), including legacy organochlorine pesticides (OCPs) and polychlorinated biphenyls (PCBs) through atmospheric transport. The exact source regions, transport pathways and time trends of POPs to the TP are not well understood. Here polystyrene–divinylbenzene copolymer resin (XAD)-based passive air samplers (PASs) were deployed at 16 Tibetan background sites from 2007 to 2012 to gain further insight into spatial patterns and temporal trends of OCPs and PCBs. The southeastern TP was characterized by dichlorodiphenyl-trichloroethane (DDT)-related chemicals delivered by Indian monsoon air masses. The northern and northwestern TP displayed the greatest absolute concentration and relative abundance of hexachlorobenzene (HCB) in the atmosphere, caused by the westerly-driven European air masses. The interactions between the DDT polluted Indian monsoon air and the clean westerly winds formed a transition zone in central Tibet, where both DDT and HCB were the dominant chemicals. Based on 5 years of continuous sampling, our data indicated declining concentrations of HCB and hexachlorocyclohexanes (HCHs) across the Tibetan region. Inter-annual trends of DDT class chemicals, however, showed less variation during this 5-year sampling period, which may be due to the ongoing usage of DDT in India. This paper demonstrates the possibility of using POP fingerprints to investigate the climate interactions and the validity of using PAS to derive inter-annual atmospheric POP time trends.

## 1 Introduction

Organochlorine pesticides (OCPs) and polychlorinated biphenyls (PCBs) are classified and regulated as persistent organic pollutants (POPs) under the Stockholm Convention (Patterson et al., 2009). Although the extensive usage of these pollutants has been forbidden for several decades, they are still ubiquitous in the global environment and the atmosphere plays a vital role in their global dispersal (Cabrerizo et al., 2011a; Brault et al., 2013; Bogdal et al., 2014).


The spatial distribution of POPs is subject to three factors. On the one hand, the global POP spatial patterns strongly related to the POP emissions with higher levels appeared in urban and agricultural sites where POPs had undergone extensively historical or current usage (Harner et al., 2004). On the other hand, long-range atmospheric transport (LRAT) is responsible for the global transport of POPs, resulting in relatively high burden of POPs in remote area (Aulagnier and Poissant, 2005; von Waldow et al., 2010). Thirdly, the global pattern of POPs could be reassigned by the secondary emissions of POPs, which is a result of “multi-hopping” between air and land–sea surfaces (Lohmann et al., 2007; Nizzetto et al., 2010). The last two factors are strongly influenced by climate variations (temperature, wind, precipitation, and others).

Temperature difference is the important factor that determines the difference in POP concentrations in air (Lamon et al., 2009; Pelley, 2004). Higher temperatures can drive increased volatilization emissions of POPs, and enhance the

POP transport (Dalla Valle et al., 2007; Lamon et al., 2009). The wind flows associated with the climate fluctuations can also exert influence on the transport strength and pathway of POPs (Pelley, 2004). Higher wind speeds can result in more efficient intercontinental transport. The Arctic Oscillation (AO), the North Atlantic Oscillation (NAO) and the Pacific North American (PNA) pattern are three major sources of climate variability in the Northern Hemisphere. During the positive phase of NAO, stronger westerly winds blow across the North Atlantic Ocean, enhancing the transport of POPs from the Canadian Prairies to Great Lakes region (Ma et al., 2004a, b). When the PNA pattern intensified, the southwesterly flow along the west coast of Canada increased and gave rise to a poleward transport of POPs to the Arctic (Gao et al., 2010; Ma et al., 2004b).

Wet deposition is the important way to scavenge pollutants from air. Modelling results suggested that maximum POP removal occurred in places with extensive wet deposition (i.e. the Intertropical Convergence Zone (ITCZ) and the region with plenty of snow) (Jurado et al., 2005). With climate change, ITCZ shifts with seasons and extreme snowy weather events frequently occur (Yancheva et al., 2007; Bednorz, 2002), which will also influence the global distribution of POPs.

Similar to the North Pole and South Pole, the Tibetan Plateau (TP) is a region of particular interest to scientists studying both climate change and POP contamination. The average elevation of the TP is 4700 m, which acts like a “wall” and splits the mid-altitude westerly into two currents (Wang et al., 2010). Moreover, the TP has an area of  $2.5 \times 10^6$  km<sup>2</sup>, which enhances the land–sea thermal contrast and strengthens the Indian monsoon (Wang et al., 2010). Climate of the TP displayed spatial difference with the north and west plateau controlled by westerlies and the south and east is dominated by the Indian monsoon (Schiemann et al., 2009). Contaminant transport over the TP is therefore complicated due to the different climate systems (transport pathways) and different source regions. For example, different perfluoroalkyl acid composition profiles were observed for snow cores from different regions of the TP (Wang et al., 2014). The seasonal variation of dichlorodiphenyltrichloroethanes (DDTs) in the southeastern TP is synchronous with the seasonality of the Indian monsoon, suggesting that the monsoon-driven transport of POPs to the TP is continuous and sustained (Sheng et al., 2013). Although connections between climate circulation and pollutants transport have been observed, the above studies are limited to few sampling points. Actually, the interactions between the westerlies and the Indian monsoon are of great concern given they will influence the moisture, heat, and anthropogenic forcing on regional and global scale. Thus, it is critical to conduct regional air sampling across the TP and investigate the linkage between spatial patterns of POPs over the TP and the interactions between climate systems.



**Figure 1.** Map showing the sampling sites of the POP monitoring programme across the Tibetan Plateau. The values in the legend are in meters.

Taking advantage of the Tibetan Observation and Research Platform (TORP) (Ma et al., 2008), a large-scale and long-term (2007–2012) atmospheric POP monitoring programme across the TP was conducted and the polystyrene–divinylbenzene copolymer resin (XAD)-2-based passive air samplers (PASs) were deployed. The result of the first year (2007–2008) has been given previously (Wang et al., 2010). In this study, all 5-year data (2007–2012) were gathered to get more reliable spatial POP patterns and to investigate the role of the climate systems in forming such patterns. The temporal trends of atmospheric POP concentrations during this 5-year period were also investigated. A better understanding of these processes will contribute to determining how the global climate systems will affect the spatial distribution of contaminants and how background POP levels over the TP vary with time increase.

## 2 Materials and methods

### 2.1 Sampling sites and programmes

An important objective of this study was to improve the knowledge of spatial patterns of POPs in background air across the TP using PAS. Therefore, 16 background sites were selected in the context of LRAT and represented a good spatial coverage of the TP (Fig. 1). Sampling sites were located from Muztaga (38° N) in the north to Mount Everest (28° N) in the south, and from Muztaga (75° E) in the west to Chayu (97° E) in the east; altitudes of sampling sites ranged from 500 to 5200 m (Table 1). Due to its high capacity and continuous accumulating over year-round exposure for POPs, XAD-PASs were chosen in this study (Wania et al., 2003). A consecutive 5-year (July 2007 to August 2012) air monitoring programme was conducted and XAD-PAS was deployed for five 1-year periods to yield annually averaged

concentrations of POPs (Gouin et al., 2008; Hayward et al., 2010). Duplicate samples were deployed at six sites (Lhasa, Xigazê, Lhazê, Mount Everest, Lulang, and Nam Co), and field blanks were taken at Golmud, Qamdo, Lulang, Lhasa, Mount Everest, and Gar for every sampling year (Fig. 1). Table 1 also gives details of the sampling programme including the date of sampling collection and a brief introduction about the sites. An active air sampler (AAS) was additionally co-deployed with XAD-PAS at Lulang (Fig. 1) to evaluate possible differences between PAS and AAS. The sampling period of AAS was from November 2008 to September 2011, and details about the sampling AAS programme were reported in a previous study (Sheng et al., 2013) and also provided in the Supplement S1.

## 2.2 Sample preparation, extraction, and analysis

Prior to deployment, XAD resin was Soxhlet extracted using in turn methanol, acetonitrile, and dichloromethane (DCM). The XAD resin (60 mL of wet XAD in methanol) was transferred to a precleaned stainless steel mesh cylinder and dried in a clean desiccator. Dry cylinders were sealed in an airtight stainless steel tube with Teflon lids. Upon completion of the sampling, all samples (XAD-cylinders) were stored at  $-20^{\circ}\text{C}$  until extraction. Chemical extraction, cleanup, and details of gas chromatographic temperature are given in S2 in the Supplement. The following compounds were measured and quantified: PCB 28, 52, 101, 138, 153 and 180; hexachlorobenzene (HCB); *o,p'*-DDE, *p,p'*-DDE, *o,p'*-DDT and *p,p'*-DDT;  $\alpha$ -HCH (hexachlorocyclohexane),  $\beta$ -HCH,  $\gamma$ -HCH, and  $\delta$ -HCH.

## 2.3 Quality assurance/quality control (QA/QC)

All analytical procedures were monitored by strict QA/QC measures. The blank concentrations and the corresponding method detection limits (MDLs) are given in Table S1 in the Supplement. MDLs were derived as the mean field blank concentration plus 3 times its standard deviation. In the present study, MDL values ranged between 0.04 and 1.2 ng per sample for OCPs, and between 0.10 and 0.32 ng per sample for PCBs (Table S1). The recoveries were between 70 and 106 % for 2,4,5,6-tetrachloro-*m*-xylene (TCmX) and between 72 and 114 % for decachlorobiphenyl (PCB-209). The full data set of the POP concentrations (in unit of ng per sample) over the five sampling years is available as Table S2. All reported values were blank-corrected but not corrected for the recoveries. If the concentration of a compound after blank correction was below the MDL, the concentration was substituted with half a MDL (Table S1). Duplicate PASs were deployed to check the repeatability, and the results showed the average relative deviation of concentrations between duplicates is generally low, which ranged from 17 to 24 % for different compounds (Table S3).

## 2.4 Uncertainties in chemical analysis

It is important to know the uncertainties of measurements when comparing the data of this study with those of other laboratories and assessing the reliable long-term trends. The factors that may introduce uncertainties include (i) loss during sample extraction and clean-up; (ii) instrumental drifts; (iii) accuracy of the employed standard; and the interferences of other compounds. The loss of analytes during extraction and clean-up is the biggest contribution for uncertainties. Estimation for the uncertainty of this step was about 25–30 % and for the whole chemical analysis was in the range of 25–35 %, varying for different compounds.

## 2.5 Sampling rate of XAD-PAS and its uncertainties

Sampling rate ( $R$ ,  $\text{m}^3 \text{day}^{-1} \text{PAS}^{-1}$ ) was estimated using the empirical equation provided in the previous study (Wang et al., 2010),

$$R = 0.16 \times \frac{T^{1.75}}{P} - 2.14. \quad (1)$$

The advantage of this empirical equation is that one can easily deduce the  $R$  of XAD-PAS by using the general parameters of temperature ( $T$ , K) and pressure ( $p$ , hpa) in a sampling site ( $R$  of each individual site was provided in Table 2). After obtaining the sequestered chemical amount per sampler and  $R$ , the PAS results can be derived to yield volumetric air concentrations ( $\text{pg m}^{-3}$ , Table S4).

Apart from the uncertainties caused by chemical analysis,  $R$  is also essential to data quality. For XAD-PAS, uncertainties associated with  $R$  can be from variable environmental conditions, such as turbulent wind and the air temperature inside the sampler shelter which can be different from the temperature recorded by weather stations. For addressing the uncertainties associated with uptake rates, field and laboratory studies have previously been carried out, comparing the air concentrations obtained on the basis of XAD-PAS and active air sampling (AAS) (Hayward et al., 2010; Gouin et al., 2008). The concentrations derived by XAD-PAS were generally lower than those of AAS by a factor of 2–3, but fell into the uncertainty range for AAS (Gouin et al., 2008; Hayward et al., 2010; Zhang et al., 2013; Melymuk et al., 2014). We refer to these studies for a better understanding of uncertainties in PAS uptake rates.

In this study, both XAD-PAS and AAS were deployed in Lulang, and details about the data comparison are provided in S3 in the Supplement. Concentration variability within a factor of 2–3 was found between the derived PAS concentrations and AAS results (Supplement S3). This concentration variability is overall in agreement with previous calibration studies (Gouin et al., 2008; Hayward et al., 2010), which indicated that the sampling rate deduced by Eq. (1) is adequate for this study.

**Table 1.** Information of sampling sites and the date of sample collection for each year.

Sampling Site	Latitude	Longitude	Elevation (m)	Temperature (°C)	Date of sample collection					
					2007	2008	2009	2010	2011	2012
Xigazê	29° 15.014' N	88° 53.319' E	3840	6.6	08/06	06/06	07/20	08/21	08/15	08/24
Nam Co	30° 46.375' N	90° 57.800' E	4740	-2.2	09/01	06/15	07/18	08/24	08/14	08/22
Lhasa	29° 38.728' N	91° 01.956' E	3660	8.1	07/07	06/03	08/06	08/15	08/05	08/20
Lhazê	29° 05.405' N	87° 38.094' E	4020	6.8	08/04	06/07	07/20	08/24	08/13	08/30
Lulang	29° 45.908' N	94° 44.246' E	3330	5.4	07/21	06/12	07/29	08/30	08/19	08/29
Mount Everest	28° 21.633' N	86° 56.948' E	4300	4.3	08/11	07/15	07/23	08/25	08/30	09/06
Naqu	31° 25.373' N	91° 58.827' E	4500	-1	09/04	05/30	07/20	08/13	08/04	08/16
Qamdo	31° 09.014' N	97° 08.624' E	3250	7.6	07/02	05/27	07/23	09/03	08/23	08/25
Saga	29° 19.889' N	85° 13.951' E	4500	6.5	08/03	06/07	08/24	08/23	08/13	08/25
Rawu	29° 22.289' N	96° 54.745' E	4540	-2	08/23	07/22	06/15	09/01	08/21	08/24
Bomi	29° 51.485' N	95° 46.167' E	2720	8.8	07/03	06/13	07/25	08/31	08/20	08/28
GBJD*	29° 53.122' N	93° 14.478' E	3420	6.2	07/04	06/16	–	08/29	08/17	08/20
Golmud	36° 23.637' N	94° 54.480' E	2830	5.3	09/06	06/23	07/17	09/15	08/02	08/14
Gar	32° 30.116' N	80° 05.654' E	4300	0.6	07/30	06/27	08/25	08/20	08/11	08/29
Chayu	28° 39.910' E	97° 27.987' N	500	14.9	–	08/04	06/20	09/01	08/24	08/22
Muztaga	38° 16.072' E	74° 50.919' N	5200	-3.6	–	–	11/15	11/20	11/30	11/15

\* GBJD is the abbreviation for Gongbujiangda.

Almost all sampling sites were in rural/remote areas. Although Lhasa is the capital city of the Tibet Autonomous Region and Golmud is the second biggest city in this plateau, samplers were installed far away (50 km) from the city centre. Due to the travel times for collection, the length of the exposure time differed slightly between different locations. Samples got lost for some sampling year at GBJD, Chayu, and Muztaga.

**Table 2.** Sampling rate ( $R$ ) for 16 sampling sites.

Site	$T$ (K)	$P$ (hpa)	$T^{1.75}/P$	$R$ ( $\text{m}^3 \text{day}^{-1}$ )
Xigazê	280	638	30.0	2.7
Nam Co	271	570	31.7	2.9
Lhasa	281	652	29.6	2.6
Lhazê	280	624	30.7	2.8
Lulang	279	680	27.9	2.3
Mount Everest	277	603	31.3	2.9
Naqu	272	587	31.1	2.8
Qamdo	281	682	28.2	2.4
Saga	280	578	33.1	3.2
Rawu	271	584	31.0	2.8
Bomi	282	732	26.5	2.1
GBJD	279	667	28.6	2.4
Golmud	278	725	26.2	2.0
Gar	274	604	30.5	2.7
Chayu	285	769	25.7	2.0
Muztaga	277	700	26.8	2.2
Average				2.6
SD				0.3

The empirical equation  $R = 0.16 \times \frac{T^{1.75}}{P} - 2.14$  (Wang et al., 2010) was used to calculate the sampling rate.

## 2.6 Statistical method


Cluster analysis is a multivariate procedure that establishes the relationship among objectives (here samples). The similarity among samples is usually defined as the Euclidean distance between the samples. Cluster analysis divides a set of samples into different groups (clusters), and each sample unambiguously falls into one of the groups. The samples in the

same group are more similar to each other due to the short Euclidean distance than to those in other clusters. The dendrogram of the cluster analysis finally provides visual evaluation on the relationships between different samples, i.e. close or distant. Using the XLSTAT software (Addinso, France), we applied cluster analysis (agglomerative hierarchical clustering) to 5-year Tibetan PAS data sets that have been obtained above. Data from XAD-resin-based PAS networks are usually reported in nanograms per sampler. However, cluster analyses were performed on compositional data, not on absolute concentrations. More details about the methodological issues are available in Liu and Wania (2014).

## 3 Results and discussion

### 3.1 Concentrations of POPs in the atmosphere of TP

The full data set for the POP concentrations ( $\text{ng sample}^{-1}$ ) in individual sampling years is available in Table S2, and the average values are provided in Table S5. The derived volumetric concentrations ( $\text{pg m}^{-3}$ , Table S4) were further averaged (Table S6). The dominant chemicals in the atmosphere of TP were HCB followed by *o,p'*-DDT and *p-p'*-DDT (Table S6). Data of this study were compared with the values reported for the previous Global Atmospheric Passive Sampling (GAPS) study (Supplement S4), which also relied on the XAD-PAS (Shunthirasingham et al., 2010) and values reported for surrounding countries of TP (Table 3). By comparison, POP concentrations in the atmosphere of TP are akin to



**Figure 2.** Spatial distribution pattern of DDTs, HCHs, HCB, and PCBs across the Tibetan Plateau.

those observed in the background sites of the world, indicating the remoteness of TP (Supplement S4). In addition, the levels of OCPs and PCBs in TP air were all much lower than the neighbouring regions, such as India (Zhang et al., 2008), Nepal (Gong et al., 2014), and Pakistan (Syed et al., 2013). Even for the background sites of India, it is 1 order of magnitude higher than those of TP (Table 3) (Zhang et al., 2008). POPs produced and consumed in these highly contaminated countries have the potential to undergo LRAT and contaminate the pristine environment of TP, especially under the favourable climatic conditions. Furthermore, the concentration range of POPs across the TP was large (Table 3), which imply some distinctions among different parts of TP, and it is necessary to figure out the factor(s) causing this spatial variability.

### 3.2 Spatial distribution of POPs across the TP

The spatial distribution of DDTs, HCHs, HCB, and PCBs has been reported in our previous study, based on 1 year (2007–2008) of data (Wang et al., 2010). This first-year study indicated some clues on the possible transport and distribution features of POPs across the TP. Yet a single year is insufficient in comprehensively understanding the POP spatial patterns. The integration of the long-term monitoring data is therefore needed. For each sampling site, the mean values of air concentrations over multi-year monitoring are more representative than concentrations for a single year. Here, the mean values of each sampling site were graphically presented in Fig. 2 to address the spatial distribution patterns of various POPs on the Tibetan Plateau. Randomized block ANOVA was further performed to test concentration differences between different sampling sites (Table S7).

The spatial distribution of DDTs shows a decreasing gradient from the southeast to the northwest of TP (Fig. 2). On the basis of the ANOVA results (Table S7), significantly

higher values of DDTs were found at Chayu, Bomi, and GBJD, which are located around the Yarlung Tsangpo Grand Canyon – a channel for receiving the pollutants from southern Asia. Similarly, significantly higher levels of HCHs were also observed at Bomi and GBJD (Table S7). Previous studies suggested that the Yarlung Tsangpo River valley is considered a “leaking wall” that contaminates the southeastern Tibetan Plateau (Sheng et al., 2013; Wang et al., 2015). Given the Indian monsoon is the distinct climate of the southeast Tibetan Plateau, and DDTs and HCHs are the dominant POPs in Indian atmosphere (Zhang et al., 2008; Gioia et al., 2012), the monsoon-driven LRAT of POPs is thus the possible reason why higher DDT/HCH contaminations occurred in the southeast of the Tibetan Plateau (Sheng et al., 2013). Besides the LRAT, scattered usage of HCHs cannot be ruled out. For example, high levels of HCHs were also found in Naqu during this 5-year period (Fig. 2 and Table S7). Naqu is located closer to the central plateau and is an agriculture and pasturage interlaced zone. Thus, the scattered HCH usage in Naqu might be possible. The spatial pattern of HCB is opposite to those observed for DDTs and HCHs, with relatively high concentrations in the northern and western TP (Fig. 2). In addition, all sampling sites in this study displayed low concentrations and a uniform spatial distribution pattern for PCBs (Table S6 and Fig. 2), and no significant difference among different sampling sites was observed by an ANOVA test (Table S7). This is a typical feature of a remote region and indicated that there is very limited primary emission of PCBs in the TP.


### 3.3 Does the soil–air exchange (secondary source) affect the spatial pattern?

Due to the remoteness of the Tibetan Plateau, both atmospheric transport and regional re-evaporation can be two important vectors that influence the spatial distribution pattern of POPs. Soil has been reported as a major environmental reservoir of POPs. Cabrerizo et al. (2011a) found that air–soil exchange controlled atmospheric concentration of OCPs in a background region. Sampling air and soil at the same sites and the determination of soil and ambient air fugacities can provide quantitative evidence to test whether soil is volatilizing POPs to the atmosphere (the direction of soil–air exchange) and how much of POPs will be evaporated (exchange fluxes).

In this study, soil data used for fugacity calculations were taken from a published study (Wang et al., 2012) in which the soils were collected from 2007. Due to its heterogeneity and slow rate of change, chemical concentrations in soil seldom change with time (Schuster et al., 2011). Therefore, we assume the soil concentrations of OCPs and PCBs during the 5-year (2007–2012) sampling period were constant. Relevant equations regarding calculation of the air ( $f_a$ ) and soil fugacity ( $f_s$ ) are provided in Supplement S5. The soil–air fugacity

**Table 3.** Concentration results of this study and the comparison with literature values reported for the surrounding countries ( $\text{pg m}^{-3}$ ).

Sampling site	TP (whole TP, this study)	India (urban)	India (rural)	India (background)	Pakistan (Punjab Province)	Nepal (southern slope of the Himalayas)
Year	2007–2012	2006	2006	2006	2011	2012
Sampler type	XAD-PAS	PUF-PAS	PUF-PAS	PUF-PAS	PUF-PAS	XAD-PAS
$\alpha$ -HCH	3.8 (0.1–17.7)	451 (22–1691)	53 (12–167)	25 (20–31)	19 (6.4–29)	–
$\gamma$ -HCH	1.6 (0.1–20.1)	909 (135–3562)	174 (31–437)	61 (34–100)	20 (5.4–45)	–
HCB	17.8 (3.0–85.0)	–	–	–	33 (13–76)	234 (128–416)
<i>o,p'</i> -DDE	0.8 (0.03–8.7)	–	–	–	63 (12–240)	10.6 (BDL–41)
<i>p,p'</i> -DDE	2.2 (0.1–18.1)	554 (26–2061)	81 (15–282)	13 (6–19)	79 (4.2–290)	154 (17–597)
<i>o,p'</i> -DDT	7.9 (0.1–44.5)	268 (23–620)	88 (BDL–307)	52 (BDL–78)	30 (3.3–77)	159 (33–509)
<i>p,p'</i> -DDT	4.4 (0.1–26.1)	110 (2–249)	79 (3–387)	25 (9–45)	34 (6.0–66)	125 (21–456)
$\sum$ PCBs	0.8 (0.1–3.9)	–	–	–	–	26.9 (3.2–78.5)
Reference		Zhang et al. (2008)	Zhang et al. (2008)	Zhang et al. (2008)	Syed et al. (2013)	Gong et al. (2014)

**Figure 3.** Air–soil fugacity fraction ( $ff$ ) of individual compounds across the TP. The boxes are defined by the 25th and 75th percentiles; whiskers mark the 10th and 90th percentiles; the median is represented by a horizontal line; the mean by a square; and outliers with an asterisk.

fraction ( $ff$ )

$$ff = f_s f_s / (f_s + f_a) \quad (2)$$

was then assessed for 13 sampling sites where both air and soil data sets were available (Table S8). A  $ff = 0.5$  indicates that soil fugacity and air fugacity are same, and the compound is at equilibrium. Due to uncertainties, fugacity fractions between 0.3 and 0.7 were not considered to differ significantly from equilibrium (Harner et al., 2004; Li et al., 2010). Figure 3 shows the box-and-whisker plot of the  $ff$  for different chemicals. From Fig. 3, HCB, PCB-28, PCB-52, and  $\alpha$ -HCH showed mean  $ff$  values slightly higher than 0.7, while mean  $ff$  values of DDTs were similar to or lower than 0.3. This suggested the air–soil exchange of volatile compounds (PCBs, HCB, and  $\alpha$ -HCH) was on the verge of equilibrium but tended to volatilization, but the air–soil exchange of less volatile compounds (DDTs) was prone to deposition.

$\gamma$ -HCH showed a small range and a high mean  $ff$  (close to 1) falling within the interquartile range, indicating a sta-


ble volatilization from soil to air (Fig. 3). Meanwhile, higher volatilization fluxes of  $\gamma$ -HCH (up to  $2.8 \text{ ng m}^{-2} \text{ h}^{-1}$ ) were also found (equations used for calculating the volatilization fluxes are given in Supplement S5 and flux values are provided in Table S9 and Fig. 4). However, HCHs are generally regarded as “swimmers”, which can easily be scavenged by precipitation (Lohmann et al., 2007) and thus their out-gassing from soil may not strongly influence their spatial distribution pattern.

With regard to other chemicals (PCB-28, PCB-52, and HCB), their  $ff$  displayed large overlap with equilibrium range (Fig. 3) and only small evaporation fluxes were derived (Fig. 4). This indicated the re-volatilization of these chemicals may not be a great contributor to their spatial distribution pattern. Given the DDT-class chemicals showed a deposition status (Fig. 3), their spatial distribution pattern was therefore less influenced by secondary emissions, but the LRAT instead. Although HCB is an example of a grasshopper chemical (“multi-hopper”) for which atmospheric concentrations are significantly influenced by the secondary emissions (Bailey, 2001), rather small evaporation fluxes were observed for HCB (average  $ff = 0.82$ ; average evaporation fluxes =  $0.32 \text{ ng m}^{-2} \text{ h}^{-1}$ ; Table S9 and Fig. 4). This may partly be due to the large uncertainties produced by both the air and soil measurements (see discussion above). Overall, given the lower soil concentrations of POPs and relatively cold environment of the TP, LRAT of chemicals instead of secondary sources was likely more important for the atmospheric concentrations and distribution patterns of POPs.

### 3.4 The spatial differences in POP sources and transport

Although the spatial POP concentrations were of primary importance, the relative composition of POPs on different spatial scales is also of interest. Cluster analysis was conducted to test whether these different samples from different sampling years displayed similarity (samples with similar POP compositions are influenced by the similar source or similar transport pathways/climate systems) and can be grouped according to their POP “fingerprints”.





**Figure 4.** The geographical distribution of the net volatilization fluxes of PCB-28, PCB-52, HCB,  $\alpha$ -HCH, and  $\gamma$ -HCH across the TP.


In the present study, we summed up levels of frequently detected compounds in each sample and normalized the level of individual chemical by this sum to yield a relative fraction (%; Table S10). The results of cluster analysis are presented in Fig. 5. In this study, all samples were classified into three groups: Group 1, Group 2, and Group 3 (from left to right in Fig. 5).

The Group 1 comprised 28 samples (Table S11). If data of the observation sites through most of sampling years can be grouped in the same cluster, this suggests that these sites have the real and consistent similarity. However, for the sites in which only one or 2 years of data can be included, these sites are doubtful sites and will not be considered as the representative of the group. In Group 1 of this study, all samples from Chayu (one sample from this site got lost and in total four samples were harvested), 80 % of samples (4 in 5 sampling years) from Lulang, and 60 % of samples (3 in 5 sampling years) from GBJD, Bomi, and Rawu were clustered in this group (Fig. 5, Table S10, 11). Chayu, Lulang, GBJD, Rawu, and Bomi were located in the southeastern TP and could be regarded as the representative sites of this group (Table S12). Although some scattered samples (e.g. Gar 1, Xigazê 1, 3 and Lhasa 3, 5) were clustered in this group, they were not likely the typical sites due to their lower frequency in this group. The dominant compounds of Group 1 were DDT-related chemicals (Fig. 5), which contributed 56.5 % (34.9–79.1 %) to the total POPs. Taking the results of spatial distribution and cluster analysis together, both higher concentrations and higher proportions of DDT class chemicals were found in the southeastern TP. Transport of the POPs strongly relied on the Indian monsoon, which has a major impact on the plateau's south side (Sheng et al., 2013) but is blocked from going further north. Based on the similarity of sampling sites that shared the same Indian source, the main contributions of cluster analysis helped to identify the influential spatial coverage of the Indian monsoon over the TP.


Regarding Group 2, it included 17 samples (Table S11); all Muztaga samples (three samples were harvested), 80 % of Golmud samples, and 60 % of Gar samples were grouped into this cluster (Table S11, 12). In this group, the representative sites (Muztaga, Golmud, and Gar) were from the northern and western TP. HCB occupied the overwhelming majority in this group, which accounted for 77.1 % (69.0–88.4 %) of the total POPs (Table S11). Meanwhile, higher HCB absolute concentrations were also observed for these sites (Table S6). It should also be noted that climate of the northern and western TP is mainly influenced by westerlies, and their upwind POP source regions include Europe and central Asia (Wang et al., 2014; Xu et al., 2009). Generally, the larger the percentage of HCB at a site, the cleaner it is (Liu and Wania, 2014). The GAPS study found that HCB is dominantly and uniformly distributed in the European air (Shunthirasingham et al., 2010). Similar results have also been observed by EMEP monitoring programme (Halse et al., 2011). Therefore, Group 2 of the present study could reflect a regional fingerprint of clean European air.

The remaining 30 samples were classified into Group 3, and samples from Lhazê, Naqu, Lhasa, Mount Everest, and Qamdo had higher frequency (> 60 %, Table S12) in this group. Due to the possible local contamination of Naqu (see discussion above), Naqu was not included as a representative site of this group. For Group 3, the dominant chemicals were HCB followed by DDTs, comprising 54.6 % (29.3–67.6 %) and 24.7 % (5.6–43.3 %) of the total POPs, respectively, and the representative sampling sites were mainly located on the central TP. This indicated the effective regional atmospheric mixing likely happened on the central TP, where both Indian and European sources could be seen. Samples of three sites (Xigazê, Nam Co, and Saga) did not specifically fall in any group but were scattered among these three groups. In part, this may be caused by the bias originating from the laboratory analysis uncertainties. On the other hand, as these sites are in closer proximity to other sites in Group 3 and are all from the central TP with relatively uniform geography, Xigazê, Nam Co, and Saga were more likely to belong to Group 3. Data of duplicate samples were brought into the cluster analysis to test whether the uncertainties of laboratory analysis could affect the cluster results (Fig. S1 in the Supplement). Nineteen pairs of duplicates appeared in the same group; only 4 of 23 pairs of duplicates were dispersed in different groups. This suggested that cluster analysis extracted real difference/similarity between sampling sites, instead of analytical variability.

Based on the classification derived from the cluster analysis, the whole Tibetan Plateau could be divided into three parts with three distinct POP fingerprints (Fig. 6): one reflecting the Indian monsoon air mass (DDTs, southeastern TP), one reflecting the clean westerly air mass (HCB, northwestern TP), and one that is just the mixture of these two air masses (mixed DDT and HCB, central TP). The formation of this spatial pattern can be attributed to the direct influ-



**Figure 5.** Dendrogram obtained by the cluster analysis (above) and the corresponding relative composition plot (bottom). The dendrogram includes a horizontal dashed line indicating where the number of clusters is stable, because moving up and down of this line along the similarity axis, the number of groups did not change. Samples in the composition plot were named by the name of sampling site and the Arabic numbers 1–5, which represent the sampling year from 2007 to 2012. For example, the first sampling year (from 2007 to 2008) is numbered as 1, and the second year (from 2008 to 2009) is numbered as 2, and so on.




**Figure 6.** The geographic groups based on the cluster analysis (Group 1: monsoon region, Group 2: westerly region, Group 3: transition region).

ences and interactions between different atmospheric circulations (Indian monsoon and westerly winds). From Fig. 6, we roughly assigned the region, south of 30° N and east of 92° E, as the monsoon region; the region, north of 35° N, as the westerly domain; and then the region located in between these two domains (from 30 to 35° N, and west of 92° E) was

regarded as the transition domain, which is under the control of a shifting climate between Indian monsoon and westerly.

Basically, precipitation oxygen isotope ratio ( $\delta^{18}\text{O}$ ) is an integrated tracer of the atmospheric processes, which has been employed to investigate the interaction between the westerlies and Indian monsoon on the TP (Tian et al., 2007). Based on long-term observations of precipitation  $\delta^{18}\text{O}$  on 20 stations of the TP, Yao et al. (2013) found that there is a transition domain located in between the westerly region and monsoon region. This is akin to our POP fingerprint patterns. Thus, outcomes of this study recommended that the POP fingerprints could also act as a tracer like  $\delta^{18}\text{O}$  to estimate the interactions between climate systems. As compared with precipitation collection, which is expensive and laborious (every single rain or snow event should be collected), PAS is cheap and simple. If XAD-PAS can be employed with sufficient spatial resolution and coverage, the POP fingerprint difference across the TP will be more distinct, and clear boundaries between monsoon region, westerly region, and transition region can be captured. Especially for the remote region like the TP, POP fingerprints obtained by the PAS would certainly help to understand the realistic synoptic atmospheric patterns. So far, researchers have paid attention to how climate change is affecting POP cycling (Bustnes et al., 2010; Dalla Valle et al., 2007). However, the opposite way of thinking allows us to use POP fingerprints as possible chemical tracers to track the climate dynamics and global pollution diffusion events.



**Figure 7.** The temporal trends of each compound over the 5 sampling years.

### 3.5 Temporal trends

Long-term air monitoring of POPs can also provide temporal patterns, which can be used to evaluate the effectiveness of regional regulations on POPs. Also, the concentrations observed over this 5-year period can be used as a benchmark for future work. Lastly, given the TP is the hinterland abutted by central, eastern, and southern Asia, the temporal patterns of POPs can also be used to test whether there is evidence of decreasing concentrations for this wide Asiatic region. Therefore, the inter-annual variation of the atmospheric POPs is given in Fig. 7 using a box-and-whisker plot. Decreasing concentrations of  $\alpha$ -HCH,  $\gamma$ -HCH, PCBs, and HCB were observed (Fig. 7). Regression analysis revealed that the concentration declines of these compounds with sampling year (from 2008 to 2012) were significant (Fig. S2), which is in good agreement with the result of the GAPS study (Shunthirasingham et al., 2010). This suggested the effectiveness of Stockholm Convention in reducing the emissions of these substances in Asian countries to the background atmosphere. As compared with other compounds, concentrations of two parent DDTs went down and up during the 5-year sampling period (Fig. 7), and the plots of DDT concentration versus sampling year did not show significant correlations (Fig. S2). Randomized block ANOVA was further performed to test differences in the concentrations of DDTs among the 5 years of sampling (2007 to 2012) in all sampling sites (Table S13). The  $p$ - $p'$ -DDT and  $p$ - $p'$ -DDE concentrations in the southeastern TP (monsoon region) did not differ significantly among the 5 years of sampling (Table S13). As mentioned above, the southeastern TP is a receptor region of Indian sources; the observed temporal pattern of DDT suggested that the regulation of DDTs in India might be less effective (Sharma et al., 2014). This suggested that ongoing DDT usage in the aspect of health and epidemic prevention and the illegal DDT ap-

plication for agricultural purpose in India needed to be better controlled and regulated.

## 4 Conclusions and implications

This study confirms that, for a remote region, the spatial distribution patterns of POPs are closely related to the variations and interactions of climate systems. This study also highlights that POP fingerprints can be used as chemical tracers to track the interactions of climate systems. This is of great significance as it indicated a simple and cost-effective PAS can yield valuable data on the realistic synoptic atmospheric interactions. Taking into account the close connections between climate fluctuations (AO, NAO, PNA and ENSO) and POP levels/fingerprints, spatial and temporal POP variations reflected by the PAS technique can provide extra evidence for understanding the process of climate change. The results obtained from this study also highlighted the feasibility of PAS to serve in identification of inter-annual trends of POPs. Long-term air monitoring of POPs using PASs can therefore be used to evaluate the effectiveness of the Stockholm Convention.

Results obtained from our investigation emphasized the need for performing more studies to better understand the secondary emission of POPs over the TP. Fugacity sampler (Cabrerizo et al., 2011b, 2013; Degrendele et al., 2016) is therefore recommended for the future studies due to the fact that it can provide accurate air–soil fugacity gradients. More studies should also be conducted to reduce the uncertainty of PAS sampling rate and get a relatively accurate air concentration, which is essential for global comparison and discerning the time trends.

**The Supplement related to this article is available online at doi:10.5194/acp-16-6901-2016-supplement.**

*Acknowledgements.* This study was supported by the National Natural Science Foundation of China (41222010 and 41071321) and Youth Innovation Promotion Association (CAS2011067). X. Wang acknowledges the staff at the Southeast Tibet Observation and Research Station for the Alpine Environment for helping with field sample collection. Details about sampling collection, analysis, detection limits, concentrations, air–soil gas exchange calculations, and regression analysis of the time trend are provided in the Supplement.

Edited by: R. Ebinghaus

## References

- Aulagnier, F. and Poissant, L.: Some Pesticides Occurrence in Air and Precipitation in Québec, Canada, *Environ. Sci. Technol.*, 39, 2960–2967, doi:10.1021/es048361s, 2005.
- Bailey, R. E.: Global hexachlorobenzene emissions, *Chemosphere*, 43, 167–182, 2001.
- Bednorz, E.: Snow cover in western Poland and macro-scale circulation conditions, *Int. J. Climatol.*, 22, 533–541, 2002.
- Bogdal, C., Müller, C. E., Buser, A. M., Wang, Z., Scheringer, M., Gerecke, A. C., Schmid, P., Zennegg, M., MacLeod, M., and Hungerbühler, K.: Emissions of Polychlorinated Biphenyls, Polychlorinated Dibenzo-p-dioxins, and Polychlorinated Dibenzofurans during 2010 and 2011 in Zurich, Switzerland, *Environ. Sci. Technol.*, 48, 482–490, 10.1021/es4044352, 2014.
- Braut, E. K., Goebel, M. E., Geisz, H. N., Canuel, E. A., and Dickhut, R. M.: Inter-Annual Variation of Persistent Organic Pollutants (POPs) in an Antarctic Top Predator *Arctocephalus gazella*, *Environ. Sci. Technol.*, 47, 12744–12752, doi:10.1021/es403577f, 2013.
- Bustnes, J. O., Gabrielsen, G. W., and Verreault, J.: Climate Variability and Temporal Trends of Persistent Organic Pollutants in the Arctic: A Study of Glaucous Gulls, *Environ. Sci. Technol.*, 44, 3155–3161, doi:10.1021/es9032919, 2010.
- Cabrerizo, A., Dachs, J., Jones, K. C., and Barceló, D.: Soil-Air exchange controls on background atmospheric concentrations of organochlorine pesticides, *Atmos. Chem. Phys.*, 11, 12799–12811, doi:10.5194/acp-11-12799-2011, 2011a.
- Cabrerizo, A., Dachs, J., Moeckel, C., Ojeda, M.-J., Caballero, G., Barceló, D., and Jones, K. C.: Factors Influencing the Soil–Air Partitioning and the Strength of Soils as a Secondary Source of Polychlorinated Biphenyls to the Atmosphere, *Environ. Sci. Technol.*, 45, 4785–4792, doi:10.1021/es200400e, 2011b.
- Cabrerizo, A., Dachs, J., Barceló, D., and Jones, K. C.: Climatic and Biogeochemical Controls on the Remobilization and Reservoirs of Persistent Organic Pollutants in Antarctica, *Environ. Sci. Technol.*, 47, 4299–4306, doi:10.1021/es400471c, 2013.
- Dalla Valle, M., Codato, E., and Marcomini, A.: Climate change influence on POPs distribution and fate: A case study, *Chemosphere*, 67, 1287–1295, doi:10.1016/j.chemosphere.2006.12.028, 2007.
- Degrendele, C., Audy, O., Hofman, J., Kučerik, J., Kukučka, P., Mulder, M. D., Přibyllová, P., Prokeš, R., Šáňka, M., Schaumann, G. E., and Lammel, G.: Diurnal Variations of Air-Soil Exchange of Semivolatile Organic Compounds (PAHs, PCBs, OCPs, and PBDEs) in a Central European Receptor Area, *Environ. Sci. Technol.*, 50, 4278–4288, doi:10.1021/acs.est.5b05671, 2016.
- Gao, H., Ma, J., Cao, Z., Dove, A., and Zhang, L.: Trend and climate signals in seasonal air concentration of organochlorine pesticides over the Great Lakes, *J. Geophys. Res.-Atmos.*, 115, D15307, doi:10.1029/2009JD013627, 2010.
- Gioia, R., Li, J., Schuster, J., Zhang, Y., Zhang, G., Li, X., Spiro, B., Bhatia, R. S., Dachs, J., and Jones, K. C.: Factors Affecting the Occurrence and Transport of Atmospheric Organochlorines in the China Sea and the Northern Indian and South East Atlantic Oceans, *Environ. Sci. Technol.*, 46, 10012–10021, doi:10.1021/es302037t, 2012.
- Gong, P., Wang, X.-p., Li, S.-H., Yu, W.-s., Kattel, D. B., Wang, W.-C., Devkota, L. P., Yao, T.-D., and Joswiak, D. R.: Atmospheric transport and accumulation of organochlorine compounds on the southern slopes of the Himalayas, Nepal, *Environ. Pollut.*, 192, 44–51, 2014.
- Gouin, T., Wania, F., Ruepert, C., and Castillo, L.: Field Testing Passive Air Samplers for Current Use Pesticides in a Tropical Environment, *Environ. Sci. Technol.*, 42, 6625–6630, doi:10.1021/es8008425, 2008.
- Halse, A. K., Schlabach, M., Eckhardt, S., Sweetman, A., Jones, K. C., and Breivik, K.: Spatial variability of POPs in European background air, *Atmos. Chem. Phys.*, 11, 1549–1564, doi:10.5194/acp-11-1549-2011, 2011.
- Harner, T., Shoeib, M., Diamond, M., Stern, G., and Rosenberg, B.: Using passive air samplers to assess urban-rural trends for persistent organic pollutants. 1. Polychlorinated biphenyls and organochlorine pesticides, *Environ. Sci. Technol.*, 38, 4474–4483, 2004.
- Hayward, S. J., Gouin, T., and Wania, F.: Comparison of Four Active and Passive Sampling Techniques for Pesticides in Air, *Environ. Sci. Technol.*, 44, 3410–3416, doi:10.1021/es902512h, 2010.
- Jurado, E., Jaward, F., Lohmann, R., Jones, K. C., Simó, R., and Dachs, J.: Wet Deposition of Persistent Organic Pollutants to the Global Oceans, *Environ. Sci. Technol.*, 39, 2426–2435, doi:10.1021/es048599g, 2005.
- Lamon, L., von Waldow, H., MacLeod, M., Scheringer, M., Marcomini, A., and Hungerbühler, K.: Modeling the Global Levels and Distribution of Polychlorinated Biphenyls in Air under a Climate Change Scenario, *Environ. Sci. Technol.*, 43, 5818–5824, doi:10.1021/es900438j, 2009.
- Li, Y.-F., Harner, T., Liu, L., Zhang, Z., Ren, N.-Q., Jia, H., Ma, J., and Sverko, E.: Polychlorinated Biphenyls in Global Air and Surface Soil: Distributions, Air–Soil Exchange, and Fractionation Effect, *Environ. Sci. Technol.*, 44, 2784–2790, doi:10.1021/es901871e, 2010.
- Liu, X. and Wania, F.: Cluster analysis of passive air sampling data based on the relative composition of persistent organic pollutants, *Environmental Science: Processes & Impacts*, 16, 453–463, 2014.
- Lohmann, R., Breivik, K., Dachs, J., and Muir, D.: Global fate of POPs: Current and future research directions, *Environ. Pollut.*, 150, 150–165, doi:10.1016/j.envpol.2007.06.051, 2007.
- Ma, J., Cao, Z., and Hung, H.: North Atlantic Oscillation signatures in the atmospheric concentrations of persistent organic pollutants: An analysis using Integrated Atmospheric Deposition Network–Great Lakes monitoring data, *J. Geophys. Res.-Atmos.*, 109, D12305, doi:10.1029/2003JD004435, 2004a.
- Ma, J., Hung, H., and Blanchard, P.: How Do Climate Fluctuations Affect Persistent Organic Pollutant Distribution in North America? Evidence from a Decade of Air Monitoring, *Environ. Sci. Technol.*, 38, 2538–2543, doi:10.1021/es0349610, 2004b.
- Ma, Y., Kang, S., Zhu, L., Xu, B., Tian, L., and Yao, T.: Roof of the world: Tibetan observation and research platform: Atmosphere-land interaction over a heterogeneous landscape, *B. Am. Meteorol. Soc.*, 89, 1487–1492, 2008.
- Melymuk, L., Bohlin, P., Šáňka, O., Pozo, K., and Klánová, J.: Current Challenges in Air Sampling of Semivolatile Organic Contaminants: Sampling Artifacts and Their Influence on Data Comparability, *Environ. Sci. Technol.*, 48, 14077–14091, doi:10.1021/es502164r, 2014.

- Nizzetto, L., Macleod, M., Borgå, K., Cabrerizo, A., Dachs, J., Guardo, A. D., Ghirardello, D., Hansen, K. M., Jarvis, A., Lindroth, A., Ludwig, B., Monteith, D., Perlinger, J. A., Scheringer, M., Schwendenmann, L., Semple, K. T., Wick, L. Y., Zhang, G., and Jones, K. C.: Past, Present, and Future Controls on Levels of Persistent Organic Pollutants in the Global Environment, *Environ. Sci. Technol.*, 44, 6526–6531, doi:10.1021/es100178f, 2010.
- Patterson, J., Donald, G., Wong, L.-Y., Turner, W. E., Caudill, S. P., DiPietro, E. S., McClure, P. C., Cash, T. P., Osterloh, J. D., Pirkle, J. L., and Sampson, E. J.: Levels in the US population of those persistent organic pollutants (2003–2004) included in the Stockholm Convention or in other long-range transboundary air pollution agreements, *Environ. Sci. Technol.*, 43, 1211–1218, 2009.
- Pelley, J.: POPs levels linked to climate fluctuations, *Environ. Sci. Technol.*, 38, 156A–156A, doi:10.1021/es040486+, 2004.
- Schiemann, R., Lüthi, D., and Schär, C.: Seasonality and interannual variability of the westerly jet in the Tibetan Plateau region, *J. Climate*, 22, 2940–2957, 2009.
- Schuster, J. K., Gioia, R., Moeckel, C., Agarwal, T., Bucheli, T. D., Breivik, K., Steinnes, E., and Jones, K. C.: Has the Burden and Distribution of PCBs and PBDEs Changed in European Background Soils between 1998 and 2008? Implications for Sources and Processes, *Environ. Sci. Technol.*, 45, 7291–7297, doi:10.1021/es200961p, 2011.
- Sharma, B. M., Bharat, G. K., Tayal, S., Nizzetto, L., Čupr, P., and Larssen, T.: Environment and human exposure to persistent organic pollutants (POPs) in India: A systematic review of recent and historical data, *Environ. Int.*, 66, 48–64, doi:10.1016/j.envint.2014.01.022, 2014.
- Sheng, J., Wang, X., Gong, P., Joswiak, D. R., Tian, L., Yao, T., and Jones, K. C.: Monsoon-driven transport of organochlorine pesticides and polychlorinated biphenyls to the Tibetan Plateau: three year atmospheric monitoring study, *Environ. Sci. Technol.*, 47, 3199–3208, 2013.
- Shunthirasingham, C., Oyiliagu, C. E., Cao, X., Gouin, T., Wania, F., Lee, S.-C., Pozo, K., Harner, T., and Muir, D. C.: Spatial and temporal pattern of pesticides in the global atmosphere, *J. Environ. Monitor.*, 12, 1650–1657, 2010.
- Syed, J. H., Malik, R. N., Liu, D., Xu, Y., Wang, Y., Li, J., Zhang, G., and Jones, K. C.: Organochlorine pesticides in air and soil and estimated air–soil exchange in Punjab, Pakistan, *Sci. Total Environ.*, 444, 491–497, 2013.
- Tian, L., Yao, T., MacClune, K., White, J., Schilla, A., Vaughn, B., Vachon, R., and Ichiyanagi, K.: Stable isotopic variations in west China: a consideration of moisture sources, *J. Geophys. Res.-Atmos.*, 112, D10112, doi:10.1029/2006JD007718, 2007.
- von Waldow, H., MacLeod, M., Jones, K., Scheringer, M., and Hungerbühler, K.: Remoteness from Emission Sources Explains the Fractionation Pattern of Polychlorinated Biphenyls in the Northern Hemisphere, *Environ. Sci. Technol.*, 44, 6183–6188, doi:10.1021/es101291q, 2010.
- Wang, X., Gong, P., Yao, T., and Jones, K. C.: Passive Air Sampling of Organochlorine Pesticides, Polychlorinated Biphenyls, and Polybrominated Diphenyl Ethers Across the Tibetan Plateau, *Environ. Sci. Technol.*, 44, 2988–2993, doi:10.1021/es9033759, 2010.
- Wang, X., Sheng, J., Gong, P., Xue, Y., Yao, T., and Jones, K. C.: Persistent organic pollutants in the Tibetan surface soil: spatial distribution, air–soil exchange and implications for global cycling, *Environ. Pollut.*, 170, 145–151, 2012.
- Wang, X., Halsall, C., Codling, G., Xie, Z., Xu, B., Zhao, Z., Xue, Y., Ebinghaus, R., and Jones, K. C.: Accumulation of perfluoroalkyl compounds in tibetan mountain snow: temporal patterns from 1980 to 2010, *Environ. Sci. Technol.*, 48, 173–181, 2014.
- Wang, X., Gong, P., Sheng, J., Joswiak, D. R., and Yao, T.: Long-Range Atmospheric Transport of Particulate Polycyclic Aromatic Hydrocarbons and the Incursion of Aerosols to the Southeast Tibetan Plateau, *Atmos. Environ.*, 115, 124–131, 2015.
- Wania, F., Shen, L., Lei, Y. D., Teixeira, C., and Muir, D. C.: Development and calibration of a resin-based passive sampling system for monitoring persistent organic pollutants in the atmosphere, *Environ. Sci. Technol.*, 37, 1352–1359, 2003.
- Xu, B., Cao, J., Hansen, J., Yao, T., Joswiak, D. R., Wang, N., Wu, G., Wang, M., Zhao, H., and Yang, W.: Black soot and the survival of Tibetan glaciers, *Proc. Natl. Acad. Sci. USA*, 106, 22114–22118, 2009.
- Yancheva, G., Nowaczyk, N. R., Mingram, J., Dulski, P., Schettler, G., Negendank, J. F., Liu, J., Sigman, D. M., Peterson, L. C., and Haug, G. H.: Influence of the intertropical convergence zone on the East Asian monsoon, *Nature*, 445, 74–77, 2007.
- Yao, T., Masson-Delmotte, V., Gao, J., Yu, W., Yang, X., Risi, C., Sturm, C., Werner, M., Zhao, H., and He, Y.: A review of climatic controls on  $\delta^{18}O$  in precipitation over the Tibetan Plateau: Observations and simulations, *Rev. Geophys.*, 51, 525–548, 2013.
- Zhang, G., Chakraborty, P., Li, J., Sampathkumar, P., Balasubramanian, T., Kathiresan, K., Takahashi, S., Subramanian, A., Tanabe, S., and Jones, K. C.: Passive Atmospheric Sampling of Organochlorine Pesticides, Polychlorinated Biphenyls, and Polybrominated Diphenyl Ethers in Urban, Rural, and Wetland Sites along the Coastal Length of India, *Environ. Sci. Technol.*, 42, 8218–8223, doi:10.1021/es8016667, 2008.
- Zhang, X., Brown, T. N., Ansari, A., Yeun, B., Kitaoka, K., Kondo, A., Lei, Y. D., and Wania, F.: Effect of Wind on the Chemical Uptake Kinetics of a Passive Air Sampler, *Environ. Sci. Technol.*, 47, 7868–7875, doi:10.1021/es401486f, 2013.