

THE
UNIVERSITY
OF RHODE ISLAND

University of Rhode Island
DigitalCommons@URI

Graduate School of Oceanography Faculty
Publications

Graduate School of Oceanography

2015

Comparative biogeochemistry–ecosystem–human interactions on dynamic continental margins

Lisa A. Levin

Kon-Kee Liu

See next page for additional authors

Creative Commons License

[Creative Commons License](#)

This work is licensed under a [Creative Commons Attribution-Noncommercial-No Derivative Works 4.0 License](#).

Follow this and additional works at: <https://digitalcommons.uri.edu/gsofacpubs>

This is a pre-publication author manuscript of the final, published article.

Citation/Publisher Attribution

Levin, L. A., Liu, K.-K., Emeis, K.-C., Breitburg, D. L., Cloern, J., Deutsch, C., Giani, M.,...Wishner, K. (2015). Comparative biogeochemistry–ecosystem–human interactions on dynamic continental margins. *Journal of Marine Systems*, 141, 3-17. doi: 10.1016/j.jmarsys.2014.04.016

Available at: <https://doi.org/10.1016/j.jmarsys.2014.04.016>

This Article is brought to you for free and open access by the Graduate School of Oceanography at DigitalCommons@URI. It has been accepted for inclusion in Graduate School of Oceanography Faculty Publications by an authorized administrator of DigitalCommons@URI. For more information, please contact digitalcommons@etal.uri.edu.

Authors

Lisa A. Levin, Kon-Kee Liu, Kay-Christian Emeis, Denise L. Breitburg, James Cloern, Curtis Deutsch, Michele Giani, Anne Goffart, Eileen E. Hofmann, Zouhair Lachkar, Karin Limburg, Su-Mei Liu, Enrique Montes, Wajih Naqvi, Olivier Ragueneau, Christopher Rabouille, Santosh Kumar Sarkar, Dennis P. Swaney, Paul Wassman, and Karen F. Wishner

46 ¹⁸ Department of Ecology and Evolutionary Biology, Cornell University, Ithaca NY
47 14853, USA, dps1@cornell.edu USA

48 ¹⁹ Dept. of Arctic and Marine Biology, Faculty of Bioscience, Fishery and Economy,
49 University of Tromsø, 9037 Tromsø, NORWAY, paul.wassmann@uit.no

50 ²⁰ University of Rhode Island, 215 S Ferry Rd, Narragansett, RI 02882, USA,
51 kwishner@gso.uri.edu

52

53 **Abstract:** The ocean's continental margins face strong and rapid change, forced by a
54 combination of direct human activity, anthropogenic CO₂-induced climate change, and
55 natural variability. Stimulated by discussions in Goa, India at the IMBER IMBIZO III,
56 we (1) provide an overview of the drivers of biogeochemical variation and change on
57 margins, (2) compare temporal trends in hydrographic and biogeochemical data across
58 different margins (3) review ecosystem responses to these changes, (4) highlight the
59 importance of margin time series for detecting and attributing change and (5) examine
60 societal responses to changing margin biogeochemistry and ecosystems. We synthesize
61 information over a wide range of margin settings in order to identify the commonalities
62 and distinctions among continental margin ecosystems. Key drivers of biogeochemical
63 variation include long-term climate cycles, CO₂-induced warming, acidification, and
64 deoxygenation, as well as sea level rise, eutrophication, hydrologic and water cycle
65 alteration, changing land use, fishing, and species invasion. Ecosystem responses are
66 complex and impact major margin services including primary production, fisheries
67 production, nutrient cycling, shoreline protection, chemical buffering, and biodiversity.
68 Despite regional differences, the societal consequences of these changes are unarguably
69 large and mandate coherent actions to reduce, mitigate and adapt to multiple stressors on
70 continental margins.

71

72 **Keywords:** anthropogenic factors, coastal biogeochemistry, climate change,
73 eutrophication, ecosystem services, time series

74

75 **Regional Terms:** Continental margins, Europe, North Atlantic, North Pacific; Arctic

76

77

78

79

80

81

82

83

84

85

86

87

88 **1. Introduction to dynamic margin ecosystems**

89

90 The oceans' continental margins extend for some 150,600 km (Jahnke, 2010) and
91 encompass estuarine, open coast, shelf, canyon, slope, and enclosed sea ecosystems.
92 They are both gateway and window to the open ocean, where water, nutrients, energy,
93 sediments, contaminants and organisms meet and are transferred through land-margin
94 and margin-open ocean interactions (Levin et al., 2001). The continental margins include
95 proximal estuaries, bays, lagoons and banks, and distal shelves, slopes and marginal seas.
96 These are susceptible to changes in biodiversity, water quality, and productivity and have
97 been increasingly perturbed by human activities.

98

99 Margin ecosystems include hard and soft-substrate habitats ranging from structurally
100 complex wetlands, kelp forests, coral reefs, rocky reefs and sand beaches, to sedimented
101 estuaries, slopes and canyons. Most of the habitat volume, however, occurs in the
102 overlying water column, with variation linked to water masses, circulation, and land and
103 atmospheric interactions. As one crosses depth contours from estuaries across the shelf
104 to the continental slope, steep gradients in nutrient concentrations, temperature, salinity,
105 oxygen, pH and suspended matter are found that impact the productivity, composition,
106 diversity, and abundance of organisms (Cloern, 1996; Hofmann et al., 2011; Levin and
107 Sibuet, 2012). Relative to their area, the margins account for a disproportionately large
108 fraction of the global primary production (10 – 15%), nutrient recycling, carbon burial (>
109 60% of total settling organic carbon), and fisheries production (Walsh et al. 1988;
110 Muller-Karger et al., 2005). They also are exceptionally dynamic systems with ecosystem
111 structures that can oscillate slowly or shift abruptly, but rarely remain static.

112

113 The current continental margin seascape has been shaped extensively by climate change
114 and human activities, yielding altered ecosystem services. Margin ecosystems provide
115 key services in the form of physical protection from waves, storms, and floods, chemical
116 buffering, food provisioning, nursery support, nutrient cycling, habitat fostering
117 biodiversity, carbon sequestration, recreation, and aesthetic value. Finely tuned
118 biogeochemical interactions drive these functions. Because human populations are
119 disproportionately concentrated in coastal cities, there is heterogeneity in the human
120 effects on margins, creating mosaics of heavily impacted and relatively pristine systems.
121 Nutrient inputs, freshwater extraction, fishing, construction, species introductions, and
122 contamination are but a few of the many ways humans alter coastal ecosystems. Also the
123 steady increase of anthropogenic CO₂ inputs to the atmosphere will result in significant
124 changes in water column temperature, oxygenation, pH, and productivity by 2100, with
125 major consequences for margin ecosystems and the over 1 billion people that depend on
126 them for food, employment and revenue (Mora et al., 2013; FAO, 2012).

127

128 While margin research has a long history among oceanographers (Banner et al., 1980.
129 Walsh et al., 1988, Biscaye et al., 1994, Duarte et al. 1999, Antia et al., 2001, Liu et al.,
130 2010), a synoptic view of dynamic coupled margin systems has emerged more slowly and
131 the linkages between human and natural biogeochemical variations, ecosystem response
132 and human social structures are only now being explored. The interactive effects of
133 remote forcing from distant inland activities, from atmospheric processes, and from

134 physical processes far out to sea are becoming more apparent. There are, however, gaps
135 in our understanding of the combined effects of multiple drivers on coastal
136 biogeochemistry and ecosystems across all continental margins. The objectives of this
137 paper are to provide an overview of sources of biogeochemical variation on margins,
138 associated ecosystem responses, and the societal and policy implications, with a focus on
139 lessons from multiple continental margin time series (Fig 1).

140

141 This paper reflects the themes and discussions of the continental margins working group
142 of IMBER IMBIZO III (Goa, India in January 2013). In this paper we examine drivers of
143 biogeochemical variation on margins, distinguishing natural from CO₂-based climate
144 variability, and more direct human drivers. We next compare temporal trends for
145 multiple physical and biogeochemical parameters at geographically contrasting locations
146 We then discuss the complex ecosystem responses to biogeochemical variation and
147 trends on margins, in particular those related to warming, deoxygenation, acidification
148 and hydrologic alterations. We subsequently identify the key roles played by continental
149 margin time-series stations (Fig. 1) in identifying and attributing drivers of change and in
150 understanding the associated ecosystem responses. Finally, we examine societal
151 responses to changing margin biogeochemistry and ecosystems, highlighting areas where
152 social and natural scientists must work together. Case studies (expanded in Supplement
153 B) are synthesized to provide insights into the sensitivity of margins to natural and human
154 perturbations, the ecological, social and economic consequences that stem from these
155 perturbations, and the policy actions needed to mitigate impacts on coastal ecosystems
156 and their resources.

157

158 **2. Natural and human-induced drivers of biogeochemical variation on margins**

159

160 Controls on biogeochemistry of margins are complex and dynamic. In this paper we
161 distinguish drivers associated with natural *variability*, anthropogenic CO₂ –driven *climate*
162 *change* and direct human (*anthropogenic*) impacts. It is often difficult to disentangle
163 these three forcing mechanisms, as well as to distinguish local change from regional to
164 global-scale pressures. Multiple factors act together – exerting top-down (often human)
165 and bottom – up (natural or human) controls on ecosystem structure simultaneously.

166

167 ***Natural Sources of Variability***

168 Natural variation in biogeochemical features affecting margin ecosystems occurs on a
169 vast range of time scales, from millions of years to hours. Direct measurements during
170 the past century in many of the systems discussed here have revealed large, abrupt,
171 persistent changes in the structure and function (or state) of an ecosystem, which were
172 sometimes interpreted as regime shifts (Mumby et al., 2007; de Young et al., 2008;
173 Barnovsky et al. 2012). These may be manifested as simultaneous changes in
174 phytoplankton, dominant consumer species, and trophic structure. Regime shifts often
175 yield major consequences for fisheries and human livelihood (McFarlane et al., 2002;
176 Zhang and Gong, 2005). Examples can be found in the North Pacific - Pacific Decadal
177 Oscillation (Wooster and Zhang, 2004), North Atlantic (Alheit et al., 2014) Caribbean
178 coral reefs (Hughes, 1994), Mediterranean Sea (Conversi et al., 2010), Northern Adriatic
179 Sea (Conversi et al., 2009) and North Sea (Beaugrand, 2004). In the Northern
180 Hemisphere, major ecosystem shifts were observed in the late 1980s to early 1990s, with

181 synchronous shifts following an overall period of high variability. These ecosystem
182 regime shifts were linked to changes in global-scale climate indices (Chavez et al., 2011;
183 Conversi et al., 2010; Möllmann et al., 2011). It is cautioned that these abrupt changes
184 are sometimes difficult to distinguish from random fluctuations or overfishing effects and
185 their true nature often remains unclear (e.g., Hsieh et al., 2005).

186

187 Margins may also be subject to basin-specific and regional influences. For example, the
188 California Cooperative Oceanic Fisheries Investigations (CalCOFI), one of the longest
189 existing oceanographic time series (63 y), reveals major natural variations in water
190 column temperature, oxygen, pH and current strength that are associated with changes in
191 the regional hydrography of the California Current Ecosystem over multiple time and
192 space scales (Checkley and Barth, 2009; McClatchie et al., 2010; Nam et al., 2011; Send
193 and Nam 2012). In addition to the Pacific Decadal Oscillation, there are decadal scale
194 ENSO cycles, seasonal and week-long upwelling events that alter productivity and/or
195 ocean biogeochemistry with marked shifts in oxygen and pH (Fig 2). Variability in these
196 environmental parameters results in changes in the regional biodiversity and ecosystem
197 structure with significant impacts on ecosystem services we depend on (Doney et al.
198 2012).

199

200 Much natural climate variability and some manifestations of climate change occur
201 abruptly over short time and space scales. Most margins experience episodic, extreme
202 events that shape their ecosystems, often through biogeochemical modification. For
203 example the Rhone River carries 80% of its solid flux during 5% of the time (Antonelli et
204 al., 2007) with large biogeochemical consequences (Cathalot et al., 2010). Extreme storm
205 events can reshape coastal systems with short-lived, dramatic changes in salinity and
206 flushing rates and through more persistent alterations of channel openings (Paerl et al.,
207 2001). Atmospheric deposition of nutrients associated with air mass outflow from the
208 Indo-Gangetic Plain to the northern Bay of Bengal is significant and most can occur over
209 4 months in association with the NE monsoon, highlighting the temporal nature of these
210 atmospheric drivers (Srinivas et al., this volume). These are likely to have direct
211 consequences for eutrophication in Bay of Bengal surface waters triggering profuse algal
212 blooms in the adjacent Sundarban wetland (Naha Biswas et al., 2013). Heat waves that
213 last for a few weeks can induce mass mortality in coastal ecosystems of the
214 Mediterranean Sea either directly (Garrabou et al., 2009; Marba and Duarte, 2010) or
215 through the spread of disease and invasive species (Lejeusne et al., 2010).

216

217 As with short time scales, small areas of the ocean can play key roles in global
218 biogeochemical fluxes on margins. For example, 1% of the ocean's water volume
219 accounts for 50% of N removal through water column denitrification and annamox in
220 oxygen deficient zones (Deutsch et al., 2011), and 60-70% of the annual denitrification
221 rate occurs in shelf sediments (Codispoti, 2007). Submarine canyons carry 80-90% of the
222 sediment and organic matter fluxes to the open seafloor sediments, with transport
223 affected by climate-driven stratification, wind regime and winter cooling (Canals et al.,
224 2006; Rabouille et al., 2013). It is proposed that the Congo River, with the second largest
225 discharge in the world, carries 50% of the river's silica through an 800-km long
226 submarine canyon to a 3000 km² deep-sea fan (Raimonet et al., this volume). The

227 functions of submarine canyons as key removal agents via deep-sea fans and deltas may
228 be diminished by climate change-driven reduction of dense water formation with less
229 cascading and deep export (Herrmann et al., 2008; Rabouille et al., 2013).

230

231 An important question to emerge is whether natural variability associated with exposure
232 to stressful conditions (such as hypoxia or hypercapnia) confers evolutionary pre-
233 adaptation to further stress from climate change or direct human activities. Evidence
234 suggests that animals in margin settings subject to upwelled, low-pH waters are resilient
235 to such conditions (e.g., Thomsen et al., 2010; Yu et al., 2011; Hoffmann et al., 2014). In
236 other coastal regions where hydrographic variability is also intense and there are multiple
237 controls from land, detection of trends, sources and biological responses including
238 adaptation can be difficult (Duarte et al., 2013). Whether hydrographic stressors that vary
239 naturally (oxygen, acidification and warming) elicit more adaptation than ‘unnatural’
240 (man made) trace organic or metal/metalloid contaminants, remains an open question.

241

242 *CO₂-driven climate drivers*

243 Rising CO₂ in the atmosphere is reshaping margin ecosystems by increasing sea level,
244 ocean warming, ocean acidification and ocean deoxygenation (Doney et al. 2012). There
245 are also climate shifts that alter patterns of heat, drought, precipitation, and flooding that
246 modify margins directly and indirectly through changes in land use, runoff, and human
247 activities.

248

249 CO₂-induced warming and enhanced stratification have been linked to declining oxygen
250 concentrations on the southern California shelf and upper slope (Bograd et al., 2008) as
251 well as increased seasonal hypoxia on the inner Oregon shelf (Chan et al., 2008). These
252 changes also involve lowered pH and high pCO₂ (Frieder et al. 2012; Alin et al. 2012),
253 with consequences for biogeochemical cycling and ecosystem structure in the California
254 Current (CC) system (Doney et al. 2012). Upwelling is intensifying and low pH (which
255 promotes aragonite undersaturation) is spreading in the northeast Pacific (Feely et al.
256 2008; Gruber et al., 2012). The observed low pH conditions in the CC system are shaping
257 characteristics of this ecosystem by affecting calcifying species and have resulted in the
258 decline of cultured bivalves (Barton et al., 2012). Whether the oxygen and pH changes
259 reflect a continuous, secular trend resulting from CO₂-driven climate changes or are part
260 of a larger (50 y) natural cycle remains controversial (McClatchie et al. 2010; Deutsch et
261 al., 2011). These changes are occurring in an ecosystem already subject to high natural
262 variability (Fig. 2). Clear understanding of this complexity is needed for forecasting
263 future conditions.

264

265 Beyond upwelling regions, perhaps the greatest manifestations of climate change are
266 found on the shelves of the Arctic Ocean. Among the most massive of inputs, a full 10%
267 of the freshwater reaching the oceans occurs in the Arctic, which has only 4 million
268 people living there. Thawing of permafrost due to warming yields increased inputs of soil
269 organic carbon and methane to the coastal ocean and atmosphere (Schurr, 2013), and will
270 influence many aspects of the Arctic coastal ecosystem (Whiteman et al., 2013). The
271 freshwater from melting sea ice combined with degradation of released organic matter is
272 causing major perturbation of low pH in the Arctic. Baseline monitoring of the W. Arctic

273 Ocean reveals that 20% of the Canada Basin surface waters exhibit aragonite
274 undersaturation (Robbins et al., 2013).

275
276 As temperatures continue to increase (Behrenfeld et al., 2006), warming is expected to
277 reduce productivity over much of the ocean (Mora et al., 2013). It is uncertain whether
278 lowered production will reduce oxygen depletion in midwater (from decomposition of
279 sinking phytoplankton and respiration of vertical migrators), counteracting the
280 deoxygenation effects of global warming (from increased stratification and reduced
281 mixing). Alternatively, intensified upwelling in a warmer world may pump more
282 nutrients into surface waters, increase respiration of microbes and other organisms, and
283 increase the rate of deoxygenation. As a direct effect or through changes in currents such
284 as the Gulf Stream, warming might increase methane emissions via dissociation of gas
285 hydrates on continental margins (Phrampus and Hornbach, 2012). Massive gas hydrate
286 deposits in the shallow Arctic Ocean are particularly susceptible and their release may
287 exacerbate acidification and oxygen depletion via aerobic methane oxidation in the water
288 column (Biajoch et al., 2011). There has yet to be exploration of modern biological
289 responses to long-term increases in methane fluxes on margins, although the geologic
290 past may hold lessons in this regard (Kennett et al., 2003).

291

292 *Direct Human Drivers*

293 Rivers are a primary conduit of nutrient loading to the shelf from terrestrial sources of
294 nutrients. Since the development in the early 20th century of the Haber-Bosch process
295 for fixing nitrogen for use in fertilizers, the global nitrogen cycle has become increasingly
296 affected by anthropogenic inputs. The net anthropogenic nitrogen inputs (NANI) to a
297 region include fertilizer application, atmospheric deposition, agricultural N fixation by
298 leguminous crops, and the nitrogen associated with food and livestock feed crossing
299 regional boundaries. Nitrogen flux in rivers is often highly correlated to the NANI of
300 their drainage basins (e.g., Howarth et al., 1996; Han and Allen, 2008; Swaney et al.,
301 2012). In areas of high population densities (e.g., coastal cities) or regions of industrial-
302 scale livestock production, as is increasingly seen in India and China, the nitrogen
303 associated with the trade of food and feed commodities may be very significant. In areas
304 of high crop production, synthetic N fertilizer is typically the dominant source of N (e.g.,
305 Yan et al., 2010). In India, use of synthetic fertilizer has grown exponentially over the
306 last fifty years, making Indian agriculture one of the most intense consumers of fertilizer
307 in the world (Swaney et al., this issue). Between 1970 and 2000, the coastal Bay of
308 Bengal has experienced massive N and P loading (50% and 35% increase, respectively)
309 causing eutrophication; 70-80% of the loading is from agricultural sources (Sattar et al.,
310 2014).

311

312 Margins play a key role in filtering nutrients and contaminants that enter the ocean via
313 runoff and rivers. Productive estuarine ecosystems, particularly wetlands, are able to
314 remove nutrients by denitrification, uptake by vascular plant, phytoplankton, and
315 microbes, by promoting flocculation and enhancing deposition and burial (Kennedy
316 1984, Howarth et al., 2006; Dähnke et al., 2008; Lassaletta et al., 2011; Howarth et al.,
317 2012). Intensive filter feeding by bivalves such as oysters and mussels can also remove
318 particulate nutrients and control eutrophication (Cloern et al. 1982; Dame 2012). The

319 filtering functions of margins have been greatly affected by massive wetland loss over the
320 past century, due largely to changing land use and sea level rise. Globally, overfishing (of
321 oysters) and species introductions (of invasive bivalves) have also had a major influence
322 on water filtration functions (Dame 2011).

323

324 Human acceleration of nutrient cycles and eutrophication are among the best studied of
325 the anthropogenic forcing factors and cause the most conspicuous adverse effects upon
326 continental margins as witnessed by diverse case studies (Table 1, Fig. 1). Intensified
327 nitrogen loading is widespread in coastal ecosystems receiving effluents from catchments
328 with dense human populations (Rabalais, 2004; Glavovic et al., submitted). This yields
329 continental margin dead zones (coastal hypoxic areas resulting from eutrophication),
330 which number over 475 and are on the rise (Diaz and Rosenberg, 2008; World Resources
331 Institute, 2013). The largest of these occur in the Baltic Sea, the Black Sea, the northern
332 Gulf of Mexico and the East China Sea (Rabouille et al., 2008; Zhu et al., 2011), where
333 historical hypoxia induced by natural climate conditions and circulation has been
334 exacerbated by human nutrient input (Zillén et al. 2008; Rabalais et al., 2010; K.-K. Liu
335 et al., this issue). The relative importance of natural and human (nutrient) drivers and
336 efficacy of nutrient legislation has been under debate in recent years (e.g., Bianchi et al.,
337 2008). For example, shrinking of hypoxic areas in the Black Sea appears to have resulted
338 from reductions in human agricultural nutrient inputs, though the extent of the human
339 impacts on this ecosystem is still not clear (Mee et al., 2005). In addition, as indicated
340 above, the balance between N, P and Si is being modified by many factors that affect
341 coastal production, both qualitatively and quantitatively (Ragueneau et al., 2005). CO₂-
342 driven changes in warming, winds, upwelling, and precipitation will inevitably influence
343 both the intensity and areal cover of hypoxia in many dead zones (Rabalais et al., 2009,
344 2014; Giani et al., 2012).

345

346 **3. Comparisons across continental margins**

347

348 To gain a broader sense of how shelf systems are responding to climatic forcing and
349 direct human activities we have compared multiple physical and biogeochemical
350 observations collected at geographically contrasting locations (Fig. 3, Table 2).
351 Consistent with the global warming trend, three out of five margins (the Cariaco Basin,
352 East China Sea and North Sea) have shown increasing temperatures over the last four
353 decades (slope = X °C yr⁻¹; $p < 0.01$), except for San Francisco Bay (slope = -0.029 °C
354 yr⁻¹; $p = 0.07$) where a cooling trend has been detected (For more detail see Supplement
355 A). The cooling trend observed in San Francisco Bay is attributed to an increase in
356 upwelling intensity across the entire California Current system resulting from increasing
357 northerly wind stress along the western coast of the US (Chavez et al., 2011). The
358 warming trend in the Cariaco Basin, in turn, is the result of the weakening of the Trade
359 Winds, and thus of upwelling intensity, along the southern Caribbean Sea (Astor et al.,
360 2013; Taylor et al., 2012).

361

362 No significant trends in sea surface salinity (SSS) are observed at the Cariaco Basin or
363 the North Sea. SSS in the Bohai Sea, however, shows a positive trend (0.0632 yr⁻¹) (Fig.

364 3b). The increasing salinity of the Bohai Sea is thought to be caused by decreasing
365 freshwater discharge from the Yellow River (Fig. 3d).

366

367 The load of dissolved inorganic nitrogen (DIN) in Changjiang (aka the Yangtze River),
368 which empties into the East China Sea, has increased by over two-fold (Liu et al., 2014)
369 between 1970 and 2002 (Fig. 3c), while this river's freshwater discharge has only
370 increased slightly (Fig. 3d). This suggests that rising DIN concentrations in the
371 Changjiang River is mainly due to the intensive use of chemical fertilizer (Yan et al.,
372 2010). By contrast, and due to EU policy change, the DIN load discharged to the North
373 Sea has decreased by 50% since 1977. Dissolved inorganic phosphorus (DIP) shows a
374 similar decreasing trend at this location (Pätsch and Lenhart, 2011)

375

376 The sea surface chlorophyll-*a* concentrations in SF Bay have increased in the last two
377 decades (Fig. 3e), which is consistent with the observed decreasing trend in SST .
378 Simultaneously, however, chlorophyll-*a* in the Cariaco Basin shows a decreasing trend
379 since the late 90's due to weaker upwelling events and stronger thermal stratification
380 (Taylor et al., 2012). The monthly mean sea surface chlorophyll-*a* in the East China Sea
381 derived from ocean color products by NASA's Sea-viewing Wide Field-of-view Sensor
382 (SeaWiFS) also exhibits a significant increasing trend since 1998, which is thought to
383 result from increased DIN loads from the Changjiang River (Fig. 3c) (K.-K. Liu et al.,
384 this issue).

385

386 In response to increasing phytoplankton growth, bottom water oxygen saturation in SF
387 Bay and the East China Sea has shown a significant decline (Fig. 3f). Oxygen saturation
388 shows a weak decline in the upper 3 m at the CARIACO Station (Fig 3f), probably due to
389 warming and possibly to lower oxygen production by phytoplankton (Fig 3e).

390

391 Different margins show markedly different responses to local stressors and to global-
392 scale change. Variations in availability and temporal coverage of different environmental
393 parameters highlight the need for comprehensive and sustained time-series observations
394 on continental margins. These are required in order to understand ecosystem responses
395 to natural, CO₂ climate-driven and direct human perturbations.

396

397 **4. Ecosystem responses to biogeochemical change on continental margins**

398

399 Some of the most apparent environmental and ecosystem consequences due to the
400 common climate and human stressors discussed in this special issue are summarized in
401 Table 1 and discussed below.

402

403 Human alteration of hydrological processes such as damming and water diversion (B3,
404 B4), drives very noticeable physical changes in margins causing loss of habitats due to
405 coastal erosion or reduced river discharge (e.g., S.M. Liu, this issue). When combined
406 with climate effects, resulting salinity increases can lead to species invasions that reshape
407 coastal ecosystems. Following massive water diversion and drought in San Francisco
408 Bay, an invasion by Asian clams altered the timing and magnitude of phytoplankton
409 availability, with cascading trophic consequences (Cloern and Jassby, 2012, B3). In the

410 Bay of Brest, introduction of an invasive limpet changed the seasonality of primary
411 production, which in turn has changed benthic biodiversity and completely modified the
412 benthic-pelagic coupling over a 30-year period (Grall and Chauvaud, 2002).

413

414 In a broad sense, land use change alters how rainfall interacts with the landscape. Some
415 land-use activities result in increased soil degradation and erosion (i.e., agriculture,
416 mining), and in eutrophication of rivers and continental margins through the use of
417 fertilizers. The coastal zone's high primary productivity and the abundant filter feeders
418 (e.g., Lotze et al., 2006) offset land-derived nutrient inputs to some extent, but are tested
419 by eutrophication and overfishing (B2). Moreover, rising sea levels will lead to flooding
420 of low-lying coastal regions like India and Bangladesh, movement of seawater farther up
421 estuaries, and intrusion of seawater into groundwater reservoirs.

422

423 Eutrophication is among the most widespread of coastal insults (Table 1; B2, B3, B4, B5,
424 B6), but the outcomes, which include productivity enhancement *and* hypoxia, can be
425 complex. For example, several cross-system comparisons indicate that increased N
426 enhances total landings of fish and mobile shellfish even in systems with hypoxia (Fig.
427 4), although individual species may decline and the overall composition of the catch can
428 be affected (Nixon and Buckley, 2002; Breitburg et al., 2009b). On the downside,
429 hypoxia - an endocrine disrupter in fish that experience chronic exposure (Thomas et al.,
430 2006) - can favor gelatinous plankton and some bivalves (Breitburg et al., 2003), and
431 create and eliminate shallow water refuges for small and juvenile fishes (Breitburg et al.
432 2009a).

433

434 Reversal of eutrophication trends have been observed in some areas such as the Danish
435 straits (Carstensen et al., 2006), the Scheldt Estuary (Soetaert et al., 2006) and other
436 continental European rivers discharging into the North Sea (Emeis et al., this volume),
437 the open Northern Adriatic (Giani et al., 2012), and the NW Black Sea (McQuatter-
438 Gollop et al., 2009). In some instances P reduction has been considered to be a primary
439 driver of these changes. In the areas subject to oligotrophication, overfishing may act
440 synergistically to diminish the trophic chain and reduce seafood resources (B5).
441 However, along the Danish and Finnish coasts, dissolved oxygen in bottom waters
442 continues to drop despite efforts to reduce nutrient discharge (Carstensen et al., 2014). In
443 the Baltic Sea, which hosts nearly 20% of the world's identified coastal hypoxic sites,
444 climate and nutrient drivers interact with regional circulation patterns and wastewater
445 treatment technologies to produce a mosaic of faunal responses (Conley et al., 2011).

446

447 While eutrophication-induced hypoxia is spreading, warming also causes the ocean to
448 lose oxygen due to the synergistic effect of reduced oxygen solubility and enhanced
449 water column stratification (Bopp et al., 2001). This has been termed deoxygenation and
450 contributes to global expansion of oxygen minimum zones (Stramma et al. 2010). Recent
451 model results demonstrate the extreme sensitivity of the volume of suboxic water in the
452 open ocean to changing climate conditions (Deutsch et al., 2011). Biological analyses
453 suggest that equator-ward species boundaries are highly sensitive to changes in ocean
454 temperature and oxygen content. Models predict a decline in metabolic scope of species
455 (energy available for maintenance and reproduction) and functional habitat loss.

456 On margins both eutrophication and intensified upwelling typically increase production
457 while drawing down oxygen and creating hypoxia at deeper water levels. Animal
458 avoidance of hypoxia acts to aggregate species around or above hypoxic zones, and leads
459 to habitat compression, both in estuarine settings and in open-ocean oxygen minimum
460 zones (OMZs). The resulting aggregations are susceptible to overfishing (Craig, 2012,
461 Breitburg et al., 2009b), but high catches may mask the consequences of ecosystem
462 stress, making detection of habitat degradation difficult (Breitburg et al., 2009b). Such
463 conditions also induce shoaling of the zooplankton biomass layer at the thermocline
464 (upper oxycline boundary) and concentration of midwater biomass in a layer at the lower
465 oxycline (Wishner et al., 2013). As oxygen declines and oxygen minima shoal in both
466 the Atlantic and Pacific, large billfish are now found at shallower depths and are
467 increasingly susceptible to overfishing (Prince and Goodyear, 2006; Prince et al., 2010;
468 Stramma et al., 2011).

469
470 Upwelling margins, which host key world fisheries, exhibit strong vertical gradients in
471 temperature, oxygen and pH associated with oxygen minimum zones (Paulmier et al.,
472 2011). Across these gradients bathyal benthic assemblages reveal shifts in diversity, body
473 size, zonation, carbon processing, bioturbation, colonization and resilience (Levin, 2003,
474 Levin et al. 2009; Gilly et al., 2013; Levin et al., 2013). Intensified upwelling is predicted
475 to result in changes in biodiversity and ecosystem functioning associated with the
476 expansion of OMZs (Stramma et al., 2010; Gilly et al., 2013). Recent onset of seasonal
477 hypoxia on the Oregon inner shelf now causes summer die-offs of fish and invertebrates
478 (Grantham et al., 2004). Responses to intensified upwelling winds and increased
479 stratification can also vary regionally. For example, comparative analyses of the
480 California and Canary systems reveal substantial differences in the responses of
481 biological production and air-sea CO₂ fluxes to upwelling intensification in these two
482 systems (Lachkar and Gruber, 2013). These differences have been attributed to various
483 drivers such as the contrasting shelf topography, eddy activity, coastal water residence
484 times and basin-scale forcing in the two regions (Marchesiello and Estrade, 2009;
485 Lachkar and Gruber, 2013). These differences also affect the vulnerability of these
486 ecosystems to global anthropogenic perturbations such as ocean acidification (Lachkar,
487 2014). Other upwelling regions have received less attention and could exhibit additional
488 (or alternative) response mechanisms.

489
490 CO₂-induced climate change is the predominant forcing on the ecosystem of the polar
491 margins (S8). Warming of the Arctic is taking place two to three times faster than global
492 rates (Trenberth et al., 2007); as a result sea-ice cover has been decreasing at a rate of
493 >10% per decade with ice-free summers expected in a few decades. Arctic ecosystems
494 are increasingly being challenged by tipping elements (Duarte et al., 2012; Naam, 2012;
495 Wassmann and Lenton, 2012). In the future Norway may experience decreased primary
496 productivity, while Russia will show increased productivity. Nowhere will adaptation be
497 a more critical element of sustainability than in the Arctic, because the Arctic shelves
498 have inordinate importance in feeding the world population. (See S8 for more details)

499
500 Among the many effects of rising atmospheric CO₂, the significant decrease of ocean pH
501 (ocean acidification) and shift in seawater carbonate chemistry (Doney et al., 2001) may

502 elicit some of the most economically significant responses from margin ecosystems.
503 Acidification alters seawater chemical speciation, most notably the lowering of calcium
504 carbonate saturation states, which impacts shell-forming marine organisms from plankton
505 to benthic molluscs, echinoderms, and corals, all of which are abundant in continental
506 margins. Ocean acidification is exacerbated in the coastal zone by increased land-derived
507 nutrient inputs, which enhance, in turn, productivity of organic matter and therefore
508 respiration and release of CO₂ (e.g., Borges and Gypens, 2010; Cai et al., 2011). A
509 serious drop of aragonite saturation state has occurred in some coastal seas, such as the
510 North Yellow Sea, threatening the aquaculture of shellfish (e.g., Zhai et al., 2014). Coral
511 reef ecosystems, which provide key fisheries, critical shoreline protection and habitats for
512 a large number of species, are highly susceptible (Andersson and Gledhill, 2013). Due to
513 ocean acidification, rates of coral calcification may decrease, whereas rates of bioerosion
514 and carbonate dissolution may increase, resulting in a transition from net accretion to net
515 erosion. Impairment of the calcifying capacity of marine organisms is therefore expected
516 to have negative impacts on coral reefs and other calcifiers (e.g., bivalves) and on the
517 ecosystem services they provide.

518
519 The complexity of ocean biogeochemical-ecosystem interactions on margins means that
520 some drivers will create responses that generate feedback – further altering a system. One
521 example occurs when acidification-induced undersaturation of carbonate minerals
522 adversely affects shell growth and settlement success of bivalves and coral polyps
523 building reefs; this is predicted to ultimately reduce oyster, mussel and clam populations
524 and coral reef building. Locally, the presence of large oyster populations buffers
525 increasing CO₂ and decreasing pH through shell dissolution and alkalinity increase. So
526 lowered pH ultimately reduces local buffering capacity, leading to further reductions in
527 pH when the mineral buffer is exhausted. In addition, mass removal of shellfish (by
528 harvest), could contribute to a deficit in the carbonate balance, as the shells form a
529 dissolution buffer needed by many animals to survive (Waldbusser et al., 2013).
530 Populations may be reduced to the point of unsustainability leading to ‘recruitment
531 overfishing’.

532
533 On some margins high-frequency climate oscillations are the dominant driver of
534 biogeochemical variation and consequently, ecosystem structure. In the Bay of Calvi in
535 the Ligurian Sea of the NW Mediterranean (Goffart et al., this issue) the biogeochemical
536 condition is very oligotrophic during mild winters and mesotrophic during moderate
537 winters (B7). During severe winters, the Bay sustains a “high nutrient - low chlorophyll”
538 situation. With little human disturbances this Bay may serve as the baseline, against
539 which ecosystem changes in the Mediterranean due to direct human impacts can be
540 detected (see B7 for more details). In the East Pacific Ocean, interannual variations
541 linked to ENSO induce low productivity (well oxygenated) El Nino and high productivity
542 (low oxygen) La Nina conditions that affect fisheries production in the Humboldt and
543 California and Benguela current ecosystems (Arntz et al., 2006).

544

545 **5. Using time series to distinguish drivers of change**

546 Hydrographic and ecological time series have provided data critical to evaluating and
547 interpreting change on margins. Koslow and Couture (2013) have referred to ecological

548 time series as the Cinderella (hard working drudges) at the climate change ball. Beyond
549 this they may provide the “Anthropocene’s canary in a coal mine” for many other forms
550 of human disturbance. Below we address the approaches, benefits and limitations of time
551 series in attribution of change on margins.

552

553 Several multi-decadal oceanographic time series measurements from a variety of coastal
554 and pelagic systems have shown how lower and intermediate trophic levels, and
555 biogeochemical cycling react to climate oscillations regionally and globally (Chavez et
556 al., 2003; Black et al., 2011; Church et al., 2013) (Table 1). Multi-decadal time series of
557 phytoplankton have been generated for many regions including San Francisco Bay
558 (Cloern and Jassby, 2013), Chesapeake Bay (Lee et al., 2013), Narragansett Bay
559 (Borkman and Smayda, 2009), the Cariaco Basin (Chavez et al., 2011, Muller-Karger et
560 al., 2013), the North Sea (Wiltshire et al., 2008) and areas of the Mediterranean Sea (e.g.
561 Goffart et al., 2002, Ninčević Gladan et al., 2010, Zingone et al., 2010, Goffart et al.,
562 submitted) including the Northern Adriatic (Bernardi-Aubry et al., 2012, Marić et al.,
563 2012, Mozetič et al., 2012) and Gulf of Naples (Ribera d’Alcalà et al., 2004). There are
564 also Arctic time series in the Bering, Chukchi, and Barents Sea. Most of these reveal
565 oscillations associated with climate variability (Borkman et al., 2009; Harrison et al.,
566 2010). Indeed, such long-term ocean time series have been fundamental for expanding
567 our knowledge about the sensitivity of marine biodiversity, ecosystems and
568 biogeochemistry to environmental change (Church et al., 2013; Koslow and Couture,
569 2013). However, moving forward an international network of time series is needed to
570 evaluate regional linkages and interpret global changes.

571

572 There are some major gaps in time series monitoring. Whereas models of nutrient fluxes
573 from watersheds abound, monitoring data to verify them do not. In the developing world,
574 the scarcity of monitoring data adequate to characterize riverine nutrient flows has
575 impeded our understanding of the relationships with human activities. Research and
576 development of monitoring in these regions, should be made a priority, and would
577 improve our management of coastal waters. It is important to add that not only the N
578 cycle should be monitored, but also changes in nutrient ratios delivered by rivers. In
579 particular the Si:N and Si:P ratios should be closely monitored as potential early warning
580 indicators of disturbances (Billen and Garnier, 2007); indeed, they are often decreasing
581 due to excessive N and P inputs and decreasing Si inputs due to damming (Humborg,
582 1997) and the proliferation of invasive species (Ragueneau et al., 2005) causing
583 replacement of diatoms by dinoflagellates. South East Asia, where anthropogenic factors
584 leading to decreasing Si:N and Si:P ratios combine, should be especially targeted for
585 monitoring (Ragueneau et al., 2006).

586

587 Modern time series gain added value when used in conjunction with paleoceanographic
588 studies (Black et al., 2011) and models (see Church et al., 2013); together these tools
589 allow researchers to discern natural sources of environmental change from variations
590 induced by climate change (warming, extreme flooding from river input or snow melt, or
591 heat waves) and direct human drivers such as eutrophication, damming and fishing
592 (Koslow and Couture, 2013). In some margin settings subject to long bouts of habitation
593 and industrialization (e.g., Chesapeake Bay, the coastal SE North Sea, northern Adriatic),

594 natural variability is a small signal relative to the influence of humans. In others (e.g.,
595 upwelling margins) natural variability produces an exceedingly strong signal and
596 irrefutably detecting CO₂-driven climate change or anthropogenic forcing is difficult.

597

598 Sediment and glacial ice core climate records often provide the long temporal perspective
599 needed to identify climate oscillations prior to high anthropogenic CO₂ (> 280 ppm)
600 conditions or resulting from long-term changes in human population density and land use
601 practices (Cooper and Brush, 1993; Emeis et al., 2000; Yasuhara et al., 2012). Some of
602 these geological climate records are now complemented by oceanographic and
603 biogeochemical time series observations, thus providing valuable insights into the effects
604 of anthropogenic perturbations on the marine environment (see Black et al., 2011).

605

606 *Time Series and Coastal Management.* The motivation underlying the establishment of
607 ecological time series varies. In California, CalCOFI was developed in the 1950s to
608 understand the boom and bust cycles of the sardine (Bograd et al., 2003; Chavez et al.,
609 2003). At its inception, the CalCOFI concept of monitoring the entire ecosystem, now
610 widely accepted in the context of ecosystem-based management, was visionary and
611 somewhat heretical. In Chesapeake Bay, a long time series of young-of-year fish
612 abundances in Maryland waters was initiated in 1954 by the state fisheries agency to aid
613 management of several anadromous species (Durrell and Weedon, 2011), a time series of
614 jellyfish abundances was begun in 1960 by a University of Maryland researcher in
615 response to the ‘Jellyfish Nuisance Act’ (Cargo and King, 1990), and bay-wide time
616 series monitoring of water quality parameters was begun in the 1980s with funding from
617 the States of Maryland and Virginia and the US EPA to aid management efforts to
618 improve water quality conditions (Boesch et al., 2001). Governments of states bordering
619 the western coasts of Europe (OSPAR) and the Baltic Sea (HELCOM) initiated
620 monitoring programs in the 1970’s to protect the marine environment from all sources of
621 pollution through intergovernmental cooperation.

622

623 Although each time series is fixed in space and provides local information, when data are
624 combined across time series they can provide a powerful synoptic understanding of the
625 link between climate variability and ocean biogeochemistry (Church et al, 2013). The
626 ICES Phytoplankton and Microbial Plankton Status Report 2009/2010 exemplifies this
627 for the North Atlantic (O’Brien et al., 2012). Records of sea ice cover and tipping points
628 in the Arctic provide another example (Carstensen and Weydmann, 2012). Under optimal
629 conditions, time series provide data prior to catastrophe (e.g., fishery collapse) so that
630 causes can be discerned. It is important, however, to recognize the value of understanding
631 regional differences and their forcing mechanisms.

632 *Time series constraints.* Spatially fixed time series may have limitations. Single-location
633 measurements typically do not reveal spatial expansions, contractions or oscillations.
634 They cannot recognize change due to relocation of organisms or features, making it
635 difficult in some cases to untangle spatial and temporal change, although spatial
636 comparisons can sometimes be used as proxies of temporal change (e.g., Wishner et al.,
637 2013). Satellite remote sensing has typically been the tool of choice for extrapolating
638 fixed time series observations to broader spatial and temporal scales. Some time-series
639 stations (i.e., Hawaiian Ocean Time-series [HOT] and the Bermuda Atlantic Time Series

640 [BATS]) have been successful at using autonomous samplers and sensors (e.g., gliders,
641 drifting profilers) for learning the regional significance of the measurements they collect.
642 These are especially important for hard-to-reach areas like the Arctic and Antarctic, and
643 can expand coverage for traditional time series in other regions. Autonomous sampling
644 platforms, however, are expensive to operate and thus are out of reach for time-series
645 programs with limited resources. Another approach is to engage platforms of opportunity.
646 Industry with a presence on the ocean margins, such as offshore wind and aquaculture,
647 fishing, fossil fuel extraction and minerals, may have a role to play in time series
648 development in the future, and should be broadly engaged to support monitoring efforts.
649 Expansion of programs like the World Ocean Council Smart Data/Smart Industries
650 (http://www.oceancouncil.org/site/smart_ocean.php) may be useful.

651 Time, funding and facilities constraints often limit time series to the upper water column
652 and basic hydrographic parameters. To link these to key resource needs and sustainable
653 management – including aquaculture, fisheries, energy and minerals – it will be necessary
654 to incorporate the sea floor and its organisms into time-series monitoring. Benthos
655 monitoring can also provide critical information about biogeochemical feedbacks from
656 the sea floor, processes often not included in large-scale climate or ecosystem models.

657
658 Most margin time series are not of sufficient duration to detect variation outside normal
659 statistical variability (especially given decadal-scale cycles emerging in the atmosphere-
660 surface ocean system). Often shifts and change are misattributed due to lack of
661 knowledge about natural variability and its sources. Paleoceanographic records in ice
662 cores, sediment cores or tree rings allow us to extend understanding of margin processes
663 back in time, prior to the establishment of *in situ* observations (Gooday et al. 2009).
664 Innovative analyses of scales, teeth, otoliths and ichnofacies may allow use of such
665 records to reconstruct complex changes in exposure histories and food web dynamics
666 (Gooday et al., 2009; Morat et al., 2014). Recent development of geochemical proxies for
667 detection of fish exposure to hypoxia offers the promise of identifying past and present
668 trends in oxygen concentration using otoliths (Limburg et al., 2011; 2014 [this volume]);
669 fish scales and other skeletal elements may also prove useful, but require testing.
670 Chronosequences from long-lived calcifying organisms (e.g., coldwater corals or
671 bivalves) may provide excellent time series of temperature or pH as a basis for
672 chronometric analyses similar to dendrochronology (e.g., Chauvaud et al., 2005, 2012;
673 Black et al., 2008). Novel functional gene microarrays may be used to evaluate the
674 diversity and composition of the denitrifying microbial community in hypoxic settings
675 like OMZs, allowing us to better understand how microbial metabolism can impact the
676 global climate through the production of N₂O, a bi-product of denitrification and a
677 powerful greenhouse gas (Jayakumar et al., 2013). We need to further develop proxy
678 variables for environmental reconstructions, expand the data bases for regional-scale
679 hindcasts, and strive to detect and interpret interannual variability from low resolution
680 archives.

681
682 While valuable in having fixed measurements, time series may also need to have an
683 adaptive observation component focused on (a) identifying changes and their underlying
684 causes and (b) monitoring parameters tied to the ecosystem services we care about, in
685 order to directly address policy concerns for effective management of these services.

686 There is often a gap between what we can realistically measure, and what we want to
687 achieve. This highlights the importance of research on basic processes and mechanisms
688 that will identify indicators of change and incorporate process-based knowledge into our
689 models. Often the early warning of degradation will come from land. For example,
690 agricultural inputs, nutrient concentrations, N:P, Si:P, Si:N ratios or multiple nutrient
691 concentration data together could be key indicators (Billen and Garnier, 2007).

692 The burgeoning number of time series and rising volumes of data highlight a need to
693 engage more scientists in the analysis phase of time series research. There is also a
694 challenge of maintaining continuity in the face of severe funding shortages; scientists
695 must advocate for continuation of monitoring programs. Key to the successful
696 continuation of time series is concise presentation of insight gained and raising the
697 awareness of the public and policy makers of their value. It is here that social scientists
698 can help natural scientists learn how to make a strong case without loss of integrity and
699 accountability.

700

701 **6. The continental margin in the Anthropocene: the convergence of** 702 **biogeochemistry, ecosystems and society**

703

704 To be effective, economic models must be able to weigh the costs of the unwanted
705 impacts of stressors and forcings on ecosystem services discussed above and associated
706 consequences against gains for society, and to attribute change to specific and
707 controllable drivers. To date scientists have failed to convey the message of the
708 overriding importance and societal consequences of CO₂ emissions (as well as other
709 greenhouse gases) in the context of global warming. Investment in collaboration efforts
710 between social and natural scientists, development of outreach and public communication
711 skills, and advocacy about the importance of individual actions are required to manage
712 margins effectively (Pidgeon and Fischhoff, 2011). A need to combine scientific with
713 traditional and local ecological knowledge, especially in settings such as the Arctic,
714 further argues for key social-natural science integration.

715

716 There is growing need for modeling of coupled human (social)-biogeochemical systems
717 on margins. A pioneering example can be found for the surfclam fishery on the Middle
718 Atlantic Bight (MAB) continental shelf (McCay et al., 2011). This million-dollar fishery
719 has been managed since the 1990s with transferable quotas, one of the first in the US to
720 do so. In recent decades the population has shifted to the north and overall abundance has
721 declined (Weinberg, 2005). Simulations of surfclam growth that use 50-year hindcasts of
722 bottom temperature obtained from an implementation of the Regional Ocean Modeling
723 System for the MAB (Kang and Curchister, 2013) show that episodic warming events
724 increase surfclam mortality and limit animal size in the southern portion of its range
725 (Narváez et al, this issue). The resulting northward movement of the stock has negative
726 economic consequences for the fishing fleet and processing plants. These studies point to
727 a key role for natural scientists in assessment of the socio-economic consequences of
728 climate change (McCay et al., 2011).

729

730 *Disasters can sometimes serve as catalysts for action.* In the United States, several
731 disaster events have caught the attention of the scientific community and policy makers,

732 and have resulted in the development of large-scale monitoring efforts that seek to
733 mechanistically understand these events and mitigate their impacts by improving our
734 predicting capabilities. One example is the unexpected failure of the oyster aquaculture
735 fishery on the US Pacific coast due to upwelling of carbonate-undersaturated waters
736 (Barton et al., 2012); this stimulated state-sponsored research programs on ocean
737 acidification (Adelsman and Binder, 2012). Highly destructive Superstorm Sandy
738 stimulated sea level rise preparedness, and massive fish kills resulting from hurricane-
739 induced release of hog waste and sewage in North Carolina (Malin et al., 1999) have
740 engendered public support for altered agricultural practices and backup treatment plants.
741 These types of events are not one-time occurrences, but are likely to become more
742 frequent. For example, low bottom-water aragonite saturation values on shelves are
743 expected to have negative effects on shellfish in the Yellow Sea (Zhai et al., 2014) and
744 off California (Gruber et al., 2012). The costs required for building community, industry
745 and ecosystem resilience are now being weighed against the massive costs of disaster
746 damage and disaster relief.

747
748 While continued and expanded time series measurements are essential to monitor status
749 and trends, scientists often know enough to make strong policy recommendations
750 regarding CO₂, nutrients and human activities. In many instances adaptation to change is
751 required and distinguishing among drivers is not essential for policy decisions. Whether
752 reduced sediment inputs to the coastal zone result from damming or from drought, it will
753 still be necessary to prepare for climate-related sea level rise and associated consequences
754 of limited land-building and net loss of coastal wetlands. For anadromous fishes, loss of
755 freshwater inputs from damming versus drought will have similar consequences, as will
756 loss of river-sea connectivity from eutrophication induced hypoxia versus upwelling-
757 induced deoxygenation. For coastal shellfish, the corrosive effects of acidification may
758 result from atmospheric CO₂ inputs, intensified upwelling, increased stratification,
759 anthropogenic nutrient loads, precipitation or sea ice melting. Acting to reduce CO₂
760 emissions and limit the now-inevitable rise in ocean temperatures, acidification, and
761 deoxygenation is critical.

762
763 There is growing consensus that direct anthropogenic stressors such as overexploitation
764 of natural resources (fisheries, mining), habitat destruction, land use/cover change,
765 alteration of river catchments, coastal construction, damming, species invasion and
766 pollution will lower the resilience of populations, species and ecosystems and make them
767 less able to cope with climate-induced stress (Bijma et al., 2013). For example, reducing
768 fishing mortality in exploited populations can also reduce total mortality and be
769 protective of declining populations, even where part of that total mortality was due to
770 hypoxia, disease or habitat degradation (Breitburg et al., 2009b). Thus policy, law and
771 management of margins must consider and address climate and direct human stressors
772 together. Relevant lessons can be drawn from regional, time series and case studies where
773 different combinations of stressors interact and their trends have been tracked over time.
774 Understanding system connectivities, seeking indicators of regime change, and
775 promoting adaptation-oriented policy to build functional resilience, are lessons from the
776 Arctic (Carmack et al., 2012) that apply well to most margin ecosystems. The recently
777 released IPCC AR5 report emphasizes the overwhelming need for societal adaptation to

778 multiple stressors associated with climate change, especially in countries where poverty
779 will exacerbate the consequences (Field et al. IPCC 2014).

780

781 The recognition of the importance of natural capital and ecosystem services to national
782 wealth has come slowly to some nations. In the USA this is now evidenced by the
783 generation of a National Ocean Policy (2004; [http://ioc-](http://ioc-unesco.org/images/stories/LawoftheSea/Documents/NationalOceanPolicy/nop.usa.pdf)
784 [unesco.org/images/stories/LawoftheSea/Documents/NationalOceanPolicy/nop.usa.pdf](http://ioc-unesco.org/images/stories/LawoftheSea/Documents/NationalOceanPolicy/nop.usa.pdf)).

785 To a large extent this policy addresses the continental margins, where most of the key
786 services and commercial resources are provided. The Marine Strategy Framework
787 Directive (Directive 2008/56/EC,

788 <http://ec.europa.eu/environment/water/marine/ges.htm>), adopted by the European
789 Commission in 2008, marks an important milestone in the development of the EU's
790 marine environmental policy and is the first framework instrument aimed expressly at
791 protecting and preserving the marine environment with a holistic approach. In 2012 The
792 EU launched its Blue Growth initiative

793 (http://ec.europa.eu/maritimeaffairs/policy/blue_growth/) that addresses three crucial
794 components of sustainable development of marine resources: gathering and channeling
795 marine knowledge to improve access to information about the continental margins of
796 Europe, maritime spatial planning to aid management of offshore resources, and
797 integrated maritime surveillance. Other nations have national ocean policies in review
798 (e.g., South Africa) or in early stages of formulation (Namibia).

799

800 *Margin management strategies must move from mono- to multiple stressor*
801 *considerations.* Most policies and research programs address only one or two factors –
802 nutrients, oxygen, ocean acidification, fishing pressure, disease, or invasive species, (e.g.,
803 Crain et al., 2008). We know that T, O₂ and CO₂ are changing simultaneously and
804 interacting (Bijma et al., 2013). There is need for scientific consensus on a) what the
805 multistressor questions are and b) how to approach the issues. How to integrate
806 laboratory studies, field observations, monitoring, modeling, and use of proxies to
807 address these questions remains a major challenge.

808

809 To incorporate natural variability and climate change into our decision making and
810 management activities *we need research that identifies, quantifies and confronts*
811 *management tradeoffs.* Stakeholder identification and finding equitable solutions is
812 critical as every decision has winners and losers. We must quantify the economic costs of
813 nutrient reduction for agriculture, fishers, and ecosystem services. Margin researchers
814 have only just begun to tackle the larger question of valuing ecosystem services and
815 biodiversity on the continental slope beyond the shelf –this is especially critical in deep
816 waters where resource extraction activities (energy, minerals and deep-water fishing) are
817 on the rise (Levin and Dayton, 2009; Jobstvogt et al., 2013).

818

819 As both top predators and guardians of the planet we face immense ocean policy
820 challenges over the next 10-50 years. The mentality of many nations is of a land-based
821 society. Managing a fluid -connected environment is fundamentally different than
822 managing land use where discrete boundaries between impacted and more pristine areas
823 can be maintained. There is a spatial disconnect between farm policies and their effects

824 on the coastal ocean and our margins. Even international climate negotiations
825 (Convergence of Parties) involve remarkably little consideration of ocean processes,
826 despite the large role the ocean plays in regulating climate. The concepts underlying
827 sustainability in ocean margins must involve an educational thrust that starts early, as
828 well as strategies to communicate at national and international levels. *We need to better*
829 *understand the process by which science is introduced to policy, and target and fast track*
830 *scientific approaches that meet those needs.* End-to-end efforts are needed that first bring
831 together the natural and human component of socio-ecosystems; and then work with
832 stakeholders and policy makers towards finding and implementing solutions.

833

834 This article focuses on the impacts of the 20th century; we recognize that the 21st century
835 may involve a different suite of primary stressors, some of which are as yet unknown.
836 The continental margins of the future will undergo further changes as the system is
837 continuously perturbed. As human populations grow, needs for fresh water, energy,
838 space, and food will create new demands of the coastal ocean including coastal
839 aquaculture, wind farms, wave energy stations and desalinization plants, intensified
840 shipping activities, and seabed mineral exploration.

841

842 Holistic consideration of margins facing the confluence of human, climate and natural
843 stressors highlights the need to integrate science with societal needs. Building on work of
844 others (e.g., Islam and Tanaka, 2004), we identify the need to:

- 845 • Formulate a clear understanding of the environmental, ecological and economic
846 value of margin ecosystems and how these vary under different climate regimes.
- 847 • Enact water quality management that recognizes land-ocean-atmosphere
848 exchanges controlled by climate and humans.
- 849 • Enact comprehensive monitoring to link policy-based changes in drivers to
850 ecosystem responses
- 851 • Improve cooperation of stakeholders, regulators, scientists and civil society
- 852 • Scale and coordinate local, regional, national and international activities to
853 maximize knowledge and promote modeling efforts
- 854 • Protect key services via ecosystem-based management
- 855 • Develop mechanisms to translate scientific knowledge into regulation and
856 legislation, and the political realities needed to achieve action.

857

858 Ultimately, we will need to set priorities, accept tradeoffs and motivate creative solutions.
859 These goals are very much in line with the approach of the Future Earth Initiative to meet
860 the grand challenge of global sustainability (Reid et al., 2010). Interactions among social
861 and natural scientists are nascent, but a growing number of national and international
862 programs recognize their importance. Achieving sufficient energy, water, food and
863 healthy margin ecosystems is a tall order, but a challenge that natural and social scientists
864 must work together to meet head on.

865

866 **6. Acknowledgements**

867

868 We are grateful for the hospitality of the National Institute of Oceanography, Goa, India,
869 which hosted the IMBIZO III. Support for the workshop was provided by IMBER (a core

870 project of IGBP), SCOR and a host of co-sponsors. We thank two anonymous reviewers
871 for their comments which have improved the manuscript. L. Levin's and D. Breitburg's
872 attendance was supported by the US Ocean Carbon Biogeochemistry program and
873 IMBER. Levin's research support is from NOAA - California Sea Grant College
874 Program Project # R/CC-04 and NSF OCE 0927445 and 1041062. Support for E.
875 Hofmann was provided by NSF grant GEO-0908939. Support for K.-K. Liu was
876 provided by the Ministry of Science and Technology, Taiwan, and SCOR.

877

878 7. References

879

- 880 Adelsman, H., and Binder, L. W. (Eds.) (2012) *Washington State Blue Ribbon Panel on*
881 *Ocean Acidification (2012): Ocean Acidification: From Knowledge to Action,*
882 *Washington State's Strategic Response*, Washington Department of Ecology,
883 Publication no. 12-01-015, Olympia, Washington.
- 884 Alheit, J. Licandro, P., Coombs, S., Garcia, A., Giráldez, A., Santamaría, M., Slotte, A.,
885 Tsikliras, A. Atlantic Multidecadal Oscillation (AMO) modulates dynamics of
886 small pelagic fishes and ecosystem regime shifts in the eastern North and Central
887 Atlantic, *Journal of Marine Systems*, Volume 131, March 2014, Pages 21-35,
888 ISSN 0924-7963, <http://dx.doi.org/10.1016/j.jmarsys.2013.11.002>.
- 889 Alin, S. R., Feely, R. A., Dickson, A. G., Hernandez-Ayon, J. M., Juranek, L. W., Ohman,
890 M. D., and Goericke, R. (2012) Robust empirical relationships for estimating the
891 carbonate system in the southern California Current System and application to
892 CalCOFI hydrographic cruise data (2005-2011), *Journal of Geophysical Research*
893 117, C05033.
- 894 Andersson, A.J., Gledhill, D., (2013). Ocean acidification and coral reefs: Effects on
895 breakdown, dissolution, and net ecosystem calcification. *Annual Review of*
896 *Marine Science*, 5: 321-348
- 897 Antia, Avan, Maaßen, J., Herman, P., Voß, M., Scholten, Jan, Groom, S. and Miller,
898 P. (2001) Spatial and Qualitative Patterns of Particle Flux at the European
899 Continental Margin: The OMEX I Project *Deep-Sea Research Part II-Topical*
900 *Studies in Oceanography*, 48 (14-15). pp. 3083-3106.
- 901 Arntz, W.E., V.A. Gallardo, D. Guteierrez, E. Isla, L.A. Levin, J. Mendo, C. Neira, G.
902 Rowe, J. Tarazona and M. Wolff. (2006) ENSO and similar perturbation effects
903 on the benthos of the Humboldt, California and Benguela Current upwelling
904 ecosystems. *Advances in Geosciences* 6, 243-265.
- 905 Astor, Y.M., Lorenzoni, L., Thunell, R., Varela, R., Muller-Karger, F., Troccoli, L.,
906 Taylor, G.T., Scranton, M.I., Tappa, E., Rueda, D., (2013) Interannual variability
907 in sea surface temperature and fCO₂ changes in the Cariaco Basin. *Deep-Sea Res.*
908 II, doi:10.1016/j.dsr2.2013.01.002.
- 909 Banner, F.T., Collins, M.E., Massey, M.S. (Eds) (1980) *The North-West European Shelf*
910 *Seas: The Sea Bed and the Sea in Motion II. Physical and Chemical*
911 *Oceanography, and Physical Resources. Elsevier Oceanography Series, Volume*
912 *24, Part B, 301-638.*
- 913 Barnosky, A.D., Hadly, E.A., Bascompte, J., Berlow, E.L., Brown, J.H., Fortelius, M.,
914 Getz, W.M., Harte, J., Hastings, A., Marquet, P.A., Martinez, N.D., Mooers, A.,
915 Roopnarine, P., Vermeij, G., Williams, J.W., Gillespie, R., Kitzes, J., Marshall,

916 C., Matzke, N., Mindell, D.P., Revilla, E., Smith, A.B., 2012. Approaching a state
917 shift in Earth's biosphere. *Nature*, 486, 52-58, doi:10.1038/nature11018.

918 Barton, A., Hales, B., Waldbusser, G.G., Langdon, C., and Feely, R.A. (2012) The
919 Pacific oyster *Crassostrea gigas*, shows negative correlation to naturally elevated
920 carbon dioxide levels: implications for near-term ocean acidification effects.
921 *Limnol. Oceanogr.* 57, 698–710

922 Beaugrand, G. (2004) The North Sea regime shift: evidence, causes, mechanisms and
923 consequences, *Progress in Oceanography* 60, 245-262.

924 Behrenfeld, M. J., R. T. O'Malley, D. A. Siegel, C. R. McClain, J. L. Sarmiento, G. C.
925 Feldman, A. J. Milligan, P. G. Falkowski, R. M. Letelier and E. S. Boss (2006)
926 "Climate-driven trends in contemporary ocean productivity." *Nature* 444, 752-
927 755.

928 Bernardi Aubry, F., Cossarini, G., Acri, F., Bastianini, M., Bianchi, F., Camatti, E., De
929 Lazzari, A., Pugnetti, A., Solidoro, C., Socal, G. (2012) Plankton communities in
930 the northern Adriatic Sea: patterns and changes over the last 30 years. *Estuarine,
931 Coastal and Shelf Science* 115, 125-137

932 Bianchi, T. S., Dimarco, S. F., Allison, M. A., Chapman, P., Cowan, J. H., Hetland, R. D.,
933 & Rowe, G. (2008) Controlling hypoxia on the US Louisiana shelf: Beyond the
934 nutrient- centric view. *Eos, Transactions American Geophysical Union*, 89(26),
935 236-237.

936 Biastoch, A., Treude, T., Rupke, L. H., Riebesell, U., Roth, C., Burwicz, E. B., Park, W.,
937 Latif, M., Boning, C. W., and Madec, G. (2011) Rising Arctic Ocean
938 temperatures cause gas hydrate destabilization and ocean acidification,
939 *Geophysical Research Letters* 38.

940 Bijma, J., Portner, H.-O., Yesson, C., and Rogers, A. D. (2013) Climate change and the
941 oceans-What does the future hold?, *Marine Pollution Bulletin* 74, 495-505.

942 Billen G., Garnier J., (2007) River basin nutrient delivery to the coastal sea: Assessing its
943 potential to sustain new production of non-siliceous algae. *Marine Chemistry* 106,
944 148-160

945 Biscaye, P. E., Flagg, C. N., and Falkowski, P. G. (1994) The shelf edge exchange
946 processes experiment, SEEP-II: an introduction to hypotheses, results and
947 conclusions, *Deep Sea Research Part II: Topical Studies in Oceanography* 41,
948 231-252.

949 Black, B.A., G.W. Boehlert, and M. Yoklavich. (2008) Establishing climate–growth
950 relationships for yelloweye rockfish (*Sebastes ruberrimus*) in the northeast Pacific
951 using a dendrochronological approach. *Fisheries Oceanography* 17, 368-379.

952 Black, D., R. Thunell, K. Wejnert, and Y. Astor (2011) Carbon isotope composition of
953 Caribbean Sea surface waters: Response to the uptake of anthropogenic CO₂,
954 *Geophys. Res. Lett.*, 38, L16609, doi:10.1029/2011GL048538.

955 Boesch, DF, RB Brinsfield and RE Magnien. (2001) Chesapeake Bay eutrophication.
956 *Journal of Environmental Quality* 30, 303-320

957 Bograd, S. J., Castro, C. G., Di Lorenzo, E., Palacios, D. M., Bailey, H., Gilly, W., and
958 Chavez, F. P. (2008) Oxygen declines and the shoaling of the hypoxic boundary
959 in the California Current, *Geophysical Research Letters* 35.

960 Bograd, S. J., Checkley, Jr, D. A., and Wooster, W. S. (2003) CalCOFI: A half century of
961 physical, chemical, and biological research in the California Current System,
962 *Deep Sea Research Part II: Topical Studies in Oceanography* 50, 2349-2353.

963 Bopp, L., Le Quere, C., Heimann, M., Manning, A.C., Monfray, P., 2002. Climate-
964 induced oceanic oxygen fluxes: Implications for the contemporary carbon budget.
965 *Global Biogeochemical Cycles*, 16(2) 10.1029/2001gb001445

966 Borges, A. V. and N. Gypens (2010). "Carbonate chemistry in the coastal zone responds
967 more strongly to eutrophication than to ocean acidification." *Limnology and*
968 *Oceanography* 55(1): 346-353.

969 Borkman, D., H. Barreta-Bekker and P. Henriksen eds. (2009) Long-term phytoplankton
970 time series. *Journal of Sea Research* 61, (1 & 2).

971 Borkman, D.G., Smayda, T. (2009) Multidecadal (1959-1997) changes in *Skeletonema*
972 abundance and seasonal bloom patterns in Narragansett Bay, Rhode Island, USA.
973 *Journal of Sea Research* 61, 84-94.

974 Breitburg, D. L., Adamack, A., Rose, K. A., Kolesar, S. E., Decker, B., Purcell, J. E.,
975 Keister, J. E., and Cowan, J. H. (2003) The pattern and influence of low dissolved
976 oxygen in the Patuxent River, a seasonally hypoxic estuary, *Estuaries* 26, 280-297.

977 Breitburg, D. L., Hondorp, D. W., Davias, L. A., and Diaz, R. J. (2009b) Hypoxia,
978 nitrogen, and fisheries: integrating effects across local and global landscapes,
979 *Annual Review of Marine Science* 1, 329-349.

980 Breitburg, DL, JK Craig, RS Fulford, KA Rose, WR Boynton, DC Brady, BJ Ciotti, RJ
981 Diaz, KD Friedland JD Hagy 111, DR Hart, AH Hines, ED Houde, SE Kolesar,
982 SW Nixon, JA Rice, DH Secor and TE Targett. (2009a) Nutrient enrichment and
983 fisheries exploitation: interactive effects on estuarine living resources and their
984 management. *Hydrobiologia* 2009, 31-47

985 Caballero-Alfonso, A. et al. (2013) Regional variability of hypoxia in the coastal Baltic
986 Sea. This volume

987 Cai, W.J. et al., (2011). Acidification of subsurface coastal waters enhanced by
988 eutrophication. *Nature Geoscience*, 4(11): 766-770

989 Canals, M., Puig, P., Durrieu de Madron, X., Heussner, S., Palanques, A., Fabres, J.,
990 (2006). Flushing submarine canyons. *Nature* 444, doi:10.1038/nature05271.

991 Cargo, D.G. and King, D. R. (1990) Forecasting the abundance of the sea nettle,
992 *Chrysaora quinquecirrha*, in the Chesapeake Bay. *Estuaries* 13, 486-491.

993 Carmack, E., McLaughlin, F., Whiteman, G., Homer-Dixon, T. (2012) Detecting and
994 coping with disruptive shocks in Arctic marine systems: a resilience approach to
995 place and people. *AMBIO* (2012) 41:56-65 DOI 10.1007/s13280-011-0225

996 [Carstensen J, Weydmann A.](#) (2012) Tipping points in the arctic: eyeballing or statistical
997 significance? [Ambio](#). 2012 Feb;41(1):34-43. doi: 10.1007/s13280-011-0223-8.

998 Carstensen, J., Conley, D.J., Andersen, J.H., Ærtebjerg, G., (2006) Coastal eutrophication
999 and trend reversal: a Danish case study. *Limnology and Oceanography* 51 (1e2),
1000 398e408.

1001 Carstensen, J., et al. (2014). "Hypoxia in the Baltic Sea: Biogeochemical Cycles, Benthic
1002 Fauna, and Management." *AMBIO* 43(1): 26-36.

1003 Cathalot, C., Rabouille, C., Pastor, L., Deflandre, B., Viollier, E., Buscail, R., Gremare,
1004 A., Treignier, C., Pruski, A.,(2010). Temporal variability of carbon recycling in

1005 coastal sediments influenced by rivers: assessing the impact of flood inputs in
1006 the Rhone River prodelta. *Biogeosciences* 7, 1187-1205.

1007 Chan, F., Barth, J., Lubchenco, J., Kirincich, A., Weeks, H., Peterson, W., and Menge, B.
1008 (2008) Emergence of anoxia in the California Current large marine ecosystem,
1009 *Science* 319, 920-920.

1010 Chauvaud L., Dunbar R., Lorrain A., Paulet Y.-M., Thouzeau G., Jean F., Guarini J. -M.,
1011 Mucciarone D. (2005) The shell of the great scallop *Pecten maximus* as a high
1012 frequency archive of paleoenvironmental change. *Geochemistry Geophysics*
1013 *Geosystems* 6, Q08001

1014 Chauvaud L., Patry Y., Jolivet A., Cam E., Le Goff C., Strand Ø., Charrier G., Thébault
1015 J., Lazure P., Gotthard K., Clavier J. (2012) Variation in size and growth of the
1016 great scallop *Pecten maximus* along the latitudinal gradient. *PLoS ONE* 7,e37717

1017 Chavez, F. P., Messie, M., and Pennington, J. T. (2011) Marine primary production in
1018 relation to climate variability and change, *Annual Review of Marine Science* 3,
1019 227-260.

1020 Chavez, F.P., Ryan, J., Lluch-Cota, S.E., Ñiquen C., M., (2003) From Anchovies to
1021 Sardines and Back: Multidecadal Change in the Pacific Ocean. *Science*
1022 299,(5604), 217-221.

1023 Checkley Jr., D. M., and Barth, J. A. (2009) Patterns and processes in the California
1024 Current System, *Progress in Oceanography* 83, 49-64.

1025 Church, M. J., M. W. Lomas, and F. E. Muller-Karger. (2013) Sea Change: Charting the
1026 course for biogeochemical ocean time series research in a new millennium. *Deep-*
1027 *Sea Research, Part II: Topical Studies in Oceanography* 93, 2-15.

1028 Cloern, J.E. (1982) Does the benthos control phytoplankton biomass in South San
1029 Francisco Bay (USA)?: *Marine Ecology-Progress Series* 9,191-202

1030 Cloern, J. E., and Jassby, A. D. (2012) Drivers of change in estuarine-coastal ecosystems:
1031 Discoveries from four decades of study in San Francisco Bay, *Reviews of*
1032 *Geophysics* 50.

1033 Cloern, J.E., (1996) Phytoplankton bloom dynamics in coastal ecosystems : a review with
1034 some general lessons from sustained investigation of San Francisco Bay,
1035 California. *Reviews of Geophysics* 34, 127-168.

1036 Conley, D.J. and 18 others. (2011) Hypoxia is increasing in the coastal zone of the Baltic
1037 Sea. *Env. Sci. and Technol. Lett.* 45, 6777-6783.

1038 Conversi, A., Fonda-Umani, S., Peluso, T., Molinero, J.C., Santojanni, A., Edward, M.
1039 (2010) The Mediterranean Sea regime shift at the end of the 1980s, and intriguing
1040 parallelisms with other European Basins. *PLoS ONE* 5 (5), e10633. [http://](http://dx.doi.org/10.1371/journal.pone.0010633)
1041 dx.doi.org/10.1371/journal.pone.0010633.

1042 Conversi, A., Peluso, T., Fonda-Umani, S. (2009) Gulf of Trieste: A changing ecosystem.
1043 *Journal of Geophysical Research*, 114, C03S90, doi:10.1029/2008JC004763,
1044 2009

1045 Cooper, S. R., and Brush, G. S. (1993) A 2,500-year history of anoxia and eutrophication
1046 in Chesapeake Bay, *Estuaries* 16, 617-626.

1047 Craig, J. K. (2012) Aggregation on the edge: effects of hypoxia avoidance on the spatial
1048 distribution of brown shrimp and demersal fishes in the Northern Gulf of Mexico.
1049 *Marine Ecology Progress Series* 445, 75-95.)

1050 Crain, C. M., Kroeker, K., and Halpern, B. S. (2008) Interactive and cumulative effects of
1051 multiple human stressors in marine systems, *Ecology Letters* 11, 1304-1315.

1052 Dähnke, K., E. Bahlmann and K.-C. Emeis (2008) A nitrate sink in estuaries? An
1053 assessment by means of stable nitrate isotopes in the Elbe estuary. *Limnol. Ocean.*
1054 **53**(4), 1504-1511.

1055 Dame, R. (2012) Bivalve Filter Feeders: in Estuarine and Coastal Ecosystem Processes.
1056 Springer Publishing Co.

1057 Dame, R. (2011). Ecology of Marine Bivalves: An Ecosystem Approach. CRC Press,
1058 Boca Raton, FL.

1059 Deutsch, C., Brix, H., Ito, T., Frenzel, H., and Thompson, L. (2011) Climate-forced
1060 variability of ocean hypoxia, *Science* 333, 336-339.

1061 deYoung, B., Barange, M., Beaugrand, G., Harris, R., Perry, R. I., Scheffer, M., and
1062 Werner, F. (2008) Regime shifts in marine ecosystems: detection, prediction and
1063 management, *Trends in Ecology & Evolution* 23, 402-409.

1064 Diaz, R.J., Rosenberg, R., (2008) Spreading dead zones and consequences for marine
1065 ecosystems. *Science* 321, 926-929.

1066 Doney, S., Ruckelshaus, M., Duffy, J.E., Barry, J.P., F. Chan, C.A. English, H.M. Galindo,
1067 J.M. grebmeier, A. B. Hollowed, N. Knowlton, J. Polovina, N. Rabalais,
1068 W.Sydeman and L. Talley. 2012. Climate change impacts on marine ecosystems.
1069 *Annu. Rev. Mar. Sci.* 2012. 4:11–37

1070 Doney, S.C., Fabry, V.J., Feely, R.A., Kleypas, J.A., (2009) Ocean Acidification: The
1071 Other CO₂ Problem. *Annual Review of Marine Science*, 1: 169-192

1072 Duarte, C. M., Hendriks, I. E., Moore, T. S., Olsen, Y. S., Steckbauer, A., Ramajo, L.,
1073 Carstensen, J., Trotter, J. A., and McCulloch, M. (2013) Is ocean acidification an
1074 open-ocean syndrome? Understanding anthropogenic impacts on seawater pH,
1075 *Estuaries and Coasts*, 1-16.

1076 Duarte, C.M., Agustí, S., Kennedy, H., Vaqué, D. (1999) The Mediterranean climate as a
1077 template for Mediterranean marine ecosystems: the example of the NE Spanish
1078 littoral. *Progress in Oceanography* 44, 245-270.

1079 Duarte, C.M., Agusti, S., Wassmann, P., Arrieta, J.M, Alcaraz, M., Coello, A., Marba, N.,
1080 Hendriks, I.E., Holding, J., Garcia-Zarandona, I., Kritzberg, E., Vaque, D. (2012)
1081 Tipping elements in the Arctic marine ecosystem, *Ambio* 41, 44-55.

1082 Durell, EQ and C Weedon. (2011) Striped bass seine survey juvenile index web page.
1083 <http://www.dnr.state.md.us/fisheries/juvinindex/index.html>. Maryland Department
1084 of Natural Resources, Fisheries Service

1085 Emeis, K., J. van Beusekom, U. Callies, R. Ebinghaus, A. Kannen, G. Kraus, I. Kröncke,
1086 H. Lenhart, I. Lorkowski, V. Matthias, C. Möllmann, J. Pätsch, M. Scharfe, H.
1087 Thomas, R. Weisse and E. Zorita (submitted) The North Sea - a shelf sea in the
1088 Anthropocene. *Journal of Marine Systems* (this volume).

1089 Emeis, K.-C., Struck, U., Leipe, T., Pollehne, F., Kundendorf, H., Christiansen, C. (2000)
1090 Changes in the C, N, P burial rates in some Baltic Sea sediments over the last 150
1091 years — relevance to P regeneration rates and the phosphorus cycle. *Marine*
1092 *Geology*, 167, (1–2), 43-59

1093 Feely, R.A., Sabine, C.L., Hernandez-Ayon, J.M., Ianson, D., Hales, B. (2008) Evidence
1094 for upwelling of corrosive “acidified” water onto the continental shelf. *Science*
1095 320, 1490-1492.

1096 Frieder, C., Nam, S., Martz, T., and Levin, L. (2012) High temporal and spatial
1097 variability of dissolved oxygen and pH in a nearshore California kelp forest,
1098 *Biogeosciences* 9, 3917-3930.

1099 Garrabou, J., Coma, R., Bensoussan, N., Bally, M., Chevaldonne, P., Cigliano, M., Diaz,
1100 D., Harmelin, J. G., Gambi, M., and Kersting, D. (2009) Mass mortality in
1101 Northwestern Mediterranean rocky benthic communities: effects of the 2003 heat
1102 wave, *Global Change Biology* 15, 1090-1103.

1103 Giani, M., Djakovac, T., Degobbis, D., Cozzi, S., Solidoro, C., and Umani, S. F. (2012)
1104 Recent changes in the marine ecosystems of the northern Adriatic Sea, *Estuarine,
1105 Coastal and Shelf Science* 115, 1-13.

1106 Gilly, W.F., Beman, J.M., Litvin, S.Y. and B.H. Robison. (2013) Oceanographic and
1107 biological effects of shoaling of the oxygen minimum zone. *Ann. Rev. Mar. Sci.* 5,
1108 21.1-21.28

1109 Glavovic, B.C. et al., 2014. Living on the Margin in the Anthropocene: Engagement
1110 Arena for Global Sustainability Research and Action. *Science*: (Submitted).

1111 Goffart, A., Hecq, J. H., and Legendre, L. (submitted) Drivers of phytoplankton bloom in
1112 the oligotrophic Bay of Calvi (NW Mediterranean) : results from a long-term
1113 study (1979-2011), *Progress in Oceanography*.

1114 Goffart, A., Hecq, J.-H., and Legendre, L. (2002) Changes in the development of the
1115 winter-spring phytoplankton bloom in the Bay of Calvi (NW Mediterranean) over
1116 the last two decades: a response to changing climate?, *Marine Ecology Progress
1117 Series* 236, 45-60.

1118 Gooday, A.J., Jorissen, F., Levin, L.A., Middelburg, J.J, Naqvi, W., Rabalais, N.,
1119 Scranton, M., Zhang, J. (2009) Historical records of coastal eutrophication and
1120 hypoxia *Biogeosciences* 6, 1-39.

1121 Grall, J; Chauvaud, L (2002) Marine eutrophication and benthos: the need for new
1122 approaches and concepts. *Global Change Biology*, 8: 813-830

1123 Grantham, B. A., F. Chan, K. J. Nielsen, D. S. Fox, J. A. Barth, A. Huyer, J. Lubchenco,
1124 and B. A. Menge (2004), Upwelling-driven nearshore hypoxia signals ecosystem
1125 and oceanographic changes in the northeast Pacific, *Nature* 429(6993), 749–754,
1126 doi:10.1038/nature02605.

1127 Gren, I.-M., 2013. The economic value of coastal waters as nutrient filters for the Baltic
1128 Sea. *Reg Environ Change*, 13: 695–703.

1129 Gruber, N., and Sarmiento, J. L. (1997) Global patterns of marine nitrogen fixation and
1130 denitrification, *Global Biogeochemical Cycles* 11, 235-266.

1131 Gruber, N., Hauri, C., Lachkar, Z., Loher, D., Frolicher, T. L., and Plattner, G.-K. (2012),
1132 Rapid progression of ocean acidification in the California Current System,
1133 *Science* 337, 220-223.

1134 Han, H., and Allan, J. D. (2008), Estimation of nitrogen inputs to catchments: comparison
1135 of methods and consequences for riverine export prediction, *Biogeochemistry* 91,
1136 177-199.

1137 Harrison, P., A. Zingone and E. Philips ed. (2010) Phytoplankton time series. *Estuaries
1138 and Coastal Ecosystems* , vol. 33,

1139 Hofmann, E.E., B. Cahill, K. Fennel, MAM Friedrichs, K Hyde, C Lee, A. Mannino, RG
1140 Najjar, JE O'Reilly, J Wilkin, J Xue, (2011) Modeling the dynamics of
1141 continental shelf carbon. *Annual Review of Marine Science* 3, 93-122.

1142 Hofmann, G.E. , T. G. Evans, M. W. Kelly, J. L. Padilla-Gamiño, C. A. Blanchette,
 1143 L. Washburn, F. Chan, M. A. McManus, B. A. Menge, B. Gaylord, T. M. Hill, E.
 1144 Sanford, M. LaVigne, J. M. Rose, L. Kapsenberg, and J. M. Dutton. 2014.
 1145 Exploring local adaptation and the ocean acidification seascape – studies in the
 1146 California Current large marine ecosystem. *Biogeosciences*, 11, 1053–1064
 1147 Howarth, R. W., Billen, G., Swaney, D., Townsend, A., Jaworski, N., Lajtha, K.,
 1148 Downing, J., Elmgren, R., Caraco, N., and Jordan, T. (1996) Regional nitrogen
 1149 budgets and riverine N & P fluxes for the drainages to the North Atlantic Ocean:
 1150 Natural and human influences, *Biogeochemistry* 35, 75-139.
 1151 Howarth, R., Swaney, D., Billen, G., Garnier, J., Hong, B., Humborg, C., Johnes, P.,
 1152 Morth, C.-M., and Marino, R. (2012) Nitrogen fluxes from the landscape are
 1153 controlled by net anthropogenic nitrogen inputs and by climate, *Frontiers in*
 1154 *Ecology and the Environment* 10, 37-43.
 1155 Howarth, R., Swaney, D., Boyer, E., Marino, R., Jaworski, N., and Goodale, C. (2006)
 1156 The influence of climate on average nitrogen export from large watersheds in the
 1157 Northeastern United States, *Biogeochemistry* 79, 163-186.
 1158 Hsieh, C. H., Glaser, S. M., Lucas, A. J., Sugihara, G. (2005) Distinguishing random
 1159 environmental fluctuations from ecological catastrophes for the North Pacific
 1160 Ocean. *Nature* 435, 336-340.
 1161 Hughes, TP. (1994) Catastrophes, phase shifts, and large-scale degradation of a
 1162 Caribbean coral reef. *Science* 265, 1547-1551
 1163 Humborg C, Ittekkot V, Cociasu A, Von Bodungen B (1997) Effect of Danube River
 1164 dam on Black Sea biogeochemistry and ecosystem structure. *Nature* 386, 385-388
 1165 IPCC (2014). Field et al. Climate Change 2014 : Impacts Adaptation and Vulnerability.
 1166 Summary for Policymakers (Copyright Pending).
 1167 Islam, S., and Tanaka, M. (2004) Impacts of pollution on coastal and marine ecosystems
 1168 including coastal and marine fisheries and approach for management: a review
 1169 and synthesis, *Marine Pollution Bulletin* 48, 624-649.
 1170 Jahnke, R. (2010), Global synthesis, in *Carbon and Nutrient Fluxes in Continental*
 1171 *Margins: A Global Synthesis*, edited by K.-K. Liu, L. Atkinson, R. Quinones and
 1172 L. Talaue-McManus, pp. 597–616.
 1173 Jayakumar, A., Peng, N., and Ward, B. (2013) Denitrifying communities in oceanic
 1174 oxygen deficient zones using microarray analyses, IMBIZO III: The future of
 1175 marine biogeochemistry, ecosystems and societies, IMBER, Goa, India.
 1176 Jobstvogt, N., Hanley, N., Hynes, S., Kenter, J., and Witte, U. (2013) Twenty thousand
 1177 sterling under the sea: Estimating the value of protecting deep-sea biodiversity,
 1178 *Ecological Economics* 97, 10-19.
 1179 Kennedy, V.S., 1984. The estuary as a filter. ????
 1180 Kennett, J. P., Cannariato, K. G., Hendy, I. L., and Behl, R. J. (2003) *Methane hydrates*
 1181 *in Quaternary climate change: The clathrate gun hypothesis*, American
 1182 Geophysical Union.
 1183 Koslow, J. A., and Couture, J. (2013) Ocean science: Follow the fish, *Nature* 502, 163-
 1184 164.
 1185 Lachkar, Z. (2014) Effects of upwelling increase on ocean acidification in the California
 1186 and Canary Current systems, *Geophys. Res. Lett.*, 40,
 1187 doi:10.1002/2013GL058726.

- 1188 Lachkar, Z., and Gruber, N. (2013), Response of biological production and air-sea CO₂
1189 fluxes to upwelling intensification in the California and Canary Current Systems,
1190 *Journal of Marine Systems* 109, 149-160.
- 1191 Lassaletta, L., Romero, E., Billen, G., Garnier, J., Garcia-Gomez, H., and Rovira, J.
1192 (2012), Spatialized N budgets in a large agricultural Mediterranean watershed:
1193 high loading and low transfer, *Biogeosciences* 9, 57-70.
- 1194 Lee, YJ, WR Boynton, M Li and Y Li. (2013) Role of late winter-spring wind influencing
1195 summer hypoxia in Chesapeake Bay. *Estuaries and Coasts* 36, 683-696
- 1196 Lejeusne, C., Chevaldonne, P., Pergent-Martini, C., Boudouresque, C. F., and Perez, T.
1197 (2010) Climate change effects on a miniature ocean: the highly diverse, highly
1198 impacted Mediterranean Sea, *Trends in Ecology & Evolution* 25, 250-260.
- 1199 Levin, L. A. (2003) Oxygen minimum zone benthos: adaptation and community response
1200 to hypoxia, in *Oceanography and Marine Biology, an Annual Review, Volume 41:*
1201 *An Annual Review*, edited by R. Gibson and R. Atkinson, pp. 1-45.
- 1202 Levin, L. A., and Dayton, P. K. (2009) Ecological theory and continental margins: where
1203 shallow meets deep, *Trends in Ecology & Evolution* 24, 606-617.
- 1204 Levin, L. A., and Sibuet, M. (2012) Understanding continental margin biodiversity: a
1205 new imperative, *Annual Review of Marine Science* 4, 79-112.
- 1206 Levin, L. A., Boesch, D. F., Covich, A., Dahm, C., Erseus, C., Ewel, K. C., Kneib, R. T.,
1207 Moldenke, A., Palmer, M. A., and Snelgrove, P. (2001) The function of marine
1208 critical transition zones and the importance of sediment biodiversity, *Ecosystems*
1209 4, 430-451.
- 1210 Levin, L., Ekau, W., Gooday, A., Jorissen, F., Middelburg, J., Naqvi, W., Neira, C.,
1211 Rabalais, N., and Zhang, J. (2009) Effects of natural and human-induced hypoxia
1212 on coastal benthos, *Biogeosciences Discussions* 6, 3563-3654.
- 1213 Levin, L.A., C. Frieder, M. Navarro, J. Gonzalez, T. Martz. (2013) Hypoxia, hypercapnia
1214 and homosapiens on upwelling margins. Oral Presentation, IMBER IMBIZO III,
1215 Goa, India, Jan. 2013.
- 1216 Limburg, K.E., C. Olson, Y. Walther, D. Dale, C. Slomp, and H. Høie (2011) Tracking
1217 Baltic hypoxia and cod migration over millennia with natural tags. *Proceedings*
1218 *of the National Academy of Sciences of the U.S.* doi:10.1073/pnas.1100684108.
- 1219 Limburg, K.E., B.D. Walther, Z. Lu, G. Jackman, J. Mohan, Y. Walther, A. Nissling,
1220 P.K. Weber, and A.K. Schmitt (2014) In search of the dead zone: use of otoliths
1221 for tracking fish exposure to hypoxia. *Journal of Marine Systems*. DOI:
1222 10.1016/j.jmarsys.2014.02.014
- 1223 Limburg, K. E., Walther, B. D., Lu, Z., Jackman, G., Mohan, J., Walther, Y., Nissling, A.,
1224 Weber, P. K., and Schmitt, A. K. (submitted), In search of the dead zone: use of
1225 otoliths for tracking fish exposure to hypoxia.
- 1226 Liu, K.-K., Yan, W., Lee, H.-J., Chao, S.-Y., Gong, G.-C., Yeh, T.-Y. (this issue) Impacts
1227 of increasing dissolved inorganic nitrogen discharge from Changjiang on primary
1228 production and sediment oxygen consumption in the East China Sea from 1970 to
1229 2002. *J. Mar. Syst.*
- 1230 Liu, K.K., Atknison, L., Quinones, R., Talaue-McManus, L. (Eds) (2010) Carbon and
1231 nutrient fluxes in continental margins: A global synthesis. Springer, Heidelberg
1232 Germany, 744 pp.

1233 Liu, S. M. (submitted, this issue) Response of nutrient transports to human activities in
1234 the ecosystem of the Bohai: under the influence of artificial floods.

1235 Liu, S., Zhang, J., Gao, H., and Liu, Z. (2008) Historic changes in flux of materials and
1236 nutrient budgets in the Bohai, *Acta Oceanologica Sinica* 27, 1-17.

1237 Mallin, M. A., Posey, M. H., Shank, G. C., McIver, M. R., Ensign, S. H., and Alphin, T.
1238 D. (1999) Hurricane effects on water quality and benthos in the Cape Fear
1239 watershed: natural and anthropogenic impacts, *Ecological Applications* 9, 350-
1240 362.

1241 Marba, N. and C.M. Duarte (2010) Mediterranean warming triggers seagrass (*Posidonia*
1242 *oceanica*) shoot mortality. *Global Change Biology*, 16, 2366–2375.

1243 Marchesiello, P., and P. Estrade (2009) Eddy activity and mixing in upwelling systems:
1244 A comparative study of Northwest Africa and California regions, *Int. J. Earth*
1245 *Sci.*, 98, 299–308, doi:10.1007/s00531-007-02356.

1246 Marić, D., Kraus, R., Godrijan, J., Supić, N., Djakovac, T., Precali, R. (2012)
1247 Phytoplankton response to climatic and anthropogenic influence in the north-
1248 eastern Adriatic during the last four decades. *Estuarine, Coastal and Shelf Science*
1249 115, 98-112.

1250 McCay, B.J., S. Brandt, C. Creed (2011) Human dimensions of climate change and
1251 fisheries in a coupled system: the Atlantic surfclam case. *ICES J. Mar. Sci.*
1252 (2011) 68 (6): 1354-1367. doi: 10.1093/icesjms/fsr044

1253 McClatchie, S., Goericke, R., Cosgrove, R., Auad, G., and Vetter, R. (2010) Oxygen in
1254 the Southern California Bight: multidecadal trends and implications for demersal
1255 fisheries, *Geophysical Research Letters* 37, L19602.

1256 McFarlane, G. A., Smith, P. E., Baumgartner, T. R., and Hunter, J. R. (2002) Climate
1257 variability and Pacific sardine populations and fisheries, *American Fisheries*
1258 *Society Symposium* 32, 195-214.

1259 McQuatter-Gollop, A., Gilbertt, A.J., Mee, L.D., Vermaat, J.E., Artioli, Y., Humborg, C.,
1260 Wulff, F., (2009) How well do ecosystem indicators communicate the effects of
1261 anthropogenic eutrophication? *Estuarine, Coastal and Shelf Science* 82, 583e596.

1262 Mee, L. D., Friedrich, J., and Gomoiu, M. T. (2005) Restoring the Black Sea in times of
1263 uncertainty, *Oceanography*, 18, 100–111

1264 Möllmann, C., Conversi A., Edwards M. (2011) Comparative analysis of European wide
1265 marine ecosystem shifts: a large-scale approach for developing the basis for
1266 ecosystem-based management. *Biology Letters* 7, 484-486.

1267 Montes, E., Lomas, M., Muller-Karger, F. E., and Lorenzoni, L. (in prep) Responses of
1268 the N cycle in the Tropical and Sub-tropical North Atlantic to modern climate
1269 variability, *Global Biogeochemical Cycles*.

1270 Mora, C., Wei, C.-L., Rollo, A., Amaro, T., Baco, A. R., Billett, D., Bopp, L., Chen, Q.,
1271 Collier, M., and Danovaro, R. (2013) Biotic and Human Vulnerability to
1272 Projected Changes in Ocean Biogeochemistry over the 21st Century, *PLoS*
1273 *Biology* 11, e1001682.

1274 Morat, F., Letourneur, Y., Dierking, J., Pecheyran, C., Bareille, G., Blamart, D.,
1275 Harmelin-Vivien, M.L., (2014). The great melting pot. Common sole population
1276 connectivity assessed by otolith and water fingerprints. *PlosOne* 9, e86585.
1277 doi:86510.81371/journal.pone.0086585.

1278 Mozetič, P., Francé, J., Kogovšek, T.; Talaber, I., Malej, A. (2012) Plankton trends and
1279 community changes in a coastal sea: bottom-up vs. top-down control in relation to
1280 local-scale and large-scale drivers. *Estuarine, Coastal and Shelf Science* 115, 138-
1281 148.

1282 Muller-Karger, F. R. Varela, R., Thunell, R. Luerssen, C. Hu and J. Walsh. (2005) The
1283 importance of continental margins in the global carbon cycle. *Geophys. Res.*
1284 *Letters*, 32, DOI: 10.1029/2004GL021346

1285 Muller-Karger, F., et al. (2013) The CARIACO Ocean Time-Series: 18 years of
1286 international collaboration in ocean biogeochemistry and ecological research,
1287 *Ocean Carbon and Biogeochemistry Newsletter, Woods Hole Oceanographic*
1288 *Institution* fall edition.

1289 Mumby, P. J.; Hastings, A.; Edwards, H.J. (2007) Thresholds and the resilience of
1290 Caribbean coral reefs". *Nature* 450, 98-101.
1291 [doi:10.1038/nature06252](https://doi.org/10.1038/nature06252) [PMID 17972885](https://pubmed.ncbi.nlm.nih.gov/17972885/).

1292 Naam, R., (2012) Arctic Sea ice: What, why and what next. *Scientific American*,
1293 [http://blogs.scientificamerican.com/guest-blog/2012/09/21/arctic-sea-ice-what-](http://blogs.scientificamerican.com/guest-blog/2012/09/21/arctic-sea-ice-what-why-and-what-next/)
1294 [why-and-what-next/](http://blogs.scientificamerican.com/guest-blog/2012/09/21/arctic-sea-ice-what-why-and-what-next/)

1295 Naha Biswas,S., Godhantaraman, N., Sarangi, R.K., B.D. Bhattacharya, B.D., Sarkar,
1296 S.K., Satpathy, K.K. (2013) Bloom of the centric diatom *Hemidiscus*
1297 *hardmannianus* (Bacillariophyceae) and its impact on water quality characteristics
1298 and plankton community structure in Indian Sundarban mangrove wetland-
1299 **CLEAN - Soil, Air and Water Pollution**, 41 (4), 333-339.

1300 Nam, S., Kim, H.-J., and Send, U. (2011) Amplification of hypoxic and acidic events by
1301 La Nina conditions on the continental shelf off California, *Geophysical Research*
1302 *Letters* 38, L22602.

1303 Narváez, D., et al., (submitted) Long-term dynamics in Atlantic surfclam (*Spisula*
1304 *Solidissima*) populations: The role of bottom water temperature. *J. Mar. Systems*
1305 (this volume).

1306 Ninčević Gladan, Ž., Marasović, I., Grbec, B., Skejić, S., Bužančić, M., Kušpilić, G.,
1307 Matijević, S., Matić, F., (2010) Inter-decadal Variability in Phytoplankton
1308 Community in the Middle Adriatic (Kaštela Bay) in Relation to the North Atlantic
1309 Oscillation. *Estuaries and Coasts* 33, 376-383.

1310 Nixon, S. W., and Buckley, B. A. (2002) "A strikingly rich zone"-nutrient enrichment
1311 and secondary production in coastal marine ecosystems, *Estuaries* 25, 782-796.

1312 O'Brien, T.D. et al. (Ed.) (2012) ICES Phytoplankton and Microbial Plankton Status
1313 Report 2009/2010. ICES Cooperative Research Report, 313. *ICES*: Denmark.
1314 ISBN 978-87-7482-115-1. 196 pp

1315 Paerl, H. W., Bales, J. D., Ausley, L. W., Buzzelli, C. P., Crowder, L. B., Eby, L. A., Fear,
1316 J. M., Go, M., Peierls, B. L., and Richardson, T. L. (2001) Ecosystem impacts of
1317 three sequential hurricanes (Dennis, Floyd, and Irene) on the United States' largest
1318 lagoonal estuary, Pamlico Sound, NC, *Proceedings of the National Academy of*
1319 *Sciences* 98, 5655-5660.

1320 Paulmier, A., Ruiz-Pino, D., and Garçon, V. (2011) CO₂ maximum in the oxygen
1321 minimum zone (OMZ), *Biogeosciences* 8, 239-252.

1322 Phrampus, B. J., and Hornbach, M. J. (2012) Recent changes to the Gulf Stream causing
1323 widespread gas hydrate destabilization, *Nature* 490, 527-530.

- 1324 Pidgeon, N., and Fischhoff, B. (2011) The role of social and decision sciences in
1325 communicating uncertain climate risks, *Nature Climate Change* 1, 35-41.
- 1326 Pätsch, J., Lenhart, H.-J. (2011) Daily nutrient loads of Nutrients, Total Alkalinity,
1327 Dissolved Inorganic Carbon and Dissolved Organic Carbon of the European
1328 continental rivers for the years 1977-2009. Berichte aus dem Zentrum für Meeres-
1329 und Klimaforschung, Reihe B: Ozeanographie, 50, Zentrum für Meeres- und
1330 Klimaforschung, Hamburg, 159 pp.
- 1331 Prince, E. D., and Goodyear, C. P. (2006) Hypoxia-based habitat compression of tropical
1332 pelagic fishes, *Fisheries Oceanography* 15, 451-464.
- 1333 Prince, E.D., Luo, J., Goodyear, C.P., Hoolihan, J.P., Snodgrass, D., Orbesen, E.S.,
1334 Serafy, J.E., Ortiz, M., Schirripa, M.J. (2010) Ocean scale hypoxia-based habitat
1335 compression of Atlantic istiophorid billfishes. *Fish. Oceanogr.* 19, 448-462.
- 1336 Rabalais, N. N. (2004), Eutrophication, in: The Global Coastal Ocean Multiscale
1337 Interdisciplinary Processes, edited by: Robinson, A. R., McCarthy, J., and
1338 Rothschild, B. J., The Sea, Vol. 13, Harvard University Press, 819–865, 2004.
- 1339 Rabalais, N. N., R. J. Diaz, L. A. Levin, R. E. Turner, D. Gilbert and J. Zhang. (2010)
1340 Dynamics and distribution of natural and human-caused coastal hypoxia.
1341 *Biogeosciences* 7: 585-619.
- 1342 Rabalais, N.N., W.-J. Cai, J. Carstensen, D.J. Conley, B. Fry, X. Hu, Z. Quiñones-Rivera,
1343 R. Rosenberg, C.P. Slomp, R.E. Turner, M. Voss, B. Wissel, and J. Zhang. 2014.
1344 Eutrophication-driven deoxygenation in the coastal ocean. *Oceanography*
1345 27(1):172–183, <http://dx.doi.org/10.5670/oceanog.2014.21>.
- 1346 Rabouille, C., Conley, D.J., Dai, M.H., Cai, W.J., Chen, C.T.A., Lansard, B., Green, R.,
1347 Yin, K., Harrison, P.J., Dagg, M., McKee, B., (2008). Comparison of hypoxia
1348 among four river-dominated ocean margins: The Changjiang (Yangtze),
1349 Mississippi, Pearl, and Rhone rivers. *Continental Shelf Research* 28, 1527-1537.
- 1350 Rabouille, O., O. Radakovitch, C. Estournel, X. Durrieu de Maron, C. Guieu, and R.
1351 Sempere. (2013) The Fate of Rhone River carbon on the mediterranean
1352 continental margin, its export to the open sea and its relation to climatic
1353 parameters. Oral Presentation, IMBER IMBIZO, Jan. 2013, Goa, India.
- 1354 Ragueneau, O., Chauvaud, L., Moriceau, B., Leynaert, A., Thouzeau, G., Donval, A., Le
1355 Loc'h, F. and Jean, F., (2005) Biodeposition by an invasive suspension feeder
1356 impacts the biogeochemical cycle of Si in a coastal ecosystem (Bay of Brest,
1357 France). *Biogeochemistry*, DOI 10.1007/s10533-004-5677-3.
- 1358 Ragueneau, O., Conley, D.J., Ni Longphuirt, S., Slomp, C. et Leynaert, A., (2006) A
1359 review of the Si biogeochemical cycle in coastal waters, II: anthropogenic
1360 perturbation of the Si cycle and responses of coastal ecosystems. Dans: Land-
1361 Ocean nutrient fluxes: silica cycle. Ittekkot, V., Humborg, C., Garnier, J. (Eds.),
1362 SCOPE Book, Island Press, pp. 197-213.
- 1363 Raimonet, M., Ragueneau, O., Jacques, V. Corvaisier, R., Moriceau, B., Khripounoff, A.,
1364 Rabouille, C. (submitted) Rapid transport and high accumulation of riverine Si in
1365 the Congo deep sea fan (this volume).
- 1366 Reid, W. V., Chen, D., Goldfarb, L., Hackmann, H., Lee, Y. T., Mokhele, K., Ostrom, E.,
1367 Raivio, K., Rockström, J., Schellnhuber, H. J., Whyte, A. (2010) Earth System
1368 Science for Global Sustainability: Grand Challenges. *Science* 330, 916-917.

1369 Ribera d'Alcalà, M., Conversano, F., Corato, F., Licandro, P., Mangoni, O., Marino, D.,
1370 Mazzocchi, M. G., Modigh, M., Montresor, M., Nardella, M., Saggiomo, V.,
1371 Sarno, D., Zingone, A. (2004) Seasonal patterns in planktonic communities in a
1372 pluriannual time series at a coastal Mediterranean site (Gulf of Naples): an attempt
1373 to discern recurrences and trends, *Scientia Marina*, **68(Suppl.1)**, 65-83,
1374 Robbins LL, Wynn JG, Lisle JT, Yates KK, Knorr PO, et al. (2013) Baseline monitoring
1375 of the Western Arctic Ocean estimates 20% of Canadian Basin surface waters Are
1376 undersaturated with respect to aragonite. *PLoS ONE* 8(9), e73796.
1377 doi:10.1371/journal.pone.0073796
1378 Sattar, M.A., C.Kroeze, and M. STrokal. 2014. The increasing impact of food production
1379 on nutrient export by rivers to the Bay of Bengal 1970-2050. *Mar. Poll. Bull.* 80:
1380 168-178.
1381 Schuur, E.A.G., (2013) High risk of permafrost thaw. Lawrence Berkeley National
1382 Laboratory. <http://dx.doi.org/doi:10.1038/480032a>
1383 Send, U., and Nam, S. (2012) Relaxation from upwelling: The effect on dissolved oxygen
1384 on the continental shelf, *Journal of Geophysical Research: Oceans (1978, 2012)*
1385 117.
1386 Soetaert, K., Middelburg, J.J., Heip, C., Meire, P., Van Damme, S., Maris, T., (2006)
1387 Long-term change in dissolved inorganic nutrients in the heterotrophic Scheldt
1388 estuary (Belgium, The Netherlands). *Limnology and Oceanography* 51 (1, Part 2),
1389 409e423.
1390 Srinivas, B., Sarin, M.M. and Sarma, S.S.V.V. (submitted) Atmospheric outflow of
1391 nutrients to the Bay of Bengal: Impact of continental sources. *Journal of Marine*
1392 *Systems* (this volume)
1393 Stramma, L., Prince, E. D., Schmidtko, S., Luo, J., Hoolihan, J. P., Visbeck, M., Wallace,
1394 D. W., Brandt, P., and Kortzinger, A. (2011) Expansion of oxygen minimum
1395 zones may reduce available habitat for tropical pelagic fishes, *Nature Climate*
1396 *Change* 2, 33-37.
1397 Stramma, L., Schmidtko, S., Levin, L. A., and Johnson, G. C. (2010) Ocean oxygen
1398 minima expansions and their biological impacts, *Deep Sea Research Part I:*
1399 *Oceanographic Research Papers* 57, 587-595.
1400 Swaney, D. P., Hong, B., Selvam, P., Howarth, R.W., Ramesh, R., Ramachandran, P.,
1401 (2013) Net anthropogenic nitrogen inputs and nitrogen fluxes from Indian
1402 watersheds: a preliminary assessment, *Journal of Marine Systems*, *submitted*.
1403 Swaney, D. P., Hong, B., Ti, C., Howarth, R. W., and Humborg, C. (2012) Net
1404 anthropogenic nitrogen inputs to watersheds and riverine N export to coastal
1405 waters: a brief overview, *Current Opinion in Environmental Sustainability* 4, 203-
1406 211.
1407 Taylor, G.T., Muller-Karger, F.E., Thunell, R.C., Scranton, M.I., Astor, Y., Varela, R.,
1408 Ghinaglia, L.T., Lorenzoni, L., Fanning, K.A., Hameed, S., Doherty, O., (2012)
1409 Ecosystem responses in the southern Caribbean Sea to global climate change. *P.*
1410 *Natl. A. Sci. USA*, doi:10.1073/pnas.1207514109
1411 Thomas, P., Rahman, M. S., Kummer, J. A., and Lawson, S. (2006) Reproductive
1412 endocrine dysfunction in Atlantic croaker exposed to hypoxia, *Marine*
1413 *environmental research* 62, S249-S252.

1414 Thomsen, J., Gutowska, M., Saphorster, J., Heinemann, A., Trubenbach, K., Fietzke, J.,
1415 Hiebenthal, C., Eisenhauer, A., Kortzinger, A., and Wahl, M. (2010) Calcifying
1416 invertebrates succeed in a naturally CO₂-rich coastal habitat but are threatened by
1417 high levels of future acidification, *Biogeosciences* 7, 3879-3891.

1418 Trenberth, K.E., Jones, P.D., Ambenje, P., Bojariu, R., Easterling, D., Klein Tank, A.,
1419 Parker, D., Rahimzadeh, F., Renwick, J. A., Rusticucci, M., Soden, B., Zhai, P.,
1420 (2007) Observations: Surface and Atmospheric Climate Change. In: Climate
1421 Change 2007: The Physical Science Basis. Contribution of Working Group I to
1422 the Fourth Assessment Report of the Intergovernmental Panel on Climate Change
1423 [eds Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K.B.,
1424 Tignor, M., Miller, H.I.]. Cambridge University Press, Cambridge, United
1425 Kingdom and New York, NY, USA.

1426 Waldbusser, G.G., E.N. Powell and R. Mann. (2013) Ecosystem effects of shell
1427 aggregations and cycling in coastal waters: an example of Chesapeake Bay oyster
1428 reefs. *Ecology* 94, 895-903.

1429 Walsh, J. J., Biscaye, P. E., and Csanady, G. T. (1988) The 1983-1984 shelf edge
1430 exchange processes (SEEP)-I experiment: hypotheses and highlights, *Continental
1431 Shelf Research* 8, 435-456.

1432 Wassmann, P. (2011) Arctic Marine ecosystems in an era of rapid climate change.
1433 *Progress in Oceanography* 90: 1-17.

1434 Wassmann, P., Lenton, T., (2012) Arctic tipping points in the Earth System perspective.
1435 *AMBIO* 41(1), 1-9.

1436 Weinberg, J. (2005) Bathymetric shift in the distribution of Atlantic surfclams: response
1437 to warmer ocean temperature. *ICES Journal of Marine Science*, 62: 1444–1453.

1438 Whiteman, G., Hope, C., Wadhams, P., (2013) Vast costs of Arctic change. *Nature*, 499:
1439 401- 403, doi:10.1038/499401a.

1440 Wiltshire, K. H., Malzahn, A. M., Wirtz, K., Greve, W., Janisch, S., Mangelsdorf, P.,
1441 Manly, B. F., Boersma, M. (2008) Resilience of North Sea phytoplankton spring
1442 bloom dynamics: An analysis of long-term data at Helgoland Roads. *Limnology
1443 and Oceanography* 54(4), 1294-1302.

1444 Wishner, K.F., Outram, D. M., Seibel, B. A., Daly, K., and Williams, R. L. (2013)
1445 Zooplankton in the Eastern Tropical North Pacific: Boundary effects of oxygen
1446 minimum zone expansion. *Deep-Sea Research I* 79, 122-140.

1447 Wooster, W. S., and Zhang, C. I. (2004) Regime shifts in the North Pacific: early
1448 indications of the 1976-1977 event, *Progress in Oceanography* 60, 183-200.

1449 World Resources Institute, (2013) Interactive map of eutrophication & hypoxia.
1450 Available online: [http://www.wri.org/resource/interactive-map-eutrophication-
1451 hypoxia](http://www.wri.org/resource/interactive-map-eutrophication-hypoxia) (accessed 1 January, 2014).

1452 Yan, W.J., Mayorga, E., Li, X.Y., Seitzinger, S.P., Bouwman, A.F., (2010) Increasing
1453 anthropogenic nitrogen inputs and riverine DIN exports from the Changjiang
1454 River basin under changing human pressures. *Global Biogeochemical Cycles*,
1455 24 Gb0a06, doi: 10.1029/2009gb003575.

1456 Yasuhara, M., Hunt, G., Breitburg, D., Tsujimoto, A., and Katsuki, K. (2012) Human-
1457 induced marine ecological degradation: micropaleontological perspectives,
1458 *Ecology and Evolution* 2, 3242-3268.

- 1459 Yu, P. C., Matson, P. G., Martz, T. R., and Hofmann, G. E. (2011) The ocean
1460 acidification seascape and its relationship to the performance of calcifying marine
1461 invertebrates: Laboratory experiments on the development of urchin larvae
1462 framed by environmentally-relevant pCO₂/pH, *Journal of Experimental Marine*
1463 *Biology and Ecology* 400, 288-295.
- 1464 Zhai, W.-D., Zheng, N., Huo, C., Xu, Y., Zhao, H.-D., Li, Y.-W., Zang, K.-P., Wang, J.-
1465 Y., and Xu, X.-M. (2013) Subsurface low pH and carbonate saturation state of
1466 aragonite on China side of the North Yellow Sea: combined effects of global
1467 atmospheric CO₂ increase, regional environmental changes, and local
1468 biogeochemical processes, *Biogeosciences Discussions* 10, 3079-3120.
- 1469 Zhai, W.D., Dai, M.H., Chen, B.S., Guo, X.H., Li, Q., Shang, S.L., Zhang, C.Y., Cai, W.J.,
1470 Wang, D.X., 2013b. Seasonal variations of sea–air CO₂ fluxes in the largest
1471 tropical marginal sea (South China Sea) based on multiple-year underway
1472 measurements. *Biogeosciences*, 10(11): 7775-7791
- 1473 Zhang, C. I., and Gong, Y. (2005) Effect of ocean climate changes on the Korean stock of
1474 Pacific saury, *Cololabis saira* (Brevoort), *Journal of Oceanography* 61, 313-325.
- 1475 Zhang, J., Yu, Z., Raabe, T., Liu, S., Starke, A., Zou, L., Gao, H., and Brockmann, U.
1476 (2004) Dynamics of inorganic nutrient species in the Bohai seawaters, *Journal of*
1477 *Marine Systems* 44, 189-212.
- 1478 Zhu, Z.-Y., Zhang, J., Wu, Y., Zhang, Y.-Y., Lin, J., Liu, S.-M., (2011) Hypoxia off
1479 the Changjiang (Yangtze River) Estuary: Oxygen depletion and organic matter
1480 decomposition. *Marine Chemistry* 125, 108–116.
- 1481 Zillén, L., D.J. Conley, T.Andren, E. Andren, S. Bjorck (2008) Past occurrences of
1482 hypoxia in the Baltic Sea and the role of climate variability, environmental
1483 change and human impact. *Earth-Science Reviews* 91, 77-92.
- 1484 Zingone, A., Dubroca, L., Iudicone, D., Margiotta, F., Corato, F., Ribera d'Alcalà, M.,
1485 Saggiomo, V., Sarno, D., (2010) Coastal phytoplankton do not rest in winter.
1486 *Estuaries and Coasts* 33, 342-361.
- 1487

1488 Table 1. Environmental drivers and ecosystem responses on continental margins: Case studies

1489

Consequences & regional responses (Supplements*)	Drivers of environmental change on continental margins from regional to global scale										Actions taken
	Land use change	Damming, diking, water diversion	Alien species	Anthropogenic nutrients & pollutants	Energy & mineral extractions from the sea	Overfishing	Changes in water cycle	Rising sea level	Warming	Climate change & oscillation	
General consequences	Changes in soil qual. & erosion in watershed	Sed. retention, decr. runoff	Change in ecosystem structure	Eutrophication, change in nutrient ratios & plankton community, HAB, hypoxia	Change in seascape & marine environments	Decr. fish stock, trophic cascades	Extreme weather, flooding, drought	Lowland flooding, salt water intrusion	Incr. stratification, drop in O ₂ , incr. vulnerability	Change in ocean circulation, upwelling,	
American margins											
Cariaco Basin (B1)						Collapse of Spanish sardine (<i>Sardinella aurita</i>)	Freshening of surface waters due to higher regional precipitation		>1°C incr. since 1995	Decr. upwelling, PP and phytoplankton biomass; incr. N* ² , change in ecosystem structure; Increasing zooplankton biomass	
Chesapeake Bay (B2)	Degrading water qual. & clarity	Blocked migratory pathways for anadromous fish	Decreased oyster populations and increased piscivory	Seasonal hypoxia, decline of SAV, fish advisories		extirpation of sturgeon, collapse of oyster fishery	Salinity drop following Tropical Storm Agnes	Rate of sea level rise much higher than average		Change in water exchange rate	Acts for reducing nutrients & restoring SAV, oysters; removing blockages to fish
San Francisco Bay (B3)		Decreased sediment supply, shrinking mudflats, incr. salinity	Decreased phytopl. PP due to Asian clams	Degrading water quality			Drought and water diversion trigger invasion			Change in bio-community	1972 Clean Water Act
Asian margins											
Bohai Sea (B4)	Changes in tidal regime	Massive drop in water discharge and sediment load, fishery collapse in 1990s		High N, low P and Si, decrease in diatom/ dinoflagellate ratio		Decrease in fish biomass; dominant fish species changed from bottom to pelagic fish			0.011oC per year increase during the 1960s-1990s		Artificially controlled water discharge in Yellow River
European margins											
Mediterranean - N. Adriatic Sea (B5)				N/P increase, anoxic events (1970s-1980s), loss of macrobenthos; trend reversed recently.		Loss of demersal fish, small pelagic fish and top predators		Reduced river flow, salinity rise			Mandates of reduction in P loading
North Sea (B6)		Altered mudflat		Massive coastal environment deterioration in 1980s	Massive wind farming				Temporally and regionally faster than global mean, provoked ecosystem shift	NAO state determines circulation mode and nutrient inventories	EU wide mitigation, Marine Strategy Framework Directive

*Note: More descriptions and references about the case studies are presented in electronic supplements listed under each case heading.

Increase in N*: Whether this is due to increased N fixation is being explored

1490
 1491
 1492
 1493
 1494

Table 2. The temporal trends of the environmental variables presented in Fig. 3. All trends are statistically significant except those shown in italics. (For more detailed information on the linear regressions see Supplement A.)

Site	SST (°C yr ⁻¹)	SSS (yr ⁻¹)	Q (km ³ yr ⁻¹)	DIN (Kt yr ⁻¹)	DIP (Kt yr ⁻¹)	Chl (mg m ⁻³ yr ⁻¹)	O ₂ saturation (% yr ⁻¹)
Bohai Sea	<i>0.0022</i>	0.0632	-0.75 ⁽¹⁾				
Cariaco Basin	0.0895	-0.0097				-0.0462	<i>-0.105</i>
East China Sea	0.0282		3.34	26.1		0.0135	-0.536
North Sea	0.0376 ⁽²⁾	<i>0.0007</i>	-7.44	-12.1	-2.3		
SF Bay	-0.0290 ⁽³⁾	<i>0.0427</i>				0.131	-0.290 ⁽³⁾

1495 Notes: (1) Water discharge of the Yellow River that empties into the Bohai Sea. (2) The
 1496 mean temperature of the top 10 m in the North Sea in winter months (DJF). (3) The
 1497 bottom temperature of the USGS Station 18 in the San Francisco Bay.

1498
 1499
 1500

1501 **Figure Captions**

1502

1503 **Figure 1.** Map showing locations of time-series observations that exemplify the interplay
1504 of natural variability, CO₂-driven climate change and human activities to shape margin
1505 ecosystems. Each area is discussed in the manuscript or summarized in Table 1.

1506 1. Arctic Sea; American margins: 2. Cariaco Basin, 3. Chesapeake Bay, 4. Gulf of
1507 Mexico off Louisiana, 5. San Francisco Bay, 6. Southern California Bight; African
1508 margins: 7. Canary Current System, 8. Congo River Submarine Canyon; Asian seas: 9.
1509 Bay of Bengal, 10. Bohai Sea, 11. East China Sea; European seas: 12. Baltic Sea, 13.
1510 Black Sea, 14. Mediterranean - Corsica (Liguran Sea), 15. Mediterranean - N. Adriatic
1511 Sea, 16. North Sea.

1512

1513 **Figure 2.** Data plots of O₂ and pH illustrating time scales of natural hydrographic
1514 variability in the nearshore southern California Bight, USA. (a) Decadal scale suggesting
1515 regime shifts (modified from McClatchie et al., 2010). (b) Interannual scale illustrating
1516 effects of ENSO at a site 6 km from Del Mar (from Nam et al., 2011) (c) Seasonal scale
1517 combining CalCOFI data at line 93 and continuous mooring measurements (from Send
1518 and Nam, 2012), (d) Event (week) scale illustrating upwelling (blue) and relaxation
1519 (green) phases (modified from Send and Nam, 2012), (e) semi-diurnal and diurnal scale
1520 variations in the La Jolla Kelp Forest during upwelling phase (blue) when there are strong
1521 semidiurnal signals and relaxation phase (green) when kelp influences the oxygen and pH
1522 variability (Frieder et al., unpublished).

1523

1524 **Figure 3.** Time series of sea surface temperature (SST) (a), sea surface salinity (SSS) (b),
1525 nutrient loads (c), riverine discharge (d), chlorophyll-*a* (e), and saturation of dissolved
1526 oxygen (f) from the Cariaco Basin and San Francisco Bay (SF Bay) in the Americas, the
1527 East China Sea (ECS) and Bohai Sea (BH) in Asia, and the North Sea (NS) in Europe
1528 (See Fig. 1 for location of time-series stations). SST time series include values obtained
1529 from satellite remote sensing monthly composites (lines without symbols) from NOAA's
1530 National Climatic Data Center (See Supplement A), and *in situ* observations (lines with
1531 symbols). Statistically significant regression ($p < 0.1$) results are shown as solid straight
1532 lines; insignificant ones are shown as dashed lines. (See text).

1533


1534 **Figure 4.** The relationship between nitrogen loading and fisheries landings as a function
1535 of hypoxic area for mobile species in estuaries and semi-enclosed seas. Modified from
1536 Breitburg et al., 2009b.

1537

1538


1539

1540 **Figure 1**
1541



1542
1543


1544 **Figure 2**
 1545
 1546



1547
 1548
 1549
 1550


1551
1552

Figure 3.



1553

1554 **Figure 4.**



1555

1556

1557

1642
1643

Table A1. Locations of margin time series observations and data sources.

Sites	Coordinates	Data sources	Remarks
Bohai Sea	37-41°N, 117-121°E	SSS: Lin et al. (2001) Discharge: Liu et al. (2012); Liu (this issue)	Domain for satellite SST data retrieval
Cariaco Basin	10.2-11.0°N, -64~66°W	SST: See text	Domain for satellite SST data retrieval
CARIACO Sta	10° 30'N 64° 40'W	(http://imars.marine.usf.edu/cariaco/cariaco-ocean-time-series-program)	<i>In situ</i> observations
East China Sea	28-32°N, 121-125°E	SST: See text Discharge, DIN load: Liu et al. (this issue)	Domain for satellite SST data retrieval
North Sea	53~57°N, 4~8.6°E	SST, SSS: (http://www.ices.dk/marine-data/dataset-collections/Pages/default.aspx) accessed on March 31, 2014 Discharge, nutrient loads: Pätzsch and Lenhart (2011)	ICES data
North Sea (L)	53-59°N, -2~8°N	SST: See text	Larger domain for satellite SST data retrieval
San Francisco Bay	38°50.8'N, 121°25.3'W	(http://sfbay.wr.usgs.gov/access/wqdata/index.html)	USGS Sta 18 (Point Blunt) Water depth = 43.0
Coastal zone adjacent to SF Bay	36.2-38.1°N, 122.5-124.7°W	SST: See text	Domain for satellite SST data retrieval

1644
1645

1646
1647

Table A2. Results of linear regression analysis of margin time series data. The statistically insignificant trends, defined as those with $p > 0.1$, are shown in italics.

Variables	Site	Period	Trend (per year)	n	R ²	p
SST (°C)	Bohai Sea	1970-2012	<i>0.0022</i>	43	0.004	0.6830
	Cariaco Basin	1970-2012	0.0303	43	0.548	<0.001
	Cariaco Basin	1995-2012	0.0351	18	0.168	0.0917
	CARIACO Sta	1995-2013	0.0895	198	0.058	<0.001
	East China Sea	1970-2012	0.0282	43	0.438	<0.001
	North Sea (winter)	1976-2013	0.0376	38	0.160	0.0129
	North Sea (L)	1970-2012	0.0301	43	0.493	<0.001
	SF Bay	1990-2013	<i>-0.0126</i>	288	0.002	0.498
	Adj. SF Bay	1970-2012	<i>-0.0087</i>	43	0.053	0.138
	SF Bay bottom water	1990-2013	-0.0290	285	0.011	0.071
SSS	CARIACO Sta	1995-2013	-0.0097	188	0.048	0.0025
	North Sea (winter)	1976-2013	<i>0.0007</i>	38	0.001	0.846
	SF Bay	1990-2013	<i>0.0427</i>	287	0.004	0.300
	Bohai Sea	1960-1999	0.0632	40	0.287	<0.001
Discharge (km ³ yr ⁻¹)	Changjiang	1970-2002	<i>3.34</i>	33	0.068	0.142
	Huanghe	1960-2009	-0.750	40	0.251	<0.001
	North Sea	1977-2010	-7.43	34	0.103	0.064
DIN load (kt yr ⁻¹)	Changjiang	1970-2002	0.0261	33	0.671	<0.001
	North Sea	1977-2010	-0.0121	31	0.601	<0.001
DIP load (Kt yr ⁻¹)	North Sea	1981-2010	-2.28	30	0.596	<0.001
Chl (mg m ⁻³)	CARIACO Sta	1995-2013	-0.0462	201	0.039	0.005
	East China Sea	1997-2010	0.0135	152	0.021	0.073
	SF Bay	1990-2013	0.131	279	0.151	<0.001
O2 satu. (%)	CARIACO Sta	1995-2013	<i>-0.105</i>	180	0.009	0.198
	East China Sea	1981-2006	-0.536	9	0.390	0.072
	SF Bay bottom	1993-2013	-0.290	251	0.042	0.001

1648
1649
1650