

THE
UNIVERSITY
OF RHODE ISLAND

University of Rhode Island
DigitalCommons@URI

Chemistry Faculty Publications

Chemistry

2011

Phenol Nitration Induced by an $\{\text{Fe}(\text{NO})_2\}^{10}$ Dinitrosyl Iron Complex

Nhut Giuc Tran

Harris Kalyvas

See next page for additional authors

Follow this and additional works at: https://digitalcommons.uri.edu/chm_facpubs

This is a pre-publication author manuscript of the final, published article.

Citation/Publisher Attribution

Tran, N. G., Kalyvas, H., Skodje, K. M., Hayashi, T., Moënne-Loccoz, P., Callan, P. E., Shearer, J., Kirschenbaum, L. J., & Kim, E. (2011). Phenol Nitration Induced by an $\{\text{Fe}(\text{NO})_2\}^{10}$ Dinitrosyl Iron Complex. *J. Am. Chem. Soc.*, 133(5), 1184-1187. doi: 10.1021/ja108313u
Available at: <https://doi.org/10.1021/ja108313u>

This Article is brought to you for free and open access by the Chemistry at DigitalCommons@URI. It has been accepted for inclusion in Chemistry Faculty Publications by an authorized administrator of DigitalCommons@URI. For more information, please contact digitalcommons@etal.uri.edu.

Authors

Nhut Giuc Tran, Harris Kalyvas, Kelsey M. Skodje, Takahiro Hayashi, Pierre Moënne-Loccoz, Jason Shearer, Louis J. Kirschenbaum, and Eunsuk Kim

Published in final edited form as:

J Am Chem Soc. 2011 February 9; 133(5): 1184–1187. doi:10.1021/ja108313u.

Phenol Nitration Induced by a $\{\text{Fe}(\text{NO})_2\}^{10}$ Dinitrosyl Iron Complex

 Nhut Giuc Tran[†], Harris Kalyvas[†], Kelsey M. Skodje[†], Takahiro Hayashi[‡], Pierre Moënne-Loccoz[‡], Paige E. Callan[¶], Jason Shearer[¶], Louis J. Kirschenbaum[§], and Eunsuk Kim^{*,†}
[†] Department of Chemistry, Brown University, Providence, Rhode Island 02912

[‡] Division of Environmental and Biomolecular Systems, Oregon Health & Science University, Beaverton, Oregon 97006

[¶] Department of Chemistry, University of Nevada, Reno, Nevada 89557

[§] Department of Chemistry, University of Rhode Island, Kingston, Rhode Island 02881

Abstract

Cellular dinitrosyl iron complexes (DNICs) have long been considered NO carriers. Although other physiological roles of DNICs have been postulated, their chemical functionality outside of NO transfer has not been demonstrated thus far. Here we report unprecedented dioxygen reactivity of an N-bound $\{\text{Fe}(\text{NO})_2\}^{10}$ DNIC, $[\text{Fe}(\text{TMEDA})(\text{NO})_2]$ (**1**). In the presence of O₂, **1** becomes a nitrating agent that converts 2,4-di-*tert*-butylphenol to 2,4-di-*tert*-butyl-6-nitrophenol via formation of a putative iron-peroxynitrite $[\text{Fe}(\text{TMEDA})(\text{NO})(\text{ONOO})]$ (**2**) that is stable below –80 °C. Iron K-edge X-ray absorption spectroscopy on **2** supports a five-coordinated metal center with a bound peroxynitrite in a cyclic bidentate fashion. The peroxynitrite ligand of **2** readily decays at increased temperature or under illumination. These results suggest that DNICs could have multiple physiological or deleterious roles, including that of cellular nitrating agents.

Dinitrosyl iron complexes (DNICs), Chart 1, are naturally occurring iron species that are generated from the reactions of nitric oxide (NO) with cellular nonheme iron species such as iron-sulfur clusters.^{1,2} A series of S- or N-bound DNICs have been reported^{2–6} since the initial discovery⁷ of a cysteine-bound DNIC formulated as $[\text{Fe}(\text{NO})_2(\text{SR})_2]^-$, an $\{\text{Fe}(\text{NO})_2\}^9$ species in the Enemark-Feltham notation.⁸ Although EPR-active S-bound $\{\text{Fe}(\text{NO})_2\}^9$ DNICs are more common, N- or O-bound DNICs have been also observed.⁹

Several physiological roles of DNICs have been suggested, including storage and transfer of NO.^{1,10} However, the nature of the chemistry that allows DNICs to play these physiological roles is not well understood. A notable recent report from the Lippard group suggests that the NO-donating ability of an N-bound DNIC might be $\{\text{Fe}(\text{NO})_2\}^9/\{\text{Fe}(\text{NO})_2\}^{10}$ redox-dependent.⁶ While such a proposal offers important chemical insights, we were intrigued by the highly reducing nature of $\{\text{Fe}(\text{NO})_2\}^{10}$ DNICs, which may suggest a rich chemistry between these motifs and oxidants such as O₂. Such reactivity could shed light on undiscovered physiological or deleterious roles of DNICs. Herein, we describe unprecedented dioxygen reactivity of an N-bound $\{\text{Fe}(\text{NO})_2\}^{10}$ DNIC, $[\text{Fe}(\text{TMEDA})(\text{NO})_2]$ (**1**), where TMEDA = *N,N,N',N'*-tetramethylethylenediamine (Chart 1), and we demonstrate the formation of an intermediate that leads to nitration of phenol.

Eunsuk_Kim@brown.edu.

 Supporting Information Available: Experimental details concerning spectroscopy, reaction product characterization and quantification, isotope labeling, electrochemistry, and nitrite detection. This material is available free of charge via the Internet at <http://pubs.acs.org>.

Compound **1** was synthesized as previously reported by Hung *et al.*^{5d} Bubbling of O₂ through a solution of **1** in dichloromethane at -80 °C leads to the formation of an EPR-silent, dark purple complex with absorption bands at 460 ($\epsilon = 420 \text{ M}^{-1}\text{cm}^{-1}$) and 560 nm ($\epsilon = 580 \text{ M}^{-1}\text{cm}^{-1}$), Figure 1A. This purple complex is stable below -80 °C, but it decomposes to an orange, insoluble precipitate upon warming.^{11,12} IR monitoring of this O₂-reaction also shows the generation of a new quasi-stable species with the appearance of two new ν_{NO} peaks at 1589 and 1805 cm^{-1} concomitant with the disappearance of the ν_{NO} of **1** at 1630 and 1687 cm^{-1} (Figure 1B).¹³ No other significant change occurs in the mid-frequency region of the IR spectra. This transformation is very different from what was observed by Tonzetich *et al.*⁶ in the case of another N-bound {Fe(NO)₂}¹⁰ DNIC with a monoanionic bidentate β -diketimate ligand. In that study, air exposure led to the one-electron oxidation of the {Fe(NO)₂}¹⁰ DNIC to an EPR-active {Fe(NO)₂}⁹ DNIC, which was also signified by >130 cm^{-1} upshifts of the two IR-active NO stretching frequencies. Here, the intermediate generated from **1**/O₂ is EPR silent and it exhibits an up- (1805 cm^{-1}) and a downshifted (1589 cm^{-1}) NO band. These data are inconsistent with the simple one-electron oxidation of **1**.¹⁴ The possibility of generating a five-coordinate superoxide (O₂⁻) adduct, [Fe(TMEDA)(NO)₂(O₂)], has been considered. Although such a species is not known, an iodide (I⁻) analogue, [Fe(TMEDA)(NO)₂(I)], has been reported to have two upshifted ν_{NO} frequencies higher than 1700 cm^{-1} ,¹⁵ which is again different from what we observe in the **1**/O₂ reaction.¹⁶ Interestingly, the ν_{NO} at 1589 cm^{-1} closely matches those from known *trans*-peroxynitrite species, O=NOOM, where M = Li, Na, K.¹⁷ These IR and EPR characteristics, along with the EXAFS data (*vide infra*), led us to consider the intermediate to be a peroxynitrite bound iron mononitrosyl species, [Fe(TMEDA)(NO)(ONOO)] (**2**).¹⁸

Iron K-edge X-ray absorption spectroscopy was used to further probe complex **2**. Comparison of the edge energies of **1** and **2** shows a shift by +1.8(4) eV upon O₂ exposure to cold solutions of **1** (Figure 2A), which is consistent with the formal oxidation of **1** by 1 electron. The Fe 1(s) \rightarrow 3(d) transition found in the pre-edge region of **2** has a peak area slightly larger than that corresponding to **1** (35(1) vs. 36(1) eV% relative to the edge height). This is most consistent with **2** having an Fe-center contained in a non-centrosymmetric coordination environment. The EXAFS region for **2** is best modeled as a five coordinate Fe-species with a coordinated bidentate O₂NO⁻ moiety (Figure 2B). We find strong multiple-scattering (MS) pathways originating from this cyclic O₂NO⁻ ligand (average N/O distance 1.91(1) Å; see inset Figure 2B). In addition, strong MS pathways are also found originating from the NO ligand (1.67(1) Å). Two additional N-scatters derived from the TMEDA ligand are also observed. The thermal decomposition product resulting from warming of **2** to room temperature yields a further shift in the edge energy of +2.6(3) eV, consistent with the production of an Fe^{2+/3+} species (Figure 2A).

Attempts at characterization of **2** by resonance Raman spectroscopy were hampered by photosensitivity. The decomposition of **2** with white light was monitored at -90 °C by UV-Vis spectroscopy but it failed to reveal new absorption features (Figure S5). In the IR, continuous white-light illumination of **2** leads to the progressive and irreversible bleaching of the 1581 and 1807 cm^{-1} bands with concomitant appearance of a single band at 1744 cm^{-1} (Figure 3). The loss of the IR band at 1581 cm^{-1} and the appearance of a new ν_{NO} at 1744 cm^{-1} suggest that illumination of **2** might lead to the formation of a new iron-mononitrosyl species after photochemistry at the peroxynitrite ligand. Like **2**, the photoproduct is EPR silent at 10 K (data not shown). While the nature of the photoproduct remains in doubt,¹⁹ the photoactive behavior of **2** is reminiscent of [Co(CN)₅(ONOO)]³⁻, where photolysis destroys the coordinated ONOO⁻.²⁰

The presence of peroxyxynitrite is often indicated by oxidation and/or nitration chemistry especially with phenolic substrates.²¹ When one equiv. of 2,4-di-*tert*-butylphenol (DBP) is added to **2** and the reaction mixture is subsequently warmed to room temperature, 2,4-di-*tert*-butyl-6-nitrophenol (NO₂-DBP) is observed along with the oxidative coupling product 2,2'-dihydroxy-3,3',5,5'-tetra-*t*-butyl-1,1'-biphenyl (Scheme 1).²² These reaction products do not form when DBP is added *after* warming the solution of **2** to room temperature, signifying that intermediate **2** is a crucial species in phenol nitration and oxidation. At room temperature, **2** is too unstable to be monitored by regular UV-Vis spectroscopy, although a glimpse of purple color can be seen momentarily. Despite the short lifetime of **2** at room temperature, when O₂ is added to a mixture of **1** and DBP (1 equiv) at room temperature nitration chemistry still occurs, yielding NO₂-DBP, while only a small amount (2%) of the oxidative coupling product is generated. This indicates that **1**/O₂ induces nitration much more efficiently than oxidation at room temperature.²³ When ¹⁸O₂ is used, approximately 50 % of ¹⁸O atom incorporates into the substrate. GC-MS analysis of NO₂-DBP displays a distribution of ¹⁸O₂N-DBP, ^{18/16}O₂N-DBP, and ¹⁶O₂N-DBP in a ratio of 30(2)%, 48(1)%, and 22(2)%.¹² Although the mechanistic investigation of phenol nitration by **2** is beyond the scope of this report, the O atom isotope distribution in NO₂-BDP warrants future mechanistic studies and likely involves the cleavage of peroxyxynitrite and participation of the other NO.²⁴

Nitration of biological phenols, such as seen in protein tyrosine nitration (PTN), is an important posttranslational modification associated with various pathological conditions including inflammatory, neurodegenerative, and cardiovascular diseases.²⁵ Although elusive, the current view^{25d} of PTN suggests that different types of cellular nitrating agents could be responsible for its specificity at various sites. Two major ways to generate PTN are known.²⁵ One is through peroxyxynitrite (ONOO⁻) that is formed from nitric oxide (NO) and superoxide (O₂⁻). The other involves reactions of heme peroxidases with hydrogen peroxide and nitrite (NO₂⁻). The reactivity of the {Fe(NO)₂}¹⁰ DNIC we report herein suggests that cellular DNICs could provide a new route to generate PTN; the DNIC derived peroxyxynitrite moiety in **2** may directly nitrate the phenol or biological tyrosine (via homolytic cleavage to •O(H) + •NO₂) or act as a •NO₂ generator (where two equiv. may lead to ArOH nitration).^{21,25} The results also agree with several literature examples of the oxidation chemistry of metal-nitrosyls,²⁶ though observation of metal-peroxyxynitrite is rare.^{26d,e} It is conceivable that small molecule metal species such as DNICs act as mobile nitrating agents in cells.

In summary, we have described unprecedented O₂ reactivity of an {Fe(NO)₂}¹⁰ iron-dinitrosyl complex [Fe(TMEDA)(NO)₂] (**1**). In the presence of O₂, **1** becomes a potent nitrating agent via formation of a putative iron-peroxyxynitrite species. The O₂ reactivity of **1** demonstrated here suggests that the physiological functions of DNICs are not limited to NO storage and transfer and deserve further studies.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported by Brown University (E.K.), the Camille and Henry Dreyfus New Faculty Award Program (E.K.), NIH (P.M.-L., GM74785; J.S. P20 RR-016464), NSF (J.S. CH-0844623), and Vertex pharmaceutical scholarship (T.H.). E.K. thanks Mr. Andrew Dineen at the Charles Stark Draper Laboratory for the initial DFT calculations. Work performed at the NSLS was supported by the DOE (98CH1086)

References

1. Vanin AF. Nitric Oxide. 2009; 21:1–13. [PubMed: 19366636]
2. Butler AR, Megson IL. Chem Rev. 2002; 102:1155–1166. [PubMed: 11942790]
3. McCleverty JA. Chem Rev. 2004; 104:403–418. [PubMed: 14871130]
4. (a) Chiang CY, Miller ML, Reibenspies JH, Darensbourg MY. J Am Chem Soc. 2004; 126:10867–10874. [PubMed: 15339171] (b) Chiang CY, Darensbourg MY. J Biol Inorg Chem. 2006; 11:359–370. [PubMed: 16520978] (c) Reginato N, McCrory CTC, Pervitsky D, Li L. J Am Chem Soc. 1999; 121:10217–10218. (d) Wang JH, Chen CH. Inorg Chem. 2010; 49:7644–7646. [PubMed: 20687531]
5. Selected examples from Liaw's group include (a) Tsai ML, Chen CC, Hsu IJ, Ke SC, Hsieh CH, Chiang KA, Lee GH, Wang Y, Chen JM, Lee JF, Liaw WF. Inorg Chem. 2004; 43:5159–5167. [PubMed: 15285693] (b) Tsai FT, Chiou SJ, Tsai MC, Tsai ML, Huang HW, Chiang MH, Liaw WF. Inorg Chem. 2005; 44:5872–5881. [PubMed: 16060642] (c) Tsai ML, Liaw WF. Inorg Chem. 2006; 45:6583–6585. [PubMed: 16903707] (d) Hung MC, Tsai MC, Lee GH, Liaw WF. Inorg Chem. 2006; 45:6041–6047. [PubMed: 16842012] (e) Tsai ML, Hsieh CH, Liaw WF. Inorg Chem. 2007; 46:5110–5117. [PubMed: 17444639] (f) Lu TT, Chen CH, Liaw WF. Chem Eur J. 2010; 16:8088–8095.
6. Tonzetich ZJ, Do LH, Lippard SJ. J Am Chem Soc. 2009; 131:7964–7965. [PubMed: 19459625]
7. (a) Commoner B, Ternberg JL. Proc Natl Acad Sci USA. 1961; 47:1374–1384. [PubMed: 13694929] (b) Vithayathil AJ, Ternberg JL, Commoner B. Nature. 1965; 207:1246–1249. [PubMed: 4287143] (c) Mallard JR, Kent M. Nature. 1964; 204:1192. [PubMed: 14264374] (d) Vanin AF, Nalbandian RM. Biofizika. 1965; 10:167–168. [PubMed: 14332030]
8. Enemark JH, Feltham RD. Coord Chem Rev. 1974; 13:339–406.
9. (a) Woolum JC, Tiezzi E, Commoner B. Biochim Biophys Acta. 1968; 160:311–320. [PubMed: 4300590] (b) D'Autreaux B, Horner O, Oddou JL, Jeandey C, Gambarelli S, Berthomieu C, Latour JM, Michaud-Soret I. J Am Chem Soc. 2004; 126:6005–6016. [PubMed: 15137765] (c) Cesareo E, Parker LJ, Pedersen JZ, Nuccetelli M, Mazzetti AP, Pastore A, Federici G, Caccuri AM, Ricci G, Adams JJ, Parker MW, Lo Bello M. J Biol Chem. 2005; 280:42172–42180. [PubMed: 16195232]
10. (a) Boese M, Mordvintcev PI, Vanin AF, Busse R, Mülsch A. J Biol Chem. 1995; 270:29244–29249. [PubMed: 7493954] (b) Ueno T, Suzuki Y, Fujii S, Vanin AF, Yoshimura T. Biochem Pharmacol. 2002; 63:485–93. [PubMed: 11853699] (c) Chen YJ, Ku WC, Feng LT, Tsai ML, Hsieh CH, Hsu WH, Liaw WF, Hung CH, Chen YJ. J Am Chem Soc. 2008; 130:10929–10938. [PubMed: 18661983]
11. The presence of nitrite (NO_2^-), Fe, and TMEDA in the precipitate has been qualitatively confirmed.¹² Further speculation concerning the nature of decomposition product(s) is not warranted.
12. See Supporting Information.
13. Synthesis of the intermediate using ^{15}NO shows clear IR shift to lower energy supporting the ν_{NO} assignments and formation of putative $[\text{Fe}(\text{TMEDA})(\text{NO})(\text{ONOO})]$ (**2**) (vide infra).¹²
14. Reversible $\{\text{Fe}(\text{NO})_2\}_9/10$ redox behavior of **1** is observed in cyclic voltammetry, with $E_{1/2} = -0.527$ V (vs. ferrocene/ferrocenium). Chemical oxidation of **1** by I_2 has been shown to yield a five-coordinate compound $[\text{Fe}(\text{TMEDA})(\text{NO})_2\text{I}]$.¹⁵ Attempts to isolate a four-coordinate counterpart, $[\text{Fe}(\text{TMEDA})(\text{NO})_2]^+$, were not successful probably due to its strong preference to be five-coordinate, as was previously discussed.¹⁵
15. $\nu_{\text{NO}} = 1773$ and 1719 cm^{-1} in diethyl ether. Chen CH, Ho YC, Lee GH. J Organomet Chem. 2009; 694:3395–3400.
16. Even more evidence to disfavor a superoxide adduct formation includes the lack of an isotope sensitive $\nu_{\text{O-O}}$ in the region of ~ 1100 cm^{-1} (Figure 1b) and the inconsistent EXAFS data for such a model. See Supporting Information for alternative EXAFS fitting results.
17. $\nu_{\text{NO}} = 1528$ – 1581 cm^{-1} for trans-MOONO, where M = K, Na, and Li. Lo WJ, Lee YP, Tsai JHM, Tsai HH, Hamilton TP, Harrison JG, Beckman JS. J Chem Phys. 1995; 103:4026–4034.

18. The two ν_{NO} of **2** at 1589 and 1805 cm^{-1} remain unchanged upon $^{18}\text{O}_2$ substitution. Characterization of **2** by ESI-MS was attempted without success due to the insufficient stability of **2**.
19. The EPR silent character of the photoproduct suggests a diamagnetic or integer spin species. This electronic structure may reflect the iron-mononitrosyl complex or the overall spin of this complex coupled with the nearby product of the photolyzed peroxyxynitrite. For an analogous situation in a metalloprotein, see the EPR studies $\{\text{FeNO}\}^6$ myoglobin complexes, where the photolyzed NO ($S=1/2$) remains spin-coupled with the high-spin iron(III) ($S=5/2$). Hori H, Ikeda-Saito M, Lang G, Yonetani T. *J Biol Chem.* 1990; 265:15028–15033. [PubMed: 2168399] Hori H, Masuya F, Dou Y, Ikeda-Saito M. *J Inorg Biochem.* 2000; 82:181–187. [PubMed: 11132625]
20. Wick PK, Kissner R, Koppenol WH. *Helv Chim Acta.* 2001; 84:3057–3062.
21. (a) Goldstein S, Lind J, Merenyi G. *Chem Rev.* 2005; 105:2457–2470. [PubMed: 15941219] (b) Gunaydin H, Houk KN. *Chem Res Toxicol.* 2009; 22:894–898. [PubMed: 19374346] (c) Ferrer-Sueta G, Radi R. *ACS Chem Biol.* 2009; 4:161–177. [PubMed: 19267456]
22. Future investigations include the reaction with Tyr as the substrate using water-soluble analogs of **1**.
23. This may also indicate that the oxidative coupling product can be generated from other low temperature stable species besides **2**.
24. We note that in a study of iron peroxycarbonate chemistry, extensive oxygen scrambling occurs. Furutachi H, Hashimoto K, Nagatomo S, Endo T, Fujinami S, Watanabe Y, Kitagawa T, Suzuki M. *J Am Chem Soc.* 2005; 127:4550–4551. [PubMed: 15796501]
25. (a) Pacher P, Beckman J, Liaudet L. *Physiol Rev.* 2007; 87:315–424. [PubMed: 17237348] (b) Szabó C, Ischiropoulos H, Radi R. *Nat Rev Drug Discov.* 2007; 6:662–680. [PubMed: 17667957] (c) Radi R. *Proc Natl Acad Sci U S A.* 2004; 101:4003–4008. [PubMed: 15020765] (d) Abello NA, Kerstjens HAM, Postma DS, Bischoff R. *J Proteome Res.* 2009; 8:3222–3238. [PubMed: 19415921]
26. (a) Roncaroli F, Videla M, Slep LD, Olabe JA. *Coord Chem Rev.* 2007; 251:1903–1930. (b) Herold S, Koppenol WH. *Coord Chem Rev.* 2005; 249:499–506. (c) Ford PC, Lorkovic I. *Chem Rev.* 2002; 102:993–1017. [PubMed: 11942785] (d) Arnold EV, Bohle DS. *Methods Enzymol.* 1996; 269:41–55. [PubMed: 8791636] (e) Park GY, Deepalatha S, Puiu SC, Lee DH, Mondal B, Narducci Sarjeant AA, del Rio D, Pau MY, Solomon EI, Karlin KD. *J Biol Inorg Chem.* 2009; 14:1301–11. [PubMed: 19662443]




Figure 1. UV-Vis (A) and IR (B) spectra of $[\text{Fe}(\text{TMEDA})(\text{NO})_2]$ (**1**) (black dashed line), and of the putative intermediate $[\text{Fe}(\text{TMEDA})(\text{NO})(\text{ONOO})]$ (**2**) (red) at -80°C in dichloromethane.




Figure 2.
 A) XANES region of the XAS for **1** (green), **2** (red) and the thermal decomposition product (black). B) Experimental (red solid line) and simulated (blue dashed line) magnitude FT k^3 EXAFS data for **2**. Best fit includes: shell #1: 2 N scatterers, $r = 2.14(1)$ Å, $\sigma^2 = 0.003(2)$ Å²; O₂NO shell: 1 O₂NO scatterer, $r_1 = 1.91(1)$ Å, $r_2 = 1.91$ Å (restrained), $\sigma^2 = 0.002(1)$ Å², $\phi = 98(7)^\circ$; NO shell: 1 NO scatterer, $r = 1.67(1)$ Å; $\sigma^2 = 0.008(2)$ Å²; $\theta = 192(2)^\circ$. $E_0 = 7120.4$ eV. $\epsilon^2 = 0.57$.





Figure 3. IR spectra of a $[\text{Fe}(\text{TMEDA})(\text{NO})(\text{ONOO})]$ (**2**) in CH_2Cl_2 at 11 K before (black) and after illumination (red). The dark minus illuminated difference spectrum is also shown (blue). Sharp IR bands are from dichloromethane.



Scheme 1.




Chart 1.
Dinitrosyl iron complexes (DNICs)