

THE  
UNIVERSITY  
OF RHODE ISLAND

University of Rhode Island  
DigitalCommons@URI

---

Biomedical and Pharmaceutical Sciences Faculty  
Publications

Biomedical and Pharmaceutical Sciences

---

2007

# Protein pyrophosphorylation by inositol pyrophosphates is a posttranslational event

Rasha Bhandari

Adolfo Saiardi

*See next page for additional authors*

Follow this and additional works at: [https://digitalcommons.uri.edu/bps\\_facpubs](https://digitalcommons.uri.edu/bps_facpubs)

Terms of Use

All rights reserved under copyright.

---

## Citation/Publisher Attribution

Bhandari, R., Saiardi, A., Ahmadibeni, Y., Snowman, A. M., Resnick, A. C., Kristiansen, T. Z., Molina, H.,...Snyder, S. H. (2007). Protein pyrophosphorylation by inositol pyrophosphates is a posttranslational event. *Proceedings of the National Academy of Sciences*, 104(39), 15305-15310. doi: 10.1073/pnas.0707338104  
Available at: <https://doi.org/10.1073/pnas.0707338104>

This Article is brought to you for free and open access by the Biomedical and Pharmaceutical Sciences at DigitalCommons@URI. It has been accepted for inclusion in Biomedical and Pharmaceutical Sciences Faculty Publications by an authorized administrator of DigitalCommons@URI. For more information, please contact [digitalcommons@etal.uri.edu](mailto:digitalcommons@etal.uri.edu).

---

**Authors**

Rasha Bhandari, Adolfo Saiardi, Yousef Ahmadibeni, Adele M. Snowman, Adam C. Resnick, Troels Z. Kristiansen, Henrik Molina, Akhilesh Pandey, J. Kent Werner Jr., Krishna R. Juluri, Yong Xu, Glenn D. Prestwich, Keykavous Parang, and Solomon H. Snyder


# Correction

## BIOCHEMISTRY


Correction for "Protein pyrophosphorylation by inositol pyrophosphates is a posttranslational event," by Rashna Bhandari, Adolfo Saiardi, Yousef Ahmadibeni, Adele M. Snowman, Adam C. Resnick, Troels Z. Kristiansen, Henrik Molina, Akhilesh Pandey, J. Kent Werner, Jr., Krishna R. Juluri, Yong Xu, Glenn D. Prestwich, Keykavous Parang, and Solomon H. Snyder, which appeared in issue 39, September 25, 2007, of *Proc Natl Acad Sci USA* (104:15305–15310; first published September 14, 2007; 10.1073/pnas.0707338104).

The authors note that Figs. 1*b*, 1*d*, 1*g*, and 4*a* have been revised to include dividing lines between lanes to show where nonessential


lanes were removed from a single original gel. Figs. 2*c*, 2*d*, and 3*c* have been revised to include boxes to indicate samples run on different gels, under the same conditions; these gels are aligned with respect to the indicated molecular weight markers. Figs. 4*b* and 4*c* have been revised to eliminate errors that occurred in the original version during file conversion. The changes were made to comply with the PNAS policy that requires dividing lines whenever entire nonessential lanes have been removed from a single original gel, and clear demarcation of samples run on separate gels. These changes do not affect the data presented nor the conclusions of the article. The corrected figures and their legends appear below.




**Fig. 1.**  $\text{IP}_7$ -mediated protein phosphorylation requires prephosphorylation by protein kinases. (a) NSR1 and SRP40 purified from *E. coli* or *S. cerevisiae* were incubated with  $5\beta[^{32}\text{P}]\text{IP}_7$  and resolved by NuPAGE; immunoblotting with a tag-specific antibody (Left) and autoradiography to determine phosphorylation (Right). (b) GST or GST-tagged NSR1 fragment (amino acids 51–166) expressed in *E. coli* and immobilized on glutathione beads were preincubated with or without *S. cerevisiae* extract or ATP, washed with PBS, and then treated with  $5\beta[^{32}\text{P}]\text{IP}_7$  and resolved by NuPAGE; Coomassie brilliant blue R250 staining (Left) and autoradiography (Right). The extra phosphorylated protein bands in lanes 3–6 are yeast proteins that bound nonspecifically to glutathione beads. (c) GST-SRP40 purified from *E. coli* was preincubated with ATP and either native *S. cerevisiae* extract (N) or extract that had been boiled for 5 min (B) and then treated with  $5\beta[^{32}\text{P}]\text{IP}_7$ , as in b. (d) GST-SRP40 purified from *E. coli* was preincubated with yeast extract and indicated nucleotides and then phosphorylated by  $5\beta[^{32}\text{P}]\text{IP}_7$ ; Coomassie brilliant blue R250 staining (Lower) and autoradiography (Upper). (e) GST or GST-tagged NSR1 fragment (amino acids 1–50) purified from *E. coli* were preincubated with or without CK2 and ATP and then treated with  $5\beta[^{32}\text{P}]\text{IP}_7$ , as in b. (f) GST-tagged NSR1 fragment (amino acids 27–50) purified from *E. coli* was incubated without or with CK2 and ATP (lanes 1 and 2) or coexpressed in *E. coli* with the catalytic A1 subunit of human CK2 (lane 3). Purified proteins were incubated with  $5\beta[^{32}\text{P}]\text{IP}_7$ , as in b. (g) GST or GST-tagged NSR1 fragments (amino acids 1–50 and 51–166) purified from *S. cerevisiae*, were preincubated in the absence or presence of  $\lambda$ -phosphatase and then phosphorylated by  $5\beta[^{32}\text{P}]\text{IP}_7$ , as in b.



**Fig. 2.** IP<sub>7</sub> phosphorylates a phosphoserine residue to generate pyrophosphoserine. (a) Extracts from *S. cerevisiae* expressing GST-tagged mouse Nopp140 fragment (amino acids 80–100) WT sequence and indicated point mutants were phosphorylated by 5β[<sup>32</sup>P]IP<sub>7</sub>; autoradiography (Upper) and immunoblotting (Lower). (b) Sequences of individual peptides derived from Nopp140 (GenBank accession no. NP\_941035). NOP, amino acids 76–100 of mouse Nopp140; NOP-2 [pS], the same sequence as NOP except with phosphoserine at positions 88 and 91; NOP-pS 88 and NOP-pS 91, single phosphoserine residues at positions 88 and 91 respectively; NOP-2[MepS], methylphosphoserine at positions 88 and 91; NOP-2[ppS], pyrophosphoserine at positions 88 and 91. (c and d) The six synthetic peptides in b were incubated with 5β[<sup>32</sup>P]IP<sub>7</sub> (c) or with CK2 and γ[<sup>32</sup>P]ATP (d) and resolved by NuPAGE. Coomassie G250 staining (Left) and autoradiography (Right). In each peptide lane, there are multiple bands in the 3- to 9-kDa range, corresponding to peptide monomers and multimers, generated possibly because of electrostatic interactions between the positively charged lysines and negatively charged acidic and phosphate-containing residues (27). Peptide dimers were also observed by MALDI-TOF analysis (data not shown).



**Fig. 3.** The properties of pyrophosphoserine differ from those of phosphoserine. (a and b) NOP-2[pS] peptide was phosphorylated by CK2 and γ[<sup>32</sup>P]ATP (a) or by 5β[<sup>32</sup>P]IP<sub>7</sub> (b), then treated with HCl and subjected to NuPAGE; Coomassie G250 staining (Left) and autoradiography (Right). (c) NOP-2[pS] peptide was phosphorylated by either CK2 and γ[<sup>32</sup>P]ATP or by 5β[<sup>32</sup>P]IP<sub>7</sub> as described in *SI Materials and Methods*. A fraction of the phosphorylated peptides was subjected to NuPAGE to ensure equivalent phosphorylation under both conditions; Coomassie G250 staining (Left) and autoradiography (Right). (d and e) The remaining CK2 phosphorylated peptide (d) and IP<sub>7</sub>-phosphorylated peptide (e) were hydrolyzed with 6 M HCl, and the resulting amino acids were resolved by HPLC. Data collected during the first 40 min of the HPLC run are presented; elution profiles of the first 3 aa (Upper) and corresponding radioactivity in each 1-ml fraction (Lower). Note that high levels of <sup>32</sup>P<sub>i</sub> in the IP<sub>7</sub>-phosphorylated sample compared with the CK2 and ATP phosphorylated sample are due to incomplete removal of 5β[<sup>32</sup>P]IP<sub>7</sub> from the peptide during desalting (see *SI Materials and Methods*).



**Fig. 4.**  $\lambda$ -phosphatase does not dephosphorylate IP<sub>7</sub>-phosphorylated proteins. (a) GST-tagged NSR1 fragment (amino acids 1–50) purified from *S. cerevisiae* was incubated without or with  $\lambda$ -phosphatase and then phosphorylated by 5β[<sup>32</sup>P]IP<sub>7</sub> (lanes 1 and 2). Alternatively, the protein was first phosphorylated by 5β[<sup>32</sup>P]IP<sub>7</sub> and then incubated without or with  $\lambda$ -phosphatase (lanes 3 and 4). The samples were resolved by NuPAGE and autoradiographed. (b) GST-tagged NSR1 fragment (amino acids 27–50) purified from *S. cerevisiae* was phosphorylated either by CK2 and γ[<sup>32</sup>P]ATP (Left) or by 5β[<sup>32</sup>P]IP<sub>7</sub> (Right) and then incubated with or without  $\lambda$ -phosphatase; Coomassie G250 staining (Left) and autoradiography (Right). (c) NOP-2[pS] peptide bound to streptavidin agarose beads was phosphorylated by either CK2 and γ[<sup>32</sup>P]ATP (Left) or by 5β[<sup>32</sup>P]IP<sub>7</sub> (Right) and then incubated with or without  $\lambda$ -phosphatase as in b. (d) *S. cerevisiae* expressing GST-tagged Nopp140 fragment (amino acids 80–100) (Left) or NSR1 fragment (amino acids 27–50) (Right) were labeled with inorganic orthophosphate [<sup>32</sup>PO<sub>4</sub>]<sub>i</sub>. Purified radiolabeled proteins were incubated with or without  $\lambda$ -phosphatase; autoradiography (Upper) and immunoblotting (Lower). (e) NOP-2[pS] peptide (Left) or NOP-2[ppS] peptide (Right) were treated with or without  $\lambda$ -phosphatase, resolved by NuPAGE, and stained with either Coomassie G250 or Pro-Q Diamond phosphoprotein gel stain, as indicated.

www.pnas.org/cgi/doi/10.1073/pnas.1218152109

# Protein pyrophosphorylation by inositol pyrophosphates is a posttranslational event

Rashna Bhandari\*, Adolfo Saiardi<sup>†</sup>, Yousef Ahmadibeni<sup>‡</sup>, Adele M. Snowman\*, Adam C. Resnick\*<sup>§</sup>, Troels Z. Kristiansen<sup>¶</sup>, Henrik Molina<sup>¶</sup>, Akhilesh Pandey<sup>¶</sup>, J. Kent Werner, Jr.\*<sup>¶</sup>, Krishna R. Juluri\*, Yong Xu<sup>||</sup>, Glenn D. Prestwich<sup>||</sup>, Keykavous Parang<sup>‡</sup>, and Solomon H. Snyder\*<sup>\*,\*\*††‡‡§§¶¶||</sup>

\*The Solomon H. Snyder Department of Neuroscience, Departments of \*\*Pharmacology and Molecular Sciences, ††Psychiatry and Behavioral Sciences, and †‡McKusick–Nathans Institute of Genetic Medicine and Department of Biological Chemistry, Johns Hopkins University School of Medicine, 725 North Wolfe Street, Baltimore, MD 21205; †Medical Research Council (MRC) Cell Biology Unit and Laboratory for Molecular Cell Biology, Department of Biochemistry and Molecular Biology, University College London, Gower Street, London WC1E 6BT, United Kingdom; ‡Department of Biomedical and Pharmaceutical Sciences, College of Pharmacy, University of Rhode Island, Kingston, RI 02881; and †‡Department of Medicinal Chemistry, University of Utah, 419 Wakara Way, Suite 205, Salt Lake City, UT 84108

Contributed by Solomon H. Snyder, August 3, 2007 (sent for review July 16, 2007)

In a previous study, we showed that the inositol pyrophosphate diphosphoinositol pentakisphosphate (IP<sub>7</sub>) physiologically phosphorylates mammalian and yeast proteins. We now report that this phosphate transfer reflects pyrophosphorylation. Thus, proteins must be prephosphorylated by ATP to prime them for IP<sub>7</sub> phosphorylation. IP<sub>7</sub> phosphorylates synthetic phosphopeptides but not if their phosphates have been masked by methylation or pyrophosphorylation. Moreover, IP<sub>7</sub> phosphorylated peptides are more acid-labile and more resistant to phosphatases than ATP phosphorylated peptides, indicating a different type of phosphate bond. Pyrophosphorylation may represent a novel mode of signaling to proteins.

inositol polyphosphate | protein phosphorylation

Inositol phosphates have diverse biologic roles with the best known, inositol-1,4,5-trisphosphate (IP<sub>3</sub>), releasing intracellular calcium (1). Inositol pyrophosphates (also designated inositol diphosphates), exemplified by diphosphoinositol pentakisphosphate (5-PP-I(1,2,3,4,6)P<sub>5</sub>, hereafter called IP<sub>7</sub>) and bis-diphosphoinositol tetrakisphosphate ([PP]<sub>2</sub>-IP<sub>4</sub>, IP<sub>8</sub>) (2–4) are formed by a group of three inositol hexakisphosphate (IP<sub>6</sub>) kinases (IP6Ks) (5–7). IP6K and inositol pyrophosphates participate in diverse physiologic activities including endocytosis (8, 9), apoptosis (10, 11), chemotaxis (12), and telomere elongation (13, 14). Recently, a novel inositol pyrophosphate, 4/6-PP-IP<sub>5</sub>, synthesized by the Vip1 protein in *Saccharomyces cerevisiae*, was identified by York and associates (15, 16) and shown to be involved in regulating cell growth, morphology, and phosphate homeostasis in yeast.

The pyrophosphate bond of IP<sub>7</sub> has a calculated phosphorylation potential that equals or exceeds that of ATP, suggesting that it could serve a similar function (4, 17). We demonstrated that IP<sub>7</sub> physiologically phosphorylates a variety of protein targets in yeast and mammals (18). Unlike phosphorylation by ATP, IP<sub>7</sub> does not require separate protein kinases but directly phosphorylates its targets. We wondered how signaling by IP<sub>7</sub> phosphorylation is distinguished from phosphorylation by ATP. We now report that protein phosphorylation by IP<sub>7</sub> involves a novel modification, pyrophosphorylation, which may provide a unique mode of signaling to proteins.

## Results and Discussion

In our earlier study, we showed that 5β[<sup>32</sup>P]IP<sub>7</sub> phosphorylates a wide range of proteins in both the yeast, *S. cerevisiae*, and in mammalian tissues (18). We characterized in detail phosphorylation of several of these proteins, such as the yeast proteins NSR1 and SRP40. To obtain large amounts of these proteins, we expressed them in *Escherichia coli* (Fig. 1a). Whereas these proteins purified from yeast are robustly phosphorylated by 5β[<sup>32</sup>P]IP<sub>7</sub>, the corresponding proteins purified from bacteria display no phosphorylation. What property of yeast but not

bacterial extracts conveys the capacity for IP<sub>7</sub> phosphorylation? One distinction is that proteins in yeast may be prephosphorylated whereas eukaryotic proteins are unlikely to be phosphorylated in *E. coli*. We examined whether yeast extracts influence IP<sub>7</sub> phosphorylation of proteins purified from *E. coli* (Fig. 1b). Although yeast extracts alone do not augment IP<sub>7</sub> phosphorylation, yeast extracts plus ATP allow IP<sub>7</sub> phosphorylation of proteins purified from bacteria comparable to phosphorylation of the same proteins purified from yeast. Boiled yeast extracts fail to prime proteins for IP<sub>7</sub> phosphorylation, indicating that the critical factor may be a protein, possibly an enzyme (Fig. 1c).

Several lines of evidence indicate that prephosphorylation of proteins primes them for IP<sub>7</sub> phosphorylation. ATP and GTP, which can donate phosphate groups, provide priming activity, whereas AMP-PNP, which is nonhydrolyzable, is inactive, as is UTP, a poor substrate for protein kinases (Fig. 1d). CK2 (formerly casein kinase-2) is one of the few protein kinases for which GTP donates phosphate to a comparable extent as ATP (19). Because GTP primes IP<sub>7</sub> phosphorylation as well as ATP, we examined the priming activity of CK2 on the IP<sub>7</sub> phosphorylation of NSR1 (1–50), a fragment of NSR1 which is an excellent substrate for IP<sub>7</sub> phosphorylation [supporting information (SI) Fig. 5] and has a strong consensus motif for CK2 phosphorylation. CK2 robustly primes IP<sub>7</sub> phosphorylation (Fig. 1e). Furthermore, the IP<sub>7</sub> substrate NSR1 (27–50) (SI Fig. 5), can be directly phosphorylated by IP<sub>7</sub> when purified from *E. coli* that coexpress the catalytic subunit of CK2 (Fig. 1f). If prephosphorylation is the critical priming event for IP<sub>7</sub> phosphorylation, then dephosphorylation should prevent such priming. Dephosphorylation of NSR1 (51–166) or NSR1 (1–50) by λ-phosphatase greatly reduces IP<sub>7</sub> phosphorylation (Fig. 1g). Protein kinase-mediated priming for IP<sub>7</sub> phosphorylation is observed for all IP<sub>7</sub> substrates tested, including yeast SRP40, NSR1, YGR130c, and

Author contributions: R.B., A.S., Y.A., A.C.R., K.P., and S.H.S. designed research; R.B., Y.A., A.M.S., A.C.R., J.K.W., and K.R.J. performed research; T.Z.K., H.M., A.P., Y.X., and G.D.P. contributed new reagents/analytic tools; R.B., A.S., and S.H.S. analyzed data; and R.B., G.D.P., K.P., and S.H.S. wrote the paper.

The authors declare no conflict of interest.

Abbreviations: IP<sub>3</sub>, inositol 1,4,5-trisphosphate; IP<sub>4</sub>, inositol tetrakisphosphate; IP<sub>5</sub>, inositol pentakisphosphate; IP<sub>6</sub>, inositol hexakisphosphate; IP<sub>7</sub>, PP-IP<sub>5</sub>, diphosphoinositol pentakisphosphate; IP<sub>8</sub>, [PP]<sub>2</sub>-IP<sub>4</sub>, bis-diphosphoinositol tetrakisphosphate; IP6K, IP<sub>6</sub> kinase; AMP-PNP, adenylyl-5'-yl imidodiphosphate; DIPP, diphosphoinositol polyphosphate phosphohydrolase.


<sup>§</sup>Present address: Division of Neurosurgery, Children's Hospital of Philadelphia, Department of Neurosurgery, University of Pennsylvania School of Medicine, Philadelphia, PA 19104.

<sup>¶</sup>To whom correspondence should be addressed at: 725 North Wolfe Street, WBSB 813, Baltimore, MD 21205. E-mail: ssnyder@jhmi.edu.

This article contains supporting information online at [www.pnas.org/cgi/content/full/0707338104/DC1](http://www.pnas.org/cgi/content/full/0707338104/DC1).

© 2007 by The National Academy of Sciences of the USA






**Fig. 1.**  $IP_7$ -mediated protein phosphorylation requires prephosphorylation by protein kinases. (a) NSR1 and SRP40 purified from *E. coli* or *S. cerevisiae* were incubated with  $5\beta[^{32}P]IP_7$  and resolved by NuPAGE; immunoblotting with a tag-specific antibody (Left) and autoradiography to determine phosphorylation (Right). (b) GST or GST-tagged NSR1 fragment (amino acids 51–166) expressed in *E. coli* and immobilized on glutathione beads were preincubated with or without *S. cerevisiae* extract or ATP, washed with PBS, and then treated with  $5\beta[^{32}P]IP_7$  and resolved by NuPAGE; Coomassie brilliant blue R250 staining (Left) and autoradiography (Right). The extra phosphorylated protein bands in lanes 3–6 are yeast proteins that bound nonspecifically to glutathione beads. (c) GST-SRP40 purified from *E. coli* was preincubated with ATP and either native *S. cerevisiae* extract (N) or extract that had been boiled for 5 min (B) and then treated with  $5\beta[^{32}P]IP_7$  as in b. (d) GST-SRP40 purified from *E. coli* was preincubated with yeast extract and indicated nucleotides and then phosphorylated by  $5\beta[^{32}P]IP_7$ ; Coomassie brilliant blue R250 staining (Lower) and autoradiography (Upper). (e) GST or GST-tagged NSR1 fragment (amino acids 1–50) purified from *E. coli* were preincubated with or without CK2 and ATP and then treated with  $5\beta[^{32}P]IP_7$  as in b. (f) GST-tagged NSR1 fragment (amino acids 27–50) purified from *E. coli* was incubated without or with CK2 and ATP (lanes 1 and 2) or coexpressed in *E. coli* with the catalytic A1 subunit of human CK2 (lane 3). Purified proteins were incubated with  $5\beta[^{32}P]IP_7$  as in b. (g) GST or GST-tagged NSR1 fragments (amino acids 1–50 and 51–166) purified from *S. cerevisiae*, were preincubated in the absence or presence of  $\lambda$ -phosphatase and then phosphorylated by  $5\beta[^{32}P]IP_7$  as in b.

APL6 and mammalian Nopp140, AP3 $\beta$ 3A, and TCOF1 (data not shown).

The requirement for priming by prephosphorylation suggests that either (i) ATP phosphorylation at certain sites allosterically facilitates  $IP_7$ -mediated phosphorylation at other sites, or (ii)  $IP_7$  directly phosphorylates amino acids that have been previously ATP-phosphorylated and hence provides a pyrophosphorylation or diphosphorylation modification. Earlier, we noted that  $IP_7$  phosphorylation occurs primarily at stretches of serines flanked by acidic amino acids and concluded that phosphorylation occurs on the serine residues (18) (also see SI Fig. 6). In Nopp140, deletion of individual serine residues in such a stretch (amino acid 80–100) decreases  $IP_7$  phosphorylation, implying that  $IP_7$  can phosphorylate several of them (Fig. 2a). Substitution of the single threonine residue in this sequence has no effect on  $IP_7$  phosphorylation. Because prephosphorylation by CK2 leads to  $IP_7$ -mediated phosphorylation of such sites, Nopp140 fragments

containing phosphates at the CK2-preferred sites should be substrates for  $IP_7$  phosphorylation in the absence of any other priming. To identify the preferred sites for CK2 phosphorylation, we phosphorylated an *E. coli*-expressed Nopp140 fragment comprising amino acids 80–100 and identified the major phosphorylation sites by mass spectrometry as serines 88 and 91 (data not shown). We synthesized a peptide with both of these serines prephosphorylated (Fig. 2b). This phosphopeptide is robustly phosphorylated by  $5\beta[^{32}P]IP_7$ , whereas the corresponding unphosphorylated peptide is resistant to  $5\beta[^{32}P]IP_7$  phosphorylation (Fig. 2c). Using two peptides with phosphates at serines 88 and 91, respectively, we observe substantially greater phosphorylation at position 91 than 88 (Fig. 2c). This differential influence on discrete sites supports the specificity of  $IP_7$ -mediated phosphorylation. Both the unphosphorylated and phosphorylated peptides are phosphorylated by  $\gamma[^{32}P]ATP$  and CK2 (Fig. 2d) on one or more of the available serine residues (data not shown).  $IP_7$



**Fig. 2.**  $\text{IP}_7$  phosphorylates a phosphoserine residue to generate pyrophosphoserine. (a) Extracts from *S. cerevisiae* expressing GST-tagged mouse Nopp140 fragment (amino acids 80–100) WT sequence and indicated point mutants were phosphorylated by  $5\beta[^{32}\text{P}]\text{IP}_7$ ; autoradiography (Upper) and immunoblotting (Lower). (b) Sequences of individual peptides derived from Nopp140 (GenBank accession no. NP\_941035). NOP, amino acids 76–100 of mouse Nopp140; NOP-2[pS], the same sequence as NOP except with phosphoserine at positions 88 and 91; NOP-pS 88 and NOP-pS 91, single phosphoserine residues at positions 88 and 91 respectively; NOP-2[MepS], methylphosphoserine at positions 88 and 91; NOP-2[ppS], pyrophosphoserine at positions 88 and 91. (c and d) The six synthetic peptides in b were incubated with  $5\beta[^{32}\text{P}]\text{IP}_7$  (c) or with CK2 and  $\gamma[^{32}\text{P}]\text{ATP}$  (d) and resolved by NuPAGE. Coomassie G250 staining (Left) and autoradiography (Right). In each peptide lane, there are multiple bands in the 3- to 9-kDa range, corresponding to peptide monomers and multimers, generated possibly because of electrostatic interactions between the positively charged lysines and negatively charged acidic and phosphate-containing residues (27). Peptide dimers were also observed by MALDI-TOF analysis (data not shown).

phosphorylation of the prephosphorylated but not the unphosphorylated Nopp140 peptide is consistent with pyrophosphorylation. This possibility is further supported by mass spectrometric analysis showing no phosphorylation by  $\text{IP}_7$  of the unphosphorylated Nopp140 peptide but addition of one or two phosphates to the prephosphorylated peptides (SI Table 1). The maximum number of phosphates accepted by a peptide from  $\text{IP}_7$  equals the number of preexisting phosphate groups on the peptide; single phosphoserine-containing peptides accept one phosphate from  $\text{IP}_7$ , whereas the peptide containing two phosphoserine residues accepts one or two phosphates.

The phosphorylated Nopp140 peptide might facilitate  $\text{IP}_7$  phosphorylation of nearby serines or might be pyrophosphorylated. To distinguish those alternatives, we synthesized the same phosphopeptide in which the two phosphates have been methylated to prevent pyrophosphorylation (Fig. 2b). The methylated phosphopeptide is not phosphorylated by  $5\beta[^{32}\text{P}]\text{IP}_7$  (Fig. 2c and SI Table 1), although it can be phosphorylated by  $\gamma[^{32}\text{P}]\text{ATP}$  and CK2 (Fig. 2d).

The failure of the methylated phosphopeptide to be  $\text{IP}_7$ -phosphorylated strongly implies that  $\text{IP}_7$  provides pyrophosphorylation. To further test this possibility, we synthesized the same Nopp140 peptide in which both serines 88 and 91 are pyrophosphorylated. This pyrophosphorylated peptide resists  $\text{IP}_7$  phosphorylation, although it can be phosphorylated by CK2 (Fig. 2c and d and SI Table 1). This further supports the notion that  $\text{IP}_7$  pyrophosphorylates its targets.

If  $\text{IP}_7$  phosphorylates proteins on serines in the same fashion as ATP, then the properties of the phosphoserine bond should be the same with  $\text{IP}_7$  and ATP phosphorylation. By contrast, the chemical properties of a pyrophosphorylated serine should differ from those of a conventional phosphoserine. We examined the acid sensitivity of the phospho-Nopp140 peptide after phosphorylation by  $\gamma[^{32}\text{P}]\text{ATP}$  and CK2 or  $5\beta[^{32}\text{P}]\text{IP}_7$  (Fig. 3a and b). Phosphorylation by  $5\beta[^{32}\text{P}]\text{IP}_7$  is more labile to treatment with 3 M HCl compared with phosphorylation by  $\gamma[^{32}\text{P}]\text{ATP}$  and CK2. Furthermore, we subjected the CK2 and ATP or  $\text{IP}_7$ -phosphorylated peptides to hydrolysis in the presence of 6 M HCl to release single amino acids. Hydrolysis liberates a prominent peak of  $[^{32}\text{P}]\text{phosphoserine}$  from the ATP-phosphorylated peptide but not from the  $\text{IP}_7$ -phosphorylated peptide (Fig. 3c–e). Thus, the pyrophosphate modification on serine is more

labile to acid than the conventional phosphoserine elicited by ATP-mediated phosphorylation.


In contrast to the greater lability to acid treatment, the  $\text{IP}_7$ -phosphorylated peptide is more stable to enzymatic dephosphorylation (Fig. 4a). As observed earlier (Fig. 1g),  $\lambda$ -phosphatase treatment before adding  $\text{IP}_7$  prevents  $\text{IP}_7$  phosphorylation of NSR1 (1–50). After  $\text{IP}_7$  phosphorylation, however, the protein is completely resistant to  $\lambda$ -phosphatase.  $\lambda$ -Phosphatase greatly reduces phosphorylation of CK2 and ATP phosphorylated NSR1 (27–50) or Nopp140 peptide but fails to affect the  $\text{IP}_7$ -phosphorylated peptides (Fig. 4b and c). The same protein fragments are resistant to  $\lambda$ -phosphatase when purified from yeast labeled with inorganic orthophosphate  $[^{32}\text{PO}_4]_i$  (Fig. 4d), implying that these proteins exist within yeast cells in a pyrophosphorylated form.

These experiments suggest that the pyrophosphoserine bond is resistant to  $\lambda$ -phosphatase. We directly tested this possibility using the Nopp140 peptide containing pyrophosphorylated serines at position 88 and 91 compared with the same peptide with phosphorylated serines at these positions.  $\lambda$ -Phosphatase abolishes phosphorylation of the phospho-Nopp140 peptide but does not affect the pyrophosphorylated Nopp140 peptide (Fig. 4e and SI Table 2).

We also examined the sensitivity of the  $\text{IP}_7$ -phosphorylated NSR1 (27–50) fragment to other protein phosphatases. The following protein phosphatases are completely inactive: calcineurin, protein phosphatase-1, and alkaline phosphatase (SI Fig. 7a). In addition, the pyrophosphatase enzymes, thermostable inorganic pyrophosphatase and tobacco acid pyrophosphatase, fail to dephosphorylate the  $\text{IP}_7$ -phosphorylated peptide (SI Fig. 7a). Because these enzymes hydrolyze inorganic pyrophosphate (PPi), it is likely that they do not recognize pyrophosphate linked to a serine residue. We also studied DIPP (diphosphoinositol polyphosphate phosphohydrolase), an enzyme that Shears and associates (20, 21) have shown to physiologically dephosphorylate  $\text{IP}_7$ . As expected, pretreatment of the reaction mix containing  $\text{IP}_7$  lowers phosphorylation, whereas treatment with DIPP after  $\text{IP}_7$  phosphorylation has no effect (SI Fig. 7b).

In summary, we report that  $\text{IP}_7$  pyrophosphorylates proteins. Evidence includes the requirement of prephosphorylation by ATP to prime proteins for  $\text{IP}_7$  phosphorylation and the selective phosphorylation by  $\text{IP}_7$  of synthetic phosphopeptides but not of






**Fig. 3.** The properties of pyrophosphoserine differ from those of phosphoserine. (a and b) NOP-2[pS] peptide was phosphorylated by CK2 and  $\gamma$ [ $^{32}$ P]ATP (a) or by  $5\beta$ [ $^{32}$ P]IP $_7$  (b), then treated with HCl and subjected to NuPAGE; Coomassie G250 staining (Left) and autoradiography (Right). (c) NOP-2[pS] peptide was phosphorylated by either CK2 and  $\gamma$ [ $^{32}$ P]ATP or by  $5\beta$ [ $^{32}$ P]IP $_7$  as described in *SI Materials and Methods*. A fraction of the phosphorylated peptides was subjected to NuPAGE to ensure equivalent phosphorylation under both conditions; Coomassie G250 staining (Left) and autoradiography (Right). (d and e) The remaining CK2 phosphorylated peptide (d) and IP $_7$ -phosphorylated peptide (e) were hydrolyzed with 6 M HCl, and the resulting amino acids were resolved by HPLC. Data collected during the first 40 min of the HPLC run are presented; elution profiles of the first 3 aa (Upper) and corresponding radioactivity in each 1-ml fraction (Lower). Note that high levels of  $^{32}$ P $_i$  in the IP $_7$ -phosphorylated sample compared with the CK2 and ATP phosphorylated sample are due to incomplete removal of  $5\beta$ [ $^{32}$ P]IP $_7$  from the peptide during desalting (see *SI Materials and Methods*).

such peptides that are “blocked” by methylation or are already pyrophosphorylated. Moreover, the properties of IP $_7$ -phosphorylated peptides differ markedly from ATP-phosphorylated peptides in terms of acid lability, which is greater with IP $_7$ -phosphorylated peptides, and sensitivity to phosphatases, which is much less with IP $_7$ -phosphorylated peptides.

We wondered whether 4/6-PP-IP $_5$ , the novel isomer of IP $_7$  identified by York and colleagues (15), and IP $_8$  are able to phosphorylate proteins in a manner similar to 5-PP-IP $_5$ . Radiolabeled 4/6-PP-IP $_5$  and IP $_8$  phosphorylate proteins in extracts obtained from *S. cerevisiae* (SI Fig. 8), the pattern of phosphorylated proteins being identical to that obtained with 5-PP-IP $_5$ .



**Fig. 4.**  $\lambda$ -phosphatase does not dephosphorylate IP $_7$ -phosphorylated proteins. (a) GST-tagged NSR1 fragment (amino acids 1–50) purified from *S. cerevisiae* was incubated without or with  $\lambda$ -phosphatase and then phosphorylated by  $5\beta$ [ $^{32}$ P]IP $_7$  (lanes 1 and 2). Alternatively, the protein was first phosphorylated by  $5\beta$ [ $^{32}$ P]IP $_7$  and then incubated without or with  $\lambda$ -phosphatase (lanes 3 and 4). The samples were resolved by NuPAGE and autoradiographed. (b) GST-tagged NSR1 fragment (amino acids 27–50) purified from *S. cerevisiae* was phosphorylated either by CK2 and  $\gamma$ [ $^{32}$ P]ATP (Left) or by  $5\beta$ [ $^{32}$ P]IP $_7$  (Right) and then incubated with or without  $\lambda$ -phosphatase; Coomassie G250 staining (Left) and autoradiography (Right). (c) NOP-2[pS] peptide bound to streptavidin agarose beads was phosphorylated by either CK2 and  $\gamma$ [ $^{32}$ P]ATP (Left) or by  $5\beta$ [ $^{32}$ P]IP $_7$  (Right) and then incubated with or without  $\lambda$ -phosphatase as in b. (d) *S. cerevisiae* expressing GST-tagged Nop140 fragment (amino acids 80–100) (Left) or NSR1 fragment (amino acids 27–50) (Right) were labeled with inorganic orthophosphate [ $^{32}$ P] $_i$ . Purified radiolabeled proteins were incubated with or without  $\lambda$ -phosphatase; autoradiography (Upper) and immunoblotting (Lower). (e) NOP-2[pS] peptide (Left) or NOP-2[ppS] peptide (Right) were treated with or without  $\lambda$ -phosphatase, resolved by NuPAGE, and stained with either Coomassie G250 or Pro-Q Diamond phosphoprotein gel stain, as indicated.

Therefore, it is likely that the principal known inositol pyrophosphates are able to pyrophosphorylate proteins.

Nonenzymatic pyrophosphorylation is thermodynamically feasible. Semiempirical calculations show that, whereas the free enthalpy ( $\Delta H$ ) of reaction for transfer of the  $\beta$ -phosphate from IP<sub>7</sub> to a simple primary alcohol is pH- and counterion-dependent,  $\Delta H$  values at pH 6.8 can reach  $-38.3$  kcal/mol (17). This strongly exothermic reaction equals or exceeds  $\Delta H$  for phosphorylation by ATP. The high phosphorylation potential is attributable to the sterically and electronically crowded environment of the IP<sub>7</sub> pyrophosphate group. The phosphoserine to pyrophosphoserine conversion by IP<sub>7</sub> will require more advanced modeling, but two important clues emerge from the protein substrates for IP<sub>7</sub> that suggest how the 3D structures of the polypeptide may organize reagent and substrate to provide an entropic driving force to contribute to the overall free energy ( $\Delta G$ ) for the reaction. First, the polyserine tracts must have Asp or Glu residues present for pyrophosphorylation to occur. Second, the pyrophosphorylation shows an absolute requirement for divalent cations, with Mg<sup>2+</sup> preferred (18). Both of these features are reminiscent of the prenyl diphosphate synthases, which feature DDXXD motifs that coordinate Mg<sup>2+</sup>, which in turn activates the pyrophosphate as a leaving group (22). For serine pyrophosphorylation, one can imagine IP<sub>7</sub> phosphates organized by H-bonds within a polySer tract, whereas the IP<sub>7</sub> pyrophosphate could be activated and targeted to a particular phosphoserine phosphate with the assistance of an Asp/Glu-chelated Mg<sup>2+</sup>.

What might be the physiologic role of pyrophosphorylation? Classic protein kinase-mediated phosphorylation modifies the conformation of proteins, enhancing or decreasing catalytic activity, determining protein localization, or altering protein-protein interactions. Presumably pyrophosphorylation also elicits such alterations in protein function. ATP-mediated phosphorylation is typically regulated by dephosphorylation, whereas pyrophosphorylated proteins are resistant to known protein phosphatases. This suggests that serine pyrophosphorylation may be a more stable modification within the cell, even though it is thermodynamically more unstable compared with phosphoserine. Such a conclusion is consistent with the fact that some of the best IP<sub>7</sub> targets, such as NSR1 and SRP40/Nopp140, exist endogenously as abundantly phosphorylated proteins (23–25). Although it is likely that pyrophosphorylation plays a central role in cellular processes modulated by inositol pyrophosphates, at this time, the biological significance of this modification remains unclear.

## Materials and Methods

**Synthesis of Inositol Pyrophosphate.** The inositol pyrophosphate used throughout these studies was 5-PP-I(1,2,3,4,6)P<sub>5</sub> (referred to as IP<sub>7</sub>). The synthesis and purification of radiolabeled 5 $\beta$ [<sup>32</sup>P]IP<sub>7</sub> was carried out as described earlier (18). By using these procedures, 5 $\beta$ [<sup>32</sup>P]IP<sub>7</sub> was obtained at a specific activity of 60 Ci/mmol (1 Ci = 37 GBq) and used at 1–2  $\mu$ Ci per reaction for protein phosphorylation. The synthesis of <sup>32</sup>P-labeled 4/6-PP-IP<sub>5</sub> and IP<sub>8</sub> is described in *SI Materials and Methods*. Unlabeled 5-PP-I(1,2,3,4,6)P<sub>5</sub> was prepared by total synthesis by using modifications of an earlier method (26). Full experimental details are provided in *SI Materials and Methods*.

**Preparation of Recombinant Proteins.** The expression and purification of all recombinant proteins are described in *SI Materials and Methods*.

**Protein and Peptide Phosphorylation Assays.** Cell lysates were obtained by resuspending yeast cells in ice-cold lysis buffer [20 mM Hepes (pH 6.8)/5 mM DTT/1 mM EGTA/1 mM EDTA/0.1% CHAPS, protease inhibitor mixture, and 200 mg/liter phenyl-

methanesulfonyl fluoride] and vortexing the sample in the presence of glass beads. The homogenates were centrifuged for 20 min at 15,000  $\times$  g, and supernatants were used in the assay. In experiments where lysates were used directly for phosphorylation, protein extracts (10–20  $\mu$ g) were incubated in IP<sub>7</sub> phosphorylation buffer [25 mM Hepes (pH 7.4)/50 mM NaCl/6 mM MgCl<sub>2</sub>/1 mM DTT] and 1  $\mu$ Ci 5 $\beta$ [<sup>32</sup>P]IP<sub>7</sub> for 15 min at 37°C. The reactions were heated at 95°C for 5 min in sample buffer before separation by NuPAGE (Invitrogen, Carlsbad, CA). The gels were either stained and dried, or transferred to a PVDF membrane. Radiolabeled proteins were detected by autoradiography. Immunoblotting using anti-Xpress (Invitrogen) or anti-GST (Sigma, St. Louis, MO) antibodies, was performed according to standard procedures.

GST- or hexahistidine-tagged proteins purified from *S. cerevisiae* were used directly for phosphorylation by 5 $\beta$ [<sup>32</sup>P]IP<sub>7</sub>. The protein (1–2  $\mu$ g) was incubated in the presence of 5 $\beta$ [<sup>32</sup>P]IP<sub>7</sub>, and the reaction was performed as described above. Where indicated, the purified proteins were preincubated with  $\lambda$ -phosphatase (400 units; New England Biolabs, Beverly, MA) at 30°C for 1 h, washed twice with PBS, and then used in a 5 $\beta$ [<sup>32</sup>P]IP<sub>7</sub> phosphorylation assay. GST fusion proteins purified from *E. coli*, were coupled to glutathione beads, and either incubated directly with 5 $\beta$ [<sup>32</sup>P]IP<sub>7</sub> as above or were first preincubated with 12  $\mu$ g of extract prepared from *S. cerevisiae* (strain DDY1810) at 30°C for 1 h in IP<sub>7</sub> phosphorylation buffer with or without 1 mM ATP or other nucleotides. Where indicated, the *S. cerevisiae* extract was incubated in boiling water for 5 min, cooled, and then used for the preincubation with purified protein. After incubation with yeast extract, the glutathione beads were washed twice with PBS and once with IP<sub>7</sub> phosphorylation buffer and used for phosphorylation by 5 $\beta$ [<sup>32</sup>P]IP<sub>7</sub> as described above. In some cases, proteins purified from *E. coli* were first preincubated with 250 units of CK2 (New England Biolabs) in the supplied CK2 reaction buffer, along with 1 mM ATP at 30°C for 1 h, and then washed and used for 5 $\beta$ [<sup>32</sup>P]IP<sub>7</sub> phosphorylation. NSR1 amino acids 27–50 purified from *E. coli* cells coexpressing human CK2A1 was used directly in 5 $\beta$ [<sup>32</sup>P]IP<sub>7</sub> phosphorylation assays.

Peptides were synthesized as described in *SI Materials and Methods*. Synthetic peptides were used for both CK2 and ATP or IP<sub>7</sub> mediated phosphorylation assays. Peptide (3–6  $\mu$ g) was incubated with 250 units of CK2 in CK2 phosphorylation buffer, 200  $\mu$ M ATP, and 1  $\mu$ Ci  $\gamma$ [<sup>32</sup>P]ATP at 30°C for 15–30 min. For IP<sub>7</sub> phosphorylation, 3–6  $\mu$ g of peptide was added to IP<sub>7</sub> phosphorylation buffer and 1–2  $\mu$ Ci of 5 $\beta$ [<sup>32</sup>P]IP<sub>7</sub> at 37°C for 15 min, followed by incubation at 95°C for 5 min. The peptides were mixed with sample buffer and resolved on a 12% gel by using the NuPAGE system. The gels were stained by using Safe Stain (Invitrogen), dried, and used for autoradiography. Although peptide phosphorylation by IP<sub>7</sub> was routinely performed at 95°C to maximize the extent of phosphorylation, a time-dependent increase in phosphorylation is observed at 37°C (*SI Fig. 9*).

**Acid-Sensitivity Assay.** To determine the sensitivity of CK2 and ATP or IP<sub>7</sub>-phosphorylated NOP-2[pS] peptide to treatment with acid, peptides were phosphorylated as described above and then incubated with either 1 M HCl at 50°C, with 3 M HCl at 25°C, or without any acid at 25°C, for 1 h. After incubation, the acid was neutralized with an appropriate volume of 10 M NaOH, and the reaction was mixed with sample buffer and subjected to NuPAGE. Phosphorylation was detected as described above. Peptide hydrolysis in the presence of 6 M HCl, followed by separation of amino acids by HPLC, was performed as described in *SI Materials and Methods*.

**$\lambda$ -Phosphatase-Sensitivity Assay.** Purified GST fusion proteins or NOP-2[pS] peptide were phosphorylated by CK2 and  $\gamma$ [<sup>32</sup>P]ATP or by 5 $\beta$ [<sup>32</sup>P]IP<sub>7</sub> as described above. After phosphorylation, the

biotinylated NOP-2[pS] peptide was bound to streptavidin agarose beads (Sigma) and washed with PBS. Protein or peptide was then incubated in the presence of  $\lambda$ -phosphatase (400 units; New England Biolabs) in supplied buffer according to the manufacturer's instructions at 30°C for 1 h, mixed with sample buffer, resolved by NuPAGE, and subjected to autoradiography. Expression and purification of  $^{32}\text{[PO}_4\text{]}_i$ -labeled GST fusion proteins and their treatment with phosphatase is described in *SI Materials and Methods*.

The Pro-Q Diamond Phosphoprotein Gel Stain (Invitrogen) was used to determine relative levels of phosphate on the NOP-2[pS] or NOP-2[ppS] peptides before or after treatment with  $\lambda$ -phosphatase. NOP-2[pS] peptide (6  $\mu\text{g}$ ) or NOP-2[ppS] peptide (24  $\mu\text{g}$ )

were incubated with 400 units of  $\lambda$ -phosphatase for 1 h at 30°C and then resolved by NuPAGE and stained with Pro-Q Diamond stain according to the manufacturer's instructions.

We thank Philip Cole, Leslyn Hanakahi, Tamara Hendrickson, Alex Huang, Michael McMahon, Ananya Majumdar, and John Honek for helpful discussions and Raghunand Tirumalai, Anutosh Chakraborty, Michael Koldobskiy, Makoto Hara, Katherine Sixt, and Asif Mustafa for valuable suggestions and comments. This work was supported by U.S. Public Health Service Grant MH18501 and Research Scientist Award DA00074 (to S.H.S.), National Institutes of Health (NIH) Grant CA106424 (to A.P.), A.S. was supported by Medical Research Council funding of the Cell Biology Unit, and G.D.P. thanks NIH Grant NS29632 for support of synthetic work.

1. Berridge MJ, Lipp P, Bootman MD (2000) *Nat Rev Mol Cell Biol* 1:11–21.
2. Bennett M, Onnebo SM, Azevedo C, Saiardi A (2006) *Cell Mol Life Sci* 63:552–564.
3. Menniti FS, Miller RN, Putney JW, Jr, Shears SB (1993) *J Biol Chem* 268:3850–3856.
4. Stephens L, Radenberg T, Thiel U, Vogel G, Khoo KH, Dell A, Jackson TR, Hawkins PT, Mayr GW (1993) *J Biol Chem* 268:4009–4015.
5. Saiardi A, Erdjument-Bromage H, Snowman AM, Tempst P, Snyder SH (1999) *Curr Biol* 9:1323–1326.
6. Saiardi A, Nagata E, Luo HR, Snowman AM, Snyder SH (2001) *J Biol Chem* 276:39179–39185.
7. Schell MJ, Letcher AJ, Brearley CA, Biber J, Murer H, Irvine RF (1999) *FEBS Lett* 461:169–172.
8. Saiardi A, Sciambi C, McCaffery JM, Wendland B, Snyder SH (2002) *Proc Natl Acad Sci USA* 99:14206–14211.
9. Dubois E, Scherens B, Vierendeels F, Ho MM, Messenguy F, Shears SB (2002) *J Biol Chem* 277:23755–23763.
10. Morrison BH, Bauer JA, Kalvakolanu DV, Lindner DJ (2001) *J Biol Chem* 276:24965–24970.
11. Nagata E, Luo HR, Saiardi A, Bae BI, Suzuki N, Snyder SH (2005) *J Biol Chem* 280:1634–1640.
12. Luo HR, Huang YE, Chen JC, Saiardi A, Iijima M, Ye K, Huang Y, Nagata E, Devreotes P, Snyder SH (2003) *Cell* 114:559–572.
13. York SJ, Armbruster BN, Greenwell P, Petes TD, York JD (2005) *J Biol Chem* 280:4264–4269.
14. Saiardi A, Resnick AC, Snowman AM, Wendland B, Snyder SH (2005) *Proc Natl Acad Sci USA* 102:1911–1914.
15. Mulugu S, Bai W, Fridy PC, Bastidas RJ, Otto JC, Dollins DE, Haystead TA, Ribeiro AA, York JD (2007) *Science* 316:106–109.
16. Lee YS, Mulugu S, York JD, O'Shea EK (2007) *Science* 316:109–112.
17. Hand CE, Honek JF (2007) *Bioorg Med Chem Lett* 17:183–188.
18. Saiardi A, Bhandari R, Resnick AC, Snowman AM, Snyder SH (2004) *Science* 306:2101–2105.
19. Tuazon PT, Traugh JA (1991) *Adv Second Messenger Phosphoprotein Res* 23:123–164.
20. Safrany ST, Caffrey JJ, Yang X, Bembenek ME, Moyer MB, Burkhart WA, Shears SB (1998) *EMBO J* 17:6599–6607.
21. Ingram SW, Safrany ST, Barnes LD (2003) *Biochem J* 369:519–528.
22. Tarshis LC, Yan M, Poulter CD, Sacchettini JC (1994) *Biochemistry* 33:10871–10877.
23. Meier UT (1996) *J Biol Chem* 271:19376–19384.
24. Meier UT, Blobel G (1992) *Cell* 70:127–138.
25. Yan C, Melese T (1993) *J Cell Biol* 123:1081–1091.
26. Reddy KM, Reddy KK, Falck JR (1997) *Tetrahedron Lett* 38:4951–4952.
27. Jackson SN, Wang HY, Yergey A, Woods AS (2006) *J Proteome Res* 5:122–126.