

Marquette University
e-Publications@Marquette

Biological Sciences Faculty Research and
Publications

Biological Sciences, Department of

1-1-1992

Modulation by cAMP of a Slowly Activating Potassium Channel Expressed in *Xenopus* Oocytes

Edward M. Blumenthal

Marquette University, edward.blumenthal@marquette.edu

Leonard K. Kaczmarek

Yale University

Published version. *Journal of Neuroscience*, Vol. 12, No. 1 (January 1992): 290-296. [Permalink](#). ©

1992 Society for Neuroscience. Used with permission.

Edward Blumenthal was affiliated with the Yale University School of Medicine at the time of publication.

Modulation by cAMP of a Slowly Activating Potassium Channel Expressed in *Xenopus* Oocytes

Edward M. Blumenthal¹ and Leonard K. Kaczmarek^{1,2}

¹Interdepartmental Neuroscience Program and ²Departments of Pharmacology and Cellular and Molecular Physiology, Yale University School of Medicine, New Haven, Connecticut 06510

When expressed in the *Xenopus* oocyte, the minK protein induces a slowly activating voltage-dependent potassium current (I_{sk}). We studied the modulation of this current by altering intracellular cAMP levels and found that the amplitude of I_{sk} is dramatically increased by treatments that raise cAMP levels and decreased by agents that lower cAMP levels. Preinjection of a protein inhibitor of the cAMP-dependent protein kinase blocked the effects of increased cAMP levels. There were no changes in the voltage dependence or kinetics of I_{sk} . Mutations that eliminate a potential phosphorylation site on the minK protein did not block the effects of activating the kinase. In addition, the membrane capacitance of the oocyte increased and decreased in parallel with I_{sk} . Our results fit a mechanism in which channel proteins are selectively inserted into and removed from the plasma membrane in response to changes in kinase activity.

Potassium channels play an important role in determining the firing pattern and the shape of action potentials in neurons and other excitable cells (Cook, 1990). In recent years, the genes coding for a number of different voltage-dependent potassium channels have been isolated. With one exception, the *minK* (minimal K) gene, all of the genes belong to the same extended family, which also includes the major subunits of the voltage-dependent sodium and calcium channels (for reviews, see Jan and Jan, 1990; Kaczmarek, 1991). The predicted structures of the proteins encoded by these genes are organized into domains containing six proposed membrane-spanning helices and a number of other highly conserved motifs. In contrast, the minK protein contains only 130 amino acids and one proposed hydrophobic transmembrane helix and has no homology with the other cloned ion channels (Takumi et al., 1988).

mRNA for the minK protein is expressed in a variety of tissues including estrogen-primed uterus, heart, and kidney proximal tubule (Folander et al., 1990; Pragnell et al., 1990; Sugimoto et al., 1990). When expressed in the oocytes of the clawed frog *Xenopus laevis*, RNA from *minK*-expressing tissue or from the cloned gene induces a voltage-dependent potassium current (I_{sk}) that activates slowly over many seconds (Boyle et al., 1987; Takumi et al., 1988).

The physiological role of the minK channel is not yet known, and speculations on its function are based solely upon its localization. Investigation of the factors that regulate I_{sk} may therefore help to clarify the role of the minK protein *in situ*.

We have now investigated the modulation of I_{sk} by the cAMP-dependent protein kinase (PKA). Modulation by second messengers has been described for a number of different classes of potassium currents in their native cellular environment (Kaczmarek and Levitan, 1987). The *Xenopus* oocyte expression system is well suited for the study of such regulation since one can record currents caused by a single species of injected RNA (Dascal, 1987). Despite this, there has been only one recent report of the modulation of a cloned potassium channel expressed in oocytes (Hoger et al., 1991), and no studies of regulation by PKA. We now report that, despite the lack of any consensus phosphorylation sites on the minK protein, the current is dramatically modulated by changes in PKA activity.

Materials and Methods

In vitro transcription. The rat *minK* clone (Takumi et al., 1988; Pragnell et al., 1990) was in pGEM2 and was linearized with HindIII (Promega or Boehringer Mannheim). *K_v1* and the human *minK* mutants (a gift of Dr. R. Swanson, Merck, Sharp and Dohme) were in pGEM-A [Swanson et al., 1990; a pGEM-9zf(-) vector with a built-in polyA tail] and were linearized with Not I (Promega or Boehringer Mannheim). Run-off transcripts were prepared in a reaction containing linearized template, reaction buffer (Promega), dithiothreitol (Promega; 10 mM), rNTPs (Promega; 500 μ M each), RNasin (Promega; 1.5 U/ μ l), mG(5')ppp(5')Gm (Pharmacia; 100 μ M), and RNA polymerase [SP6 (Promega) or T7 (gift of Dr. T. Steitz, Yale University)]. The template was then digested with DNase (Promega), and the RNA was purified by multiple ethanol precipitations and dissolved in diethyl pyrocarbonate treated sterile water at either 0.2 mg/ml (*K_v1*) or 0.05 mg/ml (others). The RNA was stored in small aliquots at -70°C .

Oocyte isolation and injection. Adult female *Xenopus laevis* (Nasco, Ft. Atkinson, WI) were anesthetized in ice-cold 0.2% 3-aminobenzoic acid ethyl ester (Sigma). Ovarian lobes were removed through a slit in the abdominal wall, which was then sutured. Animals were not operated on more than once in any 2 week period and were occasionally injected with 500 U gonadotrophin (Sigma) and left for at least 4 weeks to regenerate oocytes. The oocytes were separated by incubation at room temperature for 1.5-3 hr with 2 mg/ml collagenase (Boehringer Mannheim) in calcium-free Ringer's solution [88 mM NaCl, 1 mM KCl, 2.4 mM NaHCO₃, 0.8 mM MgSO₄, 15 mM HEPES pH 7.6 (NaOH)]. Following collagenase treatment, large healthy stage V and VI oocytes (Dumont, 1972) were transferred into OR₃ [50% L-15 (GIBCO), 15 mM HEPES pH 7.6, 50 μ g/ml gentamycin (Sigma)]. After sitting at room temperature for 2-18 hr to overnight, the oocytes were injected with 50 nl of RNA. The oocytes were incubated in OR₃ at 19°C for 3 d and then moved to 14°C. Electrophysiology was always performed at least 3 d after RNA injection.

For kinase inhibitor injection, oocytes were injected with 50 nl of a buffer containing 5 mM 2-(*N*-morpholino)-ethanesulfonic acid (MES) (CalBiochem; pH 7.0) and 1 mg/ml BSA (Sigma) with or without the

Received June 18, 1991; revised Aug. 19, 1991; accepted Aug. 26, 1991.

We thank Dr. R. Swanson for the human and mutant minK clones, Drs. S. Amara and R. Handschumacher for *Xenopus laevis*, and Todd Lubart for assistance with data analysis. This work was supported by NIH Grant HL38156 and a grant from the Cystic Fibrosis Foundation to L.K.K.

Correspondence should be addressed to Dr. Leonard Kaczmarek, Department of Pharmacology, Yale Medical School, 333 Cedar Street, New Haven, CT 06510.

Copyright © 1992 Society for Neuroscience 0270-6474/92/120290-07\$05.00/0

kinase inhibitor (Walsh et al., 1971; gift of Dr. A. Nairn, Rockefeller University) at 1.6 mg/ml. See Results for further details.

Electrophysiology. Oocyte membrane currents were recorded using standard two-electrode voltage-clamp procedures with a World Precision Instruments S-7100 voltage clamp controlled by a PDP-11/24 computer (Indec). The recording and analysis software was a modified version of bCLAMP (Indec). Electrodes were filled with 3 M KCl and had resistances of 0.2–1.5 M Ω . The current signal was filtered at 500 Hz, the time invariant leak current was subtracted, and both current and voltage records were digitized and stored on disk.

The recording solution was designed to minimize endogenous chloride currents (Barish, 1983; Boyle et al., 1987) and contained (in mM) 96 Na-aspartate, 2 K-aspartate, 0.5 CaCl₂, 5 MgCl₂, and 5 HEPES pH 7.6 (NaOH). During I_{sk} recording, the oocytes were briefly perfused before each pulse, as this seemed to give more consistent currents. All measurements of current were determined as the average of five pulses delivered every 15 sec. Except where noted, the holding potential was -50 mV, command pulse was to 50 mV for 1.5 sec, and currents were measured at the beginning of the tail current immediately following repolarization.

Membrane capacitance was measured by delivering a series of hyperpolarizing sawtooth voltage pulses, subtracting the estimated resistive current, and measuring the magnitude of the remaining capacitive current.

8-Br-cAMP (Sigma) was dissolved in recording solution at 1 mM. Progesterone (Sigma) was stored as a 1 mM stock in ethanol at -20°C and was diluted into recording solution immediately before use. Forskolin (CalBiochem) was stored as a 50 mM stock in ethanol at 4°C and was also diluted immediately before use.

Results

Modulation of I_{sk} by cAMP

Treatments that alter intracellular cAMP levels in the *Xenopus* oocytes change the amplitude of heterologously expressed I_{sk} . We used the membrane-permeable analog 8-Br-cAMP (1 mM) to raise the cAMP levels. I_{sk} amplitude consistently began to increase within 3–5 min of drug addition and continued to increase for the remainder of the experiments ($n = 9$). The increase in current after about 15 min ranged from 16% to 144% ($86 \pm 53\%$, \pm SD). In some oocytes, the current continued to increase to levels over fourfold greater than before drug treatment (Fig. 1A). We found only a weak correlation among the different injected oocytes between the final magnitudes of the current increase and the durations of 8-Br-cAMP treatment. Treatment of uninjected oocytes with 8-Br-cAMP had no consistent effects on the small endogenous outward currents (Fig. 1C).

In order to show that 8-Br-cAMP application was mimicking a rise in cAMP, we also investigated the effect of forskolin, an activator of adenylate cyclase. Forskolin (50 μM) also induced a rapid increase in I_{sk} amplitude in two oocytes (95% and 67%, respectively, within 35 min).

The steroid hormone progesterone is the physiological trigger for meiotic maturation of *Xenopus* oocytes (reviewed in Maller, 1985). It has been shown to cause a transient decrease in intracellular cAMP levels (Cicirelli and Smith, 1985). We therefore studied the effect of 1 μM progesterone on the amplitude of I_{sk} . Consistent with its effects on cAMP levels, progesterone caused a decrease in I_{sk} (Fig. 1B). This effect began within 3 min and reached a maximum reduction of $43 \pm 9\%$ ($n = 6$) 17–82 min after drug application.

A change in the amplitude of a current may result from a shift in the voltage dependence of the channels or, in the case of a very slowly activating current, a change in the kinetics of activation. We therefore analyzed these parameters for the action of 8-Br-cAMP on I_{sk} . Figure 2 shows plots of normalized current amplitude, measured at the end of a 1.5 sec command pulse, as a function of command potential. Due to the slow kinetics




Figure 1. Changes in cAMP levels alter the magnitude of I_{sk} . All current traces are averages of five 1.5 sec pulses from -50 to 50 mV. The pulses were delivered 15 sec apart to allow for complete deactivation of I_{sk} . *A*, Average traces before and 62 min after addition of 1 mM 8-Br-cAMP. *B*, Average traces before and 10 min after addition of 1 μM progesterone. *C*, 8-Br-cAMP does not alter the endogenous currents of an uninjected oocyte. Average traces are before and 17 min after 1 mM 8-Br-cAMP.

of I_{sk} , such an isochronic plot could be sensitive to alterations in the activation kinetics. Nevertheless, there was no change in the voltage dependence following 8-Br-cAMP application. We also found no change in the reversal potential of the tail currents (-92 to -96 mV; $n = 3$; data not shown), indicating that there was no change in the selectivity of the current. Finally, we observed no consistent change in I_{sk} kinetics following any of the drug treatments. Thus, the effects of cAMP appeared to be strictly confined to current amplitude.

Effects on capacitance

In addition to the changes in I_{sk} , we also observed changes in the capacitance of the oocyte plasma membrane (C_m) in response to changes in cAMP levels. The membrane capacitance of *Xenopus* oocytes, which is a function of total surface area of the plasma membrane, is known to decrease in response to phorbol




Figure 2. 8-Br-cAMP does not alter the voltage dependence of I_{sk} . Currents were measured at the end of 1.5 sec pulses to the voltages shown. The currents were normalized to the amplitude at 70 mV. The graph shows difference before 8-Br-cAMP (squares) and after 16–25 min in 1 mM 8-Br-cAMP (triangles). Each point is an average of three oocytes, each of which showed large increases in I_{sk} amplitude. Error bars represent SEs.

esters and during meiotic maturation (Kado et al., 1981; Vasilets et al., 1990). However, there have been no reports of rapid changes in C_m in response to changes in cAMP levels.

C_m increased in response to 8-Br-cAMP and decreased in response to progesterone. These changes were smaller than the changes in I_{sk} , but were temporally related. Figure 3 shows typical time courses for both effects. While the overall increases and decreases were similar, they were not identical. Notice that during certain periods of time I_{sk} increased substantially while C_m did not, and vice versa. Uninjected oocytes also underwent a capacitance increase in response to 8-Br-cAMP, indicating that this effect was not dependent upon the presence of I_{sk} or the minK protein. Because a single drug treatment changed both the current and the capacitance, we analyzed whether the magnitude of the capacitance increase in a given oocyte could account for the magnitude of the current increase. An analysis of the maximum relative current and capacitance increases among all of the oocytes treated with either 8-Br-cAMP or forskolin ($n = 11$) showed that the two effects were not significantly correlated ($r = 0.57$; $p = 0.07$).

Role of PKA

cAMP is known to influence the amplitude of ionic currents either by directly binding to the channel proteins or by activating PKA (Krebs and Beavo, 1979; DiFrancesco and Tortora, 1991). To determine the role of PKA in the regulation of I_{sk} , we injected oocytes with a heat-stable kinase inhibitor (1.55 mg/ml, 50 nl; $n = 6$; Walsh et al., 1971; Sadler and Maller, 1983). After allowing at least 2 hr for the protein to diffuse throughout the oocyte, we compared the effects of 8-Br-cAMP on inhibitor-injected and buffer-injected cells. Figure 4A shows a significant inhibition of the increase in I_{sk} amplitude in the inhibitor-injected oocytes. The small residual increase in I_{sk} was probably due to incomplete diffusion of the inhibitor protein, as it seemed to depend on the amount of time between injection and testing. We conclude that the changes in I_{sk} amplitude are due to changes in PKA activity.

Protein kinase inhibitor injection appeared to attenuate the increase in C_m , but the difference between inhibitor-injected and control oocytes was not highly significant (Fig. 4B). Because of the small size of the capacitance increase and the time-depen-




Figure 3. Time courses of the 8-Br-cAMP and progesterone effects. Typical time courses for the changes in I_{sk} (squares) and capacitance (triangles) are shown. Tail currents were measured at -50 mV following a 1.5 sec pulse to 50 mV. Drugs were added at time 0: A, 1 mM 8-Br-cAMP; B, 1 μM progesterone.

dent variability of C_m in some oocytes, we feel that this result does not rule out PKA as the cause of the capacitance changes.

While conducting the inhibitor studies, we noticed that the baseline I_{sk} amplitude was reduced in the inhibitor-injected oocytes compared with the controls. In order to test further the action of PKA on baseline I_{sk} , we compared oocytes that had been injected with either inhibitor or buffer and allowed to incubate overnight (10–13 hr). These data are summarized in Table 1. Although this treatment has been seen by others to induce oocyte maturation (Sadler and Maller, 1983), we did not consistently observe this effect. However, injection of PKA inhibitor caused a significant reduction in I_{sk} , C_m , and current density (I_{sk}/C_m) relative to oocytes injected with buffer alone. In contrast, there were no differences in the resting potentials or input resistances between the two groups. Interestingly, the magnitudes of the I_{sk} and C_m decreases (44% and 11%, respectively) were similar to those seen following progesterone treatment.

Specificity of action of cAMP

The effects of cAMP changes on I_{sk} are not due to nonspecific endocytosis and exocytosis of oocyte plasma membrane. To show this, we coexpressed I_{sk} with the delayed rectifier K_v1 , a mammalian member of the *Shaker* family (Swanson et al., 1990). K_v1 , when expressed alone, was not affected by either progesterone or 8-Br-cAMP (Fig. 5A,B). In coinjected oocytes, I_{sk} was selectively modulated by both progesterone and 8-Br-cAMP (Fig. 5C,D).




Figure 4. Effect of PKA inhibitor injection on the changes in current and capacitance. Oocytes were injected with either the inhibitor of PKA (solid bars) or buffer (shaded bars). After 2–6 hr, the oocytes were voltage clamped and treated with 1 mM 8-Br-cAMP. Tail currents were measured at -50 mV after a 1.5 sec pulse to 50 mV. The graphs show the percentage increases in tail current (A) and capacitance (B) relative to the values immediately before addition of 8-Br-cAMP ($n = 6$ oocytes). Error bars represent SEs. Currents in inhibitor-injected oocytes increased significantly less than in controls, two-tailed t test: * $p < 0.05$, ** $p < 0.005$. Injection of inhibitor did not significantly attenuate the capacitance increases: $0.15 > p > 0.10$, one-tailed t test, for last two time points.

Regulation of minK mutants

One potential mechanism for the increases in I_{sk} after activation of PKA is direct phosphorylation of the minK channel protein. Although this protein does not contain a classical consensus site for phosphorylation by PKA (R/K–R/K–X–S/T) (Krebs and Beavo, 1979), it does have one serine that is flanked by basic residues, a motif frequently found to be phosphorylated by protein kinases. We examined site-directed mutants of the human minK protein in which this putative phosphorylation site was altered (Table 2) (Murai et al., 1989). Expression of these mutants results in currents with slightly altered kinetics and voltage dependence (data not shown). Nevertheless, the currents from all three mutants, as well as the wild-type human current, increased in response to 8-Br-cAMP ($n = 2$ for each) (Fig. 6). These increases were similar in magnitude and time course to the rat I_{sk} results. We conclude that the putative phosphorylation site is not necessary for modulation of I_{sk} by PKA.

Discussion

We have shown that changes in the level of intracellular cAMP produce changes in the magnitude of I_{sk} expressed in *Xenopus*

Table 1. Inhibition of PKA affects baseline I_{sk} and C_m

Property	Oocyte	
	Buffer-injected	Inhibitor-injected
Tail current (nA)	248 ± 63	139 ± 67**
Capacitance (nF)	292 ± 10	260 ± 12**
Current density (nA/nF)	0.85 ± 0.22	0.54 ± 0.26*
Resting potential (mV)	−46 ± 2	−41 ± 6
Input resistance (MΩ)	0.9 ± 0.2	1.1 ± 0.4

Data show properties of oocytes injected with either the PKA inhibitor ($n = 7$) or buffer ($n = 9$) and allowed to incubate overnight at 14°C. Values are given as average ± SD. Inhibitor-injected differs from control, two-tailed t test: * $p < 0.05$, ** $p < 0.01$.

laevis oocytes. This modulation is due to changes in the activity of PKA. Treatments that increased PKA activity led to substantial increases in I_{sk} amplitude, while inhibition of the kinase caused the current to decrease. Modulation occurred over a relatively slow time course, tens of minutes, although a similar time course has been reported for modulation by forskolin of a heart calcium current expressed in oocytes (Dascal et al., 1986). The voltage dependence, ionic selectivity, and kinetics of I_{sk} were unaffected by manipulation of cAMP. We also found that, in addition to its effects on currents, changing the activity of PKA altered cell capacitance, a sensitive measure of membrane surface area. Activation of the kinase caused a net addition of membrane to the surface of the oocyte, while inhibition of the kinase led to a net endocytosis.

There are two general mechanisms for the modulation of ion channels that may bring about an increase in current amplitude. The first is modification of channels that are already in the plasma membrane. The second is a change in the number of channels physically present in the membrane. Although our data do not definitely distinguish between the two alternatives, they are generally more consistent with the second possibility.

For the first mechanism, the minK protein would have to exist in two states—active and inactive—which could be interconverted through the action of PKA. Since cAMP does not alter kinetics or voltage dependence, the inactive state would have to be completely silent, making this scheme different from another two-state model of channel modulation, the “willing/reluctant” model for the modulation of calcium channels by norepinephrine in dorsal root ganglion neurons (Bean, 1989). As stated earlier, there are no PKA consensus phosphorylation sites on the minK protein, and we have shown that deletion of the only likely phosphorylation site did not eliminate the PKA effect. Therefore, the conversion of minK from the inactive to the active conformation would have to involve either phos-

Table 2. Amino acid sequences of the human minK mutants

Protein	Sequence
Wild type	–Ile–Arg–Ser–Lys–Lys–
Mutant 1	–Ile–Arg–Ala–Lys–Lys–
Mutant 2	–Ile–Gln–Ser–Lys–Lys–
Mutant 3	–Ile–Arg–Ser–Gln–Gln–

This sequence lies in the proposed cytoplasmic domain of the protein between residues 66 and 70 (Murai et al., 1989).




Figure 5. Changes in cAMP levels do not affect the delayed rectifier K_v1 . *A* and *B*, Oocytes expressing K_v1 alone. Traces were taken before and 13 min after 8-Br-cAMP (*A*) and before and 24 min after progesterone (*B*). The pulse protocol was as in Figure 1, except that the holding potential was -80 mV. *C* and *D*, Oocytes coexpressing K_v1 and I_{sk} . On this time scale, the current at the beginning of the pulse represents only K_v1 while the tail currents are only I_{sk} . Traces were taken before and 29 min after 8-Br-cAMP (*C*) and before and 58 min after progesterone (*D*). The pulse protocol was as in Figure 1.

phorylation of minK by PKA at an "inhospitable" site, phosphorylation of minK by some other kinase, possibly due to the activation of a cascade of kinases, or phosphorylation of an associated endogenous oocyte protein.

The second possible mechanism for modulation of I_{sk} is a change in the amount of minK protein on the surface of the oocyte. In this scheme, changes in PKA activity would alter the kinetics of membrane trafficking to and from the surface of the oocyte. A number of assumptions are necessary for this hypothesis to explain the data. First, the effect of PKA on membrane insertion would have to be specific to membranes containing the minK protein and/or any regulatory proteins for this

channel. For example, the amplitude of the delayed rectifier K_v1 did not change along with the capacitance. Furthermore, the changes in I_{sk} amplitude were always much larger than the changes in capacitance. Therefore, we would have to assume a specific population of vesicles, to which the minK protein is targeted, whose transport is regulated by PKA activity. There is evidence that such specific membrane trafficking exists in the oocyte. Progesterone has been shown to cause the rapid formation of large numbers of coated vesicles, apparently containing the progesterone receptors (Dersch et al., 1991). This observation is consistent with our observations of progesterone's effects on capacitance. Also, activation of protein kinase C causes a re-




Figure 6. Alteration of a putative phosphorylation site does not eliminate the cAMP effect. *A*, The wild-type human I_{sk} responds to 8-Br-cAMP in the same way as rat I_{sk} . The pulse protocol was the same as in Figure 1. Traces were taken before and 9 min after drug addition. *B–D*, Pulses as above except that the command pulse was to 90 mV to compensate for the altered voltage dependence of the mutants. Currents were measured before and within 10–15 min after addition of 8-Br-cAMP. *B*, mutant 1; *C*, mutant 2; *D*, mutant 3. See Table 2 for further details.

duction in the surface area of *Xenopus* oocytes that is associated with the selective endocytosis of the Na⁺/K⁺-ATPase (Vasilets et al., 1990). Finally, a subset of ion channels is selectively removed from the membrane during the maturation of starfish oocytes (Moody and Bosma, 1985).

One potential problem with this second mechanism is the weak correlation between total changes in I_{sk} and C_m in different oocytes. It must be emphasized, however, that very few oocytes had absolutely stable baseline capacitances. It was far more common for the capacitance to vary with time, making calculations of capacitance changes inexact. The weak correlation could also be explained by assuming that the capacitance increase is due to the exocytosis of different classes of vesicles, of which only a specific subset contains the minK protein. The second, addition/subtraction mechanism more convincingly explains the lack of changes in the kinetics and voltage dependence following activation of PKA as well as the relatively slow time course of the changes in current. It is also consistent with the known ability of the oocyte to remove some proteins selectively from its surface.

Although the physiological role of the minK channel is not yet understood, there is evidence that cAMP-induced insertion and removal of plasma membrane containing ion channels occurs in the kidney, one of the tissues in which the minK protein is found (Sugimoto et al., 1990). In the principal cells of the collecting duct, vasopressin, which causes increases in cAMP, stimulates the delivery of water channels to the apical membrane (for review, see Brown, 1989). Moreover, parathyroid hormone, which elevates cAMP levels in cultured distal tubule epithelia, causes the appearance of L-type calcium channels in the cell membrane (Bacsikai and Friedman, 1990). This process is dependent on a functional exocytotic apparatus.

The *minK* gene is also expressed in cardiac muscle (Folander et al., 1990). An examination of the potassium currents found in the heart suggests a possible role for I_{sk} . The delayed rectifier of guinea pig ventricular myocytes is composed of different potassium currents, the largest and slowest of which, I_{Ks} , has kinetics and voltage dependence similar to that of I_{sk} (Sanguinetti and Jurkiewicz, 1990). In addition, this current increases in response to activation of PKA (Bennett et al., 1986; Sanguinetti et al., 1991). Walsh and Kass (1988) have shown that although PKA activation increases both the calcium current and the delayed rectifier, the two currents seem to increase by different mechanisms. The change in the delayed rectifier is highly temperature dependent, while modulation of I_{Ca} is not. It is interesting to speculate that the modulation of this I_{sk} -like potassium current might be due to exocytosis of a specific population of vesicles. Insertion and removal of channels could prove to be a widespread mechanism for cell-type-specific modulation of ionic currents.

References

- Bacsikai BJ, Friedman PA (1990) Activation of latent Ca²⁺ channels in renal epithelial cells by parathyroid hormone. *Nature* 347:388–391.
- Barish ME (1983) A transient calcium-dependent chloride current in the immature *Xenopus* oocyte. *J Physiol (Lond)* 342:309–325.
- Bean BP (1989) Neurotransmitter inhibition of neuronal calcium currents by changes in channel voltage dependence. *Nature* 340:153–156.
- Bennett P, McKinney L, Begenisich T, Kass RS (1986) Adrenergic modulation of the delayed rectifier potassium channel in calf cardiac Purkinje fibers. *Biophys J* 49:839–848.
- Boyle MB, Azhderian EM, MacLusky NJ, Naftolin F, Kaczmarek LK (1987) *Xenopus* oocytes injected with rat uterine RNA express very slowly activating potassium currents. *Science* 235:1221–1224.
- Brown D (1989) Membrane recycling and epithelial cell function. *Am J Physiol* 256:F1–F12.
- Circirelli MF, Smith LD (1985) Cyclic AMP levels during the maturation of *Xenopus* oocytes. *Dev Biol* 108:254–258.
- Cook NS ed (1990) Potassium channels: structure, classification, function, and therapeutic potential. Chichester: Horwood.
- Dascal N (1987) The use of *Xenopus* oocytes for the study of ion channels. *CRC Crit Rev Biochem* 22:317–387.
- Dascal N, Snutch TP, Lubbert H, Davidson N, Lester HA (1986) Expression and modulation of voltage-gated calcium channels after RNA injection in *Xenopus* oocytes. *Science* 231:1147–1150.
- Dersch MA, Bement WM, Larabell CA, Mecca MD, Capco DG (1991) Cortical membrane-trafficking during the meiotic resumption of *Xenopus laevis* oocytes. *Cell Tissue Res* 263:375–383.
- DiFrancesco D, Tortora P (1991) Direct activation of cardiac pacemaker channels by intracellular cyclic AMP. *Nature* 351:145–147.
- Dumont JN (1972) Oogenesis in *Xenopus laevis* (Daudin). *J Morphol* 136:153–180.
- Folander K, Smith JS, Antanavage J, Bennett C, Stein RB, Swanson R (1990) Cloning and expression of the delayed-rectifier I_{sk} channel from neonatal rat heart and diethylstilbestrol-primed rat uterus. *Proc Natl Acad Sci USA* 87:2975–2979.
- Hoger JH, Walter AE, Vance D, Yu L, Lester HA, Davidson N (1991) Modulation of a cloned mouse brain potassium channel. *Neuron* 6:227–236.
- Jan LY, Jan YN (1990) How might the diversity of potassium channels be generated? *Trends Neurosci* 13:415–419.
- Kaczmarek LK (1991) Voltage-dependent potassium channels: *minK* and *Shaker* families. *New Biol* 3:315–323.
- Kaczmarek LK, Levitan IB (1987) Neuromodulation: the biochemical control of neuronal excitability. New York: Oxford UP.
- Kado RT, Marcher K, Ozon R (1981) Electrical membrane properties of the *Xenopus laevis* oocyte during progesterone-induced meiotic maturation. *Dev Biol* 84:471–476.
- Krebs E, Beavo J (1979) Phosphorylation–dephosphorylation of enzymes. *Annu Rev Biochem* 48:923–959.
- Maller JL (1985) Regulation of amphibian oocyte maturation. *Cell Differ* 16:211–221.
- Moody WJ, Bosma MM (1985) Hormone-induced loss of surface membrane during maturation of starfish oocytes: differential effects on potassium and calcium channels. *Dev Biol* 112:396–404.
- Murai T, Kakizuka A, Takumi T, Ohkubo H, Nakanishi S (1989) Molecular cloning and sequence analysis of human genomic DNA encoding a novel membrane protein which exhibits a slowly activating potassium channel activity. *Biochem Biophys Res Commun* 161:176–181.
- Pragnell M, Snay KJ, Trimmer JS, MacLusky NJ, Naftolin F, Kaczmarek LK, Boyle MB (1990) Estrogen induction of a small, putative K⁺ channel mRNA in rat uterus. *Neuron* 4:807–812.
- Sadler SE, Maller JL (1983) The development of competence for meiotic maturation during oogenesis in *Xenopus laevis*. *Dev Biol* 98:165–172.
- Sanguinetti MC, Jurkiewicz NK (1990) Two components of cardiac delayed rectifier K⁺ current: differential sensitivity to block by class III antiarrhythmic agents. *J Gen Physiol* 96:195–215.
- Sanguinetti MC, Jurkiewicz NK, Scott A, Siegl PKS (1991) Isoproterenol antagonizes prolongation of refractory period by the class III antiarrhythmic agent E-4031 in guinea pig myocytes: mechanism of action. *Circ Res* 68:77–84.
- Sugimoto T, Tanabe Y, Shigemoto R, Iwai M, Takumi T, Ohkubo H, Nakanishi S (1990) Immunohistochemical study of a rat membrane protein which induces a selective potassium permeation: its localization in the apical membrane portion of epithelial cells. *J Membr Biol* 113:39–47.
- Swanson R, Marshall J, Smith JS, Williams JB, Boyle MB, Folander K, Luneau CJ, Antanavage J, Oliva C, Buhrow SA, Bennett C, Stein RB, Kaczmarek LK (1990) Cloning and expression of cDNA and genomic clones encoding three delayed rectifier potassium channels in rat brain. *Neuron* 4:929–939.
- Takumi T, Ohkubo H, Nakanishi S (1988) Cloning of a membrane protein that induces a slow voltage-gated potassium current. *Science* 242:1042–1045.

Vasilets LA, Schmalzing G, Madefessel K, Haase W, Schwarz W (1990) Activation of protein kinase C by phorbol ester induces downregulation of the Na⁺/K⁺-ATPase in oocytes of *Xenopus laevis*. *J Membr Biol* 118:131–142.

Walsh DA, Ashby CD, Gonzalez C, Calkins D, Fischer EH, Krebs EG

(1971) Purification and characterization of a protein inhibitor of adenosine 3':5'-monophosphate-dependent protein kinases. *J Biol Chem* 246:1977–1985.

Walsh KB, Kass RS (1988) Regulation of a heart potassium channel by protein kinase A and C. *Science* 242:67–69.