

Marquette University
e-Publications@Marquette

Biological Sciences Faculty Research and
Publications

Biological Sciences, Department of

12-1-2006

FRIGIDA LIKE 2 Is a Functional Allele in Landsberg *erecta* and Compensates for a Nonsense Allele of *FRIGIDA LIKE 1*

Michael Schläppi

Marquette University, michael.schlappi@marquette.edu

Published version. *Plant Physiology*, Vol. 142, No. 4 (December 2006): 1728-1738. DOI. © 2006

American Society of Plant Biologists. Used with permission.

Figure 1 (supplemental data) available [here](#).

FRIGIDA LIKE 2 Is a Functional Allele in Landsberg *erecta* and Compensates for a Nonsense Allele of *FRIGIDA LIKE 1*¹[W][OA]

Michael R. Schläppi

*Department of Biological Sciences, Marquette University
Milwaukee, WI*

Abstract

The Landsberg *erecta* (*Ler*) accession of *Arabidopsis* (*Arabidopsis thaliana*) has a weak allele of the floral inhibitor *FLOWERING LOCUS C* (*FLC*). *FLC-Ler* is weakly up-regulated by the active San Feliu-2 (*Sf2*) allele of *FRIGIDA* (*FRI-Sf2*), resulting in a moderately late-flowering phenotype. By contrast, the Columbia (*Col*) allele of *FLC* is strongly up-regulated by *FRI-Sf2*, resulting in a very late-flowering phenotype. In *Col*, the *FRI*-related gene *FRIGIDA LIKE 1* (*FRL1*) is required for *FRI*-mediated up-regulation of *FLC*. It is shown here that in *Ler*, the *FRL1*-related gene *FRIGIDA LIKE 2* (*FRL2*), but not *FRL1*, is required for *FRI*-mediated up-regulation of *FLC*. *FRL1-Ler* is shown to be a nonsense allele of *FRL1* due to a naturally occurring premature stop codon in the middle of the conceptual protein sequence, suggesting that *FRL1-Ler* is nonfunctional. Compared to *FRL2-Col*, *FRL2-Ler* has two amino acid changes in the conceptual protein sequence. Plants homozygous for *FRI-Sf2*, *FLC-Ler*, *FRL1-Ler*, and *FRL2-Col* have no detectable *FLC* expression, resulting in an extremely early flowering phenotype. Transformation of a genomic fragment of *FRL2-Ler*, but not of *FRL2-Col*, into a recombinant inbred line derived from these plants restores both *FRI*-mediated up-regulation of *FLC* expression and a late-flowering phenotype, indicating that *FRL2-Ler* is the functional allele of

FRL2. Taken together, these results suggest that in the two different *Arabidopsis* accessions *Col* and *Ler*, either *FRL1* or *FRL2*, but not both, is functional and required for *FRI*-mediated up-regulation of *FLC*.

The timing of reproductive development is an important decision during the life cycle of flowering plants. The coordination of flowering time is vital for self-incompatible plant species, because they strongly need their sexual partners to flower at the same time. Coordinate regulation of flowering time is also required for the reproductive success of self-compatible species such as *Arabidopsis* (*Arabidopsis thaliana*). For instance, it can determine whether a population of dormant seeds or plants in a vegetative state will overwinter, because *Arabidopsis* has evolved both naturally occurring, early flowering summer-annual ecotypes and naturally occurring, late-flowering winter-annual ecotypes (Laibach, 1937). A major difference between the two growth habits is that winter-annual types overwinter as vegetative seedlings or plants, because they require vernalization (exposure to a prolonged cold period during winter) to flower in the next spring or summer. By contrast, summer-annual types generally, but not exclusively, produce seeds that remain dormant during winter and germinate the next year for a summer-annual growth habit (Nordborg and Bergelson, 1999). The requirement for vernalization in natural populations of *Arabidopsis* is mostly controlled by the synergistic interaction of two dominant genes, *FRIGIDA* (*FRI*) and *FLOWERING LOCUS C* (*FLC*; Michaels and Amasino, 1999; Sheldon et al., 1999; Johanson et al., 2000). *FRI* encodes a plant-specific coiled-coil domain-containing protein required for the up-regulation of *FLC*, which produces a MADS domain-containing transcription factor that acts as a strong floral repressor. Vernalization is antagonistic to *FRI* and leads to the epigenetic down-regulation of *FLC* expression; that is, levels of *FLC* transcript remain low even after removal of the cold stimulus (Michaels and Amasino, 1999; Sheldon et al., 2000; Schläppi, 2001). *FRI*-mediated up-regulation of *FLC* is reset in the next generation when progeny plants become late flowering again (Amasino, 2004).

Until recently, *FRI* and *FLC* were considered the major determinants of flowering time in natural populations of *Arabidopsis*. This is because most early flowering accessions were shown to have either defects in *FRI* (Johanson et al., 2000; Le Corre et al., 2002;

Gazzani et al., 2003), weak alleles of *FLC* (Koornneef et al., 1994; Lee et al., 1994; Sanda and Amasino, 1996; Schläppi, 2001; Gazzani et al., 2003; Michaels et al., 2003) or nonfunctional *FLC* transcripts (Shindo et al., 2005; Werner et al., 2005). However, recent studies have identified late-flowering *Arabidopsis* accessions that do not fit this pattern. Those accessions have either high levels of *FLC* expression in the absence of a functional *FRI* allele or are late flowering without a functional *FLC* allele (Schläppi, 2001; Werner et al., 2005). This suggests that there is naturally occurring variation in flowering time genes other than *FRI* and *FLC*. Through mutagenesis experiments with summer- and winter-annual ecotypes, several classes of flowering time genes were identified that might be candidates for natural variation in *FRI*- or *FLC*-independent late flowering. Those are the six genes of the autonomous floral promotion pathway, *LUMINIDEPENDENS*, *FCA*, *FLOWERING LOCUS D*, *FPA*, *FY*, and *FVE*, that repress up-regulation of *FLC* expression in the absence of *FRI* (Boss et al., 2004); or the three *FLC* paralogs, *FLOWERING LOCUS M/MADS AFFECTING FLOWERING 1* (*MAF1*), *MAF2*, and *MAF3*, that together with *FLC* have an additive effect on floral repression (Y. Pan and M.R. Schläppi, unpublished data) or repress flowering when overexpressed (Ratcliffe et al., 2001, 2003; Scortecci et al., 2001). Another class of more pleiotropic suppressors of *FRI* activity and *FLC* up-regulation includes chromatin regulators such as encoded by *ACTIN-RELATED PROTEIN 6*, *PHOTOPERIOD-INDEPENDENT EARLY FLOWERING 1*, or the *VERNALIZATION INDEPENDENCE* genes (Noh and Amasino, 2003; Oh et al., 2004; Choi et al., 2005; Deal et al., 2005).

In addition to finding accessions that are late flowering in the absence of active alleles of *FRI* or *FLC*, recent mutagenesis experiments led to the identification of *FRI LIKE 1* (*FRL1*), a *FRI*-related gene that is required for the winter-annual growth habit of *Arabidopsis* (Michaels et al., 2004). A single *frl1* mutant suppresses *FRI*-mediated late flowering and up-regulation of *FLC* expression in the Columbia (Col) ecotype of *Arabidopsis*. *FRL1* is part of a gene family of six *FRI*-related genes, including *FRI LIKE 2* (*FRL2*), which has some functional redundancy with *FRL1* in the Col ecotype (Michaels et al., 2004). In this study, it is shown that there is naturally occurring variation at *FRL1* and *FRL2* between the Col and Landsberg *erecta* (*Ler*) accessions of *Arabidopsis*. In Col, *FRL1* is functional, whereas in *Ler*, *FRL2* is shown to be functional but not vice versa. This suggests

that natural variation at these two loci could potentially modify or even suppress *FLC* up-regulation and late flowering in *Arabidopsis* accessions that have functional alleles at both *FRI* and *FLC*.

Results

Genetic Identification of a Modifier Gene Required for FRI-Mediated Late Flowering in the Ler Background


In previous work, the effect of *FRI* on flowering time and its interaction with *FLC* was investigated in different genetic backgrounds of *Arabidopsis* (Schläppi, 2001). In a series of genetic experiments, control test crosses were performed between the very late-flowering Col-*FRI*-San Feliu-2 (Sf2) line (rosette leaf no. [RLN] range 51–72) and the moderately late-flowering Ler-*FRI*-Sf2 line (RLN range 12–28), both of which contained the active *FRI*-Sf2 allele in the Col or Ler background, respectively (Schläppi, 2001). While F₁ plants (RLN range 46–68) were almost as late flowering as the late Col-*FRI*-Sf2 parent, about 1/16th (17/255) very early flowering F₂ plants (RLN range 3–5) were recovered from this cross. These early flowering Col/Ler-*FRI*-Sf2 F₂ plants were considered as transgressions, because they flowered significantly earlier than the earliest Ler-*FRI*-Sf2 parent. Test crosses and mapping analyses showed that all early flowering F₂ plants were homozygous for the active and dominant *FRI*-Sf2 gene, the weak and recessive *FLC*-Ler gene, and an unlinked recessive Col-specific suppressor gene of *FRI*-mediated late flowering (Schläppi, 2001). The flowering time phenotype suggested that the naturally occurring dominant Ler variant of this suppressor gene was required for *FRI*-mediated late flowering in the Ler background. Therefore, the gene was named *ACTIVATOR OF FRI-MEDIATED LATE FLOWERING IN LER (AFL)*.

To determine the epistatic interaction between the naturally occurring Col and Ler alleles of *AFL*, three randomly selected early flowering Col/Ler-*FRI*-Sf2 plants containing *AFL*-Col (lines #1, #3, and #5; RLN range 3–4) were backcrossed with *AFL*-Ler-containing Ler-*FRI*-Sf2 (RLN range 18–27). As shown in Figure 1A, the flowering time phenotypes of F₁ plants (RLN range 5–18) from two of the three crosses were intermediate between that of the two parental lines,

whereas F_1 plants from the third cross (RLN range 16–28) flowered as late as the *Ler-FRI-Sf2* parent. As shown in Figure 1B, F_2 plants from F_1 plants with an intermediate flowering time phenotype segregated 1:2:1 ($\chi^2 = 0.0697$; $P > 0.9$) very early flowering (RLN range 3–5; 10 plants):intermediate late flowering (RLN range 7–13; 22 plants):late flowering (RLN range 16–25; 11 plants), suggesting that in this genetic background, *AFL-Ler* regulated *FRI*-mediated late flowering in a dosage-dependent manner.

Figure 1

Genetic analysis of very early flowering F_2 plants Col/*Ler-FRI-Sf2*#1, #3, and #5 from the cross of *Ler-FRI-Sf2* × Col/*FRI-Sf2*. Flowering time is expressed as the RLN at the time of bolting. Plants were grown in long-day photoperiods. Error bars indicate two sds. A, Flowering time of F_1 plants (black bars) from the cross of Col/*Ler-FRI-Sf2*#1, #3, and #5 with late-flowering *Ler-FRI-Sf2*. Flowering time of *Ler* and Col is presented for comparison. B, Frequency distribution of flowering time (RLN) for an F_2 population from the cross of Col/*Ler-FRI-Sf2*#1 × *Ler-FRI-Sf2*. Flowering time ranges and means of parent plants and F_1 hybrids are shown by horizontal lines and arrows, respectively.



AFL Is the Functional Ler Variant of FRL2


To identify the chromosomal position of *AFL*, bulked segregant analysis was done with the 17 very early flowering F_2 individuals and simple sequence length polymorphism (SSLP) molecular markers (Supplemental Fig. S1). This analysis suggested that *AFL* was linked to the *nga248* marker on chromosome 1. Individual analysis of the 17 very early flowering plants using six molecular markers on chromosome 1 indicated that *AFL* was located about 2 cM south of *UNUSUAL FLORAL ORGANS* (*UFO*; Supplemental Table S1). While mapping of *AFL* was in progress, work on *FRI*-related genes was published, describing that *FRL1* on chromosome 5 was required for *FRI*-mediated up-regulation of *FLC* in the Col-*FRI*-Sf2 background (Michaels et al., 2004). Interestingly, the *AFL* locus mapped very closely to *At1g31814*, another *FRI*-related gene on chromosome 1. *At1g31814* single mutants had no effect on flowering time; however, *frl1 At1g31814* double mutants flowered slightly earlier than *frl1* single mutants (Michaels et al., 2004). Therefore, *At1g31814* was interpreted to be functionally redundant with *FRL1* and was named *FRL2*. Thus, the map location of *FRL2* and the absence of an *frl2* single mutant phenotype in Col raised the possibility that *FRL2*-Col was identical to the suppressor allele *AFL*-Col, and conversely, that *AFL*-Ler was a functional allele of *FRL2*.

To test whether *AFL*-Ler was a functional allele of *FRL2*, a molecular-genetic approach was taken. Toward this end, a partial *FRL2*-Ler sequence was retrieved from the Monsanto Ler single-pass shotgun sequencing database (Jander et al., 2002) and compared to *FRL2*-Col. Four nonsynonymous polymorphisms were found in the coding regions between *FRL2*-Ler and *FRL2*-Col, one of which resulted in an *AluI* site in the Ler allele but not in the Col allele. As shown in Figure 2A, this *AluI* polymorphism was a useful cleaved amplified polymorphic sequence (CAPS) marker to distinguish between the Col and Ler alleles of *FRL2*. A prediction from the hypothesis that *AFL* is identical to *FRL2* was that the very early flowering Col/Ler-*FRI*-Sf2 plants had the Col-specific suppressor allele of *FRL2*. Indeed, CAPS analysis of the 17 pooled very early flowering F_2 plants from the cross of Col-*FRI*-Sf2 with Ler-*FRI*-Sf2 indicated that there was an apparent bias for *FRL2*-Col, the Col allele of *FRL2* (Fig. 2B; see result for Col/Ler

[#1–#17]). Lines Col/Ler-*FRI-Sf2*#1, #3, and #5 used for backcrosses with Ler-*FRI-Sf2* were homozygous for *FRL2*-Col, as was early flowering F₂ progeny (plants 1, 2, and 3; Fig. 2B) from the backcross shown in Figure 1B. Conversely, late-flowering F₂ progeny (plants 40, 41, and 44; Fig. 2B) from the same backcross population was homozygous for *FRL2*-Ler, whereas intermediate to late-flowering F₂ progeny (plants 14, 15, and 16; Fig. 2B) was heterozygous for both alleles. Individual CAPS analysis of the 17 very early flowering plants indicated that 16 out of 17 plants were homozygous for *FRL2*-Col and one plant was heterozygous. The heterozygous plant was subsequently shown to segregate early and late-flowering progeny, suggesting that it was misidentified during the initial selection (data not shown). Taken together, these results were an indication that the early flowering phenotype was associated with the Col allele of *FRL2*. To rule out that *FRL2*-Col was differentially expressed compared to *FRL2*-Ler, semiquantitative reverse transcription (RT)-PCR was performed on RNA isolated from Col, Ler, and *FRL2*-Col-containing, very early flowering Col/Ler-*FRI-Sf2* F₂ plants. As shown in Figure 2C, *FRL2* transcript levels were comparable in the Ler and Col genetic backgrounds, indicating that *FRL2*-Col allele was not a hypomorph.

Figure 2


A, Graphic representation of the *AluI* CAPS marker (Ala-132 site) for *FRL2* and an example of how *AluI* cleaves the Ler-specific PCR fragment. B, CAPS marker-based analysis of *FRL2* alleles in a pool of 17 and in three individual very early flowering Col/Ler-*FRI-Sf2* F₂ progeny plants from the cross of Ler-*FRI-Sf2* × Col-*FRI-Sf2* (#1–#17 and #1, #3, and #5, respectively) and in randomly selected very early flowering F₂ plants (1, 2, and 3), late-flowering F₂ plants (40, 41, and 44), and intermediate to late-flowering F₂ plants (14, 15, and 16) from the cross of Col/Ler-*FRI-Sf2*#1 × Ler-*FRI-Sf2*. C, RT-PCR analysis of *FRL2* expression levels in Col, Col-*FRI-Sf2*, Ler, Ler-*FRI-Sf2*, and Col/Ler-*FRI-Sf2*. *ACTIN2/8* (*ACT*) is shown as control for loading.



To directly show that *FRL2-Ler* was required for *FRI*-mediated late flowering, a genomic clone of *FRL2-Ler* was transformed into the very early flowering, recombinant inbred line (RIL) Col/Ler-*FRI-Sf2* #1, which had been selfed and propagated from single seeds for five generations (Supplemental Table S2). To more easily identify transgenic plants that were later flowering than Col/Ler-*FRI-Sf2* #1, but earlier flowering than Ler-*FRI-Sf2*, the flowering time of 26 T₁ plants was analyzed under less inductive 12-h photoperiods. As shown in Figure 3A, individual T₁ plants had a range of flowering time phenotypes, from flowering as early as the Col/Ler-*FRI-Sf2* #1 parent to as late as the Ler-*FRI-Sf2* control. Only six out of 26 T₁ plants flowered as early as Col/Ler-*FRI-Sf2* #1, whereas the rest flowered later (Fig. 3A). As a control, a genomic clone of *FRL2-Col* was transformed into the same very early flowering RIL, and 14 T₁ plants were recovered. All 14 plants flowered as early as the RIL control, producing only four to five rosette leaves at the time of bolting (data not shown). Taken together, these experiments demonstrated that introduction of *FRL2-Ler* into the very early flowering line Col/Ler-*FRI-Sf2* #1 restored late flowering in a majority of T₁ plants.

Figure 3

A, Frequency distribution of flowering time (RLN) for 26 Col/Ler-FRI-Sf2#1 T₁ plants transformed with a genomic copy of *FRL2-Ler*. Flowering time ranges and means of the untransformed parent Col/Ler-FRI-Sf2#1 and of Col and Ler-FRI-Sf2 control plants are shown by horizontal lines and arrows, respectively. T₁ plants and controls were grown in 12-h photoperiods. B, Late-flowering phenotype of T₂ plants from the *FRL2-Ler*-transformed T₁ line 16 and untransformed control plants grown for 36 d in long-day photoperiods.



The late-flowering time phenotype of individual T₁ plants transformed with a genomic clone of *FRL2-Ler* was generally maintained in T₂ progeny plants and cosegregated with the transgene. For instance, late-flowering T₁ line 8 had a single kanamycin resistance (*Km^R*) gene locus and thus a single *FRL2-Ler* transgene locus. Kanamycin-selected T₂ progeny plants from the selfed T₁ parent were always late flowering, whereas unselected T₂ progeny plants segregated 3:1 late flowering:very early flowering (28 late:11 early; $\chi^2 = 0.21$; $P > 0.5$). The T₃ progeny of 10 unselected T₂ plants was subsequently analyzed for *Km^R* segregation, indicating that all chosen early flowering plants were kanamycin sensitive (three plants), and that the seven chosen late-flowering plants were either homozygous

(two plants) or heterozygous (five plants) for Km^R . This suggested that the late-flowering phenotype cosegregated with the single *FRL2-Ler* transgene locus. Moreover, as shown in Figure 3B, *FRL2-Ler* transgene-containing T_2 progeny from T_1 line 16 that flowered late under 12-h photoperiods was also late flowering under long-day photoperiods. Conversely, T_2 progenies from T_1 plants that were early flowering remained early flowering. The T_2 flowering time distribution under long-day photoperiods of several transgenic plants is summarized in Table I. It is interesting to note that early flowering T_1 lines generally had a range of kanamycin-resistance cosuppression phenotypes in the T_2 generation, whereas selected T_2 progenies from late-flowering T_1 lines were fully kanamycin resistant.

Table I.

Flowering time analysis in long-day photoperiods

The flowering time of individual T_2 plants derived from the very early flowering RIL Col/*Ler-FRI-Sf2*#1, transformed with a genomic copy of *FRL2-Ler*, was determined by counting the RLN at the time of bolting. Plants were selected for Km^R (linked to the *FRL2* transgene) and grown in 16 h of light and 8 h of dark conditions. CLN, Number of cauline leaves.

Plant Lines	RLN, Mean \pm SE (<i>n</i>)	CLN, Mean \pm SE
Col/ <i>Ler-FRI-Sf2</i> #1 (RIL)	4.1 \pm 0.3 (16)	1.2 \pm 0.4
<i>Ler</i> (CS20)	4.0 \pm 0.0 (10)	1.0 \pm 0.0
Col-0	10.3 \pm 1.3 (8)	2.3 \pm 0.7
<i>Ler-FRI-Sf2</i>	36.4 \pm 4.2 (9)	6.4 \pm 0.9
T_2 -3 ^a	6.2 \pm 2.1 (15)	–
T_2 -11 ^a	10.2 \pm 2.9 (5)	–
T_2 -12 ^a	6.4 \pm 3.2 (7)	–
T_2 -8	32.1 \pm 2.7 (15)	3.8 \pm 0.6
T_2 -14	27.7 \pm 7.6 (10)	–
T_2 -15	18.5 \pm 5.4 (10)	–
T_2 -16	22.8 \pm 4.6 (10)	–
T_2 -19	24.4 \pm 5.4 (10)	–
T_2 -22	18.3 \pm 6.0 (10)	–
T_2 -23	20.3 \pm 6.0 (10)	–

^a T_2 plants with a Km^R cosuppression phenotype.

Allelic Variation at FRL2 and FRL1

When a partial genomic *FRL2-Ler* sequence obtained from the Monsanto database (Jander et al., 2002) was compared with the standard *FRL2-Col* sequence, four nonsynonymous polymorphisms were found in the 473-amino acid sequence of FRL2. However, when the genomic *FRL2-Ler* fragment used to transform plants was sequenced for this study, only the following two polymorphisms were confirmed: the functional *Ler* allele has an Ala at position 132 (Ala-132; *AluI* CAPS marker, Fig. 2) and a Leu at position 401 (Leu-401), whereas the nonfunctional *Col* allele has a Pro (Pro-132) and a Gln (Gln-401), respectively. It remains to be determined whether both Ala-132 and Leu-401 or only one of those substitutions is critical for the function of *FRL2-Ler*. An alignment of the two FRL2 proteins and their polymorphisms together with FRI is shown in Figure 4.

Figure 4

CLUSTAL alignment of FRL2 and FRI. Polymorphisms in the amino acid sequence of the FRL2 proteins are indicated with the # sign. The underlined sequences above and below the alignment indicate the predicted coiled-coil domains of FRL2 and FRI, respectively. The COILS software at http://www.ch.embnet.org/software/COILS_form.html was used to predict the coiled-coil domains in FRL2 and FRI (Lupas et al., 1991).

```

FRL2-Col      1  -----MTAAESTAAASINQIDER-----
FRL2-Ler     1  -----MTAAESTAAASINQIDER-----
FRI-Sf2      1  MSNYPPTVAAQPTTTANPLLQRHQSEQRRELPKIVETESTSMDITIGQSKQPQFLKSID

FRL2-Col     18  -----KQKLKKAFFDDLQARRSLLS-----PSFNLSWSEIDSHFSSLQSSLFNRLQSAV
FRL2-Ler     18  -----KQKLKKAFFDDLQARRSLLS-----PSFNLSWSEIDSHFSSLQSSLFNRLQSAV
FRI-Sf2     61  ELAAFSVAVETFKRQFDDIQRRHIESIENAIDSKLESNGVVLAAARNNNFHQPMLSPFRNNV

FRL2-Col     66  TSSNSGNIETPTAVTTETPVLWPELRKFCEKNDGKGLGNMYMIENSRKRLSINEELPNAIR
FRL2-Ler     66  TSSNSGNIETPTAVTTETPVLWPELRKFCEKNDGKGLGNMYMIENSRKRLSINEELPNAIR
FRI-Sf2    121  SVETTIVTSQPSQEIIVPETSNNKPEGGRMCBLMCSKGLRKYIYANISDQAKLMEETPSALK

FRL2-Col    126  #
FRL2-Ler    126  CSENPAPLVLDATIEGSYHCSSPSSSSSARAIDVKRIFVLLLEALIEIN-----ANLTD
FRI-Sf2    181  LAKEPAKFFVLDCLGKFLVQGRRAFTEKESPMSSARQVSLLESFLLMPDRGKGKVKTESW

FRL2-Col    180  LRERARTIAYDWKPNIGNK-----PSEALGFLHLVAAPELGSFPSTEEICDYIFLISK
FRL2-Ler    180  LRERARTIAYDWKPNIGNK-----PSEALGFLHLVAAPELGSFPSTEEICDYIFLISK
FRI-Sf2    241  IKDEAETAAVAWAKRRLMTEGGAAAAEKMDARGLLVLVACFGVPSNFRSTDLLDIRMS-G

FRL2-Col    233  YKQATTICKKIGLDRNRIGVLVQKFLDTGRLLVAIRFIYENEMVGEFEPVVSILKTSLKNS
FRL2-Ler    233  YKQATTICKKIGLDRNRIGVLVQKFLDTGRLLVAIRFIYENEMVGEFEPVVSILKTSLKNS
FRI-Sf2    300  SNEIAGALKRSQFLVPMVSGTVESSIKRGMHIEALEMVTTFGMEDKFSAAALVLTSPFKMS

FRL2-Col    293  REAAKRVCAGENYSLKVQNEAT-----DKELSALRAVIKVVKEKNIESEFMEEKLEECVK
FRL2-Ler    293  REAAKRVCAGENYSLKVQNEAT-----DKELSALRAVIKVVKEKNIESEFMEEKLEECVK
FRI-Sf2    360  KESFERAKRKAQSPPLAFKEAATKQLAVLSSVMQCMETHKLDPAKELPGWQIKRQIVSLEK

FRL2-Col    348  ELEDQKAQRKRATKFNSPANPQQPQEQKVDNKRPRVANGSSMEYNLTIPPLRPQQQPPLL
FRL2-Ler    348  ELEDQKAQRKRATKFNSPANPQQPQEQKVDNKRPRVANGSSMEYNLTIPPLRPQQQPPLL
FRI-Sf2    420  DTLQLDKEMEEKARSL#LMEEAALAKRMYNQOIKRPRLSPMEMPPVWSSSYSPIYRDRSF

FRL2-Col    408  PTPSQILQVNPYGLLSSIL-PGVAVPYGNPRALFGSVPAPAS-----RPVYFV
FRL2-Ler    408  PTPSQILQVNPYGLLSSIL-PGVAVPYGNPRALFGSVPAPAS-----RPVYFV
FRI-Sf2    480  PSQRDDDDDEISALVSSVYLGESTSPHRSRRSPEYMVPLRHGGLGRSVYAYEHLAPNSYS

FRL2-Col    455  QQTGYGMP-----PPQYRPPYYPQ-----
FRL2-Ler    455  QQTGYGMP-----PPQYRPPYYPQ-----
FRI-Sf2    540  PGHGHRRLHRQYSPSLVHGQRHBLQYSEPIIHQQQLPYGIQRVYRHSPSEERYLGLSNQRS

FRL2-Col
FRL2-Ler
FRI-Sf2    600  PRSNSSLDPK


```

During mapping of the Col-specific suppressor gene, it appeared that besides *FLC* (near nga249), another *Ler*-specific region linked to *FLC* on chromosome 5, near marker nga139, cosegregated with the very early flowering phenotype (Supplemental Table S1). Moreover, the previously reported complete absence of *FLC* expression in the RIL Col/*Ler-FRI-Sf2*#1 (Schläppi, 2001) was similar to the phenotype of Col-*FRI-Sf2* plants containing single *frl1* mutations (Michaels et al., 2004). This raised the possibility that the *Ler* accession had a nonfunctional allele of *FRL1*. To determine whether *FRL1-Ler* had a lesion, a partial *FRL1-Ler* sequence from the Monsanto database (Jander et al., 2002) was compared to *FRL1-Col*. Interestingly, this comparison suggested that *FRL1-Ler* contained a premature stop codon in the middle of the protein sequence; that is, the codon GAG [Glu-279] in *FRL1-Col* was changed to TAG [stop-279] in *FRL1-Ler*. Cloning and sequencing of *FRL1-Ler* confirmed this significant

polymorphism, together with two other codon changes reported by the Monsanto sequence (a Pro instead of a Thr at position 141 and a deletion of a Lys at position 387). Whether the other two amino acid changes affect FRL1-Ler function in the absence of the stop codon remains to be determined. An alignment of the two FRL1 proteins together with FRI is shown in Figure 5.

Figure 5


CLUSTAL alignment of FRL1 and FRI. Polymorphisms in the amino acid sequence of the FRL1 proteins are shown with the # sign. The stop codon in FRL1-Ler is indicated by an asterisk. The underlined sequences above and below the alignment indicate the predicted coiled-coil domains of FRL1 and FRI, respectively.



To genotype the stop codon in *FRL1-Ler*, a derived CAPS (dCAPS) marker was generated that allows cleavage of *FRL1-Ler*, but not of *FRL1-Col*, by the restriction enzyme *SpeI* (acTAGt recognition site). As shown in Figure 6A, *SpeI* indeed cleaved the dCAPS site in *FRL1-Ler*, but not in *FRL1-Col*, indicating that the dCAPS marker was functional. A dCAPS marker analysis was then done to test whether early flowering Col/*Ler-FRI-Sf2* plants had the nonfunctional *FRL1-Ler* allele, as predicted from their early flowering phenotypes. As shown in Figure 6B, all early flowering Col/*Ler-FRI-Sf2* plants assayed indeed had the *FRL1-Ler* nonsense allele. Taken together, these data suggest that in *FRI*-containing *Ler*, the functional *FRL2-Ler* allele is the major *FRI*-related gene required for *FRI*-mediated late flowering.

Figure 6

A, Graphic representation of the *SpeI* dCAPS marker and an example of how *SpeI* cleaves the *Ler*-specific PCR fragment at the internal stop codon (TGA). B, dCAPS marker-based analysis of *FRL1* alleles in very early flowering Col/*Ler-FRI-Sf2* plants (#1-#10; F₂ progeny from the cross of *Ler-FRI-Sf2* × Col/*FRI-Sf2*) and in late-flowering Col/*FRI-Sf2* and *Ler-FRI-Sf2* controls. C, LER has the same polymorphisms as *Ler* in *FRL1* and *FRL2*. CAPS marker-based analysis of *FRL2* alleles (*AluI* digestion) and dCAPS marker-based analysis of *FRL1* alleles (*SpeI* digestion) in Col, *Ler*, and LER accessions.




To determine whether the nonsense allele of *FRL1-Ler* was a result of mutagenesis in *Ler*, dCAPS analysis on LER, the unmutagenized parent of *Ler* (Rédei, 1962), was performed. The result indicated that *FRL1-LER* also has a premature stop codon and is thus a nonsense allele of *FRL1* (Fig. 6C). Moreover, a CAPS analysis of the LER allele of *FRL2* showed that it, too, contained the *AluI* site observed in *FRL2-Ler* (Fig. 6C). This suggests that the premature stop codon in *FRL1* and the *AluI* polymorphism in *FRL2* are naturally occurring in *Ler* and not the result of mutagenesis.

FRL2-Ler Promotes FRI-Mediated Activation of FLC

To determine whether introducing the genomic copy of *FRL2-Ler* into the early flowering line Col/*Ler-FRI-Sf2#1* restored *FRI*-mediated up-regulation of *FLC*, as suggested by the late-flowering phenotypes of most transformants, a semiquantitative RT-PCR analysis using RNA from late-flowering T₁ plants and untransformed control plants was done. As shown in Figure 7, the level of *FLC* transcript was very low in line Col/*Ler-FRI-Sf2#1*, comparable to *Ler* controls lacking active *FRI-Sf2*. This was consistent with previous results from RNA gel-blot analyses (Schläppi, 2001). By contrast, compared to Col/*Ler-FRI-Sf2#1*, three individual T₁ plants that were late flowering after transformation with *FRL2-Ler* (Fig. 3; Table I) had higher levels of *FLC* transcript. This demonstrated that the late-flowering phenotype, after introducing a genomic copy of *FRL2-Ler* into Col/*Ler-FRI-Sf2#1*, indeed correlated with increased levels of *FLC* transcript. Taken together, the results of these experiments were in agreement with the hypothesis that *FRL2-Ler* is a functional *FRI*-related gene and required for *FRI*-mediated up-regulation of *FLC* transcripts in *Ler*.

Figure 7

Introducing a genomic copy of *FRL2-Ler* into Col/*Ler-FRI-Sf2#1* restores *FRI*-mediated up-regulation of *FLC*. RT-PCR analysis of *FLC* expression in *FRL2-Ler*-transformed T₁ lines 19, 22, and 25 and control plants. C, Col alleles of *FRL* genes; L, *Ler* alleles of *FRL* genes. Functional alleles are shown in bold italics. *ACTIN2/8* (*ACT*) was included as loading control.



Discussion

The focus of this study was to characterize *AFL*, a gene required for *FRI*-mediated late flowering and *FLC* up-regulation in the *Ler* genetic background of *Arabidopsis*. The activity of *AFL* was discovered, because 1/16th of F_2 plants from the cross of *Col-FRI-Sf2* with *Ler-FRI-Sf2* were very early flowering, even in the presence of active *FRI-Sf2* (Schläppi, 2001). Based on phenotype and *FLC* expression, these F_2 plants were determined to be transgressions, because they flowered significantly earlier than the earliest *Ler-FRI-Sf2* parent plant and had no detectable levels of *FLC* transcript (Schläppi, 2001; Fig. 7). This suggested that at least two naturally occurring recessive genes, one derived from *Col* and one from *Ler*, recombined in those F_2 plants, which resulted both in suppression of *FRI*-mediated late flowering and *FLC* up-regulation. The *Ler* component was mapped to the vicinity of *FLC* at the top of chromosome 5, and the *Col* component to the vicinity of *UFO* at the top of chromosome 1. This initially suggested that the *Ler*-specific component was the weak *FLC-Ler* allele and that the *Col*-specific suppressor of *FRI*-mediated up-regulation of *FLC-Ler* was located near *UFO* on chromosome 1. A candidate gene for *AFL* was


FRL2, because it was previously reported that this gene was tightly linked to *UFO* (Michaels et al., 2004). Consistent with this idea, all true-breeding, very early flowering transgressions tested were homozygous for the Col allele of *FRL2*. Therefore, a *Ler*-specific genomic fragment of *FRL2* was genetically transformed into Col/*Ler-FRI-Sf2#1*, an RIL derived from one of the very early flowering transgressions, which resulted in restoration of a late-flowering phenotype and *FLC* up-regulation in a majority of T₁ plants (Figs. 3 and 7). This suggested that *AFL* is identical to the *Ler* allele of *FRL2* and was thus renamed *FRL2-Ler*. By contrast, transformation of a Col-specific genomic fragment of *FRL2* did not restore late flowering in the Col/*Ler-FRI-Sf2#1* RIL, indicating that the *Ler* allele of *FRL2* is functional and that the Col variant *FRL2-Col* is nonfunctional. This interpretation explains, at least in part, why a single *frl2* mutant does not suppress the very late-flowering phenotype of Col-*FRI-Sf2* (Michaels et al., 2004). Taken together, these results suggest that *FRL2* is an active *FRI*-related gene in the *Ler* genetic background and required for *FRI*-mediated late flowering.

The reason why *FRL2-Ler*, but not *FRL2-Col*, is an active allele of *FRL2* is not known at the moment. It does not appear that *FRL2-Col* is expressed at lower levels than the *FRL2-Ler* allele (Fig. 2), and it is thus more likely that either one or both of the amino acid substitutions in *FRL2-Col* has a negative effect on *FRL2* function. It is, however, interesting to speculate that the Ala-132 to Pro-132 change has a more dramatic effect on *FRL2* function than the other substitution. This is because Ala/Pro-132 is located between the two putative protein interacting coiled-coil domains of *FRL2* (Lupas et al., 1991; Fig. 4). As shown in Figure 8, the closest homologs of *FRL2* in Arabidopsis and other plants all have a very conserved Pro two amino acids prior to Ala/Pro-132, but none of them has an additional Pro in a Pro-X-Pro sequence as *FRL2-Col* does. It is, therefore, conceivable that an additional Pro at this position might change overall protein conformation of *FRL2-Col* and affect its ability to interact with protein partners. Because the closest *FRL2* homologs have similarity to ABI3-interacting protein 2 (AIP2) in other plants (Fig. 8), it is intriguing to speculate that the proposed Pro-induced conformational change of *FRL2-Col* compromises its ability to interact with an ABI3-type protein in Arabidopsis. These questions can be addressed in future experiments using chimeras between the two protein sequences and

might define a region or regions critical for the function of FRI-related proteins and potentially for FRI itself.

Figure 8

CLUSTAL alignment of FRL2 alleles and related proteins. The 28-amino acid sequence surrounding the Ala/Pro-132 polymorphism of FRL2 is shown (position 115–142). The site of a conserved Pro (P) is shown by an asterisk, and the site of the Ala/Pro-132 polymorphism between FRL2-Ler and FRL2-Col is shown by the # sign. AC137079, *Medicago truncatula* AIP2; AC182650, *Populus trichocarpa* clone Pop1-11704; AY109393, *Zea mays* clone CL6746_2; AY268951, *Chamaecyparis nootkatensis* AIP2 (CnAIP2); AY596574, *Saccharum officinarum* clone SCCCLR1072A04; BT013472, *Lycopersicon esculentum* clone 132133F; DQ241860, *Solanum tuberosum* clone 021F10; XM_450265, *Oryza sativa* putative AIP2; and XM_478975, *Oryza sativa* putative AIP2 (CnAIP2).



Contrary to the absence of a phenotype for single *frl2* mutants in Col-*FRI-Sf2*, mutations in the *FRL2*-related gene *FRL1* have a strong effect and significantly suppress late flowering in Col-*FRI-Sf2*, indicating that *FRL1* is an active *FRI*-related gene in the Col genetic background (Michaels et al., 2004). By contrast, the premature stop codon in the middle of *FRL1-Ler* indicates that it is a nonsense allele in the *Ler* genetic background and suggests that it is nonfunctional. It is interesting to note, however, that *FRL1-Ler* has a similar Pro substitution as *FRL2-Col* in the region between the two putative coiled-coil domains (Fig. 5), leading to a Pro-X-Pro sequence unique for *FRL2-Col* and *FRL1-Ler*. It is possible that this polymorphism leads to a nonfunctional protein even in the absence of the premature stop codon

further downstream. This possibility, or whether the deletion of a Lys in the C terminus has an additional effect on protein function, can be addressed in future studies using protein chimeras.

The *frl1* and *frl2* single mutant phenotypes thus suggest that active *FRL1* is the main requirement for *FRI*-mediated late flowering in the Col genetic background. However, *FRL2*-Col may have at least some partially overlapping function with *FRL1*, because it was reported that the *frl1frl2* double mutant was slightly earlier flowering than the *frl1* single mutant (Michaels et al., 2004). If *FRL2*-Col is indeed partially functional in Col, then its activity may not be strong enough for *FRI*-mediated up-regulation of the weak *FLC*-Ler allele. Therefore, if the strong *FLC*-Col allele requires mainly *FRL1* activity to affect its *FRI*-mediated up-regulation, is it then possible that *FLC*-Ler is so weak that it requires the cooperation of both *FRL1* and *FRL2* for its *FRI*-mediated *FLC* up-regulation? From this study, the simple answer is no, because *FRL1*-Ler has a premature stop codon at position 279 (Figs. 5 and 6) and is, therefore, an apparent null allele of *FRL1*. This suggests that *FRL2*, but not *FRL1*, is necessary for *FRI*-mediated up-regulation of *FLC* transcripts and late flowering in Ler, and, conversely, that *FRL1*, but not *FRL2*, is active in Col. It is important to note that both the Ler-type nonsense allele of *FRL1* and the *AluI* polymorphism of functional *FRL2*-Ler were also found in the LER accession (Fig. 6), the unmutagenized parent of Ler (Rédei, 1962). This indicates that the described variations at *FRL1* and *FRL2* in Ler are naturally occurring and not the result of mutagenesis.

That *FRL1* is nonfunctional in Ler is most likely the reason why about 1/16th of the F₂ progeny from the cross of Col-*FRI*-Sf2 with Ler-*FRI*-Sf2 were very early flowering, because nonfunctional *FRL1*-Ler is closely linked to weak *FLC*-Ler. Thus, early flowering transgressions homozygous for *FRL2*-Col are probably not only produced in the presence of a weak *FLC*-Ler, as assumed before (Schläppi, 2001), but rather because *FRL1* is linked to *FLC* on chromosome 5. The most likely scenario, therefore, is that *FRL1*-Ler and *FRL2*-Col are the two recessive genes with complementary gene action necessary for an early flowering transgression phenotype. However, *FLC*-Ler always cosegregated with *FRL1*-Ler in the limited number of transgressions tested here and was thus responsible for the very early flowering phenotype of those plants. If this interpretation is correct, then slightly

later-flowering transgressions with recombination events between *FLC*-Col and *FRL1*-Ler should be identified when larger populations of early flowering transgressions are analyzed in future studies.

The observation that neither Col nor Ler have fully active alleles of both *FRL1* and *FRL2* may also explain, at least in part, why the *FLC*-Col allele appeared dominant in F₁ plants from the cross of Col-*FRI*-Sf2 with Ler-*FRI*-Sf2 but semidominant in other crosses (Lee et al., 1994; Schläppi, 2001). The reason for this dominance may be that F₁ plants from the Col × Ler cross have active alleles of both *FRL1* and *FRL2*, which together might effect stronger up-regulation of either *FLC*-Col, *FLC*-Ler, or possibly both, thus compensating for the weak *FLC*-Ler copy in the F₁ hybrid. This question can be addressed in future experiments designed to determine whether *FRL1*-Col alone or a combination of *FRL1*-Col and *FRL2*-Ler enhances *FRI*-mediated up-regulation of weak *FLC*-Ler expression. Conversely, it can also be determined whether a combination of both active alleles produces very late-flowering Col-*FRI*-Sf2 plants that need a longer vernalization period to induce early flowering or whether both active alleles partially up-regulate *FLC* even in the absence of active *FRI*. It is also interesting to note that some F₁ plants from backcrosses between very early flowering transgressions and the late-flowering Ler-*FRI*-Sf2 tester had flowering times between the two parents, whereas other F₁ plants were as late as Ler-*FRI*-Sf2 (Figs. 1 and 2). One explanation for this observation is that *FRL2*-Ler may be semidominant or that the nonfunctional *FRL2*-Col protein somehow interferes with full *FRL2*-Ler activity in some crosses. This does not explain, however, why in other crosses *FRL2*-Ler can be fully dominant (Fig. 1). An alternative explanation is that accession-specific variants of other *FRI*-related genes such as *At1g14900*, *At2g22440*, *At5g27230*, and *At5g48385* (Michaels et al., 2004) or other flowering time genes interact with *FRL2* and thus regulate its activity in a dosage-dependent manner.

In summary, this study presents an example that naturally occurring variation of flowering time genes in Arabidopsis can be uncovered in very well-studied laboratory strains such as Col and Ler, which were previously used for quantitative trait loci mapping of flowering time loci (Alonso-Blanco et al., 1998; Koornneef et al., 1998; Alonso-Blanco and Koornneef, 2000). It is thus possible that *FRL1* and *FRL2* correspond to some previously identified quantitative trait loci

such as *FLG* on chromosome 5 or *AD.121C* on chromosome 1, which were identified in crosses of *Ler* to the Cape Verde Island accession of *Arabidopsis* (Alonso-Blanco et al., 1998). It is important to point out, however, that the large effect on flowering time of these naturally occurring suppressors of *FRI*-mediated late flowering in *Col* and *Ler* was uncovered in this study only because the active *FRI*-Sf2 allele had been introgressed into these laboratory stains (Lee et al., 1994). From this *Col* × *Ler* analysis it appears that there is some selection pressure to maintain an active copy of at least one *FRI*-related gene, even in the absence of an active *FRI* allele. It is thus likely that some of the previously observed, *FRI*-independent, flowering time variations (Gazzani et al., 2003; Werner et al., 2005) could be attributed to natural variation in *FRL1* and *FRL2*. This may be especially true in the case of F₂ plants from the *Ler* × *Col* cross where earliness was linked to the recessive *ms1* allele of *Ler*, which maps near *FRL1* (Koornneef et al., 1994). It is now possible to test in future studies whether the *FRL1* and *FRL2* polymorphisms identified here can be correlated with flowering time differences and the adaptation to ecological niches of the large number of available *Arabidopsis* accessions collected from the wild.

Materials and Methods

Plant Material

Early flowering *Arabidopsis* (*Arabidopsis thaliana*) accessions *Ler* and wild-type *LER* (Michaels et al., 2003) were kindly provided by T-p. Sun (Duke University) and R. Amasino (University of Wisconsin-Madison), respectively. A *Col-0* accession and lines *Col-FRI-Sf2* and *Ler-FRI-Sf2* containing the dominant *FRI*-Sf2 were kindly provided by R. Amasino and were described previously (Lee et al., 1993, 1994). *Col/Ler-FRI-Sf2*#1, #3, and #5 lines were homozygous for *FRI*-Sf2 and *FLC-Ler* and were derived from very early flowering F₂ plants of crosses between *Col-FRI-Sf2* and *Ler-FRI-Sf2*. *Col/Ler-FRI-Sf2*#1 was described previously (Schläppi, 2001). A RIL was generated with *Col/Ler-FRI-Sf2*#1 by selfing the plant and single seed propagation for five consecutive generations (Supplemental Table S2).

Plant Growth Conditions

Per sterile petri dish (90-mm plate), about 100 surface-sterilized seeds were grown on 0.8% agar-solidified medium (Difco) containing half-strength Murashige and Skoog (Gibco BRL) salts without Suc. Petri dishes were placed at 4°C for up to 2 d to break seed dormancy, then grown under cool fluorescent light with a 16-h-light/8-h-dark long-day photoperiod or a 12-h-light/12-h-dark short-day photoperiod with approximately $100 \mu\text{mol m}^{-2} \text{s}^{-1}$ photon flux and about 22°C day/night temperature. After 10 to 14 d, plantlets were transferred from petri dishes to soil (2:1:1 mix of peatmoss:vermiculite:perlite) into 2-inch pots (four plants/pot; 32 pots/flat) and grown under cool fluorescent light with a 16-h-light/8-h-dark long-day photoperiod, 20°C \pm 1°C day/night temperature, and about 60% to 70% relative humidity. Flats were watered three times per week with 0.1 g/L 15-16-17 Peters fertilizer (Grace Sierra).

Flowering Time Analysis

Flowering time of individual plants was measured as the RLN produced by the main shoot when its floral bolt was 0.5 to 1 cm high. For some plants, the number of cauline leaves on the floral bolt was also recorded (Table I).

Isolation of Genomic DNA, Cloning, Sequencing, and Plant Transformation

Genomic fragments of *FRL2* were isolated from Ler or Col DNA using the ExTaq DNA polymerase (Takara) and primers 5'-AAGAAAAGGTACCATGTCGTCGT-3' (*Kpn*I site underlined) and 5'-ATTGGCTTATTCGGATCCGTATG-3' (*Bam*HI site underlined). The primers were designed to include most of the 5' and 3' noncoding region between *FRL2* and the neighboring genes (525-bp 5' upstream from the start codon and 666-bp 3' downstream from the stop codon). The 2.6-kb PCR fragment was ligated into pGEM-T (Promega). Cloned *FRL2* DNA was sequenced using the MGW sequencing service (MGW-Biotech). The *FRL2*-Ler genomic fragment was removed as a *Kpn*I-*Bam*HI fragment from pGEM-T and ligated into the binary vector pPZP211 (Hajdukiewicz et al., 1994), cloned in *Escherichia coli*, and

introduced into *Agrobacterium tumefaciens* strain ABI using the freeze-thaw method (Chen et al., 1994). Genomic *FRL2-Ler* and *FRL2-Col* fragments were transformed into very early flowering line Col/*Ler-FRI-Sf2#1* using the floral dip method (Clough and Bent, 1998). A *FRL1-Ler* genomic fragment was isolated from *Ler* DNA using primers 5'-AGCCAAAGAAATCTTAGAGATC-3' and 5'-TAAGATCTTATTGTGCGAGATGC-3' and sequenced using the Agencourt sequencing service (Agencourt Bioscience).

RNA Isolation and Analysis

RNA was isolated from petri dish-grown whole seedlings or leaves of adult, soil-grown plants. Plant tissue was ground to a fine powder on dry ice in a mortar and pestle with added liquid N₂. Total RNA was isolated by a modified miniprep procedure as described previously (Schläppi, 2001).

For RT-PCR analysis, first-strand cDNA synthesis was performed on 1 to 2 µg of total RNA using primer 5'-GGCCACGCGTCGACTAC(T)₁₇-3' and Superscript II reverse transcriptase according to the manufacturer's instruction (Invitrogen). *FLC* was amplified using intron-spanning primers 5'-GAAATCAAGCGAATTGAGAAC-3' and 5'-TAAGATTCTCAACAAGCTTCAAC-3'; *ACT2/8* was amplified using intron-spanning primers 5'-ATGAAGATTAAGGTCGTGGCA-3' and 5'-TCCGAGTTTGAAGAGGCTAC-3'; *FRL2* was amplified using the gene-specific primer 5'-TACCGCCCACCGTACTATCCT-3' and, because the gene is intronless, the 3'-RACE primer 5'-GGCCACGCGTCGACTAC-3'. After 3 min denaturing of DNA at 95°C, hot-start PCR was performed using the *Taq* DNA polymerase (Promega) and the following cycles: 25 cycles of 95°C for 15 s, 53°C for 15 s, and 72°C for 30 s for *ACT2/8*; 28 cycles of 95°C for 15 s, 56°C for 15 s, and 72°C for 30 s for *FRL2*; and 30 cycles of 95°C for 15 s, 54°C for 15 s, and 72°C for 30 s for *FLC*. PCR fragments were separated on a 1.2% agarose gel and either visualized by ethidium bromide staining (Sambrook et al., 1989) or blotted onto Protran nitrocellulose membranes (Schleicher & Schuell), hybridized with radiolabeled probes at a concentration of 1 × 10⁶ cpm/mL as described previously (Schläppi, 2001), and visualized using a Storm PhosphorImager (Molecular Dynamics/Amersham Bioscience).

SSLP Mapping and CAPS/dCAPS Analysis

A total of 18 SSLP markers (Bell and Ecker, 1994; Lukowitz et al., 2000) were used to map *AFL*-Col to the top of chromosome I, about 2 cM south of *UFO* (Supplemental Fig. S1; Supplemental Table S1). For CAPS analysis of *FRL2* alleles, PCR primers 5'-AGCTTCCTAATGCGATTTCGAT-3' and 5'-CAAAACGAAGATCCTCTTCAC-3' were used to amplify a 136-bp product spanning a *Ler*-specific *AluI* site. After digestion with *AluI* (New England Biolabs), the *Ler*-specific product was cleaved into 99-bp and 37-bp fragments, while the Col-specific product remained intact. Cleavage products were separated on a 2% agarose gel and visualized by ethidium bromide staining. For dCAPS (Neff et al., 1998) analysis of *FRL1* alleles, dCAPS forward primer 5'-TTTTAGCAGTCAAATTCATGTAC-3' (mismatch underlined) and regular reverse primer 5'-TCTTTATCAGAGGCTTCGTTC-3' were used to amplify a 161-bp fragment. The C nucleotide at the end of the dCAPS primer introduced a *SpeI* site spanning a stop codon in *FRL1-Ler* (ACTAGT) but not in *FRL1-Col* (ACGAGT). After digestion with *SpeI*, the *Ler*-specific product was cleaved into 139-bp and 22-bp fragments, while the 161-bp Col-specific product remained intact. Cleavage products were separated on a 2% agarose gel and visualized by ethidium bromide staining.

Sequence data from this article can be found in the GenBank/EMBL data libraries under accession numbers EF052677 and EF052678.

Supplemental Data

The following materials are available in the online version of this article.

- **Supplemental Figure S1.** SSLP mapping of *AFL-Col*.


Mappig of <i>AFL-Col</i> using 17 individual very-early-flowering F2 lines from the cross of <i>Ler-FRI-Sf2</i> x <i>Col-FRI-Sf2</i>			
Chrom. I markers	# of Col chromosomes	# of Ler chromosomes	Recombination frequency
nga63	25	9	35
ciw12	30	4	12.5
nga248	31	3	9
<i>UFO</i>	32	2	6
nga280	23	11	32
nga111	16	18	53
Mappig of the early <i>Ler</i> component using 17 individual very-early-flowering F2 lines			
Chrom. V markers	# of Col chromosomes	# of Ler chromosomes	Recombination frequency
CTR1	3	31	9
nga249	1	33	3
nga139	3	31	9

-

- **Supplemental Table S1.** Recombination frequencies.

Characterization of near isogenic (F5 generation) recombinant inbred line Col/Ler-FRI-Sf2#1:		
Chromosome	Marker	Genotype (C = Columbia; L = Ler)
I	nga63	C
I	ciw12	C
I	nga248	C
I	<i>UFO</i>	C
I	<i>FRL2</i>	C
I	nga280	C
I	nga111	C
II	ciw2	L
II	ciw3	L
II	<i>ER</i>	C
II	nga163	C
III	nga172	C
III	nga162	L
III	nga6	C
IV	nga8	C
IV	nga1107	L
V	CTR1	L
V	nga249	L
V	<i>FRL1</i>	L
V	ciw9	C
V	ciw10	C

- **Supplemental Table S2.** RIL characterization.



Acknowledgments

I thank R. Amasino and S. Michaels for encouragement and insightful discussions and Y. Pan for expert advice and help with the RT-PCR experiments.

Notes

¹This work was supported in part by the National Research Initiative (competitive grant no. 2001-35100-10688) from the U.S. Department of Agriculture, and by the National Science Foundation (phosphorimager provided by instrumentation grant no. D.B.I. 0100667).

The author responsible for distribution of materials integral to the findings presented in this article in accordance with the policy described in the Instructions for Authors (www.plantphysiol.org) is: Michael R. Schläppi (ude.etteuqram@ippalhcs.leahcim).

[^W]The online version of this article contains Web-only data.

[^{OA}]Open Access articles can be viewed online without a subscription.

www.plantphysiol.org/cgi/doi/10.1104/pp.106.085571

References

- Alonso-Blanco C, El-Din El-Assal S, Coupland G, Koornneef M** (1998) Analysis of natural allelic variation at flowering-time loci in the Landsberg *erecta* and Cape Verde Island ecotypes of *Arabidopsis thaliana*. *Genetics* 149: 749–764
- Alonso-Blanco C, Koornneef M** (2000) Naturally occurring variation in *Arabidopsis*: an underexploited resource for plant genetics. *Trends Plant Sci* 5: 22–29
- Amasino R** (2004) Vernalization, competence, and the epigenetic memory of winter. *Plant Cell* 16: 2553–2559
- Bell CJ, Ecker JR** (1994) Assignment of 30 microsatellite loci to the linkage map of *Arabidopsis*. *Genomics* 19: 137–144
- Boss PK, Bastow RM, Mylne JS, Dean C** (2004) Multiple pathways in the decision to flower: enabling, promoting, and resetting. *Plant Cell* 16: S18–S31
- Chen H, Nelson RS, Sherwood JL** (1994) Enhanced recovery of transformants of *Agrobacterium tumefaciens* after freeze-thaw transformation and drug selection. *Biotechniques* 16: 664–669
- Choi K, Kim S, Kim SY, Kim M, Hyun Y, Lee H, Choe S, Kim S-G, Michaels S, Lee I** (2005) *SUPPRESSOR OF FRIGIDA3* encodes a nuclear ACTIN-RELATED PROTEIN6 required for floral repression in *Arabidopsis*. *Plant Cell* 17: 2647–2660

Clough SJ, Bent AF (1998) Floral dip: a simplified method for *Agrobacterium*-mediated transformation of *Arabidopsis thaliana*. *Plant J* 16: 735–743

Deal RB, Kandasamy MK, McKinney EC, Meagher RB (2005) The nuclear actin-related protein ARP6 is a pleiotropic developmental regulator required for the maintenance of *FLOWERING LOCUS C* expression and repression of flowering in *Arabidopsis*. *Plant Cell* 17: 2633–2646

Gazzani S, Gendall AR, Lister C, Dean C (2003) Analysis of the molecular basis of flowering time variation in *Arabidopsis* accessions. *Plant Physiol* 132: 1107–1114

Hajdukiewicz P, Svab Z, Maliga P (1994) The small, versatile *pPZP* family of *Agrobacterium* binary vectors for plant transformation. *Plant Mol Biol* 25: 989–994

Jander G, Norris SR, Rounsley SD, Bush DF, Levin IM, Last RL (2002) *Arabidopsis* map-based cloning in the post-genome era. *Plant Physiol* 129: 440–450

Johanson U, West J, Lister C, Michaels S, Amasino A, Dean C (2000) Molecular analysis of *FRIGIDA*, a major determinant of natural variation in *Arabidopsis* flowering time. *Science* 290: 344–347

Koornneef M, Alonso-Blanco C, Blankestijn-de Vries H, Hanhart CJ, Peeters AJ (1998) Genetic interactions among late-flowering mutants of *Arabidopsis*. *Genetics* 148: 885–892

Koornneef M, Blankestijn-de Vries H, Hanhart C, Soppe W, Peeters T (1994) The phenotype of some late-flowering mutants is enhanced by a locus on chromosome 5 that is not effective in the Landsberg *erecta* wild-type. *Plant J* 6: 911–919

Laibach F (1937) Über sommer- und winterannuelle Rassen von *Arabidopsis thaliana* (L.) Heynh. Ein Beitrag zur Ätiologie der Blütenbildung. *Beitr Biol Pflanz* 28: 173–210

Le Corre V, Roux F, Reboud X (2002) DNA polymorphism at the *FRIGIDA* gene in *Arabidopsis thaliana*: extensive nonsynonymous variation is consistent with local selection for flowering time. *Mol Biol Evol* 19: 1261–1271

- Lee I, Bleecker A, Amasino R** (1993) Analysis of naturally occurring late-flowering in *Arabidopsis thaliana*. *Mol Gen Genet* 237: 171–176
- Lee I, Michaels S, Masshardt A, Amasino RM** (1994) The late-flowering phenotype of *FRIGIDA* and mutations in *LUMINIDEPENDENS* is suppressed in the Landsberg *erecta* strain of *Arabidopsis*. *Plant J* 6: 903–909
- Lukowitz W, Gillmor CS, Scheible W-R** (2000) Positional cloning in *Arabidopsis*: why it feels good to have a genome initiative working for you. *Plant Physiol* 123: 795–805
- Lupas A, van Dyke M, Stock H** (1991) Predicting coiled coils from protein sequences. *Science* 252: 1162–1164
- Michaels SD, Amasino RM** (1999) *FLOWERING LOCUS C* encodes a novel MADS domain protein that acts as a repressor of flowering. *Plant Cell* 11: 949–956
- Michaels SD, Bezerra IC, Amasino RM** (2004) *FRIGIDA*-related genes are required for the winter-annual habit in *Arabidopsis*. *Proc Natl Acad Sci USA* 101: 3281–3285
- Michaels SD, He Y, Scortecci KC, Amasino RM** (2003) Attenuation of *FLOWERING LOCUS C* activity as a mechanism for the evolution of summer-annual flowering behavior in *Arabidopsis*. *Proc Natl Acad Sci USA* 100: 10102–10107
- Neff MM, Neff JD, Chory J, Pepper AE** (1998) dCAPS, a simple technique for the genetic analysis of single nucleotide polymorphisms: experimental applications in *Arabidopsis thaliana* genetics. *Plant J* 14: 387–392
- Noh YS, Amasino RM** (2003) *PIE1*, an *ISWI* family gene, is required for *FLC* activation and floral repression in *Arabidopsis*. *Plant Cell* 15: 1671–1682
- Nordborg M, Bergelson J** (1999) The effect of seed germination and rosette cold treatment on germination and flowering time in some *Arabidopsis thaliana* (Brassicaceae) ecotypes. *Am J Bot* 86: 470–475
- Oh S, Zhang H, Ludwig P, van Nocker S** (2004) A mechanism related to the yeast transcriptional regulator *Paf1c* is required for expression of

the Arabidopsis *FLC/MAF* MADS box gene family. *Plant Cell* 16: 2940–2953

Ratcliffe OJ, Kumimoto RW, Wong BJ, Riechmann JL (2003) Analysis of the Arabidopsis *MADS AFFECTING FLOWERING* gene family: *MAF2* prevents vernalization by short periods of cold. *Plant Cell* 15: 1159–1169

Ratcliffe OJ, Nadzan GC, Reuber TL, Riechmann JL (2001) Regulation of flowering in Arabidopsis by an *FLC* homologue. *Plant Physiol* 126: 122–132

Rédei GP (1962) Single locus heterosis. *Z Vererbungsl* 93: 164–170

Sambrook J, Fritsch EF, Maniatis T (1989) *Molecular Cloning: A Laboratory Manual*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY

Sanda SL, Amasino RM (1996) Ecotype-specific expression of a flowering mutant phenotype in *Arabidopsis thaliana*. *Plant Physiol* 111: 641–644

Schläppi M (2001) RNA levels and activity of *FLOWERING LOCUS C* are modified in mixed genetic backgrounds of *Arabidopsis thaliana*. *Int J Plant Sci* 162: 527–537

Scortecci K, Michaels SD, Amasino RM (2001) Identification of a MADS-box gene, *FLOWERING LOCUS M*, that represses flowering. *Plant J* 26: 229–236

Sheldon CC, Burn JE, Perez PP, Metzger J, Edwards JA, Peacock WJ, Dennis ES (1999) The *FLF* MADS box gene: a repressor of flowering in Arabidopsis regulated by vernalization and methylation. *Plant Cell* 11: 445–458

Sheldon CC, Rouse DT, Finnegan EJ, Peacock WJ, Dennis ES (2000) The molecular basis of vernalization: the central role of *FLOWERING LOCUS C (FLC)*. *Proc Natl Acad Sci USA* 97: 3753–3758

Shindo C, Aranzana MJ, Lister C, Baxter C, Nicholls C, Nordborg M, Dean C (2005) Role of *FRIGIDA* and *FLOWERING LOCUS C* in determining variation in flowering time of Arabidopsis. *Plant Physiol* 138: 1163–1173

NOT THE PUBLISHED VERSION; this is the author's final, peer-reviewed manuscript. The published version may be accessed by following the link in the citation at the bottom of the page.

Werner JD, Borevitz JO, Uhlenhaut NH, Ecker JR, Chory J, Weigel D
(2005) *FRIGIDA*-independent variation in flowering time of natural *Arabidopsis thaliana* accessions. *Genetics* 170: 1197–1207

Plant Physiology, Vol. 142, No. 4 (December 2006): pg. 1728-1738. [DOI](#). This article is © American Society of Plant Biologists and permission has been granted for this version to appear in e-Publications@Marquette. American Society of Plant Biologists does not grant permission for this article to be further copied/distributed or hosted elsewhere without the express permission from American Society of Plant Biologists.