

Marquette University
e-Publications@Marquette

Biological Sciences Faculty Research and
Publications

Biological Sciences, Department of

7-20-2007

The Exosome Subunit Rrp44 Plays a Direct Role in RNA Substrate Recognition

Claudia Schneider
University of Edinburgh

James T. Anderson
Marquette University, james.anderson@marquette.edu

David Tollervey
Marquette University

Accepted version. *Cell Biology*, Vol. 27, No. 2 (July 20, 2007): 324-331. [DOI](#). © 2007 Elsevier Inc.
Used with permission.

Marquette University

e-Publications@Marquette

Biology Faculty Research and Publications/College of Arts and Sciences

This paper is NOT THE PUBLISHED VERSION; but the author's final, peer-reviewed manuscript. The published version may be accessed by following the link in the citation below.

Molecular Cell, Vol. 27, No. 2 (July 2007): 324-331. DOI. This article is © Elsevier and permission has been granted for this version to appear in [e-Publications@Marquette](#). Elsevier does not grant permission for this article to be further copied/distributed or hosted elsewhere without the express permission from Elsevier.

The Exosome Subunit Rrp44 Plays a Direct

Claudia Schneider

Wellcome Trust Centre for Cell Biology, University of Edinburgh, Edinburgh EH9 3JR, UK

James T. Anderson

Department of Biological Sciences, Marquette University, Milwaukee, WI 53201, USA

David Tollervey

Wellcome Trust Centre for Cell Biology, University of Edinburgh, Edinburgh EH9 3JR, UK

Summary

The exosome [plays](#) key roles in RNA [maturation](#) and surveillance, but it is unclear how target RNAs are identified. We report the functional characterization of the [yeast](#) exosome component Rrp44, a member of the RNase II family. [Recombinant](#) Rrp44 and the purified TRAMP [polyadenylation](#) complex each specifically recognized tRNA^{Met} lacking a single m¹A₅₈ modification, even in the presence of a large excess of total tRNA. This tRNA is otherwise mature and functional in translation in vivo but is presumably subtly misfolded. Complete [degradation](#) of the hypomodified tRNA required both Rrp44 and the poly(A) [polymerase](#) activity of TRAMP. The intact exosome lacking only the [catalytic activity](#) of Rrp44 failed to degrade tRNA^{Met}, showing this to be a specific Rrp44 substrate. [Recognition](#) of hypomodified tRNA^{Met} by Rrp44 is genetically separable from its catalytic activity on other substrates, with the [mutations](#) mapping to distinct regions of the protein.

Introduction

The nuclear and cytoplasmic forms of the exosome share ten common components, often referred to as the “core” exosome, while the nuclear complex contains an additional 3′ [exonuclease](#) Rrp6 (reviewed in [Houseley et al. \[2006\]](#)). Recent studies proposed that Rrp44 is the only catalytically active [nuclease](#) in the [yeast](#) core exosome ([Dziembowski et al., 2007](#), [Liu et al., 2006](#)). The domain organization of Rrp44 resembles that of bacterial RNase II (see [Figure 1A](#)) and RNase R ([Vincent and Deutscher, 2006](#)), with a cold shock protein [RNA-binding domain](#) (CSD), a catalytic RNB domain, and a C-terminal S1 RNA-binding domain. However, Rrp44 has an additional [N-terminal](#) PinC domain that has no equivalent in the bacterial enzymes.




Figure 1. A [Point Mutation](#) in the RNB Motif I of Rrp44 Inactivates the Enzyme without Affecting [RNA Binding](#)

(A) Domain structures of *E. coli* RNase II and *S. cerevisiae* Rrp44. RNB motifs I–IV (adapted from [Frazao et al. \[2006\]](#)) are shown. [Positions](#) of Rrp44 point mutations are indicated.

(B) [Exonuclease](#) activity of [wild-type](#) GST-Rrp44, GST-Rrp44-20, GST-Rrp44-cat, or free GST on a 5′ end-labeled 110 nt pre-tRNA^{Ser} transcript. [Recombinant protein](#) (100 fmol) and 5 fmol RNA were incubated for 10 min at 37°C. Control: mock-digested substrate without recombinant protein. Reaction products were separated on a 12% polyacrylamide/8 M [urea](#) gel and visualized by autoradiography. The full-length RNA and ~3 nt [degradation](#) products are indicated on the right.

(C) RNA-binding assay using wild-type GST-Rrp44, GST-Rrp44-cat, GST-Rrp44-20, or free GST and 5′ end-labeled 110 nt pre-tRNA^{Ser} transcript. Substrate and reaction conditions were as described in (B). After incubation at 37°C, [samples](#) were immediately put on ice and then analyzed on a native 4% polyacrylamide gel. The positions of RNA-protein complexes, free full-length RNA, and ~3 nt degradation products are indicated on the right.

The TRAMP4 complex is a major nuclear cofactor for the exosome ([LaCava et al., 2005](#), [Vanacova et al., 2005](#)) and comprises the poly(A) [polymerase](#) Trf4, the putative ATP-dependent [RNA helicase](#) Mtr4, and either Air1 or Air2, which are zinc-knuckle putative [RNA-binding proteins](#) that appear to be functionally redundant. The TRAMP/exosome system functions in the surveillance of many nuclear RNA [species](#). In yeast strains lacking the activity of the tRNA 1-methyladenosine₅₈ [methyltransferase](#) (*trm6-504*), several tRNAs fail to be methylated, including the initiator tRNA^{Met}. This tRNA has a unique structure, which is particularly sensitive to loss of

[methylation](#) ([Basavappa and Sigler, 1991](#)). The hypomethylated tRNA_i^{Met} is stable and functional for translation at 25°C; however, transfer to 37°C leads to [degradation](#) of the [precursor](#) pre-tRNA_i^{Met} ([Kadaba et al., 2006](#)). Hypomodified pre-tRNA_i^{Met} was stabilized by [mutations](#) in the exosome components Rrp44 (*rrp44-20*) or Rrp6 (*rrp6Δ*) or in the [TRAMP complex](#) (*mtr4-20*, *trf4-20*), but it was unclear which of these factors specifically recognizes and degrades the defective pre-tRNA ([Kadaba et al., 2004](#)).

Results

Mutations in Rrp44 Alter RNA Binding and Exonuclease Activities In Vitro

The *rrp44-20* [mutation](#) was identified as a suppressor of the temperature-sensitive [lethal mutation](#) *trm6-504* ([Kadaba et al., 2004](#), [Zuo et al., 2006](#)). The genomic location of the *rrp44-20* [allele](#) was identified by [linkage analysis](#). [Sequencing](#) of the [mutant](#) allele revealed a single [nucleotide](#) substitution G₂₄₉₈ → A resulting in a Gly₈₃₃ → Asp substitution in Rrp44, within domain IV of the RNB motif (G833D; see [Figure 1A](#)).

Analysis of RNase II in *E. coli* indicated that [RNA binding](#) and [catalysis](#) involve distinct regions of the protein ([Frazao et al., 2006](#), [Zuo et al., 2006](#)). This was confirmed by the isolation of a [point mutation](#) (D209N) in RNB motif I (see [Figure 1A](#)), which inactivates the [exonuclease](#) activity of RNase II without affecting RNA binding ([Amblar and Arraiano, 2005](#)). Sequence comparisons predicted that an Asp₅₅₁ → Asn mutation (D551N: *rrp44-cat*) should abolish the [catalytic activity](#) of Rrp44 ([Figure 1A](#)) ([Dziembowski et al., 2007](#)).

[Wild-type](#) Rrp44, Rrp44-20, and Rrp44-cat were expressed in *E. coli* as GST fusions and purified on [glutathione](#) Sepharose (see [Figure S1](#) in the [Supplemental Data](#) available with this article online). These were tested in exonuclease and RNA-binding assays using in vitro-transcribed, 5' end-labeled pre-tRNA^{Ser} as a fully unmodified substrate ([Figures 1B](#) and [1C](#)). In [contrast](#) to previous analyses, the assays were performed at low [magnesium](#) concentrations (0.25 mM) similar to the reported intracellular concentration of eukaryotic cells (0.1–1 mM) ([Romani and Scarpa, 1992](#)).

In the in vitro exonuclease experiment shown in [Figure 1B](#), ~12 ng (100 fmol) protein and ~5 fmol RNA were incubated for 10 min at 37°C. The reaction products were separated on a denaturing 12% polyacrylamide/8 M [urea](#) gel and visualized by autoradiography. Both Rrp44 and Rrp44-20 degraded pre-tRNA^{Ser} in an apparently processive manner, leaving a short [oligonucleotide](#) (~3 nt) product ([Figure 1B](#)). In contrast, Rrp44-cat failed to detectably degrade pre-tRNA^{Ser} ([Figure 1B](#)) and was also inactive on a 37 nt pBS transcript, poly(A₃₀), poly(U₃₀), and tRNA_i^{Met} ([Figure 2](#), [Figure S2](#), and data not shown). This confirmed that Asp₅₅₁ is part of the catalytic site and that the activity seen for wild-type Rrp44 is not due to a contaminating *E. coli* exonuclease.

To confirm that the impaired [nuclease](#) activity of Rrp44-cat does not result from reduced substrate binding, we performed RNA band shift assays ([Figure 1C](#)). Reaction conditions were as above, except that the [samples](#) were put on ice after incubation at 37°C and separated on a precooled, native 4% polyacrylamide gel. Incubation of pre-tRNA^{Ser} with Rrp44 or Rrp44-20 generated [degradation](#) products with greater gel mobility than the free substrate ([Figure 1C](#), lanes 6 and 7). In addition, a low level of slower-migrating RNA was detected. We interpret this as showing that they bind unmodified pre-tRNA^{Ser} but then rapidly degrade the RNA. In contrast, Rrp44-cat generated only bands with slowed migration ([Figure 1C](#), lane 4), which represent stable RNA-protein complexes. The wild-type protein also formed stable RNA-protein complexes when incubated in the presence of 10 mM EDTA to inhibit nuclease activity (data not shown).

We conclude that the D551N mutation in Rrp44-cat abolishes the exonucleolytic activity of Rrp44 without affecting its ability to bind RNA, consistent with the presence of functionally separable domains that are required for either RNA binding or catalysis, as previously reported for *E. coli* RNase II ([Frazao et al., 2006](#), [Zuo et al., 2006](#)). However, Rrp44-20 was apparently unaffected in RNA binding or catalysis on a fully unmodified pre-tRNA^{Ser} substrate.

To characterize the putative defect in Rrp44-20 activity on a more biologically relevant substrate, we purified and 5' end labeled the native, hypomodified tRNA_i^{Met} from a *trm6Δ* yeast strain (Anderson et al., 2000) (Figure 2A). To facilitate analyses of RNA binding, we also generated a point mutation (Gly₉₁₆ → Glu; Rrp44-S1) in the S1 RNA-binding domain, which is predicted to disrupt its activity. This was expressed in *E. coli* as a GST fusion (Figure S1).




Figure 2. Recognition and Degradation of Native, Hypomodified tRNA_i^{Met} by Rrp44 and TRAMP

(A) Exonuclease activity of wild-type GST-Rrp44, GST-Rrp44-20, or GST-Rrp44-S1 on a 5' end-labeled, native, hypomodified tRNA_i^{Met} substrate. Recombinant protein (3, 6, 12, or 60 fmol) was incubated with 5 fmol RNA for 90 min at 30°C in the presence of 0.25 mM MgCl₂. Reaction products were separated on a 12% polyacrylamide/8 M urea gel and visualized by autoradiography. Full-length and partially degraded tRNA_i^{Met} species are indicated on the right.

(B) RNA-binding assay using wild-type or mutant GST-Rrp44 proteins and 5' end-labeled, native hypomodified tRNA_i^{Met}. Recombinant protein (3, 6, or 12 fmol) was incubated with 5 fmol RNA for 10 min at 30°C in the presence of 0.25 mM MgCl₂. Reactions were then analyzed as described in Figure 1C. The lower panel shows PhosphorImager quantification of the same experiment.

(C) Combined exonuclease/polyadenylation assay using 2 μl TAP-purified TRAMP complex, 200 fmol GST-Rrp44 fusion protein (wild-type, 44-20, or cat), and 50 fmol 5' end-labeled, native hypomodified

tRNA_i^{Met} substrate. Reactions were performed at 30°C with 0.5 mM MgCl₂ and 1 mM ATP for the times indicated. Products were separated on a 12% polyacrylamide/8 M urea gel and visualized by autoradiography. Full-length tRNA_i^{Met}, polyadenylated RNA, and degraded RNA species are indicated.

On the native, hypomodified tRNA_i^{Met} substrate, wild-type Rrp44 generated a partial degradation product (Figure 2A). The gel mobility of this would correspond to removal of the 4 nt 3' overhang comprising the CCA tail and the 3' encoded "discriminator" base. In addition, a 3 nt product was detected (lower panel). We predict that this results from complete, Rrp44-mediated degradation of tRNA fragments present in the input. In contrast, Rrp44-20 and Rrp44-S1 showed no detectable nuclease activity (Figure 2A).

To determine whether the Rrp44-20 mutation specifically impaired recognition of tRNA_i^{Met}, we performed RNA gel mobility assays (Figure 2B). Rrp44-20 showed reduced binding to the hypomodified tRNA_i^{Met} compared to the wild-type or Rrp44-cat protein (quantified in lower panel of Figure 2B). The reduction in binding of Rrp44-20 to hypomodified tRNA_i^{Met} was modest but was reproducibly observed in six independent experiments. In contrast, RNA binding was greatly reduced but not abolished in the Rrp44-S1 mutant. To analyze RNA binding in the absence of substrate degradation, the *rrp44-20* and *rrp44-S1* mutations were combined with the catalytic (D551N) mutation to generate Rrp44-20 + cat and Rrp44-S1 + cat. This abolished formation of degradation products (data not shown) but did not clearly alter the relative abilities of the wild-type, Rrp44-20, and Rrp44-S1 proteins to bind hypomodified tRNA_i^{Met} (Figure 2B).

These results suggested that the suppression phenotype of *rrp44-20* might result from impaired binding to the misfolded pre-tRNA_i^{Met}. However, subsequent analyses revealed that Rrp44-20 also showed reduced binding to other substrates tested, poly(A₃₀) (Figure S2B), and the 37 nt pBS transcript (Figure S2D). In contrast to the reduced band shift, the catalytic activity of Rrp44-20 was not clearly impaired on the 37 nt transcript (Figure S2C) and was only mildly impaired on the poly(A₃₀) substrate (Figure S2A). Rrp44-S1 showed greatly reduced catalytic activity on both substrates.

Comparing Rrp44 binding and nuclease activity on various RNA substrates, we suggest that wild-type Rrp44 can bind and completely degrade unstructured substrates but can only partially degrade the hypomodified, but otherwise mature, tRNA_i^{Met} (Figure 2). This indicates that Rrp44 can remove the presumably unstructured 3' overhang in the hypomodified tRNA_i^{Met} on its own, but it needs cofactors to completely degrade this near-native tRNA, possibly to assist in opening the stable structure of the RNA.

The Rrp44-20 protein showed marked defects in stable RNA binding on some, but not all, substrates. This reduced binding has only limited effects on the ability of the protein to degrade unstructured unmodified pre-tRNA^{Ser}, poly(A₃₀), or 37 nt pBS substrates but completely abolished the partial 3' degradation of hypomodified tRNA_i^{Met} seen for the wild-type protein (Figure 2A). From these results we conclude that the suppression phenotype of *rrp44-20* is probably not due to reduced overall binding of the hypomodified tRNA_i^{Met} but might result from impaired positioning of the unstructured 3' overhang for catalysis. This indicates that a single exosome subunit, Rrp44, is directly involved in the recognition of its cognate substrate.

TRAMP Stimulates the Degradation of Hypomodified tRNA_i^{Met} by Rrp44

The identification of mutations in the TRAMP complex as suppressors of the *trm6-504* phenotype (Kadaba et al., 2004) suggested that TRAMP could be a cofactor for Rrp44-mediated degradation of tRNA_i^{Met}.

To test this hypothesis in vitro, we incubated native, 5' end-labeled hypomodified tRNA_i^{Met} with recombinant GST-tagged Rrp44 and/or TAP-purified TRAMP complex, in the presence of 1 mM ATP (Figure 2C and Figure S3). The RNA substrate was polyadenylated by the TRAMP complex in the absence of Rrp44 (Figure 2C, lane 4), indicating that TRAMP can recognize hypomodified tRNA_i^{Met}, consistent with previous results (Vanacova et al., 2005). Addition of the TRAMP complex stimulated the activity of Rrp44 (Figure 2C, lanes 6 and 8), enhancing

both the fraction of the tRNA that was degraded (upper panel) and the yield of the 2 nt product of complete digestion (lower panel). TRAMP did not stimulate the activity of Rrp44-cat (lane 16). The activity of Rrp44-20 was slightly stimulated by TRAMP (Figure 2C, compare lanes 10 and 12), but degradation was much less robust than with the wild-type protein. The stimulation of Rrp44-mediated degradation of tRNA^{Met} by TRAMP required the presence of ATP in the reaction (Figure S3) and was inhibited by mutation of highly conserved residues (Asp_{236,238} → Ala) in the active site of Trf4 (DADA mutation). This mutation was previously shown to abolish poly(A) polymerase activity (Vanacova et al., 2005).

These data indicate that the TRAMP complex is a required cofactor in the complete, Rrp44-mediated degradation of a highly structured, native tRNA that contains all modifications but one.

Rrp44 and TAP-Purified TRAMP Complex Specifically and Independently Recognize Hypomodified tRNA_i^{Met} in Total tRNA

We next addressed whether tRNA degradation by recombinant Rrp44 and TRAMP is specific for the misfolded, hypomodified tRNA_i^{Met}. Wild-type and mutant Rrp44 proteins and/or TAP-purified TRAMP complex were incubated with total transfer RNAs isolated from either wild-type or *trm6-504* mutant yeast strains. Northern hybridization was used to analyze individual tRNAs (Figure 3A).





Figure 3. Rrp44 and TRAMP Independently Recognize Hypomodified tRNA_i^{Met} in Total tRNA

(A) Combined exonuclease/polyadenylation assay using 0.5 μ l TAP-purified [TRAMP complex](#), 100 fmol GST-Rrp44 [fusion proteins](#) (wild-type, 44-20, or cat), and total tRNAs purified from [wild-type](#) (WT) or *trm6-504* [yeast](#) strains. In order to achieve equal tRNA_i^{Met} levels in these tRNA preparations, the reaction was performed with 120 ng wild-type tRNAs and 40 ng *trm6-504* tRNAs. After a 10 min incubation at 30°C, the reaction products were separated on a 12% polyacrylamide/8 M [urea](#) gel and analyzed by northern [hybridization](#) using oligo probes specific for tRNA_i^{Met} or other tRNAs either containing (+) or lacking (-) the m¹A₅₈ modification.

(B) Western analysis of TAP-purified exosome from *GAL::rrp44* strains transformed with a [plasmid](#) expressing szz-tagged wild-type Rrp44 or the [mutant](#) Rrp44-cat protein. Strains were grown in glucose medium at 30°C. Exosomes were purified on IgG Sepharose, eluted by TEV cleavage, and analyzed by western blotting with anti-Rrp4 and anti-Rrp6 antibodies.

(C) Combined exonuclease/polyadenylation assay using 2.5 μ l TAP-purified yeast exosome, 0.5 μ l TRAMP complex, and 2.5 fmol 5' end-labeled, native, hypomodified tRNA_i^{Met} substrate in the presence of 1 mM ATP. Reactions were incubated for 90 min at 30°C at 0.75 mM MgCl₂ and analyzed as described in [Figure 2C](#).

Wild-type Rrp44 on its own degraded the hypomodified tRNA_i^{Met} from the *trm6-504* strain to an ~72 nt product but showed much lower activity with the fully modified tRNA from the wild-type strain ([Figure 3A](#), lanes 11 and 3) and no detectable degradation of other tRNAs. These included three [species](#) that also undergo Trm6-mediated m¹A modification (indicated with \oplus in [Figure 3A](#)). TAP-purified TRAMP independently recognized and polyadenylated tRNA_i^{Met} from the *trm6-504* strain ([Figure 3A](#), lane 10), but not the fully modified tRNA_i^{Met} ([Figure 3A](#), lane 2) or other tRNAs tested. Addition of the TRAMP complex strongly enhanced the degradation activity of Rrp44 ([Figure 3A](#), lanes 11 and 12) but did not stimulate Rrp44-20 or Rrp44-cat ([Figure 3A](#), lanes 13–16).

These data demonstrate that both Rrp44 and TRAMP specifically and independently recognize the misfolded hypomodified tRNA_i^{Met} in vitro, even in the presence of a large excess of other tRNA species. However, TRAMP plus Rrp44 was not sufficient to efficiently degrade the hypomodified tRNA_i^{Met} to small products in the presence of other tRNAs. This result is in contrast to the experiment shown in [Figure 2A](#), where a single, 5' end-labeled tRNA substrate was used. This suggests that the intact nuclear [exosome complex](#) might be needed for efficient and complete degradation of pre-tRNA_i^{Met} in vivo.

The Exosome Purified from a Strain Expressing Only Catalytically Inactive Rrp44 Does Not Degrade Hypomodified tRNA_i^{Met}

To determine whether Rrp44 is the only exosome component capable of degrading hypomodified tRNA_i^{Met}, the exosome was purified from the *GAL::rrp44* strain grown on glucose medium and complemented by [plasmids](#) expressing either Rrp44-szz or Rrp44-cat-szz. Complexes were purified on IgG Sepharose and eluted by [TEV protease](#) cleavage. Recovery of intact exosome complexes was confirmed by western blotting of the core component Rrp4 and the nuclear-specific component Rrp6 ([Figure 3B](#)) and by Coomassie staining (data not shown). The yield of Rrp4 was very similar for wild-type and mutant Rrp44, but ~2-fold more Rrp6 was recovered with tagged Rrp44-cat.

5' end-labeled hypomodified tRNA_i^{Met} was incubated with equivalent amounts of exosome copurified with Rrp44-szz or Rrp44-cat-szz, in the absence or presence of TAP-purified TRAMP. The degradation pattern generated by the wild-type exosome closely resembled that of recombinant Rrp44. The exosome on its own partially degraded the hypomodified tRNA_i^{Met} ([Figure 3C](#), lane 3), while the addition of the TRAMP complex

stimulated the exonuclease activity and led to a complete degradation of the substrate ([Figure 3C](#), lane 4). In contrast, the exosome copurified with Rrp44-cat-szz showed no exonuclease activity, while addition of the TRAMP complex led to [polyadenylation](#) of the RNA.

From these results we conclude that Rrp44 is the only exosome subunit that can recognize and, in the presence of the TRAMP complex, completely degrade the hypomodified tRNA_i^{Met}.

Discussion

All ten core components of the [yeast](#) exosome are essential for viability, and conditional depletion of any of these proteins leads to very similar [phenotypes](#) on [RNA metabolism](#). This has made it difficult to determine whether individual substrates are directed to, or targeted by, specific components of the complex. Here, we report that yeast Rrp44 [plays](#) a direct [role](#) in the specific [recognition](#) and [degradation](#) of defective tRNA_i^{Met}.

How Is the Hypomodified tRNA_i^{Met} Distinguished from the Fully Modified RNA?

The initiator tRNA^{Met} shows structural details that differ from other tRNAs, and [methylation](#) of [position](#) A₅₈ plays a uniquely important role in maintaining the correct [tertiary structure](#) ([Basavappa and Sigler, 1991](#)). The hypomethylated tRNA_i^{Met} can be specifically and independently recognized by both the [TRAMP complex](#) and Rrp44, even in the presence of a large excess of other tRNA [species](#) (see also [Vanacova et al. \[2005\]](#)). It seems unlikely that the specific change in base chemistry induced by the loss of adenosine methylation is the determinant for recognition. We envisage that Rrp44 and the TRAMP complex, and probably other tRNA surveillance systems ([Alexandrov et al., 2006](#)), are unable to bind correctly folded and modified tRNAs but can bind tRNAs that are misfolded. This would allow recognition of any defects in tRNA [maturation](#) that lead to altered structures. Conceptually, this might resemble the ability of [protein chaperones](#) to recognize and bind any protein that shows generic features of “misfolding” regardless of the specific [protein sequence](#).

Relating Rrp44 Structure and Function

A single point [mutation](#) in the predicted catalytic site of Rrp44 in RNB motif I abolished [nuclease](#) activity without affecting [RNA binding](#). This implies that Rrp44, like *E. coli* RNase II ([Amblar and Arraiano, 2005](#)), contains functionally separable domains that are primarily responsible for RNA binding or [catalysis](#).

Two distinct regions of RNase II, the “anchor” and “catalytic” regions, establish contacts with the RNA and act synergistically to provide processive degradation (see [Figure 4A](#)) ([Frazao et al., 2006](#)). The catalytic site lies deep within a cavity and is accessible only to single-stranded RNA, which is held in place by a “clamp” formed by two conserved aromatic residues (Tyr₂₅₃ and Phe₃₅₈). When the RNA is shortened to less than five [nucleotides](#), binding to the clamp is lost and an [oligonucleotide](#) (4 nt) product is released.

The activity of RNase II is blocked by [secondary structure](#) ([Amblar et al., 2006](#)), whereas RNase R can degrade double-stranded RNAs with single-stranded 3' overhangs of at least 7 nt ([Vincent and Deutscher, 2006](#)). Rrp44 was able to remove the unstructured 4 nt of the 3' CCA tail from the hypomodified tRNA_i^{Met}, whereas further degradation also required the TRAMP complex. We speculate that recognition of tRNA_i^{Met} by Rrp44 is permitted by the predicted wider RNA clamp, which might enable it to recognize and bind specific double-stranded RNAs with a short 3' overhang. Nuclease activity on such a substrate would, however, leave the double-stranded region of the RNA in the catalytic cavity, which is likely to prevent unwinding of the substrate. Addition of the TRAMP complex allowed complete degradation of the tRNA_i^{Met} by Rrp44, and this stimulation required ATP and the poly(A) [polymerase](#) activity of Trf4 (see [Figure S3](#)). We propose that following the specific recognition and removal of the 3' overhang by Rrp44, Trf4 adds a poly(A) tail to the substrate. The bound RNA might then “back-up,” allowing rebinding of the substrate with the longer single-stranded tail along the entire RNA-binding groove. Degradation would then resume with the double-stranded [RNA structure](#) being opened by Mtr4, possibly acting in conjunction with the S1 and CSD [RNA-binding domains](#) of Rrp44, leading to processive degradation of the remainder of the substrate. A mutation in the [S1 domain](#) (G916E) also reduced but did not completely abolish RNA binding and greatly inhibited activity on all substrates tested.

Analysis of *rrp44-20* revealed that degradation of the hypomodified tRNA_i^{Met} was inhibited by a single [amino acid substitution](#) (Gly₈₃₃ → Asp). In *E. coli* RNase II the counterpart of Rrp44 Gly₈₃₃ (Gly₄₈₆) is located in the “neck” region of the substrate-binding site of the [catalytic domain](#) ([Figures 4A and 4C](#)), which contains many amino acids conserved between RNase R and Rrp44. The neck forms specific [hydrogen bonds](#) (Hb) and van der Waals interactions (vdW) with the 3' end of the substrate (see [Figures 4A and 4C](#)) ([Frazao et al., 2006](#)). Introduction of a negatively charged side chain (Gly₄₈₆ → Asp) might reduce the affinity for RNA near the catalytic site ([Figure 4D](#)), while not affecting substrate binding by the anchor domain. Consistent with this, Rrp44-20 showed reduced binding to the hypomodified tRNA_i^{Met}, as well as poly(A₃₀) and a 37 nt transcript from the pBS polylinker. However, loss of degradation was specific for tRNA_i^{Met}. The less-structured RNA substrates may more readily enter the catalytic cavity of Rrp44, making specific interactions between the RNA and the neck of the catalytic region dispensable for catalysis. These observations imply that specific recognition and/or correct positioning of the 3' end of the hypomodified tRNA_i^{Met} takes place in the catalytic domain of Rrp44.

In understanding RNA turnover, a major limitation has been the lack of in vitro systems that faithfully recapitulate in vivo surveillance and degradation mechanisms. The results reported here should facilitate more detailed analyses of RNA surveillance.

Experimental Procedures

Expression and Purification of Recombinant Proteins

[Point mutations](#) in the expression construct for GST-Rrp44 (GST-Dis3sc) ([Noguchi et al., 1996](#)) were created using the QuikChange kit (Stratagene) and oligos Rrp44-D551N-F/R (Rrp44-cat), Rrp44-G833D-F/R (Rrp44-20), or Rrp44-G916E-F/R (S1) (see [Table S1](#)). *E. coli* strain BL21(DE3)pLysS (Stratagene) transformed with [plasmids](#) encoding GST-Rrp44 or GST alone (pGEX-4T1) was grown at 23°C to an OD₆₀₀ of 0.4, treated with IPTG at 1 mM, and incubated at 23°C to an OD₆₀₀ of 1. [Cell lysis](#) and [purification](#) were performed in buffer NB (20 mM Tris/HCl [pH 7.6], 150 mM NaCl, 5 mM MgCl₂, 8.7% [glycerol](#), and 0.1% Nonidet P40) supplemented with complete EDTA-free [protease inhibitor](#) (Roche). GST [fusion proteins](#) were affinity purified on glutathione-Sepharose beads and eluted with 20 mM [glutathione](#), which was subsequently removed by buffer exchange on NAP columns (GE Healthcare).

Exosome and TRAMP Purification

One-step purifications of exosome (at 150 mM NaCl) and [TRAMP complex](#) (at 120 mM NaCl) on IgG Sepharose columns followed by TEV [elution](#) were performed as described ([LaCava et al., 2005](#)). For exosome isolation, *GAL::rrp44* strains transformed with plasmids encoding szz-tagged Rrp44 were grown in SD –His/–Ura medium

to an OD₆₀₀ of 1.3 at 30°C. For the isolation of TRAMP complexes, an *rrp6Δ* strain transformed with a plasmid expressing zz-tagged Trf4 ([Vanacova et al., 2005](#)) was grown in SD –Leu medium to an OD₆₀₀ of 1 at 25°C. Each preparation used 2 liters of culture, producing 200–300 μl of the final TEV fraction. Enzyme concentrations were normalized by immunoblotting using anti-Rrp4 or anti-Rrp6 ([Mitchell et al., 2003](#)) antibodies.

In Vitro Assays

[Exonuclease](#) and [RNA-binding](#) assays were performed in 10 mM Tris/HCl (pH 7.6), 75 mM NaCl, 2 mM DTT, 100 μg/ml BSA, 0.8 U/μl RNasin, 4.5% glycerol, 0.05% Nonidet P40, and 0.25–0.75 mM MgCl₂. In combined exonuclease/polyadenylation assays, ATP was added to 1 mM. 5' end-labeled substrates were poly(A₃₀) and poly(U₃₀) (Dharmacon), in vitro-transcribed pre-tRNA^{Ser}, a 37 nt RNA derived from the pBS polylinker ([Mitchell et al., 1997](#)), and purified, native hypomodified tRNA_i^{Met}. In vitro [transcription](#), 5' end labeling, and gel purification of the labeled RNAs were performed as described ([LaCava et al., 2005](#)). Total tRNA was isolated from [wild-type](#) and *trm6-504* strains, and hypomodified tRNA_i^{Met} was purified from strain yJA146 (*gcd10ΔhclMT4*) as described ([Anderson et al., 2000](#)).

Acknowledgments

We thank W. Keller and J. Houseley for yeast strains and reagents and J. Houseley, S. Granneman, K. Kotovic, N. Watkins, and M. Kos for critical reading of the manuscript. This work was supported by the Wellcome Trust and a long-term HFSP fellowship to C.S.

Supplemental Data

The Exosome Subunit Rrp44 Plays a Direct Role in RNA Substrate Recognition

Table S1. Oligonucleotides Used and Strain Construction

Rrp44-D551N-F	5'-CCTCCAGGAT GTGTTGATAT TAACGATGCC CTACATGC
Rrp44-D551N-R	5'-GCATGTAGGG CATCGTTAAT ATCAACACAT CCTGGAGG
Rrp44-G833D-F	5'-CCTGACTTTA GACACTATGA TTTAGCCGTT GATATCTACA CACATTTCC
Rrp44-G833D-F	5'-GAAATGTGTG TAGATATCAA CGGCTAAATC ATAGTGTCTA AAGTCAGG
Rrp44-G916E-F	5'-GGAATAACGA GTCCACAGAA ACTGAATATG TTATTAAGGT ATTTAATAAT GG
Rrp44-G916E-R	5'-CCATTATTAA ATACCTTAAT AACATATTCA GTTTCTGTGG ACTCGTTATT CC
Rrp44-TAP-A	5'-GGATCCAATT ACTAGCAAGC GTAAGGCAGA ATTATTGTTA AAATCCATGG AAAAGAGAAG
Rrp44-TAP-B	5'-CCTTTTCGTT TTTATATCCT GATACTGAAG CATCTCCAT TACGACTCAC TATAGGG
Rrp44-genF	5'-ACGTACTION GGAATATATC CTTTTGAACT GGAGTG
Rrp44-gen(URA)R	5'-ACGTACTION GGCTGGATGG GAAGCGTACC
Anti-tRNA _i ^{Met}	5'-CGATCCGAGG ACATCAGGG
Anti-tRNA ^{Trp} (CCA)	5'-AACCTGCAAC CCTTCGA

Anti-tRNA ^{Tyr} (GΨA)	5'-AAGATTTTCGT AGTGATAA
Anti-tRNA ^{Pro} (UGG)	5'-ACCCAGGGCC TCTCG
Anti-tRNA ^{Leu} (CAA)	5'-GCATCTTACG ATACCTG
Anti-tRNA ^{Gly} (GCC)	5'-TACCACTAAA CCACTTGC

Oligonucleotides used in these analyses.

Supplemental Experimental Procedures

Strain Construction

S. cerevisiae strains used in this study were derived from the BY4741 wild-type strain (*MATa; his3Δ1; leu2Δ0; lys2Δ0; ura3Δ0*). To construct yeast expression plasmids for Rrp44, a C-terminal tag containing a streptavidin-binding peptide (Strep-tag II) (Junttila et al., 2005), TEV cleavage site and two copies of the z-domain of protein A was first added to the wild-type *RRP44* gene in BY4741. This was achieved by genomic integration of a cassette (the template plasmid (*URA3*) was derived from pBS1539 and constructed by Martin Kos, Tollervey lab) using a one-step PCR strategy and the oligonucleotides Rrp44-TAP A and B (Rigaut et al., 1999). The *RRP44-szz* gene including 500 bp of the promoter region was then PCR amplified from the genomic DNA of the tagged strain using the oligos Rrp44-genF and Rrp44-gen(URA)R and cloned into the XhoI restriction site of pRS316. The Rrp44-cat point mutation was created by site-directed mutagenesis as described above. A strain conditionally expressing HA-tagged Rrp44 from the *GAL10* promoter (*GAL::rrp44: MATa; his3Δ1; leu2Δ0; lys2Δ0; ura3Δ0, HisMX6-pGAL-3HA::rrp44* (Houseley and Tollervey, 2006)) was then transformed with an empty cloning vector (pRS316) or with constructs containing either the wild-type or the *rrp44-cat* gene (Gietz et al., 1992). Growth and handling of *S. cerevisiae* were by standard techniques. Strains were grown at 30°C in -His/-Ura dropout medium containing 0.67% nitrogen base (Difco) and either 2% glucose or 2% galactose.

The isolation of the *mtr17-1* allele was previously reported (Kadowaki et al., 1994). Subsequent analysis showed *mtr17-1* to be an allele of *RRP44* (A. Tartakoff, pers. comm.) and it was therefore designated *rrp44-1*. The strain carrying *rrp44-1* was generously provided by Alan Tartakoff (Case Western Reserve University). The *RRP44* ORF was recovered by PCR amplification and the sequence revealed two amino acid exchanges (P4S and G916E), with G916E mapping to the S1 domain of Rrp44 (see Fig. 1A). Rrp44 carrying the G916E mutation (referred to as the “S1” mutation in the text) was expressed in *S.cerevisiae* under control of the endogenous promoter. This conferred a temperature sensitive lethal phenotype following genetic depletion of the endogenous wild-type Rrp44, suggesting that the G916E mutation is responsible for the growth defect observed in *rrp44-1* strains. The GST-fusion protein containing the G916E (S1) point mutation was generated and used for *in vitro* analyses.

Supplemental References

Gietz, D., St Jean, A., Woods, R. A., and Schiestl, R. H. (1992). Improved method for high efficiency transformation of intact yeast cells. *Nucleic Acids Res* 20, 1425.

Houseley, J., and Tollervey, D. (2006). Yeast Trf5p is a nuclear poly(A) polymerase. *EMBO Rep* 7, 205-211.

Junttila, M. R., Saarinen, S., Schmidt, T., Kast, J., and Westermarck, J. (2005). Single-step Strep-tag purification for the isolation and identification of protein complexes from mammalian cells. *Proteomics* 5, 1199-1203.

Kadowaki, T., Chen, S., Hitomi, M., Jacobs, E., Kumagai, C., Liang, S., Schneiter, R., Singleton, D., Wisniewska, J., and Tartakoff, A. M. (1994). Isolation and characterization of *Saccharomyces cerevisiae* mRNA transport-defective (*mtr*) mutants. *J Cell Biol* 126, 649-659.

Rigaut, G., Shevchenko, A., Rutz, B., Wilm, M., Mann, M., and Seraphin, B. (1999). A generic protein purification method for protein complex characterization and proteome exploration. *Nat Biotechnol* 17, 1030-1032.

Schneider *et al.*, Figure S1




Figure S1. Expression of Recombinant Wild-Type and Mutant GST-Rrp44 Fusion Proteins

GST-fusion proteins were affinity-purified on glutathione-sepharose beads, separated on a denaturing 10% SDS polyacrylamide gel and stained with Coomassie Blue. The amount of wild-type and mutant Rrp44 proteins used in bandshift and exonuclease experiments were normalized to the intensity of the full-length band (marked with an arrow).




Figure S2. Activities of Mutant Forms of Rrp44 on Other RNA Substrates

(A) Exonuclease activity of wild-type GST-Rrp44, GST-Rrp44-20 or GST-Rrp44-S1 on a 5'-end labeled poly(A₃₀) RNA. Recombinant protein (3, 6 or 12 fmoles) was incubated with 10 fmoles RNA for 90 min at 30°C in the presence of 0.25 mM MgCl₂. Reaction products were analyzed as described in Fig. 2A.

(B) RNA binding assay using wild-type or mutant GST-Rrp44 proteins and 5'-end labeled poly(A₃₀) RNA. Recombinant protein (3, 6 or 12 fmoles) was incubated with 10 fmoles RNA for 10 min at 30°C in the presence of 0.25 mM MgCl₂. Reactions were analyzed as described in Fig. 2B.

(C) Exonuclease activity of wild-type GST-Rrp44, GST-Rrp44-20 or GST-Rrp44-S1 on a 5'-end labeled 37 nt substrate derived from the pBS polylinker. Recombinant protein (3, 6 or 12 fmoles) was incubated with 10 fmoles RNA for 90 min at 30°C in the presence of 0.25 mM MgCl₂. Reaction products were analyzed as described in Fig. 2A.

(D) RNA binding assays using wild-type or mutant GST-Rrp44 proteins and 5'-end labeled 37 nt RNA. Recombinant protein (3, 6 or 12 fmoles) was incubated with 10 fmoles RNA for 10 min at 30°C in the presence of 0.25 mM MgCl₂. Reactions were analyzed as described in Fig. 2B.

Schneider *et al.*, Figure S3




Figure S3. Stimulation of Rrp44-Mediated Degradation of the Hypomodified tRNA^{Met} Substrate by the TRAMP Polyadenylation Complex Requires ATP and Catalytically Active Trf4

Combined exonuclease/polyadenylation assay using 0.5 μ l TAP-purified TRAMP complex (wildtype (WT) or polyadenylation inactive DADA mutant, see Vanacova *et al.*, 2005), 100 fmoles GST-Rrp44 WT protein and 25 fmoles 5'-end labeled, native hypomodified tRNA^{Met} substrate. Reactions were performed for 60 min at 30°C with 0.625 mM MgCl₂ and in the absence or presence of 1 mM ATP.

Products were separated on a 12% polyacrylamide/ 8M urea gel and visualized by autoradiography. Full-length tRNA^{Met}, polyadenylated RNA and degraded RNA species are indicated on the right.

References

- [Alexandrov *et al.*, 2006](#) A. Alexandrov, I. Chernyakov, W. Gu, S.L. Hiley, T.R. Hughes, E.J. Grayhack, E.M. Phizicky **Rapid tRNA decay can result from lack of nonessential modifications** *Mol. Cell*, 21 (2006), pp. 87-96
- [Amblar and Arraiano, 2005](#) M. Amblar, C.M. Arraiano **A single mutation in Escherichia coli ribonuclease II inactivates the enzyme without affecting RNA binding** *FEBS J.*, 272 (2005), pp. 363-374
- [Amblar *et al.*, 2006](#) M. Amblar, A. Barbas, A.M. Fialho, C.M. Arraiano **Characterization of the functional domains of Escherichia coli RNase II** *J. Mol. Biol.*, 360 (2006), pp. 921-933
- [Anderson *et al.*, 2000](#) J. Anderson, L. Phan, A.G. Hinnebusch **The Gcd10p/Gcd14p complex is the essential two-subunit tRNA(1-methyladenosine) methyltransferase of Saccharomyces cerevisiae** *Proc. Natl. Acad. Sci. USA*, 97 (2000), pp. 5173-5178
- [Basavappa and Sigler, 1991](#) R. Basavappa, P.B. Sigler **The 3 Å crystal structure of yeast initiator tRNA: functional implications in initiator/elongator discrimination** *EMBO J.*, 10 (1991), pp. 3105-3111
- [Dziembowski *et al.*, 2007](#) A. Dziembowski, E. Lorentzen, E. Conti, B. Seraphin **A single subunit, Dis3, is essentially responsible for yeast exosome core activity** *Nat. Struct. Mol. Biol.*, 14 (2007), pp. 15-22

- [Frazao et al., 2006](#) C. Frazao, C.E. McVey, M. Amblar, A. Barbas, C. Vonrhein, C.M. Arraiano, M.A. Carrondo **Unravelling the dynamics of RNA degradation by ribonuclease II and its RNA-bound complex** *Nature*, 443 (2006), pp. 110-114
- [Houseley et al., 2006](#) J. Houseley, J. LaCava, D. Tollervey **RNA-quality control by the exosome** *Nat. Rev. Mol. Cell Biol.*, 7 (2006), pp. 529-539
- [Kadaba et al., 2004](#) S. Kadaba, A. Krueger, T. Trice, A.M. Krecic, A.G. Hinnebusch, J. Anderson **Nuclear surveillance and degradation of hypomodified initiator tRNAMet in S. cerevisiae** *Genes Dev.*, 18 (2004), pp. 1227-1240
- [Kadaba et al., 2006](#) S. Kadaba, X. Wang, J.T. Anderson **Nuclear RNA surveillance in Saccharomyces cerevisiae: Trf4p-dependent polyadenylation of nascent hypomethylated tRNA and an aberrant form of 5S rRNA** *RNA*, 12 (2006), pp. 508-521
- [LaCava et al., 2005](#) J. LaCava, J. Houseley, C. Saveanu, E. Petfalski, E. Thompson, A. Jacquier, D. Tollervey **RNA degradation by the exosome is promoted by a nuclear polyadenylation complex** *Cell*, 121 (2005), pp. 713-724
- [Liu et al., 2006](#) Q. Liu, J.C. Greimann, C.D. Lima **Reconstitution, activities, and structure of the eukaryotic RNA exosome** *Cell*, 127 (2006), pp. 1223-1237
- [Mitchell et al., 1997](#) P. Mitchell, E. Petfalski, A. Shevchenko, M. Mann, D. Tollervey **The exosome: a conserved eukaryotic RNA processing complex containing multiple 3'→5' exoribonucleases** *Cell*, 91 (1997), pp. 457-466
- [Mitchell et al., 2003](#) P. Mitchell, E. Petfalski, R. Houalla, A. Podtelejnikov, M. Mann, D. Tollervey **Rrp47p is an exosome-associated protein required for the 3' processing of stable RNAs** *Mol. Cell. Biol.*, 23 (2003), pp. 6982-6992
- [Noguchi et al., 1996](#) E. Noguchi, N. Hayashi, Y. Azuma, T. Seki, M. Nakamura, N. Nakashima, M. Yanagida, X. He, U. Mueller, S. Sazer, T. Nishimoto **Dis3, implicated in mitotic control, binds directly to Ran and enhances the GEF activity of RCC1** *EMBO J.*, 15 (1996), pp. 5595-5605
- [Romani and Scarpa, 1992](#) A. Romani, A. Scarpa **Regulation of cell magnesium** *Arch. Biochem. Biophys.*, 298 (1992), pp. 1-12
- [Vanacova et al., 2005](#) S. Vanacova, J. Wolf, G. Martin, D. Blank, S. Dettwiler, A. Friedlein, H. Langen, G. Keith, W. Keller **A new yeast poly(A) polymerase complex involved in RNA quality control** *PLoS Biol.*, 3 (2005), p. e189
- [Vincent and Deutscher, 2006](#) H.A. Vincent, M.P. Deutscher **Substrate recognition and catalysis by the exoribonuclease RNase R** *J. Biol. Chem.*, 281 (2006), pp. 29769-29775
- [Zuo et al., 2006](#) Y. Zuo, H.A. Vincent, J. Zhang, Y. Wang, M.P. Deutscher, A. Malhotra **Structural basis for processivity and single-strand specificity of RNase II** *Mol. Cell*, 24 (2006), pp. 149-156