University of Massachusetts Medical School

eScholarship@UMMS

Orthopedics and Physical Rehabilitation Publications and Presentations

Orthopedics and Physical Rehabilitation

2018-02-15

A Common Pain in Pregnancy with an Uncommon Cause

Jennifer Baima University of Massachusetts Medical School

Et al.

Let us know how access to this document benefits you.

Follow this and additional works at: https://escholarship.umassmed.edu/ortho_pp

Part of the Female Urogenital Diseases and Pregnancy Complications Commons, Musculoskeletal Diseases Commons, Obstetrics and Gynecology Commons, Orthopedics Commons, Rehabilitation and Therapy Commons, and the Women's Health Commons

Repository Citation

Baima J, Vinod AV, Most MJ. (2018). A Common Pain in Pregnancy with an Uncommon Cause. Orthopedics and Physical Rehabilitation Publications and Presentations. Retrieved from https://escholarship.umassmed.edu/ortho_pp/185

This material is brought to you by eScholarship@UMMS. It has been accepted for inclusion in Orthopedics and Physical Rehabilitation Publications and Presentations by an authorized administrator of eScholarship@UMMS. For more information, please contact Lisa.Palmer@umassmed.edu.



A common pain in pregnancy with an uncommon cause

CASE DESCRIPTION

Case: A 41-year-old woman presented with 9-months of persistent atraumatic left hip pain localized to the groin that had started in her second trimester of pregnancy.

Background: Five months after delivery at presentation to clinic, her symptoms had progressively worsened to require a cane for ambulation.


Exam: On exam, hip range of motion was limited: flexion to 90°, external rotation to 20°, and internal rotation to 10°. She had groin pain with internal and external rotation. She was neurovascularly intact without lymphadenopathy.

Imaging: Radiographs showed significant erosion of the left femoral head and acetabulum with mild protrusio. "Apple core" erosions were visualized at the femoral neck. MRI showed extensive bone edema on both sides of the joint, with erosion of the femoral head and acetabulum. There was joint effusion, synovial hypertrophy, and excessive synovial tissue.

Differential diagnoses: included PVNS, avascular necrosis, rapidly progressive osteoarthritis, inflammatory arthritis, septic arthritis, insufficiency fractures, synovial chondromatosis, and transient osteoporosis of the hip.

Diagnosis: A needle biopsy confirmed the diagnosis of PVNS.

Treatment: She was treated with synovectomy and total hip arthroplasty, and has remained pain free and without evidence of disease for the last 5 years.





2.

Baima Jennifer, Vinod Amrit, Most Mathew

Department of Orthopedics and Physical Rehabilitation, University of Massachusetts Medical School

Figure 1: T2 coronal MRI demonstrates "apple core" erosions seen in synovial chondromatosis, amyloidosis, rheumatoid arthritis, and **PVNS**




Figure 2: Histologic analysis confirmed the diagnosis of PVNS

REFERENCES

1. Uslu M, Cetik O, Atasoy P, Eksioglu F, Engin M. Localized pigmented villonodular synovitis of the knee : acute onset in pregnancy. 2006:1054-1056. doi:10.1007/s00296-006-0135-2. Ottaviani S, Ayral X, Dougados M. Pigmented Villonodular Synovitis : A Retrospective Single-Center Study of 122 Cases and Review of the Literature. Semin Arthritis Rheum. 2011;40(6):539-5doi:10.1016/j.semarthrit.2010.07.005.

Edwards MR, Tibrewal S. Patello-femoral joint pain due to unusual location of localised pigmented villonodular synovitis — a case report. Knee. 2004;11(4):327-329. doi:10.1016/S0968-0160.








Figure 3: After treatment with arthroplasty

DISCUSSION

PVNS is a disorder characterized by synovial proliferation.

There are only two previous case reports of patients who were diagnosed with PVNS during pregnancy, both of whom had monoarticular involvement of the knee.

CONCLUSIONS

- Synovectomy is the mainstay of surgical treatment of PVNS.
- Although, in patients with extensive articular involvement, synovectomy and arthroplasty may be required.
- The current understanding is that cytokines have a trophic influence leading to growth of the tumor.
- Further studies are needed to establish a definitive connection between PVNS and pregnancy.