

THE UNIVERSITY OF KANSAS  
PALEONTOLOGICAL CONTRIBUTIONS

June 1998

Number 9

PHYLOGENY OF ARENIG TO CARADOC CRINOIDS (PHYLUM  
ECHINODERMATA) AND SUPRAGENERIC CLASSIFICATION OF THE  
CRINOIDEA

William I. Ausich

Department of Geological Sciences, 155 South Oval Mall, The Ohio State University, Columbus, OH 43210,  
ausich@mps.ohio-state.edu

*Abstract.*—Stepwise, parsimony-based character analysis of Arenig to Caradoc crinoids (Ordovician) indicates that Early and middle Ordovician crinoids represent a radiation largely confined to the Ordovician. Only a few middle Paleozoic families are considered to have roots among these Early and middle Ordovician genera. Also, most genera are reinterpreted as a part of larger Ordovician clades rather than being isolated in monogeneric families.

Six subclasses recognized as a result of phylogenetic analysis are the Aethocrinea, Cladida, Flexibilia, Articulata, Camerata, and Disparida. Many aspects of the *Treatise* classification (Moore and Teichert, 1978) are followed, but significant differences exist. The zygodiplobathrid-eudiplobathrid camerate suborder distinction is not made, and disparid superfamilies are no longer recognized. The subclass Disparida is subdivided into orders rather than superfamilies. Seven Early and middle Ordovician disparid orders are recognized, and additional younger orders undoubtedly exist. Whereas disparid superfamilies in the *Treatise* were defined largely on aboral cup symmetry, disparid orders are defined here largely on the basis of presence and absence of radial plates and aboral cup architecture. The Calceocrinida, Homocrinida, Myelodactylida, Eustenocrinida, and Tetragonocrinida are new orders, having previously been regarded as superfamilies and subfamilies; the Maennilicrinida is a new order; and the order Hybocrinida is maintained in the subclass Disparida. Subfamilies are no longer recognized within the Homocrinidae or Cincinnaticrinidae.

Other changes include new families (Agostocrinidae, Columbicrinidae, and Maennilicrinidae), one family elevated from subfamily to family (Atopocrinidae), and reassignment of families to different higher categories (Acolocrinidae to the Eustenocrinida, Agostocrinidae and Colpodecrinidae to the Cladida, Cleiocrinidae to the Rhodocrinitacea, and Porocrinidae to the Cyathocrinitacea). Reassignment of 34 genera to different families is proposed (3 of these to a different subclass). A number of family-level taxa are recognized as junior synonyms and *Baerocrinus*, *Ontariocrinus*, Baerocrinidae, and Ontariocrinidae are regarded as *nomina dubia*.

INTRODUCTION

Ausich (1995, 1996a) proposed a new homology scheme for aboral-cup-plate circlets. Based on these homologies and a temporal stepwise approach, he used parsimony-based character analyses to develop a subclass-level, phylogenetic classification of the Crinoidea using Arenig and Llanvirn crinoids (Ausich, 1998a). Six subclasses were rec-

ognized: Aethocrinea, Cladida, Flexibilia, Articulata, Camerata, and Disparida. The phylogenetic trees generated by Ausich (1998a) confirm many aspects of the Moore and Teichert (1978) classification within subclasses but also indicate that several revisions are necessary, as suggested by Brower (1975), Kelly, Frest, and Strimple (1978), Kelly (1982, 1986), Donovan (1988a), Sevastopulo and Lane (1988), Simms and Sevastopulo (1993), and others.

Developed herein is a comprehensive phylogeny of Arenig to Caradoc crinoids by a continuation of this stepwise approach. This time interval includes the first substantial adaptive radiation of the Crinoidea (Eckert, 1988). During the Arenig and Llanvirn 18 and 16 genera and 19 and 23 species existed, respectively; but during the Caradoc, 76 crinoid genera were present with 259 species recognized. Immediately thereafter, crinoids were diminished in diversity during Ashgillian extinctions (Eckert, 1988; Donovan, 1988a, 1989a, 1994). Based on this phylogeny, a supra-generic classification of Arenig to Caradoc crinoids is presented herein.

The starting point of the current study is the topology of the crinoid phylogeny and subclasses of Ausich (1998a) (Fig. 1), which are based on the homologies of Ausich (1996a). These new homologies are used herein; however, the within-subclass analyses presented here are largely independent of the homology contrasts of Ausich (1996a), except perhaps for the disparids. Furthermore, this work is independent of alternative ideas on the origin of the Crinoidea (Ausich and Babcock, 1998; Ausich, 1998b), except perhaps for the Camerata.

As discussed by Ausich (1997, 1998a), the *Treatise* crinoid classification (Moore and Teichert, 1978) (see Table 1, p. 16) was largely based on work by Moore and Laudon (1943). The Moore and Laudon (1943) classification was largely derived from the competing classification schemes of Wachsmuth and Springer (1897) and Bather (1899, 1900), from a time when very little was known about Early and middle Ordovician crinoids. Therefore, underlying assumptions structuring much of the *Treatise* classification were based largely on Silurian to Mississippian crinoids. This yielded a classification in which the phylogenetic underpinning can be questioned, and, indeed, this is probably the reason for many of the proposed revisions to the *Treatise* classification (see discussions by Ausich, 1997, 1998a).

Due to new and competing ideas on aboral-cup-plate homologies, the origin of crinoids, and what is and what is not a crinoid, phylogenetic and classification analyses herein are built on many assumptions. Assumptions that most or all crinoid students support include elimination of the hemistreptocrinids and coronates from the Crinoidea (Arendt and Rozhnov, 1995; Brett *et al.*, 1983) and dissolution of the subclass Inadunata due to polyphyly (Simms and Sevastopulo, 1993, and others). Assumptions included herein that are probably not endorsed by all crinoid students are: 1. elimination of *Echmatocrinus* from the Crinoidea (Conway Morris, 1993; Ausich and Babcock, 1996, 1998; but see Sprinkle and Collins, 1995); 2. use of rhombiferans as the outgroup for the Crinoidea (Ausich, 1996b, 1998b; but see Guensburg and Sprinkle, 1997, and Sprinkle and Guensburg, 1997); 3. use of the cup-circlet-plate homology scheme of Ausich (1996a); and 4. four-circlet aboral-cup-plate condition as the primitive condition among the Crinoidea (Ausich 1996a, 1996b, 1998a).

## METHODS

The methods employed follow Ausich (1998a), who used a progressive, stepwise development of crinoid phylogeny. This method reduces the impact of convergent and iterative evolution among major crinoid clades. The topology of the initial steps of this approach (Arenig to Llanvirn crinoids from Ausich, 1998a) is accepted; and the phylogeny within each of those previously defined clades is examined herein for Arenig to Caradoc forms. Thus, cladids-flexibles, camerates, and disparids of the Arenig to Caradoc are examined independently with parsimony techniques. Furthermore, where necessary, lineage subsets of these clades were also examined. For example, analysis of all camerates yields an overall rather poorly constrained cladogram, but it clearly delineates monobathrid and diplobathrid camerates. Therefore, monobathrids and diplobathrids are analyzed independently further (see discussion of camerates below).

Parsimony-based character analyses are performed using PAUP 3.1.1 (Swofford, 1993). All crinoid stratigraphic information was correlated to the revised Ordovician stratigraphy of Fortey *et al.* (1995). No morphological evidence suggests that the recorded stratigraphic distributions could not be considered accurate at the stage level. No temporal distinctions are made within a stage, however.

For analyses presented herein, all characters are unordered and equally weighted, and all searches were run with 10 random-addition-sequence replicates, with one exception as noted. Search methods used are heuristic with random stepwise addition. Search results are presented as 50-percent majority-rule trees or as single trees. Simple consistency indices (CI), retention indices (RI), and rescaled consistency indices (RC) are given, as indicated in output from PAUP analyses. Specific details for each cladogram are presented below.

Cladograms list generic names, but characters for each genus were typically based on a single species in that genus. Most Tremadoc to Llanvirn genera are monospecific, but most of the Caradoc genera contain many species. The type species was used for character coding where appropriate or necessary. Where more than one species occurs in a genus, the oldest well-preserved species or the type species is used, unless it is poorly known (Appendix A, p. 33). Wherever possible, type specimens or casts of type specimens were examined, including material from Estonia and Russia.

An exhaustive list of morphological characters is not used because the distribution of species-level type characters is likely to add noise to an analysis seeking to uncover the underlying structure of a phylogeny. When a character set is less than comprehensive, however, biases may become a factor. Unfortunately, this is unavoidable. Character selection included those considered to represent basic architectural features of the arms, calyx, and column. Thus, 21 characters were used for cladids and the flexible; 27




Figure 1. Phylogenetic interpretation of Arenig and Llanvirn crinoids; *Caryocystites* is the outgroup (Ausich, 1998a).

characters were used for camerates; and 31 characters were used for disparids. Some characters were eliminated during preliminary analyses because they were uninformative or because character states could be determined for too few taxa due to limitations of preservation. Examples include the following: cladids—CD interray plating, anal-sac shape, and anal sac-plate sculpturing; camerates—visibility of infrabasals and characters of tegmen; disparids—relative size of C radial, aboral cup-plate sculpturing, and anal-sac shape.

Results of all character analyses and consideration of stratigraphic position, paleogeography, and crinoid mor-

phology were used to develop a final phylogenetic tree for each clade. Stratocladistic methodology (Fisher, 1988, 1994; Harvey and Ausich, 1997) is not used here because in the relatively short interval considered little stratigraphic debt can be reliably accumulated.

Results from character analyses of cladid and flexible crinoids are presented in Figures 2 and 3. In Figure 2, *Elpasocrinus* is the outgroup, and all Arenig to Caradoc cladids are used (22 genera, except *Esthonocrinus*, *Polycrinus*, and *Triboloporus*, which are poorly known). The analysis included 21 characters (Appendix B, p. 34) with a total of 67 character states. The 50-percent majority-rule tree is








Figure 3. Cladograms from character analysis of select dendrocrinid cladids. 1, Only dendrocrinid cladids considered (16 genera). *Elpasocrinus* is the outgroup. The analysis included 18 characters with a total of 54 character states. The 50-percent majority-rule tree is from 13 equally parsimonious trees; length 56; CI 0.714, RI 0.600, RC 0.429. 2, Only select dendrocrinids are considered (13 genera). *Elpasocrinus* is the outgroup. The analysis included 14 characters with a total of 47 character states. The 50-percent majority-rule tree is from 8 equally parsimonious trees; length 44; CI 0.750, RI 0.560, RC 0.420; see also Appendix B, p. 34 (new).

210, and the tree has a CI = 0.505, RI = 0.623, and RC = 0.315. The analysis is a single run and not from 10 replicates. In Figure 7.1 *Ibexocrinus* is the outgroup for analysis of homocrinid and cincinnaticrinid disparids, *sensu* Warn and Strimple, 1977 (15 genera, with the exception of *Difficilicrinus*, *Glaucocrinus*, and *Othneiocrinus*). The analysis included 17 characters (Appendix D, p. 36; A radial, C radial, D radial, E radial, radial-basal articulation, infrabasal number, lintel number, basal concavity, lintels visible, CD plating, interray plating, pore rhombs, arm number, arm habit, recumbent arms, and anal on D ray are constant) with a total of 49 character states. The 50-percent majority-rule tree is from 8 equally parsimonious trees of length 61, and the tree has a CI = 0.721, RI = 0.595, and RC = 0.429. In Figure 7.2, "*Pariocrinus*" (see Systematic Paleontology, p. 27) is the outgroup, and only iocrinid and hybocrinid disparids are considered (12 genera). The analysis included 22 characters (Appendix D, p. 36; A radial, B radial, C radial, D radial, E radial, radial-basal articulation, infrabasal number, basal concavity, interray plating, pore rhombs, and fixed brachials are constant) with a total of 66 character states. A single tree resulted with a length of 57, and the tree had a CI = 0.860, RI = 0.818, and RC = 0.703.

#### CLADID-FLEXIBLE PHYLOGENY

*Character analyses.*—Twenty-five genera of Arenig to Caradoc cladids and one flexible are recognized. Of these, three, *Elpasocrinus*, *Compagicrinus*, and *Archaetaxocrinus*, are known from Arenig strata; *Archaetaxocrinus*, *Palaeocrinus*, and a new genus (see Systematic Paleontology, p. 18) are from the Llanvirn, and the rest were first recorded from

the Caradoc. Three genera, *Esthonocrinus*, *Polycrinus*, and *Triboloporus*, are poorly known and are not included in any of the character analyses presented below. Details of the analyses are listed above and in the figure captions and appendices.

*Inferred phylogeny.*—In a comparison of *Elpasocrinus* and *Compagicrinus*, one sees that the posterior interray plating of *Elpasocrinus* is more similar to *Aethocrinus* (see Ausich, 1998a) and later cladids. Therefore, *Elpasocrinus* is taken as the base of the cladid line, having evolved from the aethocrinids through loss of the lintel circlet (Ausich, 1998a). Two cladid lineages arose during the Arenig: *Elpasocrinus* and *Compagicrinus* with the arms free above the radials and no fixed interradials or brachials; and *Archaetaxocrinus*, which had fixed brachials and interradials similar to *Aethocrinus*. In character analysis of all Arenig and Llanvirn crinoids (Ausich, 1998a), *Archaetaxocrinus* falls out as the most derived cladid, intermediate between cladids and camerates. This result is undoubtedly due to the fixed plating at the top of the calyx that is present in both *Archaetaxocrinus* and in camerates. The early age of this form, its retention of basic cladid features, and lack of camerate synapomorphies, however, indicate that it should be retained in the cladids as an early, separate lineage. Lewis (1981) argued that *Archaetaxocrinus* was the oldest flexible crinoid, but this interpretation is incompatible with the present analyses. *Archaetaxocrinus* has features such as fixed brachials and interradials that are similar to flexibles; however, it is considered here to be a cladid lineage distinct from flexibles that added plates in the distal calyx.




Figure 4. Cladogram from character analysis of all Arenig to Caradoc camerates. 1, All Arenig to Caradoc camerates are used (26 genera, except *Schizocrinus*). *Proexenocrinus* is the outgroup. The analysis included 27 characters with a total of 95 character states. The 50-percent majority-rule tree is from 1,021 equally parsimonious trees; length 134; CI 0.530, RI 0.537, RC 0.284. 2, *Proexenocrinus* is the outgroup for analysis of all monobathrid camerates. Six monobathrids are analyzed, with the exception of *Schizocrinus*. The analysis included 18 characters with a total of 44 character states. The 50-percent majority-rule tree is from 2 equally parsimonious trees; length 27; CI 0.963, RI 0.875, RC 0.843; see also Appendix C, p. 35 (new).

During the Llanvirn, the cyathocrines became established and persisted through the Caradoc. The three deeply rooted, Arenig to Llanvirn cladid lineages are therefore the dendrocrines, the cyathocrines, and the lineage begun by *Archaeataxocrinus*. Only the dendrocrines and cyathocrines persisted beyond the Caradoc. In addition, dendrocrines and cyathocrines diversified during the Caradoc, and the subclass Flexibilia evolved from the dendrocrines during the Caradoc. Separation between dendrocrines and the derived cyathocrines is evident in the character analysis that includes all cladids and the single flexible (Fig. 2). Cyathocrines became specialized with generally bowl- to globe-shaped aboral cups, narrower radial facets, narrower arms in relation to the aboral cup, and shorter anal sacs. Three early lineages of cyathocrines developed, one lineage with *Eoparisocrinus* and *Illemocrinus*, which are closely related. *Palaeocrinus*, *Porocrinus*, *Carabocrinus*, and *Triboloporus* constitute the second lineage of cyathocrines and include the conical to globose forms with a well-sutured tegmen, with pore structures, with anus on the tegmen or

the side of the cup, and with or without recumbent ambulacra. *Agostocrinus* is a third cyathocrine lineage.

From the Arenig to the Llanvirn, the fundamental dendrocrine lineage comprised *Elpasocrinus*, *Compagicrinus*, and *Brechmocrinus*, all with compound radianals. This basic lineage continued with the loss of the inferradial and diversified during the Caradoc to include *Grenprisia*, *Ottawacrinus*, *Plicodendrocrinus*, and *Esthonorcrinus*. These crinoids all have medium, cone-shaped, aboral cups; five infrabasals visible in lateral view; high infrabasals; a radianal that occupies the full proximal width of the C radial; angustary to plenary radial facets; mostly three plates in the posterior interray; rounded arms that are either isotomously branched, heterotomously branched, or isotomous proximally and heterotomous distally; and a circular or rarely pentagonal column. From these dendrocrinids a group of crinoids that added fixed plates to the aboral cup includes *Quinquecaudex*, *Cupulocrinus*, and *Protaxocrinus* (Fig. 3).

A third group of dendrocrines has low to medium, conical, aboral cups; low infrabasals; a radianal that occu-




Figure 5. Cladograms from character analysis of select diplobathrid camerates. 1, *Proexenocrinus* is the outgroup, with all diplobathrids considered (20 genera). The analysis included 25 characters with a total of 85 character states. The 50-percent majority-rule tree is from 393 equally parsimonious trees; length 108; CI 0.562, RI 0.484, RC 0.272. 2, *Proexenocrinus* is the outgroup, and all diplobathrids (except *Euptychocrinus* and *Rhaphanocrinus*) are considered (18 genera). The analysis included 24 characters with a total of 83 character states. The 50-percent majority-rule tree is from 3 equally parsimonious trees; length 103; CI 0.592, RI 0.506, RC 0.300; see also Appendix C, p. 35 (new).

pies the full width of the C radial; penepenary to plenary radial facets; one to three posterior interray plates in the cup; arms with a wide array of branching modes; and a holomeric, circular column that is either heteromorphic or xenomorphic. Moore, Lane, and Strimple (1978) placed these into numerous families: *Praecupulocrinus* into Cupulocrinidae; *Merocrinus* into Merocrinidae; *Polycrinus* into Mastigocrinidae; "*Dendrocrinus*" *acutidactylus* into Dendrocrinidae (see Systematic Paleontology, p. 21); *Eopinnocrinus* into Botryocrinidae; and *Metabolocrinus* into Metabolocrinidae. These all appear to represent one group of genera modified from the primitive dendrocrinid lineage, however (Fig. 2–3).

In the analysis of all cladids, phylogenetic relationships among *Archaeotaxocrinus*, *Colpodecrinus* (see Systematic Paleontology, p. 20), *Protaxocrinus*, and *Quinquecaudex*, all of which have fixed brachials and interradials, are unresolved. *Archaeotaxocrinus* and *Colpodecrinus* are regarded as a distinct early cladid branch (Fig. 8) that has fixed brachials and interradials probably as a derived characteristic. *Quinquecaudex* and *Protaxocrinus* evolved from *Cupulocrinus* as a convergent evolutionary trend to increase plates in the calyx. *Cupulocrinus humilis* (Billings) has some individuals with a few small fixed interradial plates. This interpretation of these cladistic results is most consistent with the hypothesis that *Protaxocrinus* is the oldest flexible, as sug-

gested by Springer (1911) and Lane (1978a), and it is recognized as the only Arenig to Caradoc member of the Flexibilia. The essential feature for this transition is the reduction to three infrabasal plates. *Quinquecaudex* is part of the same radiation, but because it retains five infrabasals, it is retained in the cladids.

Another lineage derived from the basic dendrocrinid design was the relatively minor radiation that yielded *Merocrinus*, *Praecupulocrinus*, *Polycrinus*, "*Dendrocrinus*" *acutidactylus*, *Eopinnocrinus*, and *Metabolocrinus* (Fig. 2–3, 8). These crinoids all appear to have arisen as a lineage with medium to low conical cups, very low infrabasals, one radial, plenary radial facets, one or more primibrachials, and simple to specialized arm branching.

The cyathocrinids are the most derived Arenig to Caradoc cladids. The oldest known cyathocrinid is *Palaeocrinus*, but *Eoparisocrinus* appears to represent a morphology more intermediate between dendrocrinids and cyathocrinids (Fig. 2). *Eoparisocrinus* and *Illemocrinus* represent a more primitive cyathocrinid lineage, whereas the more globular aboral cup design of *Palaeocrinus*, *Carabocrinus*, *Porocrinus*, and *Triboloporus*, with reduced arms, pore structures, and recumbent ambulacra, is more specialized in the cyathocrinid lineage. *Agostocrinus* appears to be a cyathocrinid crinoid that represents a morphological trend distinct from other Caradoc forms. *Agostocrinus* is




Figure 6. Cladograms from character analysis of all disparid crinoids. *Pogonipocrinus* is the outgroup, and all Arenig to Caradoc disparids are used (45 genera). The analysis included 31 characters with a total of 108 character states. The 50-percent majority-rule tree is from 2,027 equally parsimonious trees; length 210; CI 0.505, RI 0.623, RC 0.315; see also Appendix D, p. 36 (new).

considered a cyathocrine because of the cup shape, well-sutured tegmen plates, and apparent lack of an anal sac. Its recumbent arms and large radial in the basal cirlet make it a separate lineage.

*Discussion.*—Brower (1995a) analyzed the phylogeny of Ordovician and Early Silurian cladid crinoids on the basis of 42 characters. Methods included cluster analysis and

PAUP, and the plate homologies used were essentially the same as those of Moore and Teichert (1978). Cladids were divided into five basic groups (Brower, 1995a) (only Arenig to Caradoc crinoids are listed here): 1. stem cladids *Aethocrinus*, *Compagicrinus*, *Elpasocrinus*, *Grenprisia*, and *Ottawacrinus*; 2. *Carabocrinus*, *Palaeocrinus*, *Illemocrinus*, and *Porocrinus*; 3. *Archataxocrinus*, *Merocrinus*, *Praecupulocrinus*,




Figure 7. Cladograms from character analysis of select disparid crinoids. 1, *Ibexocrinus* is the outgroup for analysis of homocrinids, cincinnaticrinids (*sensu* Warn and Strimple, 1977), and disparids (15 genera, with the exception of *Difficilicrinus*, *Glaucoocrinus*, and *Othneiocrinus*). The analysis included 17 characters with a total of 49 character states. The 50-percent majority-rule tree is from 8 equally parsimonious trees; length 61; CI 0.721, RI 0.595, RC 0.429. 2, "*Pariocrinus*" is the outgroup, and only iocrinid and hybocrinid disparids are considered (12 genera). The analysis included 22 characters with a total of 66 character states. A single tree resulted with a length of 57; CI 0.860, RI 0.818, RC 0.703; see also Appendix D, p. 36 (new).

and *Polycrinus*; 4. *Cupulocrinus* and *Protaxocrinus*; and 5. *Dendrocrinus*, *Esthonocrinus*, *Eoparisocrinus*, *Eopinnacrinus*, *Plicodendrocrinus*, and *Quinquecaudex*.

The analyses herein and that of Brower (1995a) agree on many basic groupings of these cladids. They also agree on some potentially contentious issues: 1. *Archaetaxocrinus* is deeply rooted and not related to flexible crinoids; 2. *Ottawacrinus* is a primitive cladid closely related to *Grenprisia*; 3. *Merocrinus*, *Praecupulocrinus*, and *Polycrinus* are closely related; and 4. *Protaxocrinus* is closely related to *Cupulocrinus*. These two independent analyses disagree on several points, however, including the following: 1. *Eopinnacrinus* was considered to be a dendrocrinid by Brower (1995a) but is regarded herein as a metabolocrinid; 2. *Eoparisocrinus* was considered to be a dendrocrinid by Brower (1995a, 1995b), but it links with other cyathocrines herein; and 3. *Quinquecaudex* was considered to be a dendrocrinid by Brower (1995a) but is considered herein to be a cupulocrinid. The interpretations presented here are preferred because the choice of characters and homologies used are considered to be fundamental to crinoid designs and because the lineages identified here can be organized into well-defined superfamilies and families based on basic aspects of the aboral-cup architecture and arm characteristics (see Systematic Paleontology, p. 19–22).

#### CAMERATE PHYLOGENY

*Character analysis.*—Five Arenig to Llanvirn camerates were recognized by Ausich (1998a), with *Celtocrinus* being the only monobathrid (two-circllet) camerate and *Proexenocrinus* being the oldest and most primitive diplobathrid (three-circllet camerate). Twenty-seven Arenig to Caradoc camerates are known (Appendix C, p. 35).

Relationships among these crinoids are delineated with four character analyses (Fig. 4–5): all camerates, only monobathrids, and only diplobathrids (two analyses). For camerates, all Arenig to Caradoc camerates are analyzed with the exception of *Schizocrinus* Hall, which is very poorly known. Details of the analyses are provided above and in the figure captions and appendices.

*Inferred phylogeny.*—As discussed by Ausich (1998a), the early camerates are phylogenetically close to early dendrocrinid cladids, with cladids being more primitive. Cladograms from character analyses of these early crinoids have cladids and camerates ladderized (Ausich, 1998a), with key camerate characters (although not maintained by all taxa) being the incorporation of fixed brachials and interradials, pinnulate arms, and change in the posterior interray (Ausich, 1998a). Both three- and two-circllet camerates occur at the start of the Arenig with *Proexenocrinus* and *Celtocrinus*, respectively. Verified both by character analyses of all camerates (Fig. 4.1) and of monobathrids only (Fig. 4.2), *Celtocrinus* is linked with other monobathrids and should be regarded as the base of this monophyletic clade. Similarly, *Proexenocrinus* is considered to be the base of the diplobathrid clade (Ausich, 1998a).

*Proexenocrinus* is a simple diplobathrid, with relatively few fixed brachials and interradials, ten nonbranching arms with uniserial pinnulation, and first interradials and basals in contact in all interrays, thus separating all radial plates (Ausich, 1986b). This condition, the rhodocrinitacean arrangement, is present in all Arenig to Caradoc diplobathrids except *Euptychocrinus*.

From *Proexenocrinus* six diplobathrid lineages became established during the Llanvirn. The first was a continuation of the basic morphology of *Proexenocrinus* to yield the






Figure 8. Phylogenetic interpretation of cladid crinoids with one flexible, *Protaxocrinus*, included (new).

Archaeocrinidae-Rhodocrinitidae of Ubachs (1978c). This lineage was the principal Caradoc diplobathrid radiation. *Euptychocrinus* and *Reteocrinus* are two unusual Caradoc diplobathrids that are consistently linked peripherally to other diplobathrids in cladograms herein (Fig. 4.1, 5.1). Both were probably derived from *Neoarchaeocrinus* or a similar form, and each was an independent lineage (Fig. 9). As mentioned above, *Euptychocrinus* represented a change to an architecture with the radial circlet complete, except for the posterior interarray. Whereas this is a unique Arenig to Caradoc diplobathrid morphology, it records the base of the dimerocrinitaceans that diversified during the Silurian. *Reteocrinus* is also a very unusual Arenig to Caradoc diplobathrid because it has very small, irregular interradial plating; no large first interradial plate; and no

pinnules. This is a lineage that did not persist beyond the Ordovician.

*Trichinocrinus*, the fourth lineage, is another unusual diplobathrid because its posterior interarray is similar to the cladid posterior interarray (Ausich, Bolton, and Cumming, 1998) and may be a reversal to that condition. *Trichinocrinus* has an asymmetrical posterior interarray with a single large plate in sutural contact with the CD basal. The anitaxis and anitaxial ridge are derived from a heptagonal C radial. The essential features of this unusual posterior interarray also occur in *Pararchaeocrinus* (see Kolata, 1982), but these two forms are otherwise similar to other archaeocrinids-rhodocrinitids. In contrast, in other camerates, the posterior interarray is bilaterally symmetrical, and it begins on a single plate, the primanal, at the proximal part of the

interray. Typically either two or three plates form the first range of plates above the primanal, and if an anitaxis and/or anitaxial ridge is present its origin is from the primanal.

A fifth lineage comprises only *Cleioocrinus*. *Cleioocrinus* consistently links with *Trichinocrinus* in parsimony analyses (Fig. 4.1, 5), but this probably reflects that these two forms are significantly different from other diplobathrids rather than the fact that they are similar to one another. *Cleioocrinus* is linked to *Trichinocrinus* in analyses herein because both have a simple, small aboral cup with a basal concavity, no fixed pinnules, cuneate uniserial arms, no intrabrachials, and atomous free arms. *Cleioocrinus*, however, is so unusual that its origin from other diplobathrids is problematic. Whereas *Cleioocrinus* has a small aboral cup, it has a huge calyx composed of hypertrophied, fused brachials. Small pores penetrate the calyx at triple junctions among fused brachials, no intrabrachials are present, and the posterior interrayer is composed exclusively of the anitaxis. For the present, this unique Ordovician crinoid is regarded questionably as having been derived from *Proexenocrinus*, but no intermediate forms are known, and no Llanvirn ancestor is known. It is improbable that *Cleioocrinus* was derived from *Trichinocrinus*, because it lacks the synapomorphies of *Trichinocrinus* and *Pararchaeocrinus*.

The sixth diplobathrid lineage from *Proexenocrinus* was established from *Deocrinus* and also includes *Hercocrinus*, *Gustabilicrinus*, and *Anthracocrinus*; these are the Anthracocrinidae of Ubaghs (1978c). This lineage did not survive beyond the Caradoc.

Despite the lack of a Llanvirn monobathrid, *Celtocrinus* is certainly the root of the monobathrid clade because it has the basic characteristics of the Glyptocrinidae, which, with the exception of *Eopatelliocrinus*, represents the Arenig to Caradoc radiation of monobathrids (Fig. 4.1, 5.2, 9). *Abludoglyptocrinus* is apparently the most primitive Caradoc glyptocrinid (Fig. 4.2). *Eopatelliocrinus* is inferred to have evolved from *Abludoglyptocrinus*. Unlike glyptocrinids with a large calyx composed of several fixed brachials and interradians, *Eopatelliocrinus* has a much simplified calyx with very few fixed brachials and interradians (Brower, 1994). This began the patelliocrinacean lineage.

**Discussion.**—Early phylogeny of camerates was confused by Ubaghs (1978b) because *Proexenocrinus* was considered at that time to be a monobathrid (see Ausich, 1986b). Ubaghs (1978b) indefinitely linked major groups, although he did regard archaeocrinids as the stock from which other diplobathrids arose. This is consistent with the interpretation herein and with the interpretation of *Proexenocrinus* as a diplobathrid. Origins of monobathrids were problematic to Ubaghs (1978b), again due in part to *Proexenocrinus*. An intermediate form between diplobathrids and monobathrids (Ubaghs, 1978b), however, has still not been found.

Origins and relationships among camerates presented here differ substantially from those proposed by Donovan

(1988a) and Simms (1994a, 1994b). Donovan (1988a) used *Echmatocrinus* as the outgroup for crinoids (see Ausich and Babcock, 1996, 1998), which led to a different topology; and Simms (1994b) used very different plate-circlet homologies.

## DISPARID PHYLOGENY

**Character analysis.**—Among the earliest crinoids, the Disparida are the most diverse, with 45 genera considered here. The disparids are an interesting combination of mostly conservative, relatively simple forms with few, very distinctive, derived types. This combination is not well suited for robust parsimony character analysis. The advantages of a stepwise approach to parsimony character analysis is most evident among the disparids. Analysis of all genera (Fig. 6) is poorly resolved and cannot be interpreted clearly. Disparid phylogeny, however, can be understood from consideration first of relationships among Arenig and Llanvirn disparids (Ausich, 1998a) and then by adding analyses of Caradoc forms. Analyses of various subsets of the disparids are required to understand detailed relationships. Details of the analyses are listed above and in the figure captions and appendices.

**Inferred phylogeny.**—The initial disparids evolved through loss of the basal circlet from the four-circlet aethocrinid condition with five plates in each circlet (plus the posterior plates; Ausich, 1998a). Therefore, the most primitive disparid had five lintels, five infrabasals, and five radials, such as *Pogonipocrinus* (see Kelly and Ausich, 1978, 1979) and *Inyocrinus* (see Ausich, 1986b). The basic topology of disparid phylogeny was established during the Arenig and Llanvirn (Fig. 1). In this earliest radiation, six additional fundamental lineages became established that are identifiable on the pattern of loss of radial plates and architecture of the aboral cup. These lineages are treated here at the ordinal rank. One additional order, Calceocrinida, became established during the Caradoc. Lineages that became established include: 1. those with five infrabasals and five radials; 2. the lineage beginning with *Pultivocrinus* that lost all radials; 3. the lineage beginning with "*Pariocrinus*" that retained a single radial plate in the C ray; 4. the highly specialized hybocrinids that also retained only the C radial; 5. the homocrinid-cincinnatiocrinid lineage that began with *Ibexocrinus* with genera losing various radial plates; and 6. *Ramseyocrinus* and *Tetragonocrinus* that lost all plates in the cup except the lintels (Fig. 1). The calceocrinids were the seventh lineage, and these unique crinoids first appeared during the Caradoc. These seven basic clades are evident on the cladogram that includes all disparids (Fig. 6). This character analysis, although not particularly robust, still has most branches with 100-percent agreement and also identifies several key relationships, including the following: 1. crinoids with five (or four) infrabasals and five (or four) radials are grouped adjacent to the outgroup on a basal polytomy




Figure 9. Phylogenetic interpretation of camerate crinoids (new).

(*Pogonipocrinus*, *Ristnacrinus*, *Inyocrinus*, *Peniculocrinus*, *Eustenocrinus*, and *Virucrinus*); 2. *Ramseyocrinus* and *Tetragonocrinus* are linked; 3. *Ibexocrinus* is deeply rooted (an early branch) in the tree; and 4. even in parts of the cladogram that are poorly resolved, crinoids with one radial plate and those without radial plates are each fairly well linked (Fig. 6).

Figures 10 and 11 illustrate the inferred relationships among Arenig to Caradoc disparids. No Llanvirn disparids are known with both infrabasals and radials in every ray. It is reasonable, however, to assume that the Arenig and Caradoc crinoids with this morphology are closely related. Two unusual genera in this clade are *Virucrinus*, which lost one ray, so that it has four infrabasals and four radials, and *Acolocrinus*, which developed three lintels, radial processes, multiple armllets on radial facets, and pore structures. *Ristnacrinus* is considered to be the most primitive Caradoc form in this clade.

The extremely paedomorphic *Ramseyocrinus* and *Tetragonocrinus* had a cup composed of a single cirlet, the lintels, and left no descendants after the Llanvirn. This new interpretation of these unusual crinoids agrees, in part, with Donovan (1988b, 1989b) and Rozhnov (1988, 1989), who independently argued that *Ramseyocrinus* and *Tetragonocrinus* had an aboral cup composed of a single cirlet of plates: *Ramseyocrinus* with the radial cirlet (Donovan, 1989b) and *Tetragonocrinus* with the infrabasal cirlet (Rozhnov, 1988, 1989). The interpretation of the homology of this plate with the lintels of *Aethocrinus* is new, however.

Those disparids that lost all radials but retained all infrabasals did not survive into the Caradoc either. These include *Pultivocrinus* and *Maennilicrinus* from the Arenig and *Vosekocrinus* and *Pandorocrinus* from the Llanvirn. *Maennilicrinus* is inferred to be the ancestor of these Llanvirn forms (Fig. 10).

Various additional character analyses were also performed on the rest of the disparids in order to develop a more robust understanding of these relationships; two are presented here (Fig. 7). Figure 7.1 includes Llanvirn and Caradoc disparids that lost two to four radial plates, including those placed by Moore *et al.* (1978) in the Homocrinacea and in the Heterocrinacea (=Cincinnati crinacea of Warn and Strimple, 1977). This character analysis yields eight equally parsimonious trees. The resultant consensus-tree cladogram cannot be taken as the phylogenetic tree for these crinoids because one polytomy exists and because *Isotomocrinus* should presumably be more deeply rooted as it occurs in Llanvirn strata. This Caradoc radiation occurred entirely in North America, with the exception of *Tunguskocrinus* from the Siberian Platform, Russia. Therefore, despite the similarity of the aboral cup architecture of *Tunguskocrinus* to homocrinids, it is possible that *Tunguskocrinus* represents convergence and is not closely related to other homocrinids. Despite the fact that no Llanvirn homocrinids exist, this analysis and the phylogenetic reconstruction assume that all homocrinids evolved directly through *Ibexocrinus* rather than the homocrinids having been derived from *Isotomocrinus*, which would require first loss, then reevolution of the B radial. *Ectenocrinus* is regarded as the Caradoc homocrinid stock from which all others arose. It is thought to have evolved from *Ibexocrinus* or a similar form. From *Ectenocrinus*, six additional homocrinids arose during the Caradoc. Homocrinid-derived forms that are not shown in Figure 7.1 are the calceocrinids and *Glaucoocrinus*. The calceocrinids branch near the base on the cladogram derived from analysis of all disparids (Fig. 6). The calceocrinids *Cremacrinus*, *Calceocrinus*, and *Paracremacrinus* are considered to be a lineage that diverged early and independently from the rest of the homocrinids, as indicated in Figure 6. Harvey and Ausich (1997) recently considered the phylogeny of the calceocrinids, which is given in Figure 11. Despite the fact that Guensburg (1992) did much to clarify the morphology of *Glaucoocrinus*, questions about it remain. The sole specimen is poorly preserved, so some aspects of its unusual morphology may be a preservational artifact, or this may simply be an aberrant form. Therefore, it was not used in any analyses.

All character analyses performed on homocrinids-cincinnati crinids, regardless of the taxa, outgroup, or characters considered, consistently grouped three Cincinnati crinids, *Columbicrinus*, *Praecursoricrinus* (see Systematic Paleontology, p. 27), and *Geraocrinus*, separately from the other Cincinnati crinids, *Cincinnati crinid*, *Isotomocrinus*, *Doliocrinus*, and *Tryssocrinus*. Two alternative interpretations could explain this pattern: the loss of the B radial, the defining characteristic for the Cincinnati crinidae (=Heterocrinacea of Moore and Lane, 1978a) may have happened twice, and the character analyses presented in Figures 6 and 7.1 accurately reflect crinoid evolution; or these char-

acter analyses are flawed in some way, and the loss of the B radial among these crinoids was a unique evolutionary event. Until additional data suggest otherwise, I favor the former interpretation for three reasons: given available data separation between these taxa is very consistent; loss of radial plates was a changeable character during this early disparid diversification; and other disparids independently lost the B radial, e.g., "*Pariocrinus*" and its descendants, *Maennilicrinus* and its descendants, *Paracremacrinus*, and *Othneiocrinus*.

In summary, disparids that lost between two and four radial plates represent a single adaptive radiation from *Ibexocrinus* with five evolutionary lineages. First, *Othneiocrinus* is an unusual Arenig form that has different radial facet types in different rays and pinnulate arms. It did not give rise to descendants, as far as is known. Second, the basic homocrinid morphology supported the interpretation of a radiation of six additional genera. Third, *Isotomocrinus* became the first of the Cincinnati crinids that also included *Cincinnati crinid*, *Ohioocrinus*, *Doliocrinus*, and *Tryssocrinus*. The fourth group includes *Columbicrinus*, *Praecursoricrinus*, and *Geraocrinus*. Fifth, the calceocrinids with a hinged aboral cup and bilaterally symmetrical crown developed from the homocrinid stock.

Finally, disparids that retained a single radial plate (C radial) form two distinct clades. The oldest member of this group is "*Pariocrinus*" Rozhnov, 1988 (*non Pariocrinus* Eckert, 1984) (Fig. 1). "*Pariocrinus*" first appeared during the Arenig as the base of the iocrinid clade, and the hybocrinids began to diversify during the Llanvirn. Iocrinids include "*Pariocrinus*," *Iocrinus*, *Tornatilicrinus*, *Peltacrinus*, and *Caleidocrinus* and have cone- or bowl-shaped aboral cups, plenary to peneplenary radial facets, and arms all erect and branching. In contrast, the hybocrinids have a globose aboral cup, may have recumbent ambulacra, and have angustary radial facets; and where erect arms occur they are atomous. A single most parsimonious tree results regardless of whether "*Pariocrinus*," *Iocrinus*, or *Caleidocrinus* (*Huxleyocrinus*) is used as an outgroup. In all cases, the hybocrinids are linked most closely with *Caleidocrinus* (*H.*). "*Pariocrinus*" is stratigraphically the oldest of these crinoids; the cladogram with it as outgroup is shown in Figure 7.2. This group probably began as an early branch from *Ibexocrinus* (Fig. 1, 10) by losing all radials except the C radial. *Revalocrinus* is intermediate between iocrinids and hybocrinids, and its traditional treatment as the oldest hybocrinid (Sprinkle and Moore, 1978) is followed here.

## CLASSIFICATION

*Introduction.*—Results of these phylogenetic studies indicate that many changes are required in the suprageneric classification of Moore and Teichert (1978) (Table 1) if a phylogenetic classification is to be maintained. Unfortunately, these results do not permit a comprehensive classification of the Crinoidea, but they do provide a means




Figure 10. Phylogenetic interpretation of disparid crinoids (new).



## CALCEOCRINID AND HOMOCRINID EARLY PHYLOGENY




Figure 11. Phylogenetic interpretation of calceocrinid and homocrinid disparid crinoids (new).

to understand the classification of early forms. Completion of a comprehensive classification awaits further phylogenetic analysis of younger crinoids.

As argued by Ausich (1998a) if a hierarchical, Linnaean classification is to be developed, paraphyletic groups must be recognized, as they are herein. The basic topology of crinoid diversification and the grouping of crinoids into subclasses follows Ausich (1998a). This follows the *Treatise* (Moore and Teichert, 1978) (Table 1) with the exceptions that the Inadunata is considered to be polyphyletic, so the Cladida and Disparida are recognized as subclasses (Simms and Sevastopulo, 1993) (Table 2), the coronates are blastozoans (Brett *et al.*, 1983), and the hemistreptocrinids are regarded as columns (Arendt and Rozhonov, 1995).

In the present classification, disparids, camerates, flexibles, and articulates are monophyletic. The aethocrinids and cladids are paraphyletic (Ausich, 1998a). Whenever possible, a conservative approach has been taken regarding the grouping of crinoids into subclasses; i.e., names and ranks given by Moore and Teichert (1978) are maintained. This minimizes the number of new names but does

result in classifications within subclasses that are not always parallel. For example, the order Diplobathrida is divided into superfamilies, whereas the order Monobathrida is divided into suborders (Table 3).

*Aethocrinida*.—The least diverse but most primitive crinoid subclass is the Aethocrinea. As argued by Ausich (1998a), this subclass contains three genera grouped into two families and is included in a single order (Table 3 and in *Systematic Paleontology*, p. 19). It is regarded as a paraphyletic group that retains the most primitive crinoids.

*Cladida*.—Cladids are a subclass, as first suggested by Kelly (1982, 1986) and formally designated by Simms and Sevastopulo (1993). Moore, Lane, and Strimple (1978) subdivided the order Cladida into three suborders, the Cyathocrinina, Dendrocrinina, and Poteriocrinina. The validity of these suborders has been questioned by numerous authors (among others, McIntosh, 1986; Sevastopulo and Lane, 1988; Simms and Sevastopulo, 1993). I support recognition of the dendrocrines as the primitive cladids and of the cyathocrines as a derived cladid group (Lane, 1978b). With cladids recognized as a subclass, the

Table 1. Crinoid classification by Moore and Teichert (1978).

Class Crinoidea	
Subclass Echnatocrinea	Order Echnatocrinida
Subclass Camerata	Order Diplobathrida
	Order Monobathrida
Subclass Inadunata	Order Disparida
	Order Hybocrinida
	Order Coronata
	Order Cladida
Subclass Flexibilia	Order Taxocrinida
	Order Sagenocrinida
Subclass Articulata	Order Millerocrinida
	Order Cyrtocrinida
	Order Bourgueticrinida
	Order Isocrinida
	Order Comatulida
	Order Uintocrinida
	Order Roveocrinid
Class Hemistreptocrinoidea <sup>1</sup>	Order Hemistreptocrinida

<sup>1</sup>new class of Crinoidea proposed by Arendt (1976)

Dendrocrinida and Cyathocrinida are elevated to ordinal rank. The relationship of the pinnule-bearing poteriocrines to other cladids is beyond the scope of this study, and this important issue requires additional attention.

The difficult question for unraveling the phylogeny and classification of these early cladids is determining which taxa are morphological experiments that were convergent on later morphologies and did not radiate subsequently, and which were the beginnings of clades that radiated during the Silurian and Devonian. Moore, Lane, and Strimple (1978) and subsequent authors placed Arenig to Caradoc dendrocrines into two superfamilies, the Dendrocrininea and the Merocrininea. The principal distinction between these superfamilies was that the former has angustary radial facets and the latter has plenary radial facets (Moore, Lane, and Strimple, 1978). New taxa described since 1978 blur this single-character distinction, as genera assigned to each superfamily may have penepenary radial facets.

The Dendrocrinidae are regarded as the basic stock of cladid crinoids. The four evolutionary departures from this basic lineage are all regarded differently. The Llanvirn experimentation of adding fixed plates above the radial cirlet resulted in *Archataxocrinus* and questionably *Colpodecrinus*. These two taxa are assigned to the Colpodecrinidae, but this designation is not as strong as others below. The Caradoc experimentation of adding plates yielded two genera assigned to the Cupulocrinidae, *Cupulocrinus* and *Quinquecaudex*. *Protaxocrinus*, the first flexible crinoid, arose from this radiation. The Merocrininea include a group of dendrocrines that developed much shorter aboral cups with much less conspicuous infrabasals, including *Mero-*

*crinus*, *Praecupulocrinus*, *Polycrinus*, "*Dendrocrinus*" *acutidactylus*, *Eopinnocrinus*, and *Metabolocrinus*. These are all placed within the superfamily Merocrininea and the families Merocrinidae and Metabolocrinidae, whereas previously they were all in separate families. The Merocrinidae have three or more (rarely one) primibrachials and isotomous arm branching, whereas the Metabolocrinidae (*Metabolocrinus* and *Eopinnocrinus*) have one primibrachial and pinnulate arm branching (see Systematic Paleontology, p. 21).

Finally, the cyathocrines are recognized as an order. Llanvirn to Caradoc cyathocrines are divisible into three groups: 1. *Eoparisocrinus* and *Illemocrinus* are primitive; 2. the specialized *Palaeocrinus-Porocrinus-Carabocrinus*-type crinoids; and 3. the specialized *Agostocrinus*. Moore, Lane, and Strimple (1978) and subsequent workers have placed nearly all of the cyathocrinid genera into separate families. In contrast, the analysis herein suggests that these are closely related crinoids. For the present, all Llanvirn cyathocrines are placed into one superfamily, the Cyathocrinitacea, and three families. *Eoparisocrinus* and *Illemocrinus* are in the Euspirocrinidae. Assignment of *Eoparisocrinus* follows Ausich (1986c) and Brower (1995a), but *Illemocrinus* is reassigned from the Thalamocrinidae (Eckert, 1987) to the Euspirocrinidae. All of the specialized pore-bearing forms are placed into a single family, Porocrinidae. The Porocrinidae is reassigned to the superfamily Cyathocrinitacea. This means that the Carabocrinidae is a junior synonym of the Porocrinidae. The similarities among these crinoids and members of the Sphaerocrinidae are regarded as convergence. The new family Agostocrinidae is erected for the unusual genus *Agostocrinus*.

These revisions place all elements of this initial cyathocrinid radiation into a single superfamily. Further phylogenetic study is required to determine the relationships of these taxa to younger cyathocrines that did not radiate until the Silurian and Devonian.

*Camerates*.—Arenig to Caradoc camerates are clearly divisible into diplobathrids and monobathrids as traditionally understood, and these are regarded as ordinal in rank (Moore and Laudon, 1943; Ubaghs, 1978c). Ubaghs (1953, 1978c) subdivided the Diplobathrida into the two suborders based on the position of the radial and basal cirlets. If the basals were in the same cirlet as the radials, he regarded them as zygodiplobathrids; whereas if basals were in a typical position beneath the radials, he regarded them as eudiplobathrids. Only two genera were assigned to the zygodiplobathrids by Ubaghs (1978c), *Cleioocrinus* (middle Ordovician) and *Spyridiocrinus* (Lower Devonian). Several authors (Brower, 1975; Kelly, Frest, and Strimple, 1978; Haugh, 1979; Kolata, 1982; Ausich, 1986a) questioned this distinction; and, despite the fact that *Cleioocrinus* is unusual, phylogenetic results here fail to support these two suborders. Consequently, this subordinal distinction is abandoned. Furthermore, although beyond the scope of this study, the superfamily rank of the Nyctocrininea is ques-

Table 2. Crinoid classification by Simms and Sevastopulo (1993).

Class Crinoidea
Subclass Camerata
Order Diplobathrida
Order Monobathrida
Subclass Disparida
Subclass Cladida
"stem-group cladids"
Infraclass Cyathocrinina
Infraclass Flexibilia
Infraclass Articulata
Incertae Sedis ('Subclass') Hybocrinida

tioned. Without suborders and with consideration of the nyctocrinids as only a family, the diplobathrids are divisible into only two superfamilies, Rhodocrinitacea and Dimerocrinitacea (see Systematic Paleontology, p. 23–24).

Arenig to Caradoc rhodocrinitaceans are divided into four families. The most diverse family and the one requiring revision of Ubaghs's work (1978c) is the Rhodocrinitidae. Ubaghs (1978c) distinguished the Archaeocrinidae and Rhodocrinitidae, but Kolata (1982) combined these two families due to the lack of any clear identifying distinctions; Kolata's decision was followed by Ausich (1986a). I also suggest herein that the Arenig to Caradoc genera of these families are part of a single radiation and that it is not divided into rhodocrinitids and archaeocrinids (*sensu* Ubaghs, 1978c). Ubaghs (1978c) placed *Diaboloocrinus*, *Paradiaboloocrinus*, and *Trichinoocrinus*, as well as many younger genera, into the Rhodocrinitidae, whereas the Archaeocrinidae included *Archaeocrinus*, *Balacrinus*, *Neoarchaeocrinus*, *Pararchaeocrinus*, and *Rhaphanoocrinus*. (Note that *Proexenoocrinus* was considered a monobathrid, and *Bromidoocrinus*, *Cotylacrinna*, and *Simplococrinus* were described later.) Thus, the division of these genera by Ubaghs (1978c) does not correspond to the phylogenetic results presented herein. For the present, this family is called the Rhodocrinitidae.

Other Llanvirn to Caradoc families in the Rhodocrinitacea include the Anthracocrinidae with four genera (Fig. 9; see Systematic Paleontology, p. 24) and the monogeneric families Cleiocrinidae and Reteocrinidae. The Dimerocrinitidae, as defined by Ubaghs (1978c), is the only Arenig to Caradoc family in the Dimerocrinitacea, but this group diversified later.

With the exception of *Eopatelliocrinus*, all Arenig to Caradoc monobathrids are placed in the Glyptocrinidae. These are the oldest and most primitive members of the suborder Glyptocrinina (see Systematic Paleontology, p. 24). *Eopatelliocrinus* is the oldest member of the family Patelliocrinidae. The suborder Compsocrinina did not evolve until after the Caradoc.

*Disparids*.—Following Kelly (1982, 1986) and Simms and Sevastopulo (1993), the disparids are herein considered to be a subclass. They evolved through loss of the basal cirlet from a four-cirlet ancestor (Ausich, 1998a). Moore *et al.* (1978) subdivided disparids into superfamilies based largely on the position of the plane of bilateral symmetry through

the crown, aboral cup shape, and radial facets. As a subclass, Disparida requires subdivision into orders. Based on the results herein, Early and middle Ordovician orders are defined on the basis of the presence or absence of plate cirlets, number and position of radial plates, aboral cup symmetry and architecture, radial facets, and free arms. Accordingly, seven orders are recognized for the Ordovician disparids treated here (see Systematic Paleontology, p. 25–28). Five of these became established during the Arenig; the Hybocrinida evolved during the Llanvirn; and the Calceocrinida first appeared during the Caradoc. Undoubtedly, additional orders exist among younger disparids, but delineation of other orders is beyond the scope of the present study.

The Eustenocrinida contains crinoids with radials and infrabasals in every ray. Primitively, eustenocrinids have five radial plates and five infrabasal plates, but *Virucrinus* has only four plates in each cirlet. This order contains two families (see Systematic Paleontology, p. 25 for diagnoses and discussions of each suprageneric category). The Eustenocrinidae has a single arm on each radial facet, whereas the Acolocrinidae has multiple armllets articulating on the radial facet. The order Maennilicrinida comprises four genera in one family; and, in addition to the basal cirlet, they lost the entire radial cirlet. These four genera are all from eastern Europe and, for the present, include two poorly known genera, *Pandorocrinus*

Table 3. Ordinal classification of the Crinoidea followed herein.

Class Crinoidea
Subclass Aethocrinea
Order Aethocrinida
Subclass Cladida
Order Dendrocrinida
Order Cyathocrinida
Order Poteriocrinida <sup>1</sup>
Subclass Flexibilia
Order Taxocrinida
Order Sagenocrinida
Subclass Articulata
Order Millericrinida
Order Cyrtocrinida
Order Bourgueticrinida
Order Isocrinida
Order Comatulida
Order Uintacrinida
Order Roveacrinid
Subclass Camerata
Order Diplobathrida
Order Monobathrida
Subclass Disparida <sup>2</sup>
Order Eustenocrinida
Order Maennilicrinida
Order Tetragonocrinida
Order Homocrinida
Order Calceocrinida
Order Myelodactyla
Order Hybocrinida

<sup>1</sup>The order Poteriocrinida may not be monophyletic and requires further study.<sup>2</sup>The Disparida contains additional, younger orders not listed.

and *Vosekocrinus* (see Systematic Paleontology, p. 25–26). This order became extinct by the end of the Llanvirn (Fig. 1, 10), and these crinoids are not related to middle Paleozoic disparids that also lost all radials. The third small order is the Tetragonocrinida, which contains two genera, *Ramseyocrinus* and *Tetragonocrinus*, placed in a single family. These are exceedingly simplified, perhaps paedomorphic forms, that have only one circlet, the lintel circlet, in the aboral cup. Again, this order did not survive beyond the Llanvirn.

*Ibexocrinus* is the oldest known disparid to lose part of the radial circlet and represents the beginning of the major Early to middle Ordovician radiation of disparids, the order Homocrinida. It probably gave rise to "*Pariocrinus*," the oldest member of the Myelodactylida (Fig. 1, 10–11). It is possible that *Ibexocrinus* gave rise to the order Calceocrinida, but in Figures 10 and 11, the calceocrinids are depicted as having evolved from *Ectenocrinus* because this is most consistent with the stratigraphic occurrence of these genera. The order Homocrinida contains four families. The Homocrinidae is, as in the Homocrinacea (Moore and Lane, 1978b), composed of disparids with radial plates in the B, C, and E rays only (although one genus, *Difficilicrinus*, lost the C radial). In addition, radial facets are plenary, typically four to five primibrachials occur, and more than ten nonpinnulate arms are present. From this design, members of this order iteratively lost the B radial three times (Fig. 7.1, 10–11), and each of these separate evolutionary events is regarded as a distinct family. During the Arenig the monogeneric family Othneiocrinidae arose through loss of the B radial, change in radial facets, variable primibrachial number, and development of ten pinnulate arms. During the Llanvirn, the Cincinnaticrinidae arose with *Isotomocrinus* as the oldest known representative. These are essentially the Heterocrinacea of Moore and Lane (1978a) and the Cincinnaticrinacea of Warn and Strimple (1977), and this family is characterized by the loss of the B radial, restriction to four or five primibrachials (with one possible exception), and retention of more than ten nonpinnulate arms.

The fourth family of the order Homocrinida is the Columbocrinidae, which arose during the Caradoc by loss of the B radial, retention of plenary radial facets, limitation of one to three primibrachials, and development of ten pinnulate arms.

The order Calceocrinida is morphologically unique with no clear transitional or ancestral forms, so it is difficult to determine its origin. *Cremacrinus* with four arms and three (rather than one or two) radials is regarded as the most primitive calceocrinid, and because it contains B, C, and E radials, it has been thought to have had a homocrinid origin (Moore, 1962). Due to the deep rooting of this group in the cladogram of Figure 6 and the highly specialized morphology, these crinoids are considered to be an order. Similar to *Othneiocrinus*, the calceocrinids evolved a cup with different types of facets and unusual arms, al-

though in calceocrinids the rays affected were different, the arm branching was a bilateral heterotomy, and a movable hinge developed between the lintels and the infrabasal radial circlet.

During the Early and middle Ordovician, the order Myelodactylida had one family (Iocrinidae) represented by five genera. It is characterized principally by an erect crown, only the C radial in the radial circlet, and numerous arm branches. The order Hybocrinida probably arose from "*Pariocrinus*" and comprises six genera divided into three families (see Systematic Paleontology, p. 27–28).

## DISCUSSION

The underlying assumption herein is that the phylogeny and classification of Early and middle Ordovician crinoids should be defined by the characters and character combinations of these crinoids largely independently of younger crinoids. This approach is different from the rationale of previous studies of crinoid phylogeny and classification and yields somewhat different results. This study reveals that a substantial degree of iterative evolution occurred among crinoids during this earliest history. For example, pinnulate arms evolved in many groups; ray plates were fixed into the calyx in camerates, flexibles, and some cladids; and pore structures appeared in several groups.

In the *Treatise* classification (Moore and Teichert, 1978), many of these early crinoid genera were either placed in monogeneric families or as the oldest member of an otherwise middle Paleozoic family. Certainly, a few highly unusual monogeneric families are still recognized; but the primary perspective for interpreting this phylogeny and classification is that these Ordovician radiations may not have any relationship to younger crinoids. This must be determined by phylogenetic study of crinoid lineages through the end-Ordovician extinctions. Some of these Early and middle Ordovician genera represent the origination of a clade that diversified later, such as *Euptychocrinus* for the Dimerocrinitidae and *Protaxocrinus* for the Flexibilia. Many genera, however, especially among the cladids, are reassigned to Ordovician families.

Although I follow the revised homologies of Ausich (1996a) herein, the phylogenetic analyses within the subclasses that are presented here should be independent of conflicts about ideas concerning homologies (Ausich, 1996a; Moore and Teichert, 1978). A study of this nature may require reexamination if new Early and middle Ordovician faunas are described with markedly different morphologies.

## SYSTEMATIC PALEONTOLOGY

### Class CRINOIDEA J. S. Miller, 1821

*Discussion.*—The crinoid classification proposed here uses the subclass classification of Ausich (1998a). A summary of the systematic changes suggested here for Early and middle Ordovician crinoids is as follows. As described by Ausich (1998a), six subclasses are recognized within the



Crinoidea; and *Aethocrinus*, formerly a cladid, and *Perittocrinus* and *Tetracionocrinus*, formerly disparids, are placed in the subclass Aethocrinea.

Within the Cladida a new family of the Cyathocrinitacea is formally recognized, the Agostocrinidae, which was previously considered a disparid group. *Archaeataxocrinus* is reassigned from the flexibles to the cladids, and *Colpodecrinus* and *Colpodecrinidae* Sprinkle and Kolata are reassigned from the camerates to the cladids. *Cupulocrinidae* Moore and Laudon is reassigned to the Dendrocrinacea, and *Porocrinidae* Miller and Gurley is reassigned to the Cyathocrinitacea. Familial reassignments of genera include *Agostocrinus* to the Agostocrinidae, *Archaeataxocrinus* to the Colpodecrinidae, *Quinquecaudex* to the Cupulocrinidae, *Ottawacrinus* to the Dendrocrinidae, *Illemocrinus* to the Euspirocrinidae, *Polycrinus* and *Praecupulocrinus* to the Mero-crinidae, *Eopinnacrinus* to the Metabolocrinidae, and *Palaeocrinus* and *Carabocrinus* to the Porocrinidae. Assignment of junior synonyms include the following: *Ottawacrinidae* Moore and Laudon is a junior synonym of *Dendrocrinidae*, and *Carabocrinidae* Bather is a junior synonym of *Porocrinidae*. Finally, *Ontariocrinidae* Jaekel is considered a *nomen dubium*.

Within the Camerata the subordinal distinction of Ubaghs (1978c; zygodiplobathrids and eudiplobathrids) is not followed. The Cleiocrinidae is reassigned to the Rhodocrinitacea. Familial reassignments of genera to the Rhodocrinitidae include *Archaeocrinus*, *Balacrinus*, *Neoarchaeocrinus*, *Pararchaeocrinus*, *Rhaphanocrinus*, *Simplococrinus*, and *Spyridiocrinus*. *Celtocrinus* is assigned to the Glyptocrinidae. Assignment of junior synonyms includes the following: *Archaeocrinidae* Moore and Laudon and *Spyridiocrinidae* Jaekel are junior synonyms of the *Rhodocrinitidae*, and *Nyctocrinacea* Moore and Laudon is a junior synonym of the *Dimerocrinitacea*.

Within the Disparida is the new order Maennilicrinida; the new orders Eustenocrinida and Tetragonocrinida, having previously been regarded as subfamilies; and the new orders Homocrinida, Calceocrinida, and Myelodactylida, having previously been regarded as superfamilies. New taxa recognized herein are the following: *Maennilicrinidae* and *Columbicrinidae* as well as the *nomen translatum* *Atopocrinidae* Warn and Strimple. *Pandoracrinus* is reassigned from the cladids to the disparids. *Baerocrinus* is considered to be teratological; thus the *Baerocrinidae* is regarded as a *nomen dubium*. The *Acolocrinidae* is now in the *Eustenocrinida*; the *Tetragonocrinidae* is reassigned from the *Myelodactylacea* to the *Tetragonocrinida*; and the superfamilies *Homocrinitacea* and *Cincinnatiocrinitacea* are no longer recognized. Familial reassignments of genera include *Othneiocrinus* to the *Atopocrinidae*; *Glaucocrinus* questionably to the *Cincinnatiocrinidae*; *Columbicrinus*, *Gerocrinus*, and *Praecursoricrinus* to the *Columbicrinidae*; *Apodasmocrinus*, *Difficilicrinus*, and *Tunguskocrinus* to the *Homocrinidae*; *Tornatiliocrinus* to the *Iocrinidae*; *Maennilicrinus*, *Pandoracrinus*, *Pultivocrinus*, and *Vosekocrinus* to the

*Maennilicrinidae*; *Ramseyocrinus* to the *Tetragonocrinidae*. Assignment of junior synonyms include the following: *Apodasmocrinidae*, *Daedalocrininae*, *Homocrininae*, and *Tunguskocrinidae* are junior synonyms of *Homocrinidae*; *Ramseyocrinidae* is the junior synonym of *Tetragonocrinidae*; *Tryssocrininae* and *Atopocrininae* are junior synonyms of the *Cincinnatiocrinidae*; and *Tornatiliocrinidae* is a junior synonym of *Iocrinidae*.

#### Subclass AETHOCRINEA Ausich, 1998

*Diagnosis*.—Aboral cup composed of lintels, infrabasals, basals, and radials; aboral cup plates sutured closely; proximal brachials free or fixed; proximal interradials may be incorporated in aboral cup; tegmen not rigid; mouth probably subtegminal; anus through tegmen; posterior portion with extra plating; pore structures may be present; free arms uniserial, nonpinnulate, imperforate.

*Discussion*.—See Ausich, 1998a.

#### Order AETHOCRINIDA Ausich, 1998

*Diagnosis*.—Same as subclass.

*Included families*.—Family *Aethocrinidae* Ubaghs and *Perittocrinidae* Abel.

#### Family AETHOCRINIDAE Ubaghs, 1969

*Diagnosis*.—Aethocrinid with lintels, infrabasals, basals, and radials all large cup plates; five lintels and infrabasals; first primibrachial partially incorporated into calyx; super-radial and inferradial nearly same size; pore structures absent; column pentameric.

*Discussion*.—See Ausich, 1998a.

*Included genus*.—*Aethocrinus* Ubaghs.

#### Family PERITTOCRINIDAE Abel, 1920

*Diagnosis*.—Aethocrinid with lintels and radials as large cup plates, infrabasals and basals small cup plates; four lintels and infrabasals; first primibrachials free; inferradial much larger than superradial; pore structures may be present, column tetrameric.

*Discussion*.—See Ausich, 1998a.

*Included genera*.—*Perittocrinus* Jaekel and *Tetracionocrinus* Ubaghs.

#### Subclass CLADIDA Moore and Laudon, 1943

*Diagnosis*.—Aboral cup primitively composed of infrabasals, basals, and radials; aboral cup plates sutured closely; proximal brachials primitively free, fixed in a few derived forms; proximal interradials primitively not incorporated in aboral cup, incorporated in some derived forms; tegmen not rigid; mouth subtegminal; anus through tegmen, from anal sac, or through aboral cup; primitively posterior portion with extra plating, no extra plating in some derived forms; pore structures may be present; free arms uniserial, nonpinnulate or pinnulate, imperforate.

*Discussion*.—The elevation of the Cladida to subclass status by Kelly (1982, 1986) and Simms and Sevastopulo



(1993) is followed here. Two orders appeared early in the Ordovician, the Dendrocrinida and the Cyathocrinida. The pinnulate cladids were placed in an equally ranked category by Moore, Lane, and Strimple (1978) as the poteriocrinids. The monophyletic nature of this grouping has been widely questioned (McIntosh, 1986; Sevastopulo and Lane, 1988; Simms and Sevastopulo, 1993; and others). Resolution of the disposition of these advanced cladids awaits phylogenetic analyses of Devonian forms and cannot be addressed here.

### Order DENDROCRINIDA Bather, 1899

[*nom. transl.* Ausich, herein, *ex* Dendrocrinina Bather, 1899]

*Diagnosis.*—Aboral cup low to medium cone or low to high bowl shaped, plate sculpturing variable; radial facets angustary to plenary; anal sac tall, cylindrical or inflated, may be porous, may have conspicuous plicate plates; arms rectangular or cuneate uniserial brachials in early forms, nonpinnulate or pinnulate, isotomous or heterotomous arm branching; tegmen plates poorly sutured, presumably composed of small plates, orals not prominent; pore structures typically absent, rarely present in early forms.

*Discussion.*—The Dendrocrinida is considered to be the primitive, basic stock of cladids from which other monophyletic groups were derived (Lane, 1978b). Thus, the Dendrocrinida is a paraphyletic group with camerates, flexibles, and cyathocrines derived from them. In addition, the advanced cladids, the pinnulate poteriocrines of Moore and Teichert (1978), were derived from the dendrocrines. The phylogenetic and taxonomic status of the former is uncertain and beyond the scope of my study, as discussed above.

### Superfamily DENDROCRINACEA Bather, 1899

*Diagnosis.*—Aboral cup high to medium; infrabasals high; radial facets angustary to plenary; radianal compound or simple, if simple, not full proximal width of C radial; fixed brachials and interradials present or absent; three or more primibrachials; arm branching isotomous or heterotomous.

*Discussion.*—Early and middle Ordovician Dendrocrinina include the Dendrocrinidae, Colpodecrinidae (previously assigned to the Camerata), and the Cupulocrinidae (previously assigned to the Merocrinacea). In addition, the Ottawacrinidae, previously regarded as a merocrinacean, is considered to be a junior synonym of the Dendrocrinidae.

### Family DENDROCRINIDAE Wachsmuth and Springer, 1886

Ottawacrinidae Moore and Laudon, 1943.

*Diagnosis.*—Radial facets angustary, radianal compound or simple; pore structures, fixed brachials, interradials absent; three to eight primibrachials.

*Discussion.*—The concept of this family is essentially unchanged from that of Moore and Lane (1978c) and subsequent authors. A new cladid genus to be described by Ausich, Bolton, and Cumming (1998) is also considered

herein. The genus *Ottawacrinus*, however, is reassigned to the Dendrocrinidae, which makes the Ottawacrinidae a junior synonym of the Dendrocrinidae. Also, one Early to middle Ordovician genus, *Quinquecaudex*, is removed from this family and placed within the Cupulocrinidae.

*Included genera.*—*Dendrocrinus* Hall, *Brechmocrinus* Ausich, Bolton, and Cumming, *Compagicrinus* Jobson and Paul, *Elpasocrinus* Sprinkle and Wahlman, *Esthonocrinus* Jaekel, *Grenprisia* Moore, *Ottawacrinus* Billings, and *Plicodendrocrinus* Brower are Early and middle Ordovician genera.

### Family COLPODECRINIDAE Sprinkle and Kolata, 1982

*Diagnosis.*—Radial facets fixed peneplenary, radianal simple or compound, pore structures present or absent, fixed brachials and interradials present, three primibrachials.

*Discussion.*—This family is retained to include *Archaetaxocrinus* and *Colpodecrinus*, which are believed to represent a Llanvirn divergence from the basic dendrocrinid design. *Colpodecrinus* differs in many respects from *Archaetaxocrinus*, but it is more likely to have evolved from this genus than from any other. Of the rearrangement of genera into higher categories presented here for cladids, this is the most problematic.

Sprinkle and Kolata (1982) placed this very unusual crinoid into the Camerata; however, *Colpodecrinus* and Colpodecrinidae are reassigned herein to the Cladida. *Colpodecrinus* is considered herein to be a cladid rather than a camerate (Sprinkle and Kolata, 1982) because the posterior interray is unlike camerates, and most Caradoc camerates have biserial arms. The posterior interray of *Colpodecrinus* is reinterpreted as having four infrabasal plates, four basal plates, five radial plates with the C radial small, radianal, anal X, and fixed brachials and interradials. The radianal and anal X interrupt the radial circlet, and the hexagonal radianal supports the anal X above to the left and the small C radial above to the right. By placing *Archaetaxocrinus* in the Colpodecrinidae, it is reassigned from its previous position in the Taxocrinidae.

*Included genera.*—*Colpodecrinus* Sprinkle and Kolata and *Archaetaxocrinus* Lewis.

### Family CUPULOCRINIDAE Moore and Laudon, 1943

*Diagnosis.*—Radial facets peneplenary to plenary; radianal simple; pore structures absent; fixed brachials absent, interradials present or absent; three or four primibrachials.

*Discussion.*—The family Cupulocrinidae is redefined to include *Cupulocrinus* and *Quinquecaudex*, which represent a radiation away from the dendrocrinid lineage toward the flexibles. Two morphological changes characterize this transition: incorporation of fixed interradial plates into the calyx, and widening of radial facets from angustary to plenary. Although *Quinquecaudex* has angustary radial facets like dendrocrinids, the incorporation of fixed interradials distinguishes it from these forms. *Cupulocrinus*

has both of these advanced features. *Praecupulocrinus* is reassigned to the Merocrinidae.

*Included genera.*—*Cupulocrinus* d'Orbigny and *Quinquecaudex* Brower and Veinus.

### Superfamily MEROCRINACEA S. A. Miller, 1890

*Diagnosis.*—Aboral cup low conical; infrabasals low; radial facets peneplenary to plenary; radianal simple, full proximal width of C radial; fixed brachials and interradians absent; one to many primibrachials; arm branching isotomous, heterotomous, or endotomous.

*Discussion.*—Early and middle Ordovician families of the Merocrinacea include the Merocrinidae and Metabolo-crinidae. The Cupulocrinidae are reassigned to the Dendrocrinacea, the Ottawacrinidae is considered a junior synonym of the Dendrocrinidae, and the Ontariocrinidae is considered to be a *nomen dubium*.

### Family MEROCRINIDAE S. A. Miller, 1890

*Diagnosis.*—Three or more (rarely one) primibrachials; isotomous arm branching.

*Discussion.*—Three additional taxa are assigned to this family, which was previously monogeneric. These are *Praecupulocrinus* from the Cupulocrinidae, *Polycrinus* from the Mastigocrinidae, and “*Dendrocrinus*” *acutidactylus*. “*Dendrocrinus*” *acutidactylus* does not belong to *Dendrocrinus* and probably represents a new genus. All of these taxa share the diagnostic characters of the Merocrinidae and, thus, they are considered to be the result of a single small radiation.

*Included genera.*—*Merocrinus* Walcott, *Praecupulocrinus* Brower, *Polycrinus* Jaekel, and “*Dendrocrinus*” *acutidactylus* Billings.

### Family ONTARIOCRINIDAE Jaekel, 1918

*Discussion.*—Because the genus on which this family was founded has unknown taxonomic status, this family should be regarded as a *nomen dubium*.

*Included genus.*—*Ontariocrinus* Jaekel.

### Family METABOLOCRINIDAE Jaekel, 1918

*Diagnosis.*—One primibrachial, arms endotomously pinnulate.

*Discussion.*—This family has members with the unique combination of one primibrachial and endotomously pinnulate arm branching. *Eopinnacrinus* is reassigned to this family from the Botryocrinidae. Despite the fact that these genera are from distinct geographic areas, the unique morphology is considered to record evolutionary descent.

*Included genera.*—*Metaboloocrinus* Jaekel and *Eopinnacrinus* Brower and Veinus.

### Superfamily MASTIGOCRINACEA Jaekel, 1918

*Discussion.*—Details of this superfamily are not treated here, but *Polycrinus* is reassigned from the Mastigocrinidae to the Merocrinidae.

## Order CYATHOCRINIDA Bather, 1899

[*nom. transl.* Ausich, herein, *ex* Cyathocrinina Bather, 1899]

*Diagnosis.*—Aboral cup medium cone, bowl, or globe shaped; primitively plate sculpturing stellate; radial facets angustary; anal sac absent or short, nonporous; if absent, anal sac flush with tegmen or on side of aboral cup, arms rectangular uniserial; brachials nonpinnulate; atomous, isotomous, or heterotomous arm branching; tegmen plates well sutured, typically composed of large plates, orals prominent; pore structures present or absent in early forms, absent in later forms.

*Discussion.*—The Cyathocrinida is a monophyletic clade that deviated from the Dendrocrinida both by assuming only a restricted subset of dendrocrinid morphological variations and by evolving new morphologies. Many of these early distinctions are blurred as cyathocrines diversified later during the Paleozoic. Examples of restricted morphological variance include angustary radial facets, stellate aboral plate sculpturing, rectangular uniserial brachials, and nonpinnulate arms. New features of early cyathocrines include globe-shaped aboral cups in some, reduction of the anal sac to either a small feature or complete elimination, a tegmen with well-sutured, prominent oral plates, and pore structures in many genera.

Lane and Moore (1978) subdivided the Cyathocrinina into three superfamilies, the Cyathocrinitacea, the Gasterocomina, and the Codiocrinacea. Middle Ordovician genera were placed in both the Cyathocrinitacea and the Gasterocomina, and the principal distinction between these two superfamilies was that the former had an anal sac and the latter had an anal opening flush on the tegmen or cup. Accordingly, *Porocrinus*, *Palaeocrinus*, *Carabocrinus*, and *Triboloporus*, which lack an anal sac, were placed in the Gasterocomina. There are no known crinoids that link these Ordovician crinoids to younger gasterocominids, and the one gasterocominid that is alleged to be similar, *Sphaerocrinus* (Lane & Moore, 1978), is actually quite different. My interpretation is that the anal sac was lost in two different lineages of Cyathocrinitacea, once in the Ordovician porocrinids, also characterized by pore structures, and a second time in Silurian and Devonian forms. A better, revised diagnosis of the Gasterocomina requires phylogenetic analysis of Silurian and Devonian cyathocrines.

### Superfamily CYATHOCRINITACEA Bassler, 1938

*Diagnosis.*—Aboral cup cone, bowl, or globe shaped, five or three infrabasals, five radials; radianal simple, compound, or absent; anal X large; tegmen with slender anal sac or anal opening flush on tegmen; pore structures present or absent; arm branching isotomous, rarely heterotomous.

### Family EUSPIROCRINIDAE Bather, 1890

*Diagnosis.*—Aboral cup cone or bowl shaped; three or five infrabasals; radial facets large; radianal simple; radial facets narrow to wide angustary; no pore structures; two or

three anal plates in cup, radial tetragonal or pentagonal; anal sac high; arms erect, isotomous arm branching.

*Discussion.*—Two Early and middle Ordovician genera are assigned to this family. Previously, *Illemocrinus* was assigned to the Thalamocrinidae.

*Included genera.*—*Eoparisocrinus* Ausich and *Illemocrinus* Eckert are Early and middle Ordovician members.

#### Family THALAMOCRINIDAE Miller and Gurley, 1895

*Discussion.*—*Illemocrinus* Eckert is reassigned to the Euspirocrinidae, which restricts this family to early Silurian and younger representatives.

#### Family AGOSTOCRINIDAE new family

*Diagnosis.*—Aboral cup medium, cone shaped; three infrabasals; radial plates much reduced; aboral cup basally composed of infrabasals, basals, and radianal; radianal simple, in contact with infrabasals; radial facet fixed; no pore structures; anal X and anal opening on tegmen; recumbent arms, unusual heterotomous branching.

*Discussion.*—Kesling and Paul (1971) and Moore and Strimple (1978) considered *Agostocrinus* to be a highly unusual disparid and assigned it to the Catillocrinidae. This crinoid, however, has recumbent arms, not multiple facets on radial plates, so it is not similar to other catillocrinids. Termier and Termier (1972) suggested that *Agostocrinus* and *Acolocrinus* should be assigned to a single family and suggested that that family be the Agostocrinidae. *Acolocrinus* is now assigned to a distinct family, the Acolocrinidae (Brett, 1980), and is treated here as a disparid; but *Agostocrinus* is considered as a very unusual cladid. *Agostocrinus* is unusual because of the very reduced radials, recumbent arms, arm branching, and anal X. Although cladids with simple radianals, as interpreted for *Agostocrinus*, do not have the radianal in contact with the infrabasals, cladids with compound radianals do. As a cladid, *Agostocrinus* is interpreted as an unusual, early cyathocrine that left no descendants. The Agostocrinidae was never formally described by Termier and Termier (1972), and it is erected here for this unusual cladid.

*Included genera.*—*Agostocrinus* Kesling and Paul.

#### Family POROCRINIDAE Miller and Gurley, 1894

Carabocrinidae Bather, 1899.

*Diagnosis.*—Aboral cup cone to globe shaped; five infrabasals; anal X, radial plates large; radianal simple or compound; radial facets angustary; pore structures; no anal sac, anal opening in tegmen; arms erect, isotomous or atomous arm branching.

*Discussion.*—All of the conical to globose Llanvirn to Caradoc cyathocrinids with pore structures, well-sutured oral plates, and the anus on the tegmen or the side of the cup are regarded as a single evolutionary cluster of species. Accordingly, they belong to the same family rather than being divided among three families. As in other taxa, the presence of a compound radianal in *Carabocrinus* was con-

sidered by Lane and Moore (1978) to be sufficient to define a separate family, but in the lineage herein, this feature is considered to be only a generic-level character. With this interpretation, Carabocrinidae is regarded as a junior synonym of Porocrinidae. Also, *Palaeocrinus* is reassigned from the Sphaerocrinidae to the Porocrinidae. Furthermore this family is reassigned to the Cyathocrinitacea from the Gasterocomacea.

*Included genera.*—*Porocrinus* Billings, *Carabocrinus* Billings, *Palaeocrinus* Billings, and *Triboloporus* Kesling and Paul.

#### Superfamily GASTEROCOMACEA Roemer, 1854

[*nom. transl.* Moore and Lane in Moore and Strimple, 1973, p. 18, *ex* Gasterocomidae Roemer, 1854, p. 299]

*Diagnosis.*—Aboral cup bowl to globe shaped; five, three, or one infrabasal; five radials; anal X present or absent; radianal simple if present; anus flush with tegmen and on tegmen or on side of aboral cup, no anal sac; pore structures absent; arm branching isotomous or atomous.

*Discussion.*—The Porocrinidae is reassigned from this superfamily to the Cyathocrinitacea. Thus, Gasterocomina is now confined to Devonian and younger forms.

#### Family GASTEROCOMIDAE Roemer, 1854

*Discussion.*—Details of this family are not treated here.

#### Family SPHAEROCRINIDAE Jaekel, 1895

*Diagnosis.*—Details of this family are not treated here.

*Discussion.*—With removal of *Palaeocrinus* from this family, it is restricted to the Devonian.

#### Family CROTALOCRINITIDAE Bassler, 1938

*Discussion.*—Details of this family are not treated here.

#### Superfamily CODIACRINACEA Bather, 1890

*Discussion.*—The details of this superfamily are not treated here.

#### Order POTERIOCRINIDA Jaekel, 1918

[*nom. transl.* Ausich, herein, *ex* Poteriocrinina Jaekel, 1918]

*Discussion.*—Details of these advanced cladids are not treated here because they are younger than the Caradoc. This grouping of crinoids, as organized by Moore, Lane, and Strimple (1978) is undoubtedly polyphyletic to some extent, as suggested by McIntosh (1986), Sevastopulo and Lane (1988), and Simms and Sevastopulo (1993), among others.

#### Subclass FLEXIBILIA von Zittel, 1895

*Diagnosis.*—Aboral cup composed of infrabasals (primitively three, fewer in some derived forms), basals, and radials; aboral cup plates sutured loosely; proximal brachials fixed primitively or free in some derived forms; proximal interradians incorporated primitively in aboral cup, not incorporated in some derived forms; tegmen flexible; mouth on tegmen; anus through tegmen; posterior portion with extra plating primitively, no extra plates in some



derived forms; pore structures absent; free arms uniserial, nonpinnulate, imperforate.

*Discussion.*—*Archaetaxocrinus* is removed from the Flexibilia to the Cladida, an interpretation that follows traditional views on the origins of flexible crinoids from *Cupulocrinus* during the Caradoc (Springer, 1911). See discussion in the phylogeny section above, p. 5.

#### Order TAXOCRINIDA Springer, 1913

*Discussion.*—Only *Taxocrinus* and *Protaxocrinus* are treated here.

#### Superfamily TAXOCRINACEA Angelin, 1878

##### Family TAXOCRINIDAE Angelin, 1878

*Diagnosis.*—See Moore, 1978.

*Included genera.*—*Taxocrinus* Phillips and *Protaxocrinus* Springer. Further phylogenetic analyses are required to determine whether the remaining Taxocrinidae of Moore (1978) belong with these genera or elsewhere. *Archaetaxocrinus* is reassigned to the Colpodecrinidae.

#### Order SAGENOCRINIDA Springer, 1913

*Discussion.*—Details of this order are not treated here.

##### Subclass ARTICULATA von Zittel, 1879

*Diagnosis.*—Aboral cup composed primitively of infrabasals, basals, and radials, few in derived forms; aboral cup plates sutured closely; proximal brachials free primitively, fixed in some derived forms; proximal interradians not incorporated primitively in aboral cup, incorporated in some derived forms; tegmen not rigid; mouth on tegmen; anus through tegmen; posterior portion with no extra plating; pore structures absent; free arms uniserial, pinnulate, perforate.

*Discussion.*—Details of this subclass are not treated here.

##### Subclass CAMERATA Wachsmuth and Springer, 1885

*Diagnosis.*—Aboral cup composed of infrabasals, basals, and radials or of basals and radials; aboral cup plates sutured rigidly; proximal brachials fixed primitively, free in some derived forms; proximal interradians incorporated in aboral cup primitively, not incorporated in some derived forms; tegmen rigid; mouth subtegmental; anus through tegmen or anal tube; posterior portion with extra plating primitively, no extra plates in some derived forms; pore structures absent; free arms uniserial, nonpinnulate, imperforate primitively, in derived forms biserial and pinnulate.

#### Order DIPLOBATHRIDA Moore and Laudon, 1943

*Diagnosis.*—Camerates with aboral cup composed primitively of infrabasals, basals, and radial plates.

*Discussion.*—As discussed above, the suborder distinction (zygodiplobathrids and eudiplobathrids) of Ubaghs (1978c) is not followed. *Cleiocrinus* is placed in its own

family in the Rhodocrinitacea, and *Spyridiocrinus* is probably also a Rhodocrinitidae.

#### Superfamily RHODOCRINITACEA Roemer, 1855

*Diagnosis.*—Diplobathrid with radials separated in all rays by sutural contact between the first plate in interray and the basal plate.

*Discussion.*—A distinction between the Rhodocrinitidae and the Archaeocrinidae is not made; see discussion below.

*Included families.*—Rhodocrinitidae Roemer, Anthracocrinidae Strimple and Watkins, Reteocrinidae Wachsmuth and Springer, and Cleiocrinidae S. A. Miller are Early and middle Ordovician families supported by phylogenetic interpretations herein. Verification of Opsiocrinidae Kier and Anthemocrinidae Jaekel that were recognized in this superfamily by Ubaghs (1978c) requires phylogenetic analysis of younger genera.

##### Family RHODOCRINITIDAE Roemer, 1855

Archaeocrinidae Moore and Laudon, 1943.

Spyridiocrinidae Jaekel, 1918.

*Diagnosis.*—Calyx globular, conical, or bowl shaped; basal concavity shallow or absent; interradian plates large, regular, in contact with tegmen; anitaxis primitively absent in some derived forms; no pores through calyx; some fixed arm plates; weblike extension of fixed brachials and pinnulars absent; arms uniserial or biserial, pinnulate.

*Discussion.*—Kolata (1982) combined the Rhodocrinitidae and Archaeocrinidae to form one family ranging from the Early Ordovician to the early Mississippian. Until further study of younger genera demonstrates otherwise, I follow Kolata's (1982) classification, and the Rhodocrinitidae has priority. It is quite possible, however, that these Ordovician genera, perhaps with some Silurian genera, represent a single clade that is distinct from later forms and that these should be separated at the family level from younger genera in the Rhodocrinitidae as constituted here. If so, Archaeocrinidae would be an appropriate family for the older forms. The following reassignments are made: *Archaecrinus*, *Balacrinus*, *Neoarchaecrinus*, *Pararchaecrinus*, *Rhaphanocrinus*, and *Simplococrinus* are reassigned from the Archaeocrinidae to the Rhodocrinitidae; *Spyridiocrinus*, a Devonian zygodiplobathrid (Ubaghs, 1978c) is also reassigned to the Rhodocrinitidae. At present, two families are regarded as junior synonyms of Rhodocrinitidae. These are Archaeocrinidae, as discussed above, and Spyridiocrinidae, because *Spyridiocrinus* is assigned to the Rhodocrinitidae.

*Included genera.*—Arenig to Caradoc genera that belong in the Rhodocrinitidae are *Archaecrinus* Wachsmuth and Springer, *Balacrinus* Ramsbottom, *Bromidocrinus* Kolata, *Cotylacrinna* Brower, *Crinocrinus* Kolata, *Diabolocrinus* Wachsmuth and Springer, *Neoarchaecrinus* Strimple and Watkins, *Pararchaecrinus* Strimple and Watkins, *Paradiabolocrinus* Brower and Veinus, *Proexenocrinus* Strimple and McGinnis,

*Rhaphanocrinus* Wachsmuth and Springer, *Simplococrinus* Frest, Strimple, and Kelly, and *Trichinocrinus* Moore and Laudon. Twenty-one additional genera are currently assigned to this family (Ubaghs, 1978c; Kolata, 1982; Ausich, 1986a), and *Spyridiocrinus* should also be assigned here.

**Family ANTHRACOCRINIDAE**  
Strimple and Watkins, 1955

*Diagnosis.*—Calyx conical to bowl shaped, deep basal concavity; interrational plates large, regular; not in contact with tegmen; anitaxis absent; no pores through calyx; some fixed arm plates; weblike extension at the base of arms formed by fixed brachials and pinnulars; arms uniserial or biserial, pinnulate.

*Included genera.*—*Anthracocrinus* Strimple and Watkins, *Deocrinus* Hudson, *Gustabilicrinus* Guensburg, and *Herocrinus* Hudson.

**Family RETEOCRINIDAE**  
Wachsmuth and Springer, 1885

*Diagnosis.*—Calyx conical; basal concavity absent; interradial plates small, irregular, in contact with tegmen; anitaxis present; no pores through calyx; some fixed arm plates; weblike extension of fixed brachials and pinnulars absent; arms uniserial, pinnulate or nonpinnulate.

*Included genera.*—*Reteocrinus* Wachsmuth and Springer is the only genus demonstrated herein to belong to this family. Further phylogenetic analyses are required to determine whether *Gaurocrinus* S. A. Miller belongs in this family or elsewhere.

**Family CLEIOCRINIDAE S. A. Miller, 1890**

*Diagnosis.*—Calyx conical, deep basal concavity; interrational plates absent; anitaxis present; pores through calyx, extremely numerous fixed brachials in weblike extension; arms uniserial, pinnulate.

*Discussion.*—This family is moved to the Rhodocrinitacea from the Zygodiplobathrida, the latter being dissolved.

*Included genera.*—*Cleioocrinus* Billings.

**Family OPSIOCRINIDAE Kier, 1952**

*Discussion.*—Details of this family are not treated here.

**Family ANTHEMOCRINIDAE Jaekel, 1918**

*Discussion.*—Details of this family are not treated here.

**Superfamily DIMEROCRINITACEA von Zittel, 1879**

Nyctocrinacea Moore and Laudon, 1943.

*Diagnosis.*—Diplobathrid with radials primitively separated only in the CD interray where primanal and the CD basal plate in sutural contact, in derived forms radials in contact in all interrays.

*Discussion.*—The definition of this superfamily follows Ubaghs (1978c) with the exception that the Nyctocrinidae is included within rather than being a separate superfamily.

ily. Thus Nyctocrinacea Moore and Laudon is considered to be a junior synonym of Dimerocrinitacea.

*Included families.*—Dimerocrinitidae Zittel, Lamptero-crinidae Bather, Gazacrinidae S. A. Miller, Orthocrinidae Jaekel, and Nyctocrinidae Moore and Laudon.

**Family DIMEROCRINITIDAE von Zittel, 1879**

*Diagnosis.*—See Ubaghs, 1978c.

*Included genera.*—*Dimerocrinites* Phillips and *Euptychocrinus* Brower. Further phylogenetic analyses are required to determine whether the remaining Dimerocrinitidae of Ubaghs (1978c) belong with these genera or elsewhere.

**Family LAMPTEROCRINIDAE Bather, 1899**

*Discussion.*—Details of this family are not treated here.

**Family GAZACRINIDAE S. A. Miller, 1892**

*Discussion.*—Details of this family are not treated here.

**Family ORTHOCRINIDAE Jaekel, 1918**

*Discussion.*—Details of this family are not treated here.

**Family NYCTOCRINIDAE Moore and Laudon, 1943**

*Discussion.*—Details of this family are not treated here.

**Order MONOBATHRIDA Moore and Laudon, 1943**

*Diagnosis.*—Camerates with aboral cup composed of basals and radial plates.

**Suborder COMPSOCRININA Ubaghs, 1978c**

*Diagnosis.*—Monobathrid with hexagonal basal circlet; radials separated only in CD interray where primanal and the CD basal plate are in sutural contact.

*Discussion.*—Details of this suborder are not treated here.

**Suborder GLYPTOCRININA Moore, 1952**

*Diagnosis.*—Monobathrid with pentagonal basal circlet; radials adjoining in all interrays because primanal in sutural contact with C and D basal plates.

**Superfamily GLYPTOCRINACEA von Zittel, 1879**

*Diagnosis.*—See Ubaghs, 1978c.

*Included family.*—Glyptocrinidae von Zittel.

**Family GLYPTOCRINIDAE von Zittel, 1879**

*Diagnosis.*—See Ubaghs, 1978c.

*Discussion.*—*Celtocrinus*, previously left unassigned within the monobathrids (Donovan and Cope, 1989), is assigned to the Glyptocrinidae.

*Included genera.*—*Glyptocrinus* Hall, *Abludoglyptocrinus* Kolata, *Celtocrinus* Donovan and Cope, *Periglyptocrinus* Wachsmuth and Springer, *Pynocrinus* S. A. Miller, and *Schizocrinus* Hall.

**Superfamily PATELLIOCRINACEA Angelin, 1878**

*Diagnosis.*—See Ubaghs, 1978c.



**Family PATELLIOCRINIDAE Angelin, 1878**

*Diagnosis.*—See Ubaghs, 1978c.

*Included genera.*—*Patelliocrinus* Angelin and *Eopatelliocrinus* Brower. Further phylogenetic analyses are required to determine whether the remaining Patelliocrinidae of Ubaghs (1978c) belong with these genera or elsewhere.

**Family STELIDIOCRINIDAE Angelin, 1878**

*Discussion.*—Details of this family are not treated here.

**Superfamily MELOCRINITACEA d'Orbigny, 1852**

*Discussion.*—Details of this superfamily are not treated here.

**Superfamily PLATYCRINITACEA  
Austin and Austin, 1842**

*Discussion.*—Details of this superfamily are not treated here.

**Subclass DISPARIDA Moore and Laudon, 1943**

*Diagnosis.*—Aboral cup composed of lintels, infrabasals, and a varying number of radials, primitively five but varies from five to none; aboral cup plates typically sutured rigidly; proximal brachials free; proximal interradians not incorporated in aboral cup; tegmen not rigid; mouth subtegmenal; anus through tegmen or on anal sac; posterior portion with extra plating primitively, no extra plating in some derived forms; pore structures may be present; free arms uniserial, nonpinnulate, imperforate primitively, pinnulate in some derived forms.

*Discussion.*—The interpretation of homology of aboral cup plates used here follows Ausich (1996a); therefore the lowest two circllets of the disparid aboral cup are the lintels and infrabasals. If superradials (*sensu* Ubaghs, 1978a) are present, these are considered to be homologous to the radial circllet of other crinoids. With elevation of the Disparida to a subclass, superfamilies of Moore *et al.* (1978) are regarded as orders: Homocrinida, Calceocrinida, and Myelodactylida. In addition, other new orders include the Eustenocrinida, Tetragonocrinida, and Maennilicrinida; and the order Hybocrinida is maintained. The following higher-level groups are not retained: Tunguskocrinidae and Cincinnaticrinacea.

**Order EUSTENOCRINIDA new order**

*Diagnosis.*—Disparids with aboral cup composed of five lintels, five infrabasals, and five radials primitively, derived form with four infrabasals and radials; plate circllets symmetrical about oral-aboral axis; aboral cup medium cone shaped, radial facets peneplenary or plenary, and free arms branch.

*Discussion.*—The Eustenocrinidae was placed in the Myelodactylacea by Moore *et al.* (1978) along with the Iocrinidae and Myelodactylidae because all shared a crown bilateral symmetry on the A-CD crinoidal plane. In the

present view of disparid evolution, however, the presence of radials rather than the plane of bilateral symmetry is key. The eustenocrinids are considered as primitive and represent a separate order with two families.

*Included families.*—Eustenocrinidae Ulrich and Acolocrinidae Brett.

**Family EUSTENOCRINIDAE Ulrich, 1925**

*Diagnosis.*—Five lintel plates; one arm per radial, arms may branch; radial processes and pore structures absent.

*Discussion.*—This family remains basically the same as defined by Moore *et al.* (1978) and as used by subsequent authors. *Pultivocrinus* is reassigned to the new order Maennilicrinida.

*Included genera.*—*Eustenocrinus* Ulrich, *Inyocrinus* Ausich, *Peniculocrinus* Moore, *Pogonipocrinus* Kelly and Ausich, *Ristnacrinus* Öpik, and *Virucrinus* Rozhnov.

**Family ACOLOCRINIDAE Brett, 1980**

*Diagnosis.*—Three lintel plates; multiple armlets per radial, arms atomous; radial processes present; pore structures present.

*Discussion.*—Brett (1980) recognized *Acolocrinus* as a distinctive crinoid and described the family Acolocrinidae to comprise *Acolocrinus* and *Paracolocrinus*. He considered *Acolocrinus* to be distinct from *Agostocrinus*, to which it had commonly been linked. Brett (1980) also considered *Acolocrinus* to be most probably an allagecrinacean but did not assign his new family to a superfamily. The fact that the acolocrinids have five radials and five infrabasals clearly aligns them with the order Eustenocrinida.

*Included genera.*—*Acolocrinus* Kesling and Paul and *Paracolocrinus* Brett.

**Order MAENNILICRINIDA new order**

*Diagnosis.*—Disparids with aboral cup composed of five lintels and five infrabasals; plate circllets symmetrical about oral-aboral axis; aboral cup cone or bowl shaped, radial facets peneplenary or plenary, and free arms branch.

*Discussion.*—This new order is designated for disparids that lost the entire radial circllet during the Arenig radiation. This order contains one family with four genera.

*Included family.*—Maennilicrinidae new family.

**Family MAENNILICRINIDAE new family**

*Diagnosis.*—Same as for order.

*Discussion.*—This new family unites genera that were previously assigned to several families. *Maennilicrinus* and *Vosekocrinus* were previously in the Iocrinidae, and *Pultivocrinus* was previously in the Eustenocrinidae. *Pandoracrinus* was questionably considered a cladid by Moore and Lane (1978c, p. 615), but it was linked with the disparid *Vosekocrinus*. Both of these crinoids are poorly known, but they appear to be similar and are placed within the Maennilicrinidae.

*Included genera.*—*Maennilicrinus* Rozhnov, *Pandoracrinus* Jaekel, *Pultivocrinus* Rozhnov, and *Vosekocrinus* Jaekel.

#### Order TETRAGONOCRINIDA new order

*Diagnosis.*—Disparids with aboral cup composed of a single circlet of four or five lintels; plate circlets symmetrical about oral-aboral axis; aboral cup low cone shaped, radial facets peneplenary or plenary, and free arms branch.

*Discussion.*—This order includes two very paedomorphic crinoids interpreted here to retain only one circlet, the lintel circlet, in the aboral cup.

*Included family.*—Tetragonocrinidae Stukalina.

#### Family TETRAGONOCRINIDAE Stukalina, 1980

Ramseyocrinidae Donovan, 1984.

*Diagnosis.*—Same as for order.

*Discussion.*—With *Tetragonocrinus* and *Ramseyocrinus* combined into a single family, Tetragonocrinidae Stukalina, 1980 is the senior synonym of Ramseyocrinidae Donovan, 1984. Previously, the Tetragonocrinidae was assigned to the superfamily Myelodactylacea, and the Ramseyocrinidae was unassigned to a superfamily.

*Included genera.*—*Ramseyocrinus* Bates and *Tetragonocrinus* Yeltyshwa.

#### Order HOMOCRINIDA new order

*Diagnosis.*—Disparids with aboral cup composed primitively of five lintels, five infrabasals, and three radials (B, C, and E rays), derived forms may lose radials; plate circlets symmetrical about oral-aboral axis; aboral cup cone or bowl shaped, radial facets primitively plenary, and free arms branch.

*Discussion.*—The Homocrinida is erected for a large number of Early and middle Ordovician crinoids that lost either two or three radial plates. This new order contains four families. Previously, the Homocrinidae was of superfamily rank. The Cincinnaticrinacea is no longer regarded to have rank above the family level and is in the Homocrinida. The Columbocrinidae is new and the Atopocrinidae is a *nomen translatum* from the subfamilial level.

*Included families.*—Homocrinidae Kirk, Cincinnaticrinidae Warn and Strimple, Columbocrinidae new family, and Atopocrinidae Warn and Strimple.

#### Family HOMOCRINIDAE Kirk, 1914

Apodasmocrinidae Frest, Strimple, and McGinnis, 1979.

Daedalocrininae Warn and Strimple, 1977.

Homocrininae Kirk, 1914.

Tunguskocrinidae Arendt, 1963.

*Diagnosis.*—B, C, and E radials primitive, C may be lost; radial facets plenary, all similar on an individual; two to nine primibrachials; more than ten nonpinnulate arms.

*Discussion.*—Genera assigned to the Homocrinidae were previously assigned to a variety of families and superfamilies. Reassignments include the following: *Apodasmocrinus* and *Difficilicrinus* were in the Apodasmocrinidae, *Daedalo-*

*crinus* was in the homocrinid subfamily Daedalocrininae, and *Tunguskocrinus* was in the Tunguskocrinidae within the Pisocrinacea. Previously, the Homocrinidae were divided into subfamilies (Warn and Strimple, 1977), but that distinction is not followed herein. Families and subfamilies now considered to be junior synonyms of the Homocrinidae are the Apodasmocrinidae, Daedalocrininae, Homocrininae, and Tunguskocrinidae.

*Included genera.*—*Apodasmocrinus* Frest, Strimple, and McGinnis, *Cataraquicrinus* Kolata, *Daedalocrinus* Ulrich, *Difficilicrinus* Frest, Strimple, and McGinnis, *Ectenocrinus* S. A. Miller, *Ibexocrinus* Lane, *Penicillicrinus* Warn, and *Tunguskocrinus* Arendt are Early and middle Ordovician genera considered herein as part of the Homocrinidae. Further phylogenetic analyses are required to determine whether the remaining Homocrinidae of Moore and Lane (1978b) belong with these genera or elsewhere, including *Homocrinus* Hall.

#### Family CINCINNATICRINIDAE

Warn and Strimple, 1977

Atopocrininae Warn and Strimple, 1977.

Tryssocrininae Guensburg, 1984.

*Diagnosis.*—C and E radials; radial facets primitively plenary, peneplenary in derived forms, all similar on an individual; four to five primibrachials (one genus questionably assigned has one); more than ten arms.

*Discussion.*—Warn and Strimple (1977) described the Cincinnaticrinacea as containing one family and two subfamilies, and Guensburg (1984) added a third subfamily. As discussed above, the cincinnaticrinids are regarded as belonging to the order Homocrinida. These genera are still aligned as a single family, but subdivision into subfamilies is not considered necessary. Hence, the subfamilies Atopocrininae Warn and Strimple and Tryssocrininae Guensburg are synonymized with the Cincinnaticrinidae. *Columbicrinus* and *Praecursoricrinus* are reassigned to the Columbocrinidae, and *Othneiocrinus* is reassigned to the Atopocrinidae. *Glaucocrinus* is a poorly understood genus (Guensburg, 1992) and is questionably assigned to this family; it had previously been assigned to the Anomalocrinidae.

*Included genera.*—Arenig to Caradoc genera assigned to this family are *Cincinnaticrinus* Warn and Strimple, *Dollocrinus* Warn, *Isotomocrinus* Ulrich, *Ohioocrinus* Wachsmuth and Springer, *Tryssocrinus* Guensburg, and questionably *Glaucocrinus* Parks and Alcock. Additional younger taxa should also be assigned but those designations await further phylogenetic analyses.

#### Family ATOPOCRINIDAE Warn and Strimple, 1977

[*nom. transl.* Atopocrinidae Ausich, herein, *ex* Atopocrininae Warn and Strimple, 1977]

*Diagnosis.*—C and E radials; radial facets plenary and angustary on same individual; two or five primibrachials; ten pinnulate arms.

*Discussion.*—*Othneiocrinus* is a morphologically unusual crinoid that diverged early from *Ibexocrinus* but lacked descendants. This is regarded as a separate evolutionary event from the evolution of the cincinnaticrinids or the columbicrinids, so designation of a monogeneric family is necessary.

*Included genus.*—*Othneiocrinus* Lane.

#### Family COLUMBICRINIDAE new family

*Diagnosis.*—C and E radials; radial facets plenary, all same on an individual; one to three primibrachials; ten pinnulate arms.

*Discussion.*—The Columbicrinidae is designated for three Caradoc disparid crinoids that retained only the C and E radials and pinnulate arms. Based on phylogenetic analyses presented here, these three crinoids represent an independent lineage from *Isotomocrinus*.

Previously, *Columbicrinus* and *Praecursoricrinus* were assigned to the Cincinnaticrinidae, and *Geraocrinus* was assigned to the Anomalocrinidae.

Note that Guensburg (1984) synonymized *Praecursoricrinus* with *Columbicrinus*. Whereas this may indeed be the correct interpretation, the type species of these two genera differ in aboral cup shape, and the number of primibrachials and stem construction is variable. At least for the analyses presented here, these two genera are treated as distinctive.

*Included genera.*—*Columbicrinus* Ulrich, *Geraocrinus* Ulrich, and *Praecursoricrinus* Frest, Strimple, and McGinnis.

#### Order CALCEOCCRINIDA new order

*Diagnosis.*—Disparids with aboral cup composed primitively of four lintels, five infrabasals, and three radials (B, C, and E rays), derived forms lose some lintels and radials; plate circlets articulated by a synarthrial ridge, asymmetrical about oral-aboral axis; aboral cup flattened or conical, radial facets peneplenary, and free arms branch.

*Discussion.*—The Calceocrinidae have previously been recognized as a distinctive lineage at the superfamilial level (Moore *et al.*, 1978). Here, the calceocrinids are elevated to an order because of the distinctiveness of morphology of these crinoids and the deeply rooted position of the Caradoc forms when analyzed with all other Arenig to Caradoc disparids (Fig. 6).

*Included family.*—Calceocrinidae Meek and Worthen.

#### Family CALCEOCCRINIDAE Meek and Worthen, 1869

*Diagnosis.*—Same as for order.

*Discussion.*—Three middle Ordovician genera of this family exist, *Cremacrinus*, *Calceocrinus*, and *Paracremacrinus*. Harvey and Ausich (1997) interpreted the phylogeny of this family and agreed with Moore (1962) that *Cremacrinus* is the most primitive genus.

*Included genera.*—*Calceocrinus* Hall, *Anulocrinus* Ramsbottom, *Catatonocrinus* Brett, *Charactocrinus* Brett, *Chirocrinus* Angelin, *Chiropinna* Moore, *Cremacrinus* Ulrich, *Cunctocrinus*

Kesling and Sigler, *Deltacrinus* Ulrich, *Diaphorocrinus* Eckert, *Dolerocrinus* Prick, *Eohalysiocrinus* Prokop, *Epihalysiocrinus* Arendt, *Espanocrinus* Webster, *Grypocrinus* Strimple, *Halysiocrinus* Ulrich, *Minicrinus* Prokop, *Paracremacrinus* Brower, *Senariocrinus* Schmidt, *Stibarocrinus* Ausich, *Synchirocrinus* Jaekel, and *Trypherocrinus* Ausich.

#### Order MYELODACTYLIDA new order

*Diagnosis.*—Disparids with aboral cup composed of five lintels, five infrabasals, and one radial (C ray); plate circlets symmetrical or asymmetrical about oral-aboral axis; aboral cup cone or bowl shaped, radial facets peneplenary or plenary, and free arms branch.

*Discussion.*—This order corresponds to the superfamily Myelodactylacea of Moore and Lane (1978d) without the Eustenocrinidae. Thus, this order is restricted to disparids with a single radial plate; one radial only in the C ray.

*Included families.*—Myelodactylidae S. A. Miller and Iocrinidae Moore and Laudon.

#### Family MYELODACTYLIDAE S. A. Miller, 1883

*Discussion.*—Details of this family not treated here.

#### Family IOCRINIDAE Moore and Laudon, 1943

Tornatiliocrinidae Guensburg, 1984.

*Diagnosis.*—Myelodactylan with a large crown; crown erect on the column; numerous isotomous arm branches.

*Discussion.*—*Tornatiliocrinus* is reassigned from Tornatiliocrinidae Guensburg to the Iocrinidae, thus the former is a junior synonym of the latter.

*Included genera.*—*Iocrinus* Hall, *Caleidocrinus* (*Caleidocrinus*) Waagen and Jahn, *Caleidocrinus* (*Huxleyocrinus*) Donovan, *Pariocrinus* Rozhnov, 1988 (*non* Eckert, 1984), *Peltacrinus* Warn, and *Tornatiliocrinus* Guensburg.

#### Order HYBOCRINIDA Jaekel, 1918

[*nom. transl. et correct.* Hybocrinida Moore, 1952, p. 613, *ex suborder* Hybocrinites Jaekel, 1918, p. 90]

*Diagnosis.*—Disparids with aboral cup composed of five lintels, five infrabasals, and one radial (C ray); plate circlets symmetrical about oral-aboral axis; aboral cup cone to globe shaped; radial facets angustary; free arms atomous.

*Discussion.*—Except for the removal of *Baerocrinus* and the Baerocrinidae, I follow herein the classification by Sprinkle and Moore (1978). Rozhnov (personal communication, 1996) considered *Baerocrinus* to be teratological.

Rozhnov (1985) argued that the hybocrinids should be elevated to subclass rank, equivalent to the Inadunata of Moore and Teichert (1978). Although the hybocrinids are morphologically distinct and unusual crinoids, this is not necessarily the case in the context of Early and middle Ordovician forms. During the initial crinoid radiation, a great disparity of morphologies developed that included, among others, the one-circlet ramseyocrinids, the calceocrinids with a hinged aboral cup, and the cleiocrinids with fused, hypertrophied brachials. In hybocrinids, the

basic architecture is regarded as that of a disparid composed of five lintels, five infrabasals, and one radial plate, similar to iocrinids. Many of the seemingly unique features of hybocrinids represent either a lack of development or a reversal of characters to an ancestral condition, which is considered here to be that of rhombiferans (Ausich, 1996b, 1998b). On the basis of cryptic pentameres in *Hybocrinus nitidus* Sinclair, Sprinkle (1982) suggested the possibility that hybocrinids were pseudomonocyclic. Rozhnov (1985) used this condition as a criterion to designate the hybocrinids as a distinct subclass. If true, this would be strong evidence for separating the hybocrinids from the disparids. Ausich (1996), however, demonstrated that hybocrinids are similar to other disparids by considering the orientation of lumen angles, which was argued as the most reliable character for determining the symmetry relationship between the calyx and column.

*Included families.*—Hybocrinidae von Zittel, Hybocystitidae Jaekel, and Cornucrinidae Regnéll.

#### **Family HYBOCRINIDAE von Zittel, 1879**

*Diagnosis.*—Hybocrinids with five relatively short, erect arms; one in each ray.

*Included genera.*—*Hybocrinus* Billings, *Hoplocrinus* Grewingk, *Revalocrinus* Jaekel.

#### **Family HYBOCYSTITIDAE Jaekel, 1918**

*Diagnosis.*—Hybocrinids with three relatively short, erect arms with long, recurved ambulacral grooves in A, C, and

D rays; two long, recumbent ambulacra, one each in B and E rays.

*Included genus.*—*Hybocystites* Wetherby.

#### **Family CORNUCRINIDAE Regnéll, 1948**

*Diagnosis.*—Hybocrinids with no erect arms; three recumbent arms, one each in A, C, and D rays; B and E ray with appendages or ambulacra absent.

*Included genera.*—*Cornucrinus* Regnéll and *Tripatocrinus* Sprinkle.

#### **ACKNOWLEDGMENTS**

I am most grateful to M. G. Bassett, National Museum of Wales; D. Collins, Royal Ontario Museum; J. R. Nudds, Manchester Museum, University of Manchester, England; T. E. Bolton, Geological Survey of Canada, Ottawa, Canada; E. Pietrzeniuk, Museum für Naturkunde zu Berlin; P. Sutherland, University of Oklahoma; J. Thompson, United States National Museum of Natural History; and S. Rozhnov, Russian Academy of Sciences, for providing loans and allowing access to specimens used herein. S. M. Bergström aided in questions of Ordovician stratigraphy; S. Rozhnov, B. E. Bodenbender, and N. G. Lane helped by providing valuable discussion on several issues; and J. C. Brower and S. A. Marcus improved earlier drafts of this manuscript. B. Heath helped with typing, and T. Gray and K. Tyler aided with the illustrations. This research was supported by the National Science Foundation (EAR-9404752).



## REFERENCES

- Abel, Othenio. 1920. Lehrbuch der Paläozoologie. Gustav Fisher. Jena. 500 p.
- Angelin, N. P. 1878. Iconographia crinoideorum in Stratis Sueciae Siluricus fossilium. Samson and Wallin. Holmiae. 62 p.
- Arendt, Y. A. 1963. Krona morskoi lilii iz srednego ordovika r. Podkamennoi tunguski [The crown of a crinoid from the Middle Ordovician of the Podkamennaya Tunguska River]. *Paleontologicheskii Zhurnal* 2:117–121.
- . 1976. Ordovikskie iglokozhiie Gemistreptokrinoidei [Ordovician Echinodermata: Hemistreptocrinida]. *Moskovskoe Obshchestvo Ispytatelei Prirody, Byulletin Otdel Geologicheskii* 51:63–84.
- Arendt, Y. A., and S. V. Rozhnov. 1995. Concerning hemistreptocrinoids. *Paleontological Journal* 29:161–166.
- Ausich, W. I. 1986a. Early Silurian rhodocrinitacean crinoids (Brassfield Formation, Ohio). *Journal of Paleontology* 60:84–106.
- . 1986b. The crinoids of the Al Rose Formation (Early Ordovician, Inyo County, California, USA). *Alcheringa* 10:217–224.
- . 1986c. Early Silurian inadunate crinoids (Brassfield Formation, Ohio). *Journal of Paleontology* 60:719–735.
- . 1995. The primitive crinoid had four circlets of aboral cup plates—revised homologies for the class Crinoidea (Echinodermata). *Geological Society of America Abstracts with Program* 27(6):367.
- . 1996a. Crinoid plate circlet homologies. *Journal of Paleontology* 70:955–964.
- . 1996b. Phylogeny and classification of Ordovician Crinoidea (Echinodermata). *Geological Society of America Abstracts with Program* 28(7):292.
- . 1997. Calyx plate homologies and early evolutionary history of the Crinoidea. In J. A. Waters and C. G. Maples, eds., *Geobiology of Echinoderms*. *Paleontological Society Papers* 3:289–304.
- . 1998a. Early phylogeny and subclass division of the Crinoidea (Phylum Echinodermata). *Journal of Paleontology* 72:499–510.
- . 1998b. Origin of the Crinoidea. In Richard Mooi and Malcolm Telford, eds., *Echinoderms*: San Francisco. Balkema Press. Rotterdam. p. 127–132.
- Ausich, W. I. and L. E. Babcock. 1996. Phylogenetic affinities of *Echmatocrinus brachiatus* (Middle Cambrian, Canada). Sixth North American Paleontological Convention Abstracts of Papers, *Paleontological Society Special Publication* 8:16.
- . 1998. Phylogenetic position of *Echmatocrinus brachiatus* Sprinkle 1973. *Palaeontology* 41:193–202.
- Ausich, W. I., T. E. Bolton, and L. M. Cumming. 1998. Whiterockian (Ordovician) crinoid fauna from the Table Head Group, western Newfoundland (Canada). *Canadian Journal of Earth Sciences* 35:121–130.
- Austin, Thomas, Sr., and Thomas Austin, Jr. 1842. Proposed arrangement of the Echinodermata, particularly as regards the Crinoidea, and a subdivision of the class Adelostella (Echinidae). *Annals and Magazine of Natural History (series 1)* 10:106–113.
- Bassler, R. S. 1938. *Pelmatozoa Palaeozoica*. In Werner Quenstedt, ed., *Animalia, Fossilium Catalogus*. W. Junk. s'Gravenhag. p. 1–194.
- Bassler, R. S., and M. W. Moodey. 1943. Bibliographic and faunal index of Paleozoic pelmatozoan echinoderms. *Geological Society of America Special Papers* 45:734 p.
- Bather, F. A. 1890. British fossil crinoids. *Annals and Magazine of Natural History (series 6)* 5:373–388, 458–486.
- . 1899. A phylogenetic classification of the Pelmatozoa. *British Association for the Advancement of Science Report (1898)*:916–923.
- . 1900. The Crinoidea. In E. R. Lankester, ed., *A Treatise on Zoology*. Adam and Charles Black. London. p. 94–204.
- Brett, C. E. 1980. *Paracolocrinus*, a new inadunate crinoid genus from the Rochester Shale (Silurian, Wenlockian) of New York. *Journal of Paleontology* 54:913–922.
- Brett, C. E., T. J. Frest, James Sprinkle, and C. R. Clement. 1983. Coronioidea: A new class of blastozoan echinoderms based on taxonomic reevaluation of *Stephanocrinus*. *Journal of Paleontology* 57:627–651.
- Brower, J. C. 1975. Silurian crinoids from the Pentland Hills, Scotland. *Palaeontology* 18:631–656.
- . 1994. Camerate crinoids from the middle Ordovician (Galena Group, Dunleith Formation) of northern Iowa and southern Minnesota. *Journal of Paleontology* 68:570–599.
- . 1995a. Dendrocrinid crinoids from the Ordovician of northern Iowa and southern Minnesota. *Journal of Paleontology* 69:939–960.
- . 1995b. Eoparisocrinid crinoids from the middle Ordovician (Galena Group) of northern Iowa and southern Minnesota. *Journal of Paleontology* 69:351–366.
- Conway Morris, Simon. 1993. The fossil record and the early evolution of the Metazoa. *Nature* 361:219–225.
- Donovan, S. K. 1984. *Ramseyocrinus* and *Ristnacrinus* from the Ordovician of Britain. *Palaeontology* 27:623–634.
- . 1988a. The early evolution of the Crinoidea. In C. R. C. Paul and A. B. Smith, eds., *Echinoderm Phylogeny and Evolutionary Biology*. Oxford University Press. Oxford. p. 236–244.
- . 1988b. *Ramseyocrinus* (Crinoidea) from the Arenig of Morocco. *Journal of Paleontology* 62:283–285.
- . 1989a. The significance of the British Ordovician crinoid fauna. *Modern Geology* 13:243–255.
- . 1989b. More about *Ramseyocrinus* Bates (Crinoidea). *Journal of Paleontology* 63:124–125.
- . 1994. Extinct sea lilies. *National Geographic Research and Exploration* 10:72–79.
- Donovan, S. K., and J. C. W. Cope. 1989. A new camerate crinoid from the Arenig of South Wales. *Palaeontology* 32:101–107.
- Eckert, J. D. 1984. Early Llandovery crinoids and stelleroids from the Cataract Group (Lower Silurian) in southern Ontario, Canada. *Royal Ontario Museum, Life Sciences Contributions* 137:83 p.
- . 1987. *Illemocrinus amphiatius*, a new cladid inadunate crinoid from the middle Ordovician of Ontario. *Canadian Journal of Earth Sciences* 24:860–865.
- . 1988. Late Ordovician extinction of North American and British crinoids. *Lethaia* 21:147–167.
- Fisher, D. C. 1988. Stratocladistics: integrating stratigraphic and morphologic data in phylogenetic inference. *Geological Society of America Abstracts with Program* 20:A186.
- . 1994. Stratocladistics: morphological and temporal patterns and their relation to phylogenetic process. In Lance Grande and Olivier Rieppel, eds., *Interpreting the Hierarchy of Nature; from Systematic Patterns of Evolutionary Process and Theories*. Academic Press. San Diego. p. 133–171.
- Fortey, R. A., D. A. T. Harper, J. K. Ingham, A. W. Owen, and A. W. A. Rushton. 1995. A revision of Ordovician series and stages from the historical type area. *Geological Magazine* 132:15–30.
- Frest, T. J., H. L. Strimple, and M. R. McGinnis. 1979. Two new



- crinoids from the Ordovician of Virginia and Oklahoma, with notes on pinnulation in the Disparida. *Journal of Paleontology* 53:399–415.
- Guensburg, T. E. 1984. Echinodermata of the middle Ordovician Lebanon Limestone, central Tennessee. *Bulletins of American Paleontology* 86(319):100 p.
- . 1992. *Glaucoocrinus falconeri* Parks and Alcock, 1912 (Crinoidea) and its systematic status. *Royal Ontario Museum Life Sciences Occasional Paper* 39:7 p.
- Guensburg, T. E., and James Sprinkle. 1997. Rhombiferans are not the ancestors of crinoids. *Geological Society of America Abstracts with Program* 29:A341.
- Harvey, E. W., and W. I. Ausich. 1997. Phylogeny of calceocrinid crinoids (Paleozoic: Echinodermata): Biogeography and mosaic evolution. *Journal of Paleontology* 71:229–305.
- Haugh, B. N. 1979. Late Ordovician channel-dwelling crinoids from southern Ontario, Canada. *American Museum Novitates* 2665:25 p.
- Jaekel, Otto. 1895. Beiträge zur Kenntnis der palaeozoischen Crinoiden Deutschlands. *Palaeontologische Abhandlungen (new series)* 3:1–116.
- . 1918. Phylogenie und System der Pelmatozoen. *Paläontologische Zeitschrift* 3:1–128.
- Kelly, S. M. 1982. Origin of the crinoid orders Disparida and Cladida: possible inadunate cup plate homologies. *Third North American Paleontological Convention (Montreal), Proceedings, vol. I*. p. 285–290.
- . 1986. Classification and evolution of class Crinoidea. *Fourth North American Paleontological Convention, Abstracts*. University of Colorado, Boulder. p. A23.
- Kelly, S. M., and W. I. Ausich. 1978. A new Lower Ordovician disparid crinoid from Utah. *Journal of Paleontology* 52:916–920.
- . 1979. A new name for the Lower Ordovician crinoid *Pogocrinus* Kelly and Ausich. *Journal of Paleontology* 53:1433.
- Kelly, S. M., T. J. Frest, and H. L. Strimple. 1978. Additional information on *Simplococrinus persculptus*. *Journal of Paleontology* 52:1227–1232.
- Kesling, R. V., and C. R. C. Paul. 1971. *Agostocrinus* and *Acolocrinus*, two new Ordovician crinoids with peculiar ray and respiratory structures. *University of Michigan Museum of Paleontology Contributions* 23:221–237.
- Kier, P. M. 1952. Echinoderms of the Middle Devonian Silica Formation of Ohio. *University of Michigan Museum of Paleontology Contribution* 10(4):59–81.
- Kirk, Edwin. 1914. Notes on the fossil crinoid genus *Homocrinus* Hall. *United States National Museum Proceedings* 46:473–483.
- Kolata, D. R. 1982. Camerates. In James Sprinkle, ed., *Echinoderm faunas from the Bromide Formation (middle Ordovician) of Oklahoma*. The University of Kansas Paleontological Contributions Monograph 1:170–205.
- Lane, N. G. 1978a. Evolution of flexible crinoids. In R. C. Moore and Curt Teichert, eds., *Treatise on Invertebrate Paleontology*. Part T, Echinodermata 2(2). The Geological Society of America and The University of Kansas. Boulder and Lawrence. p. 301–302.
- . 1978b. Cladida Cyathocrinina and Dendrocrinina. In R. C. Moore and Curt Teichert, eds., *Treatise on Invertebrate Paleontology*. Part T, Echinodermata 2(2). The Geological Society of America and The University of Kansas. Boulder and Lawrence. p. 294–298.
- Lane, N. G., and R. C. Moore. 1978. Suborder Cyathocrinina. In R. C. Moore and Curt Teichert, eds., *Treatise on Invertebrate Paleontology*. Part T, Echinodermata 2(2). The Geological Society of America and The University of Kansas. Boulder and Lawrence. p. 578–600.
- Lewis, R. D. 1981. *Archaeataxocrinus*, new genus, the earliest known flexible crinoid (Whiterockian) and its phylogenetic implications. *Journal of Paleontology* 55:227–238.
- McIntosh, G. C. 1986. Phylogeny of the dicyclic inadunate order Cladid. *Fourth North American Paleontological Convention Abstracts*. University of Colorado, Boulder. p. A31.
- Meek, F. B., and A. H. Worthen. 1869. Descriptions of new Crinoidea and Echinoidea from the Carboniferous rocks of the western states, with a note on the genus *Onychaster*. Philadelphia Academy of Natural Sciences Proceedings 21:67–83.
- Miller, J. S. 1821. A natural history of the Crinoidea or lily-shaped animals, with observations on the genera *Asteria*, *Euryale*, *Comatula*, and *Marsupites*. Bryan and Co. Bristol. 150 p.
- Miller, S. A. 1883. The American Palaeozoic fossils: a catalogue of the genera and species, with names of authors, dates, places of publication, groups of books in which found, and the etymology and significance of the words, and an introduction devoted to the stratigraphical geology of the Palaeozoic rocks, 2nd edition, Echinodermata. Cincinnati, Ohio. p. 247–334.
- . 1890. The structure, classification, and arrangement of American Palaeozoic crinoids into families. *American Geologist* 6:275–286.
- . 1892. North American geology and paleontology, first appendix. Western Methodist Book Concern. Cincinnati, Ohio. p. 665–718.
- Miller, S. A., and W. F. E. Gurley. 1894. New genera and species of Echinodermata. *Illinois State Museum Natural History Bulletin* 5:53 p.
- . 1895. New and interesting species of Palaeozoic fossils. *Illinois State Museum Natural History Bulletin* 7:89 p.
- Moore, R. C. 1952. Crinoids. In R. C. Moore, C. G. Lalicker, and A. G. Fischer, *Invertebrate Fossils*. McGraw-Hill. New York. p. 604–652.
- . 1962. Revision of Calceocrinidae. *The University of Kansas Paleontological Contributions, Echinodermata, Article* 3:40 p.
- . 1978. Flexibilia. In R. C. Moore and Curt Teichert, eds., *Treatise on Invertebrate Paleontology*. Part T, Echinodermata 2(2). The Geological Society of America and The University of Kansas. Boulder and Lawrence. p. 759–812.
- Moore, R. C., and N. G. Lane. 1978a. Superfamily Heterocrinacea Zittel, 1879. In R. C. Moore and Curt Teichert, eds., *Treatise on Invertebrate Paleontology*. Part T, Echinodermata 2(2). The Geological Society of America and The University of Kansas. Boulder and Lawrence. p. 549–550.
- . 1978b. Superfamily Homocrinacea Kirk, 1914. In R. C. Moore and Curt Teichert, eds., *Treatise on Invertebrate Paleontology*. Part T, Echinodermata 2(2). The Geological Society of America and The University of Kansas. Boulder and Lawrence. p. 522–524.
- . 1978c. Superfamily Dendrocrinacea Wachsmuth and Springer, 1886. In R. C. Moore and Curt Teichert, eds., *Treatise on Invertebrate Paleontology*. Part T, Echinodermata 2(2). The Geological Society of America and The University of Kansas. Boulder and Lawrence. p. 607–613.
- . 1978d. Superfamily Myelodactylacea S. A. Miller, 1883. In R. C. Moore and Curt Teichert, eds., *Treatise on Invertebrate Paleontology*. Part T, Echinodermata 2(2). The Geological Society of America and The University of Kansas. Boulder and Lawrence. p. 550–555.
- Moore, R. C., N. G. Lane, and H. L. Strimple. 1978. Order

- Cladida Moore and Laudon, 1943. In R. C. Moore and Curt Teichert, eds., *Treatise on Invertebrate Paleontology*. Part T, Echinodermata 2(2). The Geological Society of America and The University of Kansas. Boulder and Lawrence. p. 578–759.
- Moore, R. C., N. G. Lane, H. L. Strimple, and James Sprinkle. 1978. Order Disparida Moore and Laudon, 1943. In R. C. Moore and Curt Teichert, eds., *Treatise on Invertebrate Paleontology*. Part T, Echinodermata 2(2). The Geological Society of America and The University of Kansas. Boulder and Lawrence. p. 520–574.
- Moore, R. C., and L. R. Laudon. 1943. Evolution and classification of Paleozoic crinoids. *Geological Society of America Special Papers* 46:153 p.
- Moore, R. C., and H. L. Strimple. 1973. Lower Pennsylvanian (Morrowan) crinoids from Arkansas, Oklahoma, and Texas. *University of Kansas Paleontological Contributions*, Article 60, Echinodermata Article 12:1–84, fig. 1–7, pl. 1–23.
- . 1978. Superfamily Allagecrinacea Carpenter and Etheridge, 1881. In R. C. Moore and Curt Teichert, eds., *Treatise on Invertebrate Paleontology*. Part T, Echinodermata 2(2). The Geological Society of America and The University of Kansas. Boulder and Lawrence. p. 537–548.
- Moore, R. C., and Curt Teichert, eds. 1978. *Treatise on Invertebrate Paleontology*. Part T, Echinodermata 2. The Geological Society of America and The University of Kansas. Boulder and Lawrence. 1027 p.
- d'Orbigny, A. D. 1852. *Prodrome du paléontologie stratigraphique universelle des animaux mollusques et rayonnés faisant suite au cours élémentaire de paléontologie et de géologie stratigraphique*, vol. 3. V. Masson. Paris. 196 p.
- Regnéll, Gerhard. 1948. Swedish Hybocrinida (Crinoidea Inadunata Disparata; Ordovician–Lower Silurian). *K. Svenska Vetenskapsakad., Arkiv for Zoologi* 40A:27 p.
- Roemer, C. F. 1854–1855. *Erste Periode, Kohlen-Gebrige*. In H. G. Bronn, *Lethaea Geognostica*, 3rd ed., 1851–1856, vol. 2. E. Schweizerbart. Stuttgart. 788 p.
- Rozhnov, S. V. 1985. Hybocrinea—novyy podklass morskich liliy [Hybocrinea—a new subclass of crinoid]. *Doklady Akademii Nauk SSSR* 280:1012–1015.
- . 1988. Morfologiya i sistematskoye polozheniye nizhneordovikskikh morskich liliy. *Paleontologicheskii Zhurnal* 2:67–79.
- . 1989. The morphology and systematic position of Lower Ordovician sea lilies. *Paleontological Journal* 2:62–75.
- Sevastopulo, G. D., and N. G. Lane. 1988. Ontogeny and phylogeny of disparid crinoids. In C. R. C. Paul and A. B. Smith, eds., *Echinoderm Phylogeny and Evolutionary Biology*. Oxford University Press. Oxford. p. 245–253.
- Simms, M. J. 1994a. Reinterpretation of thecal plate homology and phylogeny in the class Crinoidea. *Lethaia* 26:303–312.
- . 1994b. A new interpretation of crinoid thecal plate homology and phylogeny. In Bruno David, Alain Guille, J.-P. Feral, and Michel Roux, eds., *Echinoderms Through Time*. A. A. Balkema. Rotterdam. p. 257–263.
- Simms, M. J., and G. D. Sevastopulo. 1993. The origin of articulate crinoids. *Palaeontology* 36:91–109.
- Springer, Frank. 1911. On the Trenton echinoderm fauna. *Geological Survey of Canada Memoir* 15-P:69 p.
- . 1913. Crinoidea. In K. A. von Zittel, *Text-book of paleontology*, 2nd ed. Macmillan and Co., Ltd. London. p. 173–243.
- Sprinkle, James. 1982. *Hybocrinus*. In James Sprinkle, ed., *Echinoderm faunas from the Bromide Formation (middle Ordovician) of Oklahoma*. The University of Kansas Paleontological Contributions Monograph 1:119–128.
- Sprinkle, James, and Desmond Collins. 1995. *Echmatocrinus* revisited: still an echinoderm and probably the earliest crinoid. *Geological Society of America Abstracts with Program* 27:A113–A114.
- Sprinkle, James, and T. E. Guensburg. 1997. How the crinoid got its cup plating: Arm brachials down plus stem (penta) meres up. *Geological Society of America Abstracts with Program* 29:A341.
- Sprinkle, James, and D. R. Kolata. 1982. “Rhomb-bearing” camerate. In James Sprinkle, ed., *Echinoderm faunas from the Bromide Formation (middle Ordovician) of Oklahoma*. The University of Kansas Paleontological Contributions Monograph 1:206–211.
- Sprinkle, James, and R. C. Moore. 1978. Hybocrinida. In R. C. Moore and Curt Teichert, eds., *Treatise on Invertebrate Paleontology*. Part T, Echinodermata 2(2). The Geological Society of America and The University of Kansas. Boulder and Lawrence. p. 564–574.
- Strimple, H. L., and W. T. Watkins. 1955. New Ordovician echinoderms. 1, Three new genera. *Washington Academy of Science Journal* 45:347–353.
- Stukalina, G. A. 1980. New species of quadrilateral from the Ordovician of Kazakhstan, Urals, and eastern European platform. In G. A. Stukalina, ed., *New species of ancient plants and invertebrates of the USSR*, 5. Akademiia Nauk SSSR, Paleontologicheskii Institut. Moscow. p. 88–95.
- Swofford, D. L. 1993. PAUP: phylogenetic analysis using parsimony, version 3.1.1. Computer program distributed by the Illinois Natural History Survey. Champaign, Illinois.
- Termier, Henri, and Geneviève Termier. 1972. Sur les affinités des Herpétocystidés (Echinodermata Cystoïdes). *Compte Rendu de l'Académie des Sciences, Paris* 274:1645–1647.
- Ubaghs, Georges. 1953. Classe des Crinoïdes. In Jean Piveteau, ed., *Traité de paléontologie*, Volume 3. Masson and Cie. Paris. p. 658–773.
- . 1969. *Aethocrinus moorei* Ubaghs, n. gen., n. sp., le plus ancien crinoïde dicyclique connu. *The University of Kansas Paleontological Contributions Paper* 38:25 p.
- . 1978a. Skeletal morphology of fossil crinoids. In R. C. Moore and Curt Teichert, eds., *Treatise on Invertebrate Paleontology*. Part T, Echinodermata 2(2). The Geological Society of America and The University of Kansas. Boulder and Lawrence. p. 58–216.
- . 1978b. Evolution of camerate crinoids. In R. C. Moore and Curt Teichert, eds., *Treatise on Invertebrate Paleontology*. Part T, Echinodermata 2(2). The Geological Society of America and The University of Kansas. Boulder and Lawrence. p. 281–292.
- . 1978c. Subclass Camerata Wachsmuth and Springer, 1885. In R. C. Moore and Curt Teichert, eds., *Treatise on Invertebrate Paleontology*. Part T, Echinodermata 2(2). The Geological Society of America and The University of Kansas. Boulder and Lawrence. p. 410–519.
- Ulrich, E. O. 1925. New classification of the “Heterocrinidae.” In A. F. Foreste, *Upper Ordovician faunas of Ontario and Quebec*. Canada Geological Survey Memoir 138:82–104.
- Wachsmuth, Charles, and Frank Springer. 1885. Revision of the Palaeocrinoidea, part 3, sec. 1. Discussion of the classification and relations of the brachiata crinoids, and conclusion of the generic descriptions. *Philadelphia Academy of Sciences Proceedings*. p. 223–364 (1–162).
- . 1886. Revision of the Palaeocrinoidea, part 3, sec. 2. Discussion of the classification and relations of the brachiata crinoids, and conclusion of the generic descriptions.

- Philadelphia Academy of Sciences Proceedings. p. 64–226 (140–302).
- . 1897. North American Crinoidea Camerata. Museum of Comparative Zoology Memoirs 20–21:897 p.
- Warn, John, and H. L. Strimple. 1977. The disparid inadunate superfamilies Homocrinacea and Cincinnaticrinacea (Echinodermata: Crinoidea), Ordovician–Silurian, North America. *Bulletins of American Paleontology* 72(296):138 p.
- Webster, G. D. 1973. Bibliography and index of Paleozoic crinoids 1942–1968. *Geological Society of America Memoir* 137:341 p.
- . 1977. Bibliography and index of Paleozoic crinoids 1969–1973. *Geological Society of America Microform Publication* 8:235 p.
- . 1986. Bibliography and index of Paleozoic crinoids 1974–1980. *Geological Society of America Microform Publication* 16:405 p.
- . 1988. Bibliography and index of Paleozoic crinoids 1981–1985. *Geological Society of America Microform Publication* 18:236 p.
- von Zittel, K. A. 1879. *Handbuch der Palaeontologie, Band 1, Palaeozoologie, Abt. 1.* R. Oldenbourg. München. 557 p.
- . 1895. *Grundzüge der Palaeontologie (Palaeozoologie)*, 1st ed. R. Oldenbourg. München. 971 p.

Appendix A. Species from which morphological data were used to characterize genera for Arenig to Caradoc character analyses. Illustrations and descriptions of these species are by Moore and Teichert (1978) or can be located in the work of Bassler and Moodey (1943) or Webster (1973, 1977, 1986, 1988).

## AETHOCRINIDS

*Aethocrinus moorei* Ubahgs (type species)  
*Perritocrinus radiatus* (Beyrich) (type species)  
*Tetracionocrinus transitor* (Jaekel) (type species)

## CLADIDS

*Agostocrinus xenus* Kesling and Paul (type species)  
*Archaeataxocrinus burfordi* Lewis (type species)  
*Carabocrinus huronensis* Foerste (one of oldest species)  
*Colpodeocrinus quadrifidus* Sprinkle and Kolata (type species)  
*Compagicrinus fenestratus* (Jobson and Paul) (type species)  
*Cupulocrinus humilis* (Billings) (one of oldest species)  
*Dendrocrinus acutidactylus* (Billings) (one of oldest species)  
*Elpasocrinus radiatus* Sprinkle and Wahlman (type species)  
*Eoparisocrinus crossmani* Brower (one of oldest species)  
*Eopinnacrinus pinnulatus* Brower and Veinus (type species)  
*Esthocrinus laevior* Jaekel (type species)  
*Grenprisia billingsi* Springer (type species)  
*Illemocrinus ampiatus* Eckert (type species)  
*Merocrinus typus* Walcott (type species)  
*Metabolocrinus rossicus* Jaekel (type species)  
*Ottawacrinus typus* Billings (type species)  
 for *Palaeocrinus* used two well-preserved species for determination of characters: *Palaeocrinus planobasalis* (Brower and Veinus) and *Palaeocrinus hudsoni* (Sinclair) new cladid genus and species (type species)  
*Plicodendrocrinus proboscidiatus* (Billings) (one of oldest species)  
*Polycrinus ramulatus* Jaekel (type species)  
*Porocrinus conicus* Billings (type species)  
*Praecipulocrinus conjugans* (Billings) (type species)  
*Quinquecaudex glabellus* Brower and Veinus (type species)  
*Triboloporus cryptoplicatus* Kesling and Paul (type species)  
 new cladid genus from Newfoundland Ausich, Bolton, and Cumming (1998)

## FLEXIBLE

*Protaxocrinus laevis* (Billings) (one of oldest species)

## CAMERATES

*Abludoglyptocrinus laticostatus* Kolata (one of oldest species)  
*Anthraxocrinus primitivus* Strimple and Watkins (type species)  
*Archaeocrinus lacunosus* Billings (type species)  
*Balacrinus basalis* (M'Coy) (type species)  
*Bromidocrinus nodosus* Kolata (type species)  
*Celtocrinus ubahgsi* Donovan and Cope (type species)  
*Cleioocrinus regius* Billings (type species)  
*Cotylacrinna sandra* Brower (type species)  
*Crinocrinus parvicostatus* Kolata (type species)  
*Deocrinus asperatus* (Billings) (type species)  
*Diabolocrinus perplexus* Wachsmuth and Springer (type species)  
*Eopatellioocrinus scyphogracilis* Brower (type species)  
*Euptychoocrinus skopaios* (Shumard) (one of oldest species)  
*Glyptocrinus decadaactylus* Hall (type species)  
*Gustabilicrinus plektanikaulos* Guensburg (type species)  
*Hercocrinus elegans* Hudson (type species)  
*Neoarchaeocrinus pyriformis* (Billings) (type species)  
*Paradiabolocrinus irregularis* Brower and Veinus (type species)  
*Pararchaeocrinus decoratus* Strimple and Watkins (type species)  
*Periglyptocrinus billingsi* Wachsmuth and Springer (type species)

*Proexenocrinus inyoensis* Strimple and McGinnis (type species)  
*Pycnocrinus dyeri* (Meek) (well-known species)  
*Reteocrinus alveolatus* Miller and Gurley (one of oldest species)  
*Rhaphanocrinus sculptus* (S. A. Miller) (well-known species)  
*Schizocrinus nodosus* Hall (type species)  
*Simplococrinus persculptus* Frest, Strimple, and Kelly (type species)  
*Trichinocrinus terranovicus* Moore and Laudon (type species)

## DISPARIDS

*Acolocrinus hydraulicus* Kesling and Paul (type species)  
*Apodasmocrinus daubei* Warn and Strimple (type species)  
*Calceocrinus longifrons* Brower (one of oldest species)  
*Caleidocrinus* (*Caleidocrinus*) *multiramous* (Waagen and Jahn) (type species)  
*Caleidocrinus* (*Huxleyocrinus*) *turgidulus* (Ramsbottom) (type species)  
*Cataraquicrinus elongatus* Kolata (type species)  
*Cincinnatiocrinus varibrachialis* Warn and Strimple (type species)  
*Columbicrinus crassus* Ulrich (type species)  
*Cornucrinus mirus* Regnéll (type species)  
*Cremacrinus punctatus* Ulrich (type species)  
*Daedalocrinus bellewillensis* (W. R. Billings) (one of older species)  
*Difficilicrinus coneyi* Frest, Strimple, and McGinnis (type species)  
*Doliocrinus pustulatus* Warn (type species)  
*Ectenocrinus simplex* Hall (type species)  
*Eustenocrinus springeri* Ulrich (type species)  
*Geraocrinus sculptus* Ulrich (type species)  
*Glaucoocrinus falconeri* Parks and Alcock (type species)  
*Hoplocrinus dipentus* (Leuchtenberg) (type species)  
*Hybocrinus conicus* Billings (type species)  
*Hybocystites problematicus* Wetherby (type species)  
*Ibexocrinus leptus* Lane (type species)  
*Inyocrinus strimplei* Ausich (type species)  
*Iocrinus brithdirenensis* Bates (one of oldest species)  
*Isotomocrinus typus* (type species)  
*Maennilicrinus concinnus* Rozhnov (type species)  
*Ohiocrinus brauni* Ulrich (well-known species)  
*Othneiocrinus priscus* (Lane) (type species)  
*Pandorocrinus pinnulatus* Jaekel (type species)  
*Paracremacrinus laticardinalis* Brower (type species)  
*"Pariocrinus" ladogensis* Rozhnov (type species)  
*Peltacrinus sculptatus* Warn (type species)  
*Penicillicrinus parvus* Warn (type species)  
*Peniculocrinus miller* (Wetherby) (type species)  
*Pogonipocrinus antiquus* Kelly and Ausich (type species)  
*Praecursoricrinus sulphurensis* Frest, Strimple, and McGinnis (type species)  
*Pultivocrinus fundatus* Rozhnov (type species)  
*Ramseyocrinus cambriensis* (Hicks) (type species)  
*Revalocrinus costatus* Jaekel (type species)  
*Ristnacrinus marinus* Öpik (type species)  
*Tetragonocrinus pygmaeus* (Eichwald) (type species)  
*Tomatilicrinus longicaudis* Guensburg (type species)  
*Tripatocrinus pustulatus* Sprinkle (type species)  
*Tryssocrinus endotomous* Guensburg (type species)  
*Tunguskocrinus ivanovae* Arendt (type species)  
*Virucrinus kegelensis* (Yeltyscheva) (type species)  
*Vosekocrinus granulatus* Jaekel (type species)



Appendix B. Characters, character states, and character matrix for Arenig to Caradoc cladids-flexibles for PAUP character analyses cited herein.

1. C radial smaller than others: 0, yes; 1, no
2. Radial facets: 0, fixed; 1, angustary; 2, peneplenary; 3, plenary; 4, fixed-recumbent
3. Number of infrabasal plates: 0, 5; 1, 4; 2, 3
4. Aboral cup shape: 0, medium cone; 1, medium globe; 2, low bowl; 3, low cone; 4, medium bowl; 5, high bowl
5. Radial circlet interrupted: 0, CD interrays; 1, no interrays; 2, all
6. Basal concavity: 0, no; 1, yes
7. Pore rhombs: 0, no; 1, yes
8. Plate sculpturing (in addition to ray ridges, if present): 0, stellate; 1, smooth
9. Radial in contact with: 0, infrabasals; 1, basals; 2, C radial; 3, lintels
10. Regular interradii with plate(s) in aboral cup: 0, yes; 1, at corners; 2, no
11. Brachials (most advanced state, typically most distal): 0, rectangular uniserial; 1, cuneate uniserial
12. Brachials fixed: 0, yes; 1, no
13. Erect arm number: 0, 5; 1, 0
14. Arm habit: 0, erect; 1, recumbent
15. Arm branching: 0, isotomous; 1, atomous; 2, endotomous ramules; 3, pinnulate; 4, heterotomous-recumbent; 5, isotomous-heterotomous
16. Number of primibrachials: 0, 7–11; 1, 1; 2, 2; 3, 3–6; 4, all
17. Patelloid processes: 0, no; 1, yes
18. Stem construction: 0, pentameric; 1, holomeric; 2, tetrameric
19. Columnal shape: 0, pentalobate; 1, circular; 2, pentagonal; 3, tetralobate
20. Lumen shape: 0, pentalobate; 1, circular; 2, pentagonal; 3, tetralobate
21. Column (based on proximal part): 0, heteromorphic; 1, xenomorphic; 2, homeomorphic

CHARACTER MATRIX FOR CLADIDS-FLEXIBLES.

<i>Aethocrinus</i>	00000	00000	00000	00000	0
<i>Agostocrinus</i>	14202	10032	00114	10022	?
<i>Archaetaxocrinus</i>	00000	00100	00005	300(12)0	0
<i>Brechmocrinus</i>	02000	00112	01000	00012	2
<i>Carabocrinus</i>	11040	01002	01000	20020	0
<i>Colpodecrinus</i>	00150	01010	00002	30233	0
<i>Compagicrinus</i>	01000	00002	01000	3000?	0
<i>Cupulocrinus</i>	03000	00111	01000	3(01)110	0
" <i>Dendrocrinus</i> "	01000	00112	01000	1011?	0
<i>Elpasocrinus</i>	01000	00002	01000	30011	0
<i>Eoparisocrinus</i>	01000	00012	01000	30(01)(12)0	0
<i>Eopinnacrinus</i>	02040	00112	11003	1011?	0
<i>Esthonocrinus</i>	01000	00012	01000	30?10	?
<i>Grenprisia</i>	01000	00111	01000	30021	0
<i>Illemocrinus</i>	01040	00012	01000	20(01)(12)0	0
<i>Merocrinus</i>	13031	00122	01000	30111	0
<i>Metabolocrinus</i>	03020	00012	11002	10110	0
<i>Ontariocrinus</i>	11000	00012	01000	2011?	0
<i>Ottawacrinus</i>	03000	00112	0100?	30022	0
<i>Palaeocrinus</i>	11010	01012	01000	(123)0010	0
<i>Plicodendrocrinus</i>	01000	00012	01005	30102 0	
<i>Polycrinus</i>	?3000	00112	01005	30?1?	0
<i>Porocrinus</i>	01010	01012	01001	40110	1
<i>Praecupulocrinus</i>	02030	00112	01000	(23)0112	1
<i>Protaxocrinus</i>	00200	00110	00000	21110	0
<i>Quinquecaudex</i>	02000	00110	0(01)000	30000	0
<i>Triboloporus</i>	11010	01012	0100?	30110	?

## Appendix C. Characters, character states, and character matrix for Arenig to Caradoc camerates for PAUP character analyses cited herein.

	CHARACTER MATRIX FOR CAMERATES						
1. Number of basal plates: 0, 5; 1, 3;	<i>Ablduglyptocrinus</i>	01400	02000	13011	11111	11111	?0
2. Number of infrabasal plates: 0, 5; 1, 0	<i>Aethocrinus</i>	00000	00000	00000	00000	00000	00
3. Calyx shape: 0, medium cone; 1, high bowl; 2, low bowl; 3, low cone; 4, medium bowl; 5, high cone; 6, medium globe; 7, low globe	<i>Anthracocrinus</i>	00011	03010	0(24)061	11313	11111	00
4. Radial circlet interrupted: 0, CD interrays; 1, all interrays; 2, no interrays	<i>Archaeocrinus</i>	00411	00010	030(41)1	02111	01111	00
5. Basal concavity: 0, no; 1, yes	<i>Balacrinus</i>	00610	01110	03002	11111	11111	10
6. Ray ridges: 0, yes; 1, no	<i>Bromidocrinus</i>	00611	04100	07023	02111	11100	00
7. Plate sculpturing (in addition to ray ridges, if present): 0, stellate; 1, smooth; 2, pustulose; 3, fine reticulate; 4, vermiform; 5, gently convex; 6, very convex; 7, knobby; 8, coarsely pitted; 9, ray ridges	<i>Celtocrinus</i>	015?0	00???	??011	00112	21111	?0
8. Extra plates in CD interrays: 0, yes; 1, no	<i>Cleiocrinus</i>	00311	05010	012(40)5	01405	11102	21
9. Anitaxis of plates: 0, no; 1, yes	<i>Cotylacrinna</i>	00211	06010	02011	12101	41101	00
10. Anitaxial ridge: 0, no; 1, yes	<i>Crinocrinus</i>	00711	00010	03022	0?214	??1??	??
11. Primal in contact with: 0, CD basal; 1, C+D radials; 2, C radial	<i>Deocrinus</i>	00611	17000	05014	11211	111?1	1?
12. Posterior interradius plating: 0, iRA, sRA, X; 1, P-1-1; 2, P-3-3 or 2; 3, P-3-4 or 5; 4, X-P-3 or 2; 5, irregular; 6, P-1+1+; 7, 2-3-3; 8, P-2-2	<i>Diaboloocrinus</i>	00211	10000	04051	02101	11111	00
13. Interradial plating: 0, large plates; 1, small plates; 2, none	<i>Eopattellioocrinus</i>	11020	00011	13011	11111	11111	00
14. first primibrachial in contact with: 0, ibr1; 1, ibr 1+2; 2, ibr 2+3; 3, radial and ibr 1; 4, radial; 5, ibr 2; 6, ibr 4+5	<i>Euptychoocrinus</i>	00000	05010	0301(12)	01111	11111	20
15. Interradii plating (AB, BC, DE, EA): 0, I only; 1, I - 2; 2, I - 3; 3, ii - 3 or 4; 4, irregular; 5, none	<i>Glyptocrinus</i>	01120	00011	12011	01214	11111	00
16. Fixed pinnules: 0, no; 1, yes	<i>Gustabilicrinus</i>	00411	11100	18011	11214	11111	20
17. Brachials (most advanced state, typically most distal): 0, rectangular uniserial; 1, cuneate uniserial; 2, biserial	<i>Hercocrinus</i>	00611	18100	05054	12101	?1111	20
18. Highest fixed brachials: 0, primibrachials; 1, secundibrachials; 2, tertibrachials; 3, quartibrachials; 4, sexibrachials	<i>Neoarchaeocrinus</i>	00410	01???	0?011	02111	01111	00
19. Intrabrachials: 0, no; 1, yes	<i>Paradiaboloocrinus</i>	00711	00000	0?023	0?111	??1??	??
20. Erect arm number: 0, 5; 1, 10; 2, 10–15; 3, 15–20; 4, 20; 5, 85–90	<i>Pararchaeocrinus</i>	00211	00011	24022	12111	01112	00
21. Arm branching: 0, isotomous; 1, atomous; 2, isotomous-heterotomous; 3, 1 isotomous	<i>Periglyptocrinus</i>	01120	00011	12011	02214	11111	10
22. Free-arm character: 0, nonpinnulate; 1, pinnulate	<i>Proexenocrinus</i>	00411	09000	08011	01101	11111	?1
23. Number of primibrachials: 0, 7–11; 1, 2; 2, 4–5	<i>Pycnocrinus</i>	01420	00011	12011	11111	01111	00
24. Stem construction: 0, pentameric; 1, holomeric	<i>Reteocrinus</i>	00410	09011	06164	00101	00201	30
25. Columnal shape: 0, pentalobate; 1, circular; 2, pentagonal	<i>Rhaphanocrinus</i>	00410	000??	02021	00111	11111	10
26. Lumen shape: 0, pentalobate; 1, circular; 2, pentagonal; 3, complex pentalobate	<i>Schizocrinus</i>	01420	01???	1?011	01101	01211	20
27. Column (based on proximal part): 0, heteromorphic; 1, homeomorphic	<i>Simplococrinus</i>	00410	00000	03021	12214	111?1	??
	<i>Trichinocrinus</i>	00311	19011	25001	01101	11112	20

## Appendix D. Characters, character states, and character matrix for Arenig to Caradoc disparids for PAUP character analyses cited herein.

	CHARACTER MATRIX FOR DISPARIDS						
1. Radial plate in A ray: 0, yes; 1, no	<i>Acolocrinus</i>	00000	40023	01011	11400	10611	?1014 0
2. Radial plate in B ray: 0, yes; 1, no	<i>Apodasmocrinus</i>	10010	30003	10021	01000	7020(01)	00002 0
3. Radial plate in C ray: 0, yes; 1, no	<i>Calceocrinus</i>	10110	21011	11022	00000	00211	21102 0
4. Radial plate in D ray: 0, yes; 1, no	<i>Caleidocrinus (C.)</i>	11011	20002	10122	00000	0021(12)	12131 0
5. Radial plate in E ray: 0, yes; 1, no	<i>Caleidocrinus (H.)</i>	11011	10003	10021	00000	0021(12)	12132 0
6. Radial or infrabasal facets: 0, fixed; 1, angustary; 2, penepenary; 3, plenary; 4, multiple small; 5, none	<i>Cataraquicrinus</i>	10010	30000	10021	00000	00(04)02	0001? 0
7. Articulation between lintels and infrabasals: 0, absent; 1, present	<i>Cincinnatiocrinus</i>	11010	30000	10021	00000	70402	22011 1
8. Number of infrabasal plates: 0, 5; 1, 4; 2, 3; 3, 0	<i>Columbicrinus</i>	11010	30003	10021	01000	30(23)(01)(12)	00013 0
9. Number of lintels: 0, 5; 1, 4; 2, 3; 3, 0	<i>Cornucrinus</i>	11011	50010	100?2	02312	20711	01124 0
10. Aboral cup shape: 0, medium cone; 1, cylindrical; 2, low cone; 3, medium bowl; 4, high cone; 5, medium globe	<i>Cremacrinus</i>	10010	21011	11021	00100	60211	20102 0
11. Radial cirlet interrupted: 0, yes; 1, no	<i>Daedalocrinus</i>	10010	30002	10021	00000	50(34)00	00012 0
12. Basal concavity: 0, no; 1, yes	<i>Difficilicrinus</i>	10110	30002	10011	0?000	???00	0?03? 0
13. Lintels visible: 0, yes; 1, no	<i>Doliocrinus</i>	11010	30000	10021	0?000	?0?01	20011 0
14. Anal X in contact with: 0, CD basal; 1, C+D infrabasal; 2, C radial; 3, primibrachial 1; 4, none	<i>Ectenocrinus</i>	10010	30000	10021	01000	70231	00012 1
15. Posterior interradius plating: 0, iRA, sRA, X; 1, X only; 2, none	<i>Eustenocrinus</i>	00000	30000	10022	00100	01001	20110 1
16. Respiratory structure: 0, no; 1, yes	<i>Geraocrinus</i>	11010	(23)0002	10021	00000	40(12)01	00003 0
17. Brachials (most advanced state): 0, rectangular uniserial; 1, cuneate uniserial; 2, none	<i>Glaucoocrinus</i>	11010	10003	10021	00000	70100	10102 0
18. Erect arm number: 0, 5; 1, 4; 2, 3; 3, 0; 4, 40+	<i>Hoplocrinus</i>	11011	10003	10011	00000	10611	0?014 0
19. Arm habit: 0, erect; 1, recumbent	<i>Hybocrinus</i>	11011	10000	10011	00000	10611	01014 0
20. Recumbent arm number: 0, 0; 1, 2; 2, 3	<i>Hybocystites</i>	11011	10005	10011	002(01)3	10611	01034 0
21. Arm branching: 0, isotomous; 1, atomous; 2, none; 3, 1 isotomous-pinnulate; 4, endotomous pinnulation; 5, endotomous ramulate; 6, bilateral heterotomy; 7, 1 isotomous-ramulate; 8, 1 isotomous-heterotomous	<i>Ibexocrinus</i>	10010	30000	10021	00000	80(0234)01	01000 1
22. Brachials fixed between adjacent proximal rays: 0, no; 1, yes	<i>Inyocrinus</i>	00000	20000	?00??	00000	00011	?1?00 0
23. Number of primibrachials: 0, 7-11; 1, 1; 2, 2; 3, 3; 4, 4-6; 5, >11; 6, all; 7, none	<i>Iocrinus</i>	11011	30000	10022	00000	00(04)1(12)	20110 0
24. Stem construction: 0, pentameric; 1, holomeric; 2, tetrameric; 3, trimeric	<i>Isotomocrinus</i>	11010	30000	10021	00000	00402	20010 0
25. Columnal shape: 0, pentalobate; 1, circular; 2, pentagonal; 3, tetralobate; 4, tetragonal	<i>Maennilicrinus</i>	11111	20004	10022	00000	004??	??113 0
26. lumen shape: 0, pentalobate; 1, circular; 2, pentagonal; 3, tetralobate; 4, tetragonal; 5, elliptical	<i>Ohiocrinus</i>	11010	30000	10021	00000	80412	?0001 0
27. Column (based on proximal part): 0, heteromorphic; 1, homeomorphic; 2, xenomorphic	<i>Othneiocrinus</i>	11010	(23)0000	10032	01000	30(24)01	2?103 0
28. Proximal anal on D radial: 0, yes; 1, no.	<i>Paracremacrinus</i>	11110	21011	11021	01100	60211	?0132 0
29. Lintel cirlet width to height ratio: 0, 2.2 to 2.8; 1, 1.1 to 2.0; 3, 0.5; 4, 3.0 to 5.0.	<i>"Pariocrinus"</i>	11011	20000	10022	00000	00301	20110 1
30. Secundibrachial number: 0, many; 1, 4-5; 2, 2-3; 3, all; 4, none.	<i>Peltacrinus</i>	11011	20000	10021	00000	00(234)00	20131 0
31. Aboral cup higher than wide: 0, no; 1, yes.	<i>Penicillicrinus</i>	10010	30000	10021	00000	002?1	?0011 0
	<i>Peniculocrinus</i>	00000	30000	10032	00000	01(04)12	2000? 1
	<i>Pogonipocrinus</i>	00000	30000	?00??	00000	00212	?0?00 0
	<i>Praecursoricrinus</i>	11010	30000	10021	01000	3021(12)	00013 0
	<i>Pultivocrinus</i>	11111	30000	10022	0?000	?0???	??11? 0
	<i>Ramseyocrinus</i>	11111	30012	1?142	00000	00523	3010? 1
	<i>Revalocrinus</i>	11011	10000	10022	00000	?0???	??0?? ?
	<i>Ristnacrinus</i>	00000	30030	10021	00000	01(12)11	500?1 0
	<i>Tetragonocrinus</i>	11111	30312	10042	00200	00(234)24	4011? 1
	<i>Tornatilicrinus</i>	11011	30000	10021	00000	7000(12)	20100 0
	<i>Tripatocrinus</i>	11011	50005	00001	02312	107??	??014 0
	<i>Tryssocrinus</i>	11010	?0000	00011	00000	714(01)(01)	00012 1
	<i>Tunguskocrinus</i>	10010	30003	10021	00000	001?1	1?013 0
	<i>Virucrinus</i>	10000	30100	10021	00100	0000(01)	20113 1
	<i>Vosekocrinus</i>	11111	10003	?00??	00000	002?0	??132 0