

SUPPLEMENT TO ECHINODERMATA ARTICLES 8-10

RAYMOND C. MOORE, RUSSELL M. JEFFORDS, and THEO. H. MILLER

COLLECTING LOCALITIES

Localities are designated by assigned letters (e.g., Ogz, as recorded in lists of the University of Kansas Paleontological Institute—UKPI) or by letter and numbers (e.g., E 733, recorded in lists of the Esso Production Research Company, Houston). The localities are arranged in groups according to the age of the fossil-bearing strata, beginning with Ordovician (oldest represented) and proceeding upward in stratigraphic succession.

ORDOVICIAN

- | LOC. | DESCRIPTION |
|------|---|
| Ocb | <i>Fernvale Formation, Richmond Group, Cincinnati, Upper Ordovician</i> ; near Decaturville, Camden County, Missouri. Moore coll., Univ. Kansas Paleont. Inst. |
| Ogg | <i>Catheys Limestone, Trentonian, Middle Ordovician</i> ; bottom of Reservoir Hill in Nashville, Davidson County, Tennessee. Moore coll., Univ. Kansas Paleont. Inst. |
| Ogq | <i>Blanchester Formation (Waynesville), Richmond Group, Cincinnati, Upper Ordovician</i> ; near Clarks-ville, Clinton County, Ohio, U. S. Natl. Museum. |
| Ogr | <i>Maquoketa Shale, Richmond Group, Cincinnati, Upper Ordovician</i> ; near Spring Valley, Fillmore County, Minnesota, U. S. Natl. Museum. |
| Ogv | <i>Wilmington Limestone, Richmond Group, Cincinnati, Upper Ordovician</i> ; near Savannah, Carroll County, Illinois. U. S. Natl. Museum. |
| Ogw | <i>Fernvale Limestone, Richmond Group, Cincinnati, Upper Ordovician</i> ; near Cape Girardeau, Cape Girardeau County, Missouri. Ulrich coll., U. S. Natl. Museum. |
| Ogx | <i>Wilmington Limestone, Richmond Group, Cincinnati, Upper Ordovician</i> ; near Wilmington, Wills County, Illinois, U. S. Natl. Museum. |
| Ogz | <i>Cincinnati, Upper Ordovician</i> ; unspecified locality in Cincinnati, Hamilton County, Ohio. O. C. Marsh coll., Yale Univ. Peabody Museum 17110. |
| Oha | <i>Cincinnati, Upper Ordovician</i> ; Cincinnati, Hamilton County, Ohio. U. S. Natl. Museum. |
| Ohg | <i>Maquoketa Shale, Cincinnati, Upper Ordovician</i> ; near Clermont, Fayette County, Iowa. Slocom coll. |

SILURIAN

- | LOC. | DESCRIPTION |
|------|--|
| Sad | <i>Waldron Clay Member, Waynesville Formation, Niagaran, Upper Silurian</i> ; quarry of Lambert Bros. Stone Company 1.5 miles west-southwest of junction of U. S. 70N and 70S 11 miles west-southwest of center of Nashville, Davidson County, Tennessee, Moore and Jeffords coll., Univ. Kansas Paleont. Inst.—Esso Prod. Research Co. |
| Sdg | <i>Brownsport Formation, Niagaran, Upper Silurian</i> ; Blue Mound Glade (Amsden, 1949, p. 38, loc. 18) 0.8 mile northeast of Cedar Grove Church and 0.2 mile north of road to Mt. Carmel Church, Perryville quad-range, near Decaturville, Decatur County, Tennessee. Moore and Jeffords coll., Univ. Kansas Paleont. Inst.—Esso Prod. Research Co. |
| Sfr | <i>Waldron Shale, Niagaran, Upper Silurian</i> ; south mouth of tunnel at Tunnel Mill, about 0.1 mile west of Ind. 3, 0.5 mile south of Vernon, Jennings County, Indiana (SE $\frac{1}{4}$ sec. 11, T. 6 S., R. 8 E.). Moore, Jeffords, and Hattin coll., Univ. Kansas Paleont. Inst.—Esso Prod. Research Co. |
| Shd | <i>Waldron Shale, Niagaran, Upper Silurian</i> ; on county road 2.5 miles east of Dupont, Jefferson County, Indiana. Moore and Jeffords coll., Univ. Kansas Paleont. Inst.—Esso Prod. Research Co. |
| | county road 2.5 miles east of Dupont, Jefferson County, Indiana. Moore and Jeffords coll., Univ. Kansas Paleont. Inst.—Esso Prod. Research Co. |
| Shg | <i>Waldron Shale, Niagaran, Upper Silurian</i> ; pit in west part of Waldron, Shelby County, Indiana. Moore and Jeffords coll., Univ. Kansas Paleont. Inst.—Esso Prod. Research Co. |

DEVONIAN

- | LOC. | DESCRIPTION |
|------|---|
| Ddv | <i>Rockport Limestone (lower part), Traverse Group, Erian, Middle Devonian</i> ; quarry at Rockport, Alpena County, Michigan, Laudon coll., Esso Prod. Research Co. |
| Ddx | <i>Wanakah Shale Member, Ludlowville Formation, Hamilton Group, Erian, Middle Devonian</i> ; on 18-mile Creek near bridge on old Lakeshore Road near Lake Erie southwest of Buffalo, Erie County, New York. Jeffords coll., Esso Prod. Research Co. |
| Dh | <i>Birdsong Shale, Linden Group, Helderbergian, Lower Devonian</i> ; along Turkey Creek and valley wall on old Love farm east of Ky. 69 about 9.5 miles south of underpass at south edge of Camden, Benton County, Tennessee. Moore and Jeffords coll., Univ. Kansas Paleont. Inst.—Esso Prod. Research Co. |
| Di | <i>Birdsong Shale, Linden Group, Helderbergian, Lower Devonian</i> ; road cut on Tenn. 13 about 4 miles north of Linden, Perry County, Tennessee. Moore and Jeffords coll., Univ. Kansas Paleont. Inst.—Esso Prod. Research Co. |
| Dia | <i>Wanakah Shale Member, Ludlowville Formation, Hamilton Group, Erian, Middle Devonian</i> ; shale pit on Big Tree road about 0.5 mile southeast of N. Y. 5 near Bay View, about 10 miles south of downtown Buffalo, Erie County, New York. W. F. Berry coll., Univ. Kansas Paleont. Inst. |
| Dib | <i>Wanakah Shale Member (upper part), Ludlowville Formation, Hamilton Group, Erian, Middle Devonian</i> ; cut on N. Y. 63 about 2 miles southeast of East Bethany, Genesee County, New York. W. F. Berry coll., Univ. Kansas Paleont. Inst. |
| Did | <i>Centerfield Limestone, Darien Member, Ludlowville Formation, Erian, Middle Devonian</i> ; 2 miles northeast of Darien, Genesee County, New York, W. F. Berry coll., Univ. Kansas Paleont. Inst. |
| Dii | <i>Birdsong Shale, Linden Group, Helderbergian, Lower Devonian</i> ; near Swaynes Mill, Benton County, Tennessee. Springer coll., U. S. Natl. Museum. |

MISSISSIPPIAN

- | LOC. | DESCRIPTION |
|------|---|
| Ma | <i>New Providence Shale, Borden Group, Osagian, Lower Mississippian</i> ; pit of Coral Ridge Brick and Tile Company on South Park road 0.3 mile southeast of junction with old National Turnpike, 1.1 miles southeast of Coral Ridge and about 3 miles west of Kentucky Turnpike, also 1.2 miles northwest of summit of Button Mould Knob, about 15 miles south of downtown Louisville, Bullitt County, Kentucky. Moore and Jeffords coll., Univ. Kansas Paleont. Inst.—Esso Prod. Research Co. |
| Maa | <i>New Providence Shale, Borden Group, Osagian, Lower Mississippian</i> ; northwest flank of Button Mould |

- | | |
|-----|--|
| | Knob, 2.3 miles southeast of Coral Ridge and about 15 miles south of downtown Louisville, Bullitt County, Kentucky. Moore and Jeffords coll., Univ. Kansas Paleont. Inst.—Esso Prod. Research Co. |
| Mc. | <i>Brodhead Formation, Borden Group, Osagian, Lower Mississippian</i> ; hill side at northwest edge of Knob Creek Church on Ky. 44 about 2.7 miles west of Shepherdsville, Bullitt County, Kentucky. Moore and Jeffords coll., Univ. Kansas Paleont. Inst.—Esso Prod. Research Co. |
| Md | <i>Brodhead Formation, Borden Group, Osagian, Lower Mississippian</i> ; north side of U. S. 62 at point 0.75 west of junction with Kentucky 583 east of Elizabethtown, Hardin County, Kentucky. Moore and Jeffords coll., Univ. Kansas Paleont. Inst.—Esso Prod. Research Co. |
| Mdp | <i>"Fort Payne Limestone" (Borden Group equivalent), Osagian, Lower Mississippian</i> ; weathered road cut along Eagle Creek, 8 miles northeast of Livingston, Overton County, Tennessee. Moore and Jeffords coll., Univ. Kansas Paleont. Inst.—Esso Prod. Research Co. |
| Me | <i>Floyds Knob Formation, Borden Group, Osagian, Lower Mississippian</i> ; large cut on U. S. 31W on Muldraugh Hill, southeast of Muldraugh, Bullitt County, Kentucky. Moore and Jeffords coll., Univ. Kansas Paleont. Inst.—Esso Prod. Research Co. |
| Mec | <i>Burlington Limestone, Osagian, Lower Mississippian</i> ; in and near Burlington, Des Moines County, Iowa. Springer coll., U. S. Natl. Museum. |
| Mey | <i>Paint Creek Formation, Chesteran, Upper Mississippian</i> ; along creek near Vogel School, SE $\frac{1}{4}$ SW $\frac{1}{4}$ sec. 3, T. 2 S., R. 9 W., about 1.5 miles west of Floraville, St. Clair County, Illinois. Moore and Van Sant coll., Univ. Kansas Paleont. Inst.—Esso Prod. Research Co. |
| Mfm | <i>Edwardsville Formation ("Crawfordsville beds"), Borden Group (middle part), Osagian, Lower Mississippian</i> ; Montgomery County Poor Farm in NE cen. SW $\frac{1}{4}$ sec. 29, T. 19 N., R. 4 W., northeast of Crawfordsville, Indiana, along creek bank below hog fields. Moore, Jeffords, and Gary Lane coll., Univ. Kansas Paleont. Inst.—Esso Prod. Research Co. |
| Mfn | <i>Edwardsville Formation, Borden Group, Osagian, Lower Mississippian</i> ; cut on north side of Indiana 46 approximately 6 miles east of Bloomington, Monroe County, Indiana ("Stobo bioherm," in sec. 4, T. 84 N., R. 1 E.). Moore, Jeffords, and Hattin coll., Univ. Kansas Paleont. Inst.—Esso Prod. Research Co. |

PENNSYLVANIAN

- | LOC. | DESCRIPTION |
|------|---|
| Pak | <i>Mingus Shale Member, Garner Formation, Strawn Group, Desmoinesian, Middle Pennsylvanian</i> ; cut on south side of U. S. 80 just east of Eastland-Erath County line, 3.5 miles west of junction with Texas |

- 108, Erath County, Texas. Moore and Jeffords coll., Univ. Kansas Paleont. Inst.—Esso Prod. Research Co.
- Paw *Chaffin Limestone Member, Thrifty Formation, Cisco Group, Virgilian, Upper Pennsylvanian*; north side of FM 1850 1.8 miles west of Texas 279, Brown County, Texas. Moore and Jeffords coll., Univ. Kansas Paleont. Inst.—Esso Prod. Research Co.
- Pba *South Bend Shale Member, Graham Formation, Cisco Group, Virgilian, Upper Pennsylvanian*; side of stock tank at northeast base of prominent sandstone-capped hill in southwest part of South Bend, east of FM 701 with Main Street, Young County, Texas. Moore and Jeffords coll., Univ. Kansas Paleont. Inst.—Esso Prod. Research Co.
- Pbd *Blach Ranch Limestone Member, Thrifty Formation, Cisco Group, Virgilian, Upper Pennsylvanian* (fossil bed "B" called Breckenridge Limestone by Roth and Kauffman); stock tank dam 5.6 miles northwest of U. S. 281 junction with Texas 199, about 12 miles northwest of Jacksboro, Jack County, Texas. Moore, Jeffords, and Roth coll., Univ. Kansas Paleont. Inst.—Esso Prod. Research Co.
- Pca *Pumpkin Creek Limestone Member, Dornick Hills Formation, Atokan, Middle Pennsylvanian*; on Daube Ranch road about 4.75 miles east of Baum, near center sec. 32, T. 3 S., R. 4 E., Johnston County, Oklahoma. Moore and Jeffords coll., Univ. Kansas Paleont. Inst.—Esso Prod. Research Co.
- Pdq *Belknap Limestone Member, Thrifty Formation, Cisco Group, Virgilian, Upper Pennsylvanian*; on FM 578 and U. S. 183 at point 6.1 miles north of Birch Hotel in Breckenridge, Stephens County, Texas. Jeffords coll., Univ. Kansas Paleont. Inst.—Esso Prod. Research Co.
- Pef *Coffeyville Formation, Missourian, Upper Pennsylvanian*; old shale pit in west part of Coffeyville, Montgomery County, Kansas. Moore coll., Univ. Kansas Paleont. Inst.
- Peg *Marble Falls Limestone, Atokan, Middle Pennsylvanian*; near Turkey Roost Creek, Jym Sloan ranch, 1¼ miles south of Maxwell Crossing on San Saba River, 10 miles south of Richland Springs and southwest of San Saba, San Saba County, Texas. Moore coll., Univ. Kansas Paleont. Inst.
- Phu *Caprock of Mineral coal near middle of Cabaniss Formation, Cherokee Group, Desmoinesian, Middle Pennsylvanian*; strip pits about 5 miles west of Pittsburg, Crawford County, Kansas, Moore coll., Univ. Kansas Paleont. Inst.
- Piw *Minturn Formation, Desmoinesian, Middle Pennsylvanian*; near McCoy, Eagle County, Colorado. Univ. Kansas Paleont. Inst.
- Pr *Millsap Lake Formation, Strawn Group, Desmoinesian, Middle Pennsylvanian*; cattle tank 250 feet west of fence located 0.3 mile north of ranch road at point 0.75 mile south of U. S. 190, which is joined by ranch road 2.7 miles east of railroad overpass at Rochelle, McCulloch County, Texas. Moore and Jeffords coll., Univ. Kansas Paleont. Inst.—Esso Prod. Research Co.
- Pt *Wayland Shale Member, Graham Formation, Cisco Group, Virgilian, Upper Pennsylvanian*; scarp about 50 yards south of road at point 1.25 miles east of Fife, McCulloch County, Texas. Moore and Jeffords coll., Univ. Kansas Paleont. Inst.—Esso Prod. Research Co. [=loc. E 5].
- E 5 *Wayland Shale Member, Graham Formation, Cisco Group, Virgilian, Upper Pennsylvanian*; scarp about 50 yards south of road at point 1.25 miles east of Fife (or east of U.S. 283 road junction about 3 miles south of Colorado River), McCulloch County, Texas [=loc. Pt of Moore & Jeffords, 1967]. Collectors: R. M. Jeffords, R. G. Todd, R. C. Moore, T. H. Miller, and E. K. Martin, 1958-66.
- E 672 *Wayland Shale Member (upper part), Graham Formation, Cisco Group, Virgilian, Upper Pennsylvanian*; stock-tank area just west of road, 2 miles southeast of Thrifty; locality is reached by going 1.1 miles southeast of Thrifty on F.M. 2492 and turning south on gravel road (U. S. Geol. Survey loc. 16020), Brown County, Texas. Collectors: R. M. Jeffords, T. H. Miller, and E. K. Martin, 1963-64.
- E 675 *Wayland Shale Member, Graham Formation, Cisco Group, Virgilian, Upper Pennsylvanian*; unrecorded locality in McCulloch County, Texas. Collector: E. K. Martin, 1963.
- E 682 *Wayland Shale Member, Graham Formation, Cisco Group, Virgilian, Upper Pennsylvanian*; one mile due south of Gunsight, Stephens County, on Perry Farm. Follow road south, taking west fork at 0.9 mile and park at wire gate 0.3 mile from road fork. Exposure is in second scarp to east. Collectors: R. C. Moore and R. M. Jeffords, 1962.
- E 705 *Wayland Shale Member, Graham Formation, Cisco Group, Virgilian, Upper Pennsylvanian*; scarp north of stock tank on Gary ranch which is east of U. S. 283 at a point 1.7 miles north of junction with FM 504, McCulloch County. Drive past ranch house and barn on caliche road to stock tank; walk north behind tank approximately 50 yards to shale scarp. Collectors: R. M. Jeffords, T. H. Miller, and E. K. Martin, 1964.
- E 706 *Gunsight Limestone*. Flat below scarp on Cornill Ranch near locality 705. From locality 705, walk north 50 yards, cross fence to Cornill Ranch, and go 25 yards to northeast. Collectors: R. M. Jeffords, T. H. Miller, and E. K. Martin, 1964.
- E 722 Same as locality 672.
- E 732 *Wayland Shale Member, Graham Formation, Cisco Group, Virgilian, Upper Pennsylvanian*; along bluff west of road and in road ditch, about 4 miles south of Whon, Coleman County. Collectors: R. M. Jeffords and T. H. Miller, 1964.

- E 733 *Wayland Shale Member, Graham Formation, Cisco Group, Virgilian, Upper Pennsylvanian*; base of road cut and in farm road to north, north side of U. S. 80, 4.15 miles west of Eastland, Eastland County. Collectors: R. C. Moore and R. M. Jeffords, 1962.
- E 734 *Wayland Shale (upper part)*; bluff 400 yards west of road, 4.5 miles south of Whon, northeast of Parks Mountain, County. Collectors: R. M. Jeffords and T. H. Miller, 1964.
- E 790 *Wayland Shale*; scarp on Cornill Ranch just north of localities 705 and 706, McCulloch County. From locality 706, go about 300 yards north past second stock tank to prominent scarp. Collectors: T. H. Miller and E. K. Martin, 1964.

PERMIAN

- | | |
|------|-------------|
| LOC. | DESCRIPTION |
|------|-------------|
- Qry *Waldrip Shale Member (Waldrip Limestone 3), Pueblo Formation, Wichita Group, Wolfcampian, Lower Permian*; cut on FM 765 about 350 yards west of bridge over Saddle Creek, about 7 miles west of Fife, McCulloch County, Texas. Moore and Jeffords coll., Univ. Kansas Paleont. Inst.—Esso Prod. Research Co.

CRETACEOUS

- | | |
|------|-------------|
| LOC. | DESCRIPTION |
|------|-------------|
- Kix *San Felipe Formation (upper part), Senonian, Upper Cretaceous*; Las Flores and Chocoy haciendas, about 65 km. northwest of Tampico, Tamaulipas, Mexico. Springer coll., U. S. Natl. Museum.
- Kja *Prairie Bluff Chalk, Maastrichtian, Upper Cretaceous*; along Josey Creek about 4 miles northwest of Starkville, Oktibbeha County, Mississippi. P. H. Dunn coll., Univ. Kansas Paleont. Inst.
- Kjc *Ripley Formation, Maastrichtian, Upper Cretaceous*; near Livingston, Sumter County, Alabama. Cincinnati Soc. Nat. History coll.
- Kji *Prairie Bluff Chalk, Maastrichtian, Upper Cretaceous*; at roadside and cut on Houston Road (sec. 10, T. 15 S., R. 3 E.), 1.25 miles north of Sparta, Chickasaw County, Mississippi, (U. S. Geol. Survey loc. 17235). L. W. Stephenson and W. H. Monroe coll., U.S. Natl. Museum.

PALEOGENE-NEOGENE

- | | |
|------|-------------|
| LOC. | DESCRIPTION |
|------|-------------|
- Aig *Universidad Formation, Lower Eocene (upper part)*; 2 km. northwest of Radas, Las Villas Province, Cuba. Bermúdez coll., Univ. Kansas Paleont. Inst.

REFERENCES

Given here is a consolidated list of references to literature cited in one or more of the preceding articles which are issued together in this publication. Repetition thus is avoided.

- AGASSIZ, J. L. R., 1835 [1836], *Prodrome d'une monographie des radiaires ou echinodermes*: Soc. Sci. Nat. Neuchâtel, Mém., v. 1, p. 168-199.
- ANGELIN, N. P., 1878, *Iconographia crinoideorum in stratis Sueciae Siluricis fossilium*: Samson & Wallin, Stockholm, Holmiae, 62 p., 29 pl.
- ARENDT, YU. A., & GEKKER [HECKER], R. F., 1964 [1965], *Klass Crinoidea, Morskije lilii, Sistematicheskaya chasty* [Class Crinoidea, sea lilies, systematic part], in Orlov, Yu. A. (ed.), *Osnovy paleontologii, Iglokoynhie, Gemikhordovye, Pogonofory, i Shchetinkochelyustnye* [Treatise on paleontology, echinoderms, hemichordates, pogonophores, chaetognaths]: p. 80-105, fig. 115-140, pl. 8-16.
- AUSTIN, THOMAS, 1875, *Observations on the genus Platycrinus*: Annals and Mag. Nat. History, ser. 4, v. 16, p. 90-91.
- , & AUSTIN, THOMAS, JR., 1842, XVII. *Proposed arrangement of the Echinodermata, particularly as regards the Crinoidea, and a subdivision of the class Adelostella (Echinidae)*: Annals and Mag. Nat. History, ser. 1, v. 10, no. 63, p. 106-113.
- , & ——, 1843, XXXIII. *Description of several new genera and species of Crinoidea*: Annals and Mag. Nat. History, ser. 1, v. 11, no. 69, p. 195-207.
- AVROV, D. P., & STUKALINA, G. A., 1964, *Novye dannye o siluriyskikh oilozheniyakh yuga gornogo Altaya* [New data on Silurian deposits in southern Altai range], in *Materialy po geologii i poleznykh iskopayemykh Altaya i Kazakhstana* [Information on the geology and useful fossils of Altai and Kazakhstan]: Vses. Nauchno-Issled. Geol. Inst., Trudy [VSEGEI], v. 3, p. 25-29, fig. 1.
- BASSLER, R. S., 1938, *Pelmatozoa Palaeozoica*, in Quenstedt, Werner (ed.), *Fossilium catalogus, I, Animalia*, pt. 83: W. Junk, s'Gravenhage, 194 p.
- , & MOODEY, M. W., 1943, *Bibliographic and faunal index of Paleozoic pelmatozoan echinoderms*: Geol. Soc. America, Spec. Paper 45, 734 p.
- BATHER, F. A., 1893, *The Crinoidea of Gotland, Part 1, The Crinoidea Inadunata*: K. Svenska Vetenskapsakad., Handl., v. 25, no. 2, 182 p., 10 pl.
- , 1895, *On Uintacrinus, a morphological study*: Zool. Soc. London, Proc. 1895, p. 974-1004, pl. 54-56.

- , 1898, *Petalocrinus (Weller and Davidson)*: Geol. Soc. London, Quart. Jour., v. 54, p. 401-441, 15 fig.
- , 1898-99, *Wachsmuth and Springer's monograph on crinoids*: Geol. Mag., n. ser., dec. 4, v. 5 (1898), 1st notice, p. 276-283; 2nd notice, p. 318-329; 3rd notice, p. 419-428; 4th notice, p. 522-527; v. 6 (1899), 5th notice, p. 32-44; 6th notice, p. 117-127.
- , 1899, *A phylogenetic classification of the Pelmatozoa*: British Assoc. Rept. for 1898, p. 916-822.
- , 1900, *The Echinoderma*, in Lankester, E. R. (ed.), *A treatise on zoology*, pt. 3, Adam & Charles Black, London, 344 p., 309 fig. [Crinoidea, p. 94-204, fig. 1-127].
- , 1909, *Triassic echinoderms of Bakony*: Wiss. Erforsch. Balatonsees Result., Paläont. Anhang, v. 1, pt. 1, p. 24-30.
- BERRY, W. F., (1952), *A descriptive study of crinoid columnals from the Hamilton Group of New York*: Univ. Massachusetts, unpubl. M. A. dissert.
- BIESE, WALTER, 1927, *Ueber die Encriniten der unteren Muschelkalkes von Mitteleuropa*: Preuss. geol. Landesanstalt, Abh., new ser., no. 103, 119 p., 6 fig., 4 pl.
- , 1934, *Crinoidea triadica*: in Quenstedt, Werner (ed.), *Fossilium catalogus*, I, Animalia, pt. 66; W. Junk, s'Gravenhage, 255 p.
- , 1935-37, *Crinoidea jurassica I*: in Quenstedt, Werner (ed.), *Fossilium catalogus*, I, Animalia, pts. 70, 73, 76; W. Junk, s'Gravenhage, 739 p.
- , & SIEVERTS-DORECK, HERTHA, 1937, *Crinoidea cretacea*: in Quenstedt, Werner (ed.), *Fossilium catalogus*, I, Animalia, pt. 77; W. Junk, s'Gravenhage, 254 p.
- , & ———, 1939a, *Crinoidea caenozoica*: in Quenstedt, Werner (ed.), *Fossilium catalogus*, I, Animalia, pt. 80; W. Junk, s'Gravenhage, 151 p.
- , 1939b, *Supplementum ad Crinoidea triadica, jurassica, cretacea et caenozoica*: in Quenstedt, Werner (ed.), *Fossilium catalogus*, I, Animalia, pt. 88; W. Junk, s'Gravenhage, 81 p.
- BLAINVILLE, H. M. D. DE, 1834-37, *Manuel d'actinologie et de zoophytologie*: F. G. Levrault, Paris, 695 p., 100 pl.
- BLUMENBACH, J. F., 1802-04, *Abbildungen naturhistorischer Gegenstände*: Göttingen, H. 7, no. 70, 4 p., 1 pl.
- BREIMER, ALBERT, 1962, *A monograph on Spanish Palaeozoic Crinoidea*: Leidse Geol. Mededeel., v. 27, 89 p., 39 fig., 16 pl.
- BRONN, H. G., 1840, *Ctenocrinus, ein neues Krinoiden-Geschlecht der Grauwacke*: Neues Jahrb. Mineralogie, Geologie u. Palaeontologie, p. 542-548, pl. 8B.
- , 1848, *Index palaeontologicus, unter Mitwirkung der Herren Prof. H. R. Göppert und H. von Meyer*: Handbuch einer Geschichte der Natur, Stuttgart, v. 5, Abt. 1, no. 1, 2, pt. 3, A. Nomenclator palaeontologicus; A-M, p. 1-775, N-Z, p. 776-1381.
- BUCKLAND, WILLIAM, 1837, *Geology and mineralogy; the Bridgewater treatise*: London, v. 6, pt. 1, 599 p.; pt. 2, 110 p., 86 pl.
- BULLARD, F. M., & CUYLER, R. H., 1936, *The Upper Pennsylvanian and Lower Permian section of the Colorado River valley, Texas*: Texas Univ., Bull. 3501, p. 191-258, 2 fig., pl. 4.
- BYRNE, F. E., & SEEBERGER, EVELYN, 1942, *Fragmentary crinoids from the Lower Permian of the Manhattan area [Kansas]*: Kansas Acad. Sci., Trans., v. 45, p. 225-228, pl. 1.
- CARPENTER, P. H., 1884, *Report upon the Crinoidea collected during the voyage of H.M.S. Challenger, during the years 1873-1876. Pt. I—General morphology, with descriptions of the stalked crinoids*: Rept. Scientific Results Explor. Voyage H.M.S. Challenger, Zoology, v. 11, pt. 26, 442 p., 21 fig., 62 pl.
- CASEY, RAYMOND, 1960, *A Lower Cretaceous gastropod with fossilized intestines*: Palaeontology, v. 2, p. 270-276, 1 fig., pl. 41.
- CLARK, A. H., 1913, *Articulata*: in Zittel, K. A., *Textbook of paleontology*, Eastman, C. R. (ed.), Macmillan & Co., London, 2nd ed., p. 226-241, fig. 332-346.
- , 1915-47, *A monograph of the existing crinoids*: U.S. Natl. Mus., Bull. 82, v. 1, The comatulids, pt. 1 (1915), 406 p., 513 fig., 17 pl., pt. 2 (1921), 795 p., 949 fig., 57 pl., pt. 3 (1931), 816 p., 82 pl.; pt. 4a (1941), 582 p., 61 pl.; pt. 4b (1947), 459 p., 43 pl.
- CLARK, W. B., & TWITCHELL, M. W., 1915, *The Mesozoic and Cenozoic Echinodermata of the United States*: U.S. Geol. Survey, Mon. 54, 341 p., 108 pl.
- CRONEIS, CAREY, 1938, *Utilitarian classification for fragmentary fossils*: Jour. Geology, v. 46, p. 975-984.
- CUÉNOT, LUCIEN, 1948, *Classe des crinoïdes*: in Cuénot, Lucien, & Dawydoff, C., *Embranchement des échinodermes, Traité de Zoologie*, Grassé, P.-P. (ed.), v. 11, p. 30-74, fig. 38-98.
- CUMBERLAND, GEORGE, 1826, *Reliquiae conservatae, "From the primitive materials of our present globe with popular descriptions of the prominent characters of some remarkable fossil Encrinites and their connecting links"*: Bristol, 45 p., 13 pl.
- DAMES, W. B., 1885, *Petrefakten aus dem Daghestan und der Turkmenensteppe*: Deutsch. Geol. Gesell., Zeitschr., v. 37, p. 218-220.
- DAVILA, P. F., & ROMÉ DE L'ISLE, J. B. L. DE, 1767, *Catalogue systématique et raisonné des curiosités de la nature et de l'art qui composent le cabinet de M. Davila*: Paris, v. 3, 290 p., 8 pl.
- DELAGE, YVES, & HÉROUARD, E. J. E., 1903 (1904), *Traité de zoologie concrète*: Paris, v. 3, Echinodermes.
- DESOR, EDOUARD, 1847, *Résumé de ses études sur les crinoïdes fossiles de la Suisse*: Soc. Sci. Nat. Neuchâtel, Bull., v. 1, p. 211-222.

- DEVONSHIRE, P. F. P. C., 1954, *A faunal study of the Pumpkin Creek Limestone Member of the Big Branch Formation in the Ardmore [Oklahoma] area*: Shale Shaker, v. 5, p. 5-16, 21-22, 3 fig., 11 pl. [Republished with other papers from Shale Shaker, ?1962, p. 314-327].
- DÖDERLEIN, LUDWIG, 1907, *Die gestielten Crinoiden der Siboga-Expedition: Siboga Expeditie*: Uitkomsten op zoologisch, botanisch, oceanographisch en geologisch gebied, verzameld in Nederlandsch Oost-Indië 1899-1900, v. 42a, p. 1-54, 12 fig., pl. 1-23 (Leiden).
- , 1912, *Die gestielten Crinoiden der deutschen Tiefsee-Expedition*: Gustav Fischer, Jena, Wiss. Ergebnisse der deutschen Tiefsee-Exped. auf dem Dampfer "Valdivia" 1898-99, v. 17, no. 1, 34 p., 9 fig., 12 pl.
- DUBATOLOVA, YU. A., 1964, *Morskikh liliy devona Kuzbassa* [Devonian crinoids from Kuznetsk Basin]: Akad. Nauk SSSR, Sibirskoye Otdeleniye, Inst. Geologii i Geofizik, Trudy, 152 p., 16 fig., 14 pl.
- , 1967, *Devonskie krinoidei Khrebia Tas-khayaktakh (Severo-vostok SSSR)* [Devonian crinoids from Tas-khayaktakh Range (Northeastern USSR)]: Akad. Nauk SSSR, Sibirskoye Otdeleniye, Inst. Geologii i Geofiziki, Novye dannye po biostratigrafii Devona i Verkhnego Paleozoya Sibiri, p. 32-41, fig. 1-6, pl. 6.
- , & SHAO, TSZE, 1959, *Stebli morskih liliy Kamenougolnykh, Permskih, i Triasovykh otzheniy yuzhnogo Kitaya* [Carboniferous, Permian, and Triassic crinoid stems from Yuzhna Kitaya region (China)]: Acta Palaeontologica Sinica, v. 7, no. 1, p. 41-83, 4 fig., pl. 1-2.
- , & YELTSHEVA, R. S., 1960 (1961), *Paleontologicheskaya kharakteristika Devona Sarno-Altayskoy oblasti (morskije liliy)* [Paleontological characteristics of Devonian in Sarno-Altayskoy region (central Kazakhstan) (crinoids)]: in L. L. Khalfin *et al.*, Biostratigrafiya Paleozoya Sayano-Altayskoy Gornoy Oblasti, v. 2, Sredniy Paleozoy, Sibirskkogo Nauchno-Issledovatel'skogo Instituta Geologii, Geofiziki i Mineralnogo Siroya (SNIIGGIMS), Minsisterstva Geologii i Okhrany Nedr SSSR, pt. 20, 850 p., 104 fig., 133 pl. [crinoids, p. 294-296, 552-560, pl. D86-D87].
- EARGLE, D. H., 1960, *Stratigraphy of Pennsylvanian and Lower Permian rocks in Brown and Coleman Counties, Texas*: U.S. Geol. Survey, Prof. Paper 315-D, p. 55-77, 2 fig., 6 pl.
- EASTON, W. H., 1962, *Carboniferous formations and faunas of central Montana*: U.S. Geol. Survey, Prof. Paper 348, 126 p., 1 fig., 14 pl.
- EHRENBERG, KURT, 1929, *Pelmatozoan root-forms (fixation)*: Am. Mus. Nat. History, Bull., v. 59, art. 1, 76 p., 42 fig.
- EICHWALD, EDUARD D', 1840, *Ueber das silurische Schichtensysteme Estlands*: Acad. Sci. St. Petersburg, Bull., v. 7, 210 p. [In German and French.]
- , 1856, *Beitrag zur geographischen Verbeitung der fossilen Thiere Russlands*: Soc. Imp. Nat. Moscou (Moskov. Obshch. Ispytateley Prirody Byull.), v. 29, p. 88-127.
- , 1859-68, *Lethaea rossica ou paleontologie de la Russie, vol. 1, Premiere section de l'ancienne periode*: E. Schweizerbart, Stuttgart, v. 1, 681 p., 59 pl. [Atlas of plates: Ancienne periode, 59, 1859; Mayenne periode, 40, 1868; Moderne periode, 14, 1859.]
- ELTSYSHEVA, R. S. (see YELTSYSHEVA, R. S.)
- ÉTALLON, C. A., 1859, *Etudes paléontologiques sur le Haut-Jura; Rayonnés du Corallien*: Soc. Emul. Doubs, Mém., ser. 3, n. 3, p. 401-553.
- FENTON, M. A., 1929, *Notes on several forms of Lichenocrinus from Black River formations*: Am. Midland Naturalist, v. 11, p. 494-499, pl. 37.
- FISCHER DE WALDHEIM, GOTTHELF, 1811, *Recherches sur les encrinites les polycères et les ombellulaires etc.*: Notice des fossiles du Gouv. de Moscou, servant de Progr. l'invitation à la séance publique de la Soc. Imp. Moscou, 32 p., 2 pl.
- FOERSTE, A. F., 1919, *Echinodermata of the Brassfield (Silurian) formation of Ohio*: Denison Univ., Bull., Sci. Lab. Jour., v. 19, p. 3-31, 7 pl.
- GEINITZ, H. B., 1846, *Grundriss des Versteinerungskunde*: Arnold, Dresden and Leipzig, 815 p., 28 pl.
- GIRTY, G. H., 1915, *Fauna of the Wewoka Formation of Oklahoma*: U.S. Geol. Survey, Bull. 544, 335 p., 35 pl.
- GISLÉN, TORSTEN, 1924, *Echinoderm studies*: Zool. Bidrag fran Uppsala, v. 9, 330 p., 349 fig.
- , 1934, *A reconstruction problem: Analysis of fossil comatulids from North America, with a survey of all known types of comatulid arm ramifications*: Univ. Arssk. (Acta Univ. Lundensis), neue Folge, Band 30, Avd. 2 (K. Fysiog. Sällsk. Lund Handl., neue Folge, Bd. 45, Nr. 11), 59 p., 63 fig.
- GOLDFUSS, G. A., 1826-44, *Petrefacta Germaniae, tam ea, Quae in Museo Universitatis Regiae Borussiae Fridericiae Wilhelmae Rhenanae, servantur, quam alia quaecunque in Museis Hoeninghusiano Muensteriano aliisque, extant, iconibus et descriptionibus illustrata.—Abbildungen und Beschreibungen der Petrefacten Deutschlands und der Angränzenden Lander, unter Mitwirkung des Herrn Grafen Georg zu Münster, herausgegeben von August Goldfuss*: Arnz & Co., Düsseldorf, v. 1 (1826-33), Divisio prima, Zoophytorum reliquiae (p. 1-114); Divisio secunda, Radiariorum reliquiae (p. 115-221) [Echinodermata]; Divisio tertia, Annulatorium reliquiae (p. 222-242); v. 2 (1834-40), Divisio quarta, Molluscorum acephaliorum reliquiae. I. Bivalvia (p. 65-286); II. Brachiopoda (p. 287-303); v. 3 (1841-44); Divisio quinta,

- Molluscorum gasteropodum reliquiae (p. 1-121); atlas of plates (1-199).
- , 1839, *Beiträge zur Petrefaktenkunde*: K. Leopold. Carolin. Akad. Naturf., Verh., v. 19, p. 329-364, pl. 30-33.
- GOLDRING, WINIFRED, 1923, *The Devonian crinoids of the state of New York*: New York State Mus., Mem. 16, 670 p., 60 pl.
- , 1935, *New and previously known Middle Devonian crinoids of New York*: Carnegie Mus. Annals, v. 24, p. 349-368, pl. 25-27.
- , 1936, *Some Hamilton (Devonian) crinoids from New York*: Jour. Paleont., v. 10, p. 14-22, 1 fig., pl. 6-7.
- , 1942, *Crown of Ancyrocrinus bulbosus Hall*: Buffalo Soc. Nat. Sci., Bull., v. 17, p. 13-18, 2 fig., pl. 4.
- GRANT, R. E., 1963, *Unusual attachment of a Permian linoproductid brachiopod*: Jour. Paleont., v. 37, p. 134-140, 1 fig., pl. 19.
- GREGORIO, ANTONIO DE, 1930, *Sul Permiano di Sicilia*: Annales Géologie et Paléontologie, v. 52, p. 1-70, pl. 1-21.
- HAECKEL, E. H. P. A., 1896, *Die Amphorideen und Cystoideen; Beiträge zur morphologie und Phylogenie der Echinodermen*: W. Engelmann, Leipzig, 179 p., 5 pl. [Separate: "Festschrift für Carl Gegenbaur."]
- HAGENOW, F. V., 1840, *Monographie der rügenschen Kreide-Versteinerungen, Abt. 2. Radiarien und Annulaten*: Neues Jahrb. Mineralogie, Jahrg. 1840, p. 631-672, 1 pl.
- HALL, JAMES, 1843, *Geology of New York. Part IV. Comprising the survey of the fourth geological district*: Nat. History of New York, Albany, 683 p., 74 fig., 19 pl.
- , 1852, *Containing descriptions of the organic remains of the lower middle division of the New York system*: New York State Geol. Survey, Palaeontology of New York, v. 2, 362 p., 84 pl.
- , 1858a, *Crinoids of New York*: Am. Jour. Sci. Arts, ser. 2, v. 25, p. 277-279.
- , 1858b, *Palaeontology of Iowa*: Iowa Geol. Survey, v. 1, pt. 2, p. 473-724 [Crinoids, p. 479-483, 524-596, 608-634, 654-655, 669-673, 678-690; pl. 1, 8-11, 15-18, 25].
- , 1859a, *Contributions to the paleontology of Iowa; being descriptions of new species of Crinoidea and other fossils*: Iowa Geol. Survey, supplement to v. 1, pt. 2 of Geol. Rept. of Iowa, 94 p., 3 pl.
- , 1859b, *Descriptions and figures of the organic remains of the Lower Helderberg group and the Oriskany sandstone*: New York Geol. Survey, v. 3, 532 p., 120 pl.
- , 1860, *Contributions to paleontology, 1858 and 1859*: New York State Cab. Nat. History, Ann. Rept. 13, p. 95-112, 121-124, 8 fig.
- , 1861, *Descriptions of new species of Crinoidea from the Carboniferous rocks of the Mississippi valley*: Boston Soc. Nat. History, Jour., v. 7, p. 261-328, pl. 1-7. [Plates 1-7 published 1872 and privately distributed.]
- , 1862, *Preliminary notice of some of the species of Crinoidea, known in the Upper Helderberg and Hamilton groups of New York*: New York State Cab. Nat. History, Ann. Rept. 15, p. 115-153, 2 pl.
- , 1866a (Nov.), *Descriptions of new species of Crinoidea and other fossils from the Lower Silurian strata of the age of the Hudson-River Group and Trenton Limestone*: Albany, 17 p. [Printed in advance without plates=Hall, 1872.]
- , 1866b (1867), *Account of some new or little known species of fossils from rocks of the age of the Niagara group*: New York State Cab. Nat. History, Ann. Rept. 20, p. 305-401, pl. 10-23 (revised edit. 1870, p. 347-438, pl. 10-25).
- , 1868 (1870), *Descriptions of new or little-known species of fossils from rocks of the age of the Niagara Group*: New York State Cab. Nat. History, Ann. Rept. 20, p. 347-438, pl. 10-25.
- , 1872, *Descriptions of some new species of Crinoidea, etc.* [title same as Hall, 1866a]: New York State Cab. Nat. History, Ann. Rept. 24, p. 205-224, pl. 5-8.
- , 1879 (1880), *Notice of some new and remarkable forms of Crinoidea from the Lower Helderberg group of New York and Tennessee*: revised and reprinted edition of paper from Albany, New York State Mus. Nat. History, Ann. Rept. 28 for 1879, p. 1-8, pl. 35-37.
- , 1882, *Descriptions of the species of fossils found in the Niagara group at Waldron, Indiana*: Indiana Dept. Geology and Nat. History, 11th Ann. Rept., p. 217-345, 36 pl.
- HOFER, JOHANN, 1760, *Tentaminis lithologici de Polyporitis*: Acta Helvetica, v. 4, p. 169-213, pl. 6-9.
- INTERNATIONAL COMMISSION ON ZOOLOGICAL NOMENCLATURE, 1961, *International code of zoological nomenclature adopted by the XV International Congress of Zoology*: International Trust for Zoological Nomenclature, London, 176 p.
- JAEKEL, OTTO, 1895, *Beiträge zur Kenntnis der palaeozoischen Crinoiden Deutschlands*: Palaeont. Abh., new ser., v. 3 (der Ganzen Reihe, Bd. 7), p. 1-116, fig. 1-29, pl. 1-10.
- , 1898, *Über einen neuen devonischen Pentacrinoiden*: Deutsch. Geol. Gesell., Zeitschr., Verh. 1898, v. 50, p. 28-32.
- , 1918, *Phylogenie und System der Pelmatozoen*: Paläont. Zeitschr., v. 3, p. 1-128, 114 fig.
- JEFFORDS, R. M., & MILLER, T. H., 1967, *Ontogenetic development in Late Pennsylvanian crinoid columnals*:

- Univ. Kansas, Paleont. Contrib., Echinodermata, Art. 10, p. 1-14, fig. 1-5, pl. 1-4.
- KAUFFMAN, A. E., & ROTH, R. I., 1966, *Upper Pennsylvanian and Lower Permian fusulinids from north-central Texas*: Cushman Found. Foram. Research, Spec. Publ. 8, 49 p., 2 fig., 11 pl.
- KESLING, R. V., 1965, *Proctothylacocrinus esseri, a new crinoid from the Middle Devonian Silica Formation of northwestern Ohio*: Univ. Michigan, Mus. Paleont., Contrib., v. 20, no. 4, p. 75-87, 1 fig., 5 pl.
- KIER, P. M., 1952, *Echinoderms of the Middle Devonian Silica Formation of Ohio*: Univ. Michigan, Mus. Paleont., Contrib., v. 10, no. 4, p. 59-81, 4 pl.
- KIRK, EDWIN, 1933, *Syndetocrinus, a new crinoid genus from the Silurian of Canada*: Am. Jour. Sci., 5th ser., v. 26, p. 344-354, 8 fig.
- , 1941, *Four new genera of Mississippian Crinoidea Inadunata*: Jour. Paleont., v. 15, p. 82-88, pl. 18-19.
- KONINCK, L. G. DE, & LEHON, H. S., 1854, *Recherches sur les crinoïdes du terrain carbonifère de la Belgique*: Acad. Royale Belgique, Mém., v. 28, 217 p., 8 pl.
- LAMBE, L. M., 1896, *Description of a supposed new genus of Polyzoa from the Trenton limestone of Ottawa*: Canadian Rec. Sci., v. 7, p. 1-3.
- LANE, N. G., 1963, *The Berkeley crinoid collection from Crawfordsville, Indiana*: Jour. Paleont., v. 37, p. 1001-1008, 2 fig., pl. 128.
- , & WEBSTER, G. D., 1966, *New Permian crinoid fauna from southern Nevada*: Univ. California, Publ. Geol. Sci., v. 63, p. 1-60, 19 fig., pl. 1-13.
- LEE, WALLACE, et al., 1938, *Stratigraphic and paleontologic studies of the Pennsylvanian and Permian rocks in north-central Texas*: Texas Univ., Bull. 3801, 252 p., 9 fig., 11 pl.
- LINNÉ, KARL [LINNAEUS, CAROLUS], 1758 [1759], *Systema naturae per regna tria naturae, secundum classes, ordines, genera, species cum characteribus, differentiis, synonymis locis*: Laurentii Salvii, Holmiae, 10th edit., 2 v., 824 p.
- LORIOU, PERCEVAL DE, 1882, *Description of a new species of Bourgueticrinus*: Cincinnati Soc. Nat. History, Jour., v. 5, p. 118, pl. 5.
- , 1882-89, *Paléontologie française, ou description des fossiles de la France*, d'Orbigny, A.D. (ed.): Première série, Animaux invertébrés, Terrain jurassique, tome 2, première partie, crinoïdes (1882-84), 627 p., pl. 1-121; deuxième partie, crinoïdes (1884-89), 580 p., pl. 122-229.
- , 1904, *Notes pour servir à l'étude des échinodermes*: Basel, Genève, & Berlin, sér. 2, 68 p., 4 pl.
- LYON, S. S., 1857, *Paleontological report*: Kentucky Geol. Survey, Rept. 3, p. 465-498, 5 pl.
- , & CASSEDAY, S. A. 1859 (1860), *Description of nine new species of Crinoidea from the Subcarboniferous rocks of Indiana and Kentucky*: Am. Jour. Sci., ser. 2, v. 28, p. 233-246. [No illus.]
- MAREZ-OYENS, F. A. H. W. DE, 1940, *Neue permische Krinoiden von Timor*: in Geological expedition to the Lesser Sunda Islands under leadership of H. A. Brower; Amsterdam, v. 1, p. 287-348, 11 fig., pl. 1-4.
- MARSHALL, A. F. GRAF VON, 1873, *Nomenclator zoologicus continens nomina systematica generum animalium tam viventium quam fossilium secundum ordinem alphabeticum disposita, sub auspiciis et sumptibus C. R. Societatis zoologico-botanicae conscriptus a comite Augusto de Marschall*: M. Salzar, Vindobonae, 482 p.
- MARTIN, E. K., 1965, *Biostratigraphy of the Upper Pennsylvanian Wayland Shale in McCulloch-Coleman Counties, central Texas*: Rice Univ., unpubl. dissert., 99 p., 20 fig., 9 pl.
- , 1966, *Biostratigraphy of the Upper Pennsylvanian Wayland Shale in McCulloch-Coleman Counties, central Texas* [Abst.]: Houston Geol. Soc., Bull., v. 8, no. 3, p. 19, 20.
- MEEK, F. B., 1871 (1872), *Preliminary list of the fossils collected . . . in Utah and Wyoming Territories, with descriptions of a few new species*: in Hayden, F. V., Preliminary report U. S. Geol. Survey Montana and portions of adjacent territories (fifth ann. rept.); p. 373-377.
- , 1872, *Report on the paleontology of eastern Nebraska with some remarks on the Carboniferous rocks of that district*: in Hayden, F. V., Final report U. S. Geol. Survey Nebraska . . .; U. S. 42nd Congress, 1st sess., House Exec. Doc. 19, p. 83-264, pl. 1-11.
- , 1873, *Descriptions of invertebrate fossils of the Silurian and Devonian systems*: Ohio Geol. Survey, v. 1, pt. 2, 246 p., 23 pl.
- , & HAYDEN, F. V., 1858, *Descriptions of new organic remains collected in Nebraska Territory . . . together with some remarks on the geology of the Black Hills and portions of the surrounding country*: Acad. Nat. Sci. Philadelphia, Proc. (1858), v. 10, p. 41-59.
- , & WORTHEN, A. H., 1860, *Descriptions of new species of Crinoidea and Echinoidea from the Carboniferous rocks of Illinois, and other western states*: Acad. Nat. Sci. Philadelphia, Proc., ser. 2, v. 4, p. 379-397.
- , & ——, 1865a, *Description of new species of Crinoidea, etc., from the Paleozoic rocks of Illinois and some of the adjoining states*: Acad. Nat. Sci. Philadelphia, Proc., ser. 1, v. 17, p. 143-166.
- , & ——, 1865b, *Description of new species of Crinoidea, etc., from the Palaeozoic rocks of Illinois and some of the adjoining states*: Acad. Nat. Sci. Philadelphia, Proc., v. 17, no. 3, p. 143-155.
- , & ——, 1866, *Descriptions of invertebrates from*

- the Carboniferous system*: Illinois Geol. Survey, v. 2, sec. 2, p. 143-411, pl. 14-32.
- , & ———, 1868a, *Paleontology* [of Illinois]: Illinois Geol. Survey, v. 3, pt. 2, p. 289-565, pl. 1-20.
- , & ———, 1868b, *Remarks on some types of Carboniferous Crinoidea with descriptions of new genera and species of the same, and of one echinoid*: Acad. Nat. Sci. Philadelphia, Proc. (Dec. 1869), v. 20, p. 335-359.
- , & ———, 1873, *Palaeontology. Descriptions of invertebrates from Carboniferous system*: Illinois Geol. Survey, v. 5, pt. 2, p. 323-619, pl. 1-32.
- MILLER, J. S., 1821, *A natural history of the Crinoidea or lily-shaped animals, with observation on the genera Asteria, Euryale, Comatula, and Marsupites*: Bryan & Co., Bristol, 150 p., 50 pl.
- MILLER, S. A., 1874, *Lichenocrinus tuberulatus*: Cincinnati Quart. Jour. Sci., v. 1, p. 346-347, fig. 38.
- , 1883, *The American Palaeozoic fossils* [Echinodermata]: Cincinnati, edit. 2, p. 276-288.
- , 1888, *A new genus of crinoids from the Niagara Group*: Am. Geologist, v. 1, p. 263-264.
- , 1889, *North American geology and paleontology*: Cincinnati, p. 1-664, fig. 1-1194, First appendix, 1892, p. 665-718, fig. 1195-1265.
- , 1890, *The structure, classification, and arrangement of American Paleozoic crinoids into families*: Am. Geologist, v. 6, p. 275-286, 340-357.
- , & GURLEY, W. F. E., 1890, *Description of some new genera and species of Echinodermata, from the Coal Measures and Subcarboniferous rocks of Indiana, Missouri, and Iowa*: Cincinnati Soc. Nat. History, Jour., v. 13, p. 3-25, pl. 1-4.
- , & ———, 1894, *New genera and species of Echinodermata*: Illinois Mus. Nat. History, Bull. 5, 53 p., 5 pl.
- MOORE, R. C., 1939a, *Platycrinid columnals in Lower Permian limestone of western Texas*: Jour. Paleont., v. 13, p. 228-229, 1 fig.
- , 1939b, *The use of fragmentary crinoidal remains in stratigraphic paleontology*: Denison Univ., Bull., Sci. Lab. Jour., v. 33 (1938), p. 165-250, 14 fig., 4 pl.
- , 1939c, *New crinoids from Upper Pennsylvanian and Lower Permian rocks of Oklahoma, Kansas, and Nebraska*: Denison Univ., Bull., Sci. Lab. Jour., v. 34, p. 171-279, 39 fig., pl. 5-9.
- , 1948, *Evolution of the Crinoidea in relation to major paleogeographic changes in earth history*: Internat. Geol. Cong., 18th, Great Britain, rept. 1948, pt. 12, p. 27-53, 18 fig.
- , 1952, *Crinoids*: in Moore, R. C., Lalicker, C. G., and Fischer, A. G., *Invertebrate fossils*, McGraw-Hill, New York, Toronto, London, 766 p. [Crinoids p. 574-714, fig. 16-21.]
- , 1962a, *Revision of Calceocrinidae*: Univ. Kansas, Paleont. Contrib., Echinodermata, Art. 4, 40 p., 21 fig., 3 pl.
- , 1962b, *Ray structures of some inadunate crinoids*: Univ. Kansas, Paleont. Contrib., Echinodermata, Art. 5, 47 p., 17 fig., 4 pl.
- , 1967, *Unique stalked crinoids from Upper Cretaceous of Mississippi*: Univ. Kansas, Paleont. Contrib., Paper 17, 35 p., 8 fig., 8 pl.
- , & JEFFORDS, R. M., 1967, *Classification and nomenclature of fossil crinoids based on dissociated parts of columns*: Univ. Kansas, Paleont. Contrib., Echinodermata, Art. 9, p. 1-86, fig. 1-6, pl. 1-28.
- , & MILLER, T. H., 1967, *Morphological features of crinoid columns*: Univ. Kansas, Paleont. Contrib., Echinodermata, Art. 8, p. 1-30, fig. 1-5, pl. 1-4.
- , & LAUDON, L. R., 1943, *Evolution and classification of Paleozoic crinoids*: Geol. Soc. America, Spec. Paper 46, 153 p., 18 fig., 14 pl.
- , & ———, 1944, *Crinoidea*: in Shimer, H. W., and Shrock, R. R., *Index fossils of North America*, John Wiley & Sons, Inc., New York, p. 137-211, pl. 52-79.
- , & PLUMMER, F. B., 1938, *Upper Carboniferous crinoids of the Morrow Subseries of Arkansas, Oklahoma and Texas*: Denison Univ., Bull., Sci. Lab. Jour., v. 32, p. 209-313, pl. 12-16.
- , & ———, 1940, *Crinoids from the Upper Carboniferous and Permian strata in Texas*: Texas Univ., Bull. 3945, 468 p., 78 fig., 21 pl.
- , & SYLVESTER-BRADLEY, P. C., 1957a, *Proposed insertion in the "Règles" of provisions recognizing "parataxa" as a special category for the classification and nomenclature of discrete fragments or of life-stages of animals which are inadequate for identification of whole-animal taxa, with proposals for procedure for the nomenclature of "parataxa"*: Bull. Zool. Nomenclature, v. 15, quadruple pt. ¼, Sept. 1957, p. 5-13.
- , & ———, 1957b, *First supplemental application for a ruling of the International Commission directing that the classification and nomenclature of discrete conodonts be in terms of "parataxa"*: Bull. Zool. Nomenclature, v. 15, quadruple pt. ¼, p. 14-33.
- , & ———, 1957c, *Second supplemental application for a ruling by the International Commission that the classification and nomenclature of ammonoid aptychi (Class Cephalopoda) be in terms of "parataxa"*: Bull. Zool. Nomenclature, v. 15, quadruple pt. ¼, p. 35-70.
- , & VOKES, H. E., 1953, *Lower Tertiary crinoids from northwestern Oregon*: U. S. Geol. Survey, Prof. Paper 233-E, p. 111-147, fig. 27-39, pl. 14-24.
- MORRIS, JOHN, 1843, *A catalogue of British fossils. Comprising all the genera and species hitherto described*;

- with reference to their geological distribution and to the localities in which they have been found: John Van Voorst, London, 222 p. (1st edit.).
- MÜLLER, JOHANNES, 1855, *Ueber die Echinodermen in der Umgegend von Coblenz und in dem Eifeler Kalke*: in Zeiler, F., and Wirtgen, P. W., *Bemerkungen ueber die Petrofacten der ältern devonischen Gebirge am Rheine, insbesondere, ueber die der Umgegend von Coblenz vorkommenden Arten*, Rheinland Nat. Ver. Verh., v. 12, p. 79-85, pl. 10-12.
- MÜNSTER, G. G. ZU, 1839, *Beschreibung einiger neuen Crinoideen aus der Uebergangsformation*: Beiträge zur Petrefaktenkunde, v. 1, p. 1-124, pl. 1-19.
- MURCHISON, R. I., 1839, *The Silurian System, Part 1. Founded on geological researches in the counties of Solop, Hereford, Radnor, Montgomery, Caermarthen, Brecon, Pembroke, Monmouth, Gloucester, Worcester, and Stafford; with descriptions of the coal-fields and overlying formations*: John Murray, London, p. 1-578; Part 2. Organic remains, p. 579-768, 37 pl.
- MYERS, D. A., 1960, *Stratigraphic distribution of some Pennsylvanian Fusulinidae from Brown and Coleman Counties, Texas*: U. S. Geol. Survey, Prof. Paper 315-C, 53 p., 2 fig., 10 pl.
- NIELSEN, K. B., 1913, *Crinoiderne i Danmarks Kridtafleveringer*: Danmarks Geol. Undersøgelse, ser. 2, no. 26 (Univ. Copenhagen, Mus. Mineralogy and Paleontology, Paleont. Communications 11), 120 p., 34 fig., 12 pl.
- OOSTER, W. A., 1865, *Petrifications remarquables des Alpes Suisses; Synopsis des echinodermes des Alpes Suisses*: H. Georg, Genève and Bâle, 131 p., 29 pl.
- ORBIGNY, A. D. D', 1840, *Histoire naturelle générale et particulière des crinoïdes, vivants et fossiles, comprenant description géologique et zoologique de ces animaux*: The author, Paris, 98 p., 18 pl.
- , 1849-51, *Cours élémentaire de paléontologie et géologie stratigraphiques*: Victor Masson, Paris, v. 1 (1849), 299 p., 165 fig.; v. 2 (1851), 847 p., 628 fig.
- PHILIP, G. M., 1961, *Lower Devonian crinoids from Toongabbie, Victoria, Australia*: Geol. Mag., v. 98, p. 143-160, 10 fig., pl. 8.
- PHILLIPS, JOHN, 1836, *Illustration of the geology of Yorkshire, or a description of the strata and organic remains, Pt. 2. The Mountain limestone districts*: John Murray, London, 253 p., 25 pl.
- , 1839, in Murchison, R. I., *The Silurian system*: John Murray, London, p. 670-675, pl. 17-18.
- , 1841, *Figures and descriptions of the Paleozoic fossils of Cornwall, Devon, and west Somerset*: Longman, Brown, Green, & Longmans, London, 232 p., 6 pl.
- , 1843, *Crinoids*: in Morris, John, *A catalogue of British fossils*, 222 p.
- PLUMMER, F. B., & MOORE, R. C., 1922, *Stratigraphy of the Pennsylvanian formations of north-central Texas*: Texas Univ., Bull. 2132, 236 p., 19 fig., 27 pl.
- PORKORNÝ, VLADIMIR, 1958, *Echinoderma-Stachelhäuter*: in Grundzüge der zoologischen Mikropalaontologie, Band II, Veb Deutscher Verlag der Wiss. Berlin, p. 323-351, fig. 1046-1070.
- PUSCH, G. G., 1837, *Polens Palaeontologie*: Stuttgart, 218 p., 16 pl.
- QUENSTEDT, F. A., 1852, *Handbuch der Petrefactenkunde*: H. Laupp Verlag, Tübingen, 1st edit., 792 p., 62 pl. (text and plates bound separately). [Descriptions and figures of numerous crinoids, including isolated columnals and pluricolumnals.]
- , 1867, *Handbuch der Petrefactenkunde*: H. Laupp Verlag, Tübingen, 2nd edit. (essentially same as 1st edit., crinoids, p. 714-758).
- , 1874-76, *Petrefactenkunde Deutschlands*: Fues's Verlag (R. Reisland), Leipzig, v. 4, Abt. 1, Echinodermen, p. 1-742, pl. 90-114.
- , 1882 (1885), *Handbuch der Petrefactenkunde*: H. Laupp Verlag, Tübingen, 3rd edit., Abt. 4, crinoids, p. 912-965, pl. 63-80.
- RAMSBOTTOM, W. H. C., 1961, *A monograph on British Ordovician Crinoidea*: Palaeontographical Soc., Mon., v. 114, 37 p., 11 fig., 8 pl.
- RASMUSSEN, H. W., 1953, *Cretaceous Crinoidea; preliminary report on the species found in Denmark*: Dansk Geol. Foren. Medd., v. 12, p. 415-419.
- , 1961, *A monograph on the Cretaceous Crinoidea*: K. Vidensk. Selsk. Biol. Skr., v. 12, no. 1, 428 p., 60 pl.
- RINGUEBERG, E. N. S., 1886, *New genera and species of fossils from the Niagara shales*: Buffalo Soc. Nat. Sci., Bull. 5, p. 1-22, pl. 1.
- ROEMER, C. F., 1860, *Die silurische Fauna des westlichen Tennessee*: E. Trewendt, Breslau, 100 p., 5 pl.
- ROEMER, F. A., 1836, *Die Versteinerungen des norddeutschen Oolithengebirges*: Hahn, Hannover, 218 p., 16 pl.
- , 1840-41, *Die Versteinerungen des norddeutschen Kreidegebirges*: Hahn, Hannover, 145 p., 16 pl.
- , 1852-54, *II. Erste Periode, Kohlen-Gebirge*: in Bronn, H. G., 1851-56, *Lethaea Geognostica*, E. Schweizerbart, Stuttgart, 3rd edit., v. 2, 788 p.
- ROVERETO, G., 1914, *Nuovi studi sulla stratigraphia e sulla fauna dell' Oligocene Ligure*: Gênes, 180 p., 1 pl.
- RUSCONI, CARLOS, 1955, *Fosiles Cambricos y Ordovicios al oeste de San Isidro, Mendoza*: Mus. Historia Nat. Mendoza, Rev., v. 8, p. 3-64, 5 pl.
- SANDBERGER, GUIDO & SANDBERGER, FRIDOLIN, 1850-56, *Die Versteinerungen des rheinischen Schichtensystems in Nassau*: xiv + p. 1-564, pl. 1-41, Kreidel & Niedner, Wiesbaden, Verlagshandlung; Atlas (sep.), 41 pl. [Crinoidea p. 383-403, pl. 35].

- SARDESON, F. W., 1908, *Discoid crinoidal roots and Camarocrinus*: Jour. Geology, v. 16, p. 239-254, 31 fig.
- SARS, MICHAEL, 1863 (1864), *Om en ny Art Brachiolaria*: Vidensk. Selsk. Förh. Christiania, p. 126-137.
- SCHMIDT, W. E., 1930, *Die Echinodermen des deutschen Unterkarbons in die Fauna des deutschen Unterkarbons*: Preuss. geol. Landesanstalt, Abh., new ser., no. 122, pt. 1, 92 p., 20 fig., 3 pl.
- , 1931, *Crinoideen und Blastoideen aus dem jüngsten Unterdevon Spaniens*: Paleontographica, v. 76, p. 1-34, pl. 1-4.
- , 1934, *Die Crinoideen des rheinischen Devons, Teil I, Die Crinoideen des Hunsrückschiefers*: Preuss. Geol. Landesanstalt, Abh., new ser., no. 163, 149 p., 34 pl.
- , 1942, *Die crinoideen des Rheinischen Devons, Teil II, A. Nachtrag zu die crinoideen des Hunsrückschiefers; B. Die crinoideen des Unterdevons bis zur Cultrijugatus-zone (mit Ausschluss des Hunsrückschiefers)*: Reichstelle f. Bodenforschung Abh., new ser., no. 182, 253 p., 26 pl.
- SCHULTZE, LUDWIG, 1867, *Monographie der Echinodermen des Eifler Kalkes*: Kgl. Akad. Wiss., math.-Naturw. Cl., v. 26, p. 113-230, 19 fig., 13 pl.
- SCHULZE, C. F., 1760, *Betrachtung der versteinerten See Sterne und ihrer Theile*: Groll, Warsaw and Dresden, 59 p., 3 pl. [Rejected publication for purposes of zoological nomenclature, ICZN (Bull. Zool. Nomencl., v. 19, p. 399).]
- SHEVCHENKO, T. V., 1964, *Nizhnesiluriyskije krinoidei Tsentralnogo Tadzhikistana* [Lower Silurian crinoids of central Tadzhikistan]: Akad. Nauk Tadzhikskoy SSR, Tadzhikskoe Otdelenie Vsesoyuznogo Paleont. Obschestva, Paleontologiya Tadzhikistana [Dushanbe], p. 8-20, pl. 1-4.
- , 1966, *Morskije lilii iz Verkhnesiluriyskikh i Nizhnedevonskikh otzheniy lugo-zapadnogo Tian-Shanya i ikh stratigraficheskoe znachenie* [Crinoids from Upper Silurian and Lower Devonian deposits of southwestern Tian-Shan and their stratigraphic significance]: Upravleniya Geologii Soveta Ministrov Tadzhikskoy SSR, Trudy, Paleontologiya i Stratigrafiya, v. 2, p. 123-188, fig. 1-41, pl. 1-8.
- SHROCK, R. R., & TWENHOFEL, W. H., 1953, *Echinoderma*: in Principles of Invertebrate Paleontology, 2nd edit., McGraw-Hill, New York, p. 642-735, fig. 1-57.
- SIEVERTS-DORECK, HERTHA, 1939, *Jura-und Kreide Crinoideen aus Deutsch-ostafrika*: Paleontographica, suppl. 7, Reich pt. II, no. 3, p. 217-231, pl. 16. [Dissociated pluricolumnals referred to *Apiocrinus* sp. aff. *A. crassus* (d'Orbigny) described and figured.]
- , 1942, *Crinoiden aus dem Perm Tasmaniens*: Zentralbl. Mineralogie, Geologie u. Palaeontologie, Jahrg. 1942, Abt. B, no. 7, p. 222-231, 7 fig. [*Camptocrinus? tasmaniensis*, n. sp., based on dissociated columnals and pluricolumnals.]
- , 1943, *Armgleider von Roveacrinus aus einem norddeutschen Senongeschiebe*: Z. Sonderabdruck aus Zeitschr. für. Geschiebeforsch. und Flacklandsgeologie, v. 18, p. 96-100, fig. 1-24, 1 pl.
- , 1951a, *Crinoiden aus dem Unterkarbon des Oberharzes*: Neues Jahrb. Geologie u. Palaeontologie, Abh., v. 93, p. 117-144, 8 fig., pl. 8-9. [Describes *Orocrinus* and new species based on dissociated stem parts.]
- , 1951b, *Ein Isocrinus aus dem obersten Kreide Bulgariens*: Paläont. Zeitschr., v. 24, p. 72-75, 3 fig. [*Isocrinus gocevi*, n. sp., based on dissociated columnals and pluricolumnals, figured articularia compared with typical *Balanocrinus* and *Austinocrinus*.]
- , 1953a, *Ueber Austinocrinus im norddeutschen Senon mit einem Beitrag zur Gliederung, Stammesgeschichte und Verbreitung der Gattung*: Hamburg Geol. Staatsinst., Mitt., no. 22, p. 102-118, 6 fig., pl. 17-18.
- , 1953b, *Ueber einige inadunate Crinoiden aus dem rheinischen Devon*: Hesse Landesamt. Bodenforschung Notizbl., v. 81, p. 75-87, 8 fig., pl. 4. [Includes morphological features of *Myelodactylus canaliculatus* (Goldfuss).]
- , 1954, *Weitere Mitteilungen ueber Myelodactylus aus dem Mittel-Devon der Eifel*: Hesse Landesamt. Bodenforschung, Notizbl., v. 82, p. 46-49, 1 pl. [Morphology of *Myelodactylus canaliculatus* (Goldfuss) described and figured.]
- , 1957, *Bemerkungen uber altpaläozoische Crinoiden aus Argentinien*: Neues Jahrb. Geologie u. Palaeontologie, Monatsh. Abt. B, p. 151-156, 4 fig. [Discusses pentamerally divided column of *Salagastiana* Rusconi.]
- , 1960, *Spezielle Arbeitsgebiete der Mikropaläontologie 3. Echinodermen*: Handbuch der Mikroskopie in der Technik, Band II, Teil 3, Mikroskopie in der Geologie sedimentärer Lagerstätten (Mikropalaeontologie), p. 239-264, pl. 1-6.
- SIZOVA, E. N., 1960, *Znachenie iskopaemykh stebley morskikh liliy dlya stratigrafii Devona i Karbona tsentralnogo Kazakhstana* [Significance of fossil crinoid stems in Devonian and Carboniferous stratigraphy of central Kazakhstan]: Materialy po geologii i poleznykh iskopaemykh Altaya i Kazakhstana, Vses. Nauchno-Issled. Geol. Inst., Trudy [VSEGEI], new ser., no. 33, p. 51-65.
- SLOCOM, A. W., 1907, *New crinoids from the Chicago area*: Field Columbian Mus., Pub. 123, v. 2, no. 10, p. 273-306, 11 fig., pl. 82-87.
- , & FOERSTE, A. F., 1924, *New echinoderms from the Maquoketa beds of Fayette County, Iowa*: Iowa Geol. Survey, v. 29, Ann. Rept. 1919 and 1920, p. 315-384, pl. 29-32.

- SPRINGER, FRANK, 1901, *Uintacrinus, its structure and relations*: Harvard College Mus. Comp. Zoology, Mem. 25, p. 1-89, 8 pl.
- , 1909, *A new American Jurassic crinoid*: U.S. Natl. Mus., Proc., v. 36, p. 179-190, pl. 4.
- , 1911, *Some new American fossil crinoids*: Harvard College, Mus. Comp. Zoology, Mem. 25, no. 3, p. 117-161, 6 pl.
- , 1913, *Crinoidea*: in Zittel, K. A., Text-book of paleontology, transl. & ed. by C. R. Eastman, 2nd edit., Macmillan & Co., Ltd., London, v. 1, 839 p., 1594 fig. [Crinoids, p. 173-243.]
- , 1917, *On the crinoid genus Scyphocrinus and its bulbous root, Camarocrinus*: Smithsonian Inst., Pub. 2440, 74 p., illus.
- , 1920, *The Crinoidea Flexibilia*: Smithsonian Inst., Pub. 2501, 486 p., 51 fig., 79 pl.
- , 1921, *The fossil crinoid genus Dolatocrinus and its allies*: U. S. Natl. Mus., Bull. 115, 78 p., 6 fig., 16 pl.
- , 1922a, *Balanocrinus in America*: Pan-Am. Geologist, v. 38, p. 262-263.
- , 1922b, *Crinoids from the Upper Cretaceous of Tamaulipas, Mexico*: U. S. Natl. Mus., Proc., v. 61, art. 5, 4 p., 1 pl.
- , 1926a, *American Silurian crinoids*: Smithsonian Inst., Pub. 2871, 239 p., 33 pl.
- , 1926b, *Upper Devonian crinoids from the Mackenzie River valley*: Canada Geol. Survey, Mus. Bull. 42, p. 127-132, pl. 24.
- , 1926c, *Unusual forms of fossil crinoids*: U. S. Natl. Mus., Proc., v. 67, art. 9, 137 p., 26 pl.
- STEININGER, JOHANN, 1837, *No title [Notes from meeting in which 2 new genera, with their types are published]*: Soc. Géol. France, Bull., v. 8, p. 230-232, 2 fig. [*Haplocrinites*, n. gen., p. 231; *Haplocrinites sphaeroideus*, n. sp., p. 232 (small figure p. 232); *Lichas antiquus* n. gen. & n. sp., p. 231, with figure.]
- , 1853, *Geognostische Beschreibung der Eifel*: Fr. Lints, Trier, 143 p., 9 pl.
- STEINMANN, GUSTAV, 1908, *Die geologischen Grundlagen der Abstammungslehre*: Leipzig, 284 p.
- STEVENS, C. H., JONES, D. H., & TODD, R. G., 1960, *Ultrasonic vibrations as a cleaning agent for fossils*: Jour. Paleont., v. 34, p. 727-730, pl. 93-94.
- STRIMPLE, H. L., 1961, *Late Desmoinesian crinoid faunule from Oklahoma [Pennsylvanian]*: Oklahoma Geol. Survey, Bull. 93, 189 p., 23 fig., 19 pl.
- , 1962, *Platycrinid columnals from the Pumpkin Creek Limestone*: Oklahoma Geology Notes, v. 22, p. 3-5, 7 fig.
- , 1963, *Crinoids of the Hunton Group (Devonian-Silurian) of Oklahoma*: Oklahoma Geol. Survey, Bull. 100, 169 p., 30 fig., 12 pl.
- STUKALINA, G. A., 1960, *Kompleks llanoveriyskikh stebly morskikh liliy, khr. chingiz* [Crinoid stems from Llandoverly complex of Khr. Chingiz]: Vses. Nauchno-Issled. Geol. Inst., Trudy [VSEGEI], no. 35, p. 95-110, 7 fig.
- , 1961, *Stebli krinoidey iz otlozheniy verknego Silura gor, Akhsar'lyi (Tsentralnyy Kazakhstan)* [Crinoid stems from Upper Silurian deposits of Akhsar'ly region (central Kagakhstan)]: Vses. Nauchno-Issled. Geol. Inst., Trudy, no. 42, p. 31-42, pl. 1-2.
- , 1964a, *K metodike izuchniya i sborov stebly morskikh liliy* [On methods of studying and collecting crinoid stems]: in Materialy po geologii i paleznykh iskopaenykh Altaya i Kazakhstana [Information on the geology and useful fossils of Altai and Kazakhstan], Vses. Nauchno-Issled. Geol. Inst., Trudy [VSEGEI], v. 3, p. 31-35, 1 fig. [General discussion, no systematics.]
- , 1965a, *Morskije lilii Karaespinskogo gorizonta* [Crinoids of the Karaspinsk horizon]: in Stratigrafiya nizhnepaleozoyskikh i Siluriyskikh otlozheniy tsentralnogo Kazakhstana [Stratigraphy of lower Paleozoic and Silurian deposits of central Kazakhstan], Vses. Nauchno-Issled. Geol. Inst., Trudy [VSEGEI], p. 134-141, 7 fig., 2 pl.
- , 1965b, *Novye vidy Hexacrinites(?) tsentralnogo Kazakhstana* [New species of *Hexacrinites?* central Kazakhstan]: Ezhegodnik Vses. Paleont. Obsch., v. 17, p. 188-193, 1 pl.
- , 1965c, *O taksonomicheskoy znachenii stebly drevnykh morskikh liliy* [On taxonomic significance of ancient crinoid stems]: Vses. Nauchno-Issled. Geol. Inst., Trudy [VSEGEI], Biostrat. sbornik, n. ser., v. 115, p. 210-217, 2 fig.
- , 1966, *O printsipakh klassifikatsii stebly drevnykh morskikh liliy* [Principles of classification of ancient crinoid stems]: Palaeont. Zhurn., no. 3, p. 94-102, 2 fig.
- , 1967, *O taksonomicheskikh proznamenakh segmentirovannykh stebly morskikh liliy* [On taxonomic indications in segmentation of crinoid stems]: Vsesoyuz. Nauchno-Issled. Geol. Inst. (VSEGEI), Trudy, nov. ser., v. 129, Biostratigraficheskii sbornik, pt. 3, p. 200-206, fig. 1-6.
- , & TUYUTYAN, YU. A., 1967, *Novye Ordovikskie morskie lilii Kazakhstana* [New Ordovician crinoid stems from Kazakhstan]: Akad. Nauk Kazakhskoy SSR, Izvestiya, ser. geol. 4, p. 73-76, fig. 1-12.
- SUTTON, A. H., & WINKLER, V. D., 1940, *Mississippian Inadunata—Eupachycrinus and related forms*: Jour. Paleont., v. 14, p. 544-567, pl. 66-68.
- TERMIER, HENRI, & TERMIER, GENEVIÈVE, 1949, *Hiérarchie et corrélations des caractères chez les crinoïdes fossiles*: Serv. Carte Géol. l'Algérie, Bull., sér. 1, Paléontologie, no. 10, p. 1-90, 8 pl.
- , & ———, 1958, *Les échinodermes permien de Djebel Tebaga (extrême sud Tunisien)*: Soc. Géol. France, Bull., sér. 6, v. 8, p. 51-64, 7 pl.

- THOMAS, A. O., 1923, *Some fossils from an outcrop in Des Moines*: Iowa Acad. Sci., Trans., v. 30, p. 471-483, 2 pl.
- TIEN, C. C., 1926, *Crinoids from the Taiyuan series of north China*: Palaeontologia Sinica, ser. B., v. 5, p. 5-51, pl. 3.
- TRAUTSCHOLD, HERMANN, 1859, *Recherches géologiques aux environs de Moscou*: Soc. Imp. Nat. Moscou, Bull. (Moskov. Obshch. Ispytateley Prirody Byull.), v. 32, p. 109-121, 2 pl.
- , 1867, *Einige Crinoiden und andere Tierreste des jüngeren Bergkalks in Gouvernement Moskau*: Soc. Imp. Nat. Moscou, Bull. (Moskov. Obshch. Ispytateley Prirody Byull.), v. 40, pt. 2, no. 3, 49 p., 5 pl.
- , 1879, *Die Kalkbrüche von Mjatschkowa; eine Monographie des oberen Bergkalks Schlusslieferung*: Soc. Imp. Nat. Moscou (Moskov. Obshch. Ispytateley Prirody Byull.), nouv. mém., v. 14, p. 101-180, pl. 12-18.
- , 1880 [1881], *Ueber Synphocrinus*: Soc. Imp. Nat. Moscou, Bull., (Moskov. Obshch. Ispytateley Prirody Byull.), v. 55, no. 4, ann. 1880, p. 390-396, 1 fig., pl. 6.
- TROOST, GERARD, 1849 [1850], *A list of the fossil crinoids of Tennessee*: Am. Jour. Sci. Arts, Proc., ser. 2, v. 8, p. 59-64.
- UBAGHS, GEORGES, 1953, *Classe des crinoïdes*: in Piveteau, Jean, *Traité de paléontologie*, Masson et Cie, Paris, v. 3, p. 658-773, fig. 1-166.
- VALETTE, D. A., 1934, *Le Permien marin de l'extrême-sud Tunisien; pt. 3, Les crinoïdes permien du sud de la Tunisie*: Serv. Carte Géol. Tunisie, Mém., n. sér., no. 1, pt. 3, p. 91-101, 1 pl.
- VAN SANT, J. F., 1964, *Crawfordsville crinoids*: in Van Sant, J. F., & Lane, N. G., *Crawfordsville (Indiana) crinoid studies*, Univ. Kansas, Paleont. Contrib., Art. 7, Echinodermata, p. 34-136, fig. 10-41, pl. 1-8.
- VYALOV, O. S., 1953, *K voprosy o klassifikatsii stebelkov morskikh-liliy* [Problem of classification of crinoid columns]: Akad. Nauk SSSR, Doklady, v. 89, no. 6, p. 1087-1090. [Lvov. Geol. Obshch., paleont. ser., Trudy, no. 2.]
- WAAGEN, WILHELM, & JAHN, J. J., 1899, *Recherches paléontologiques, v. 7, Classe des échinodermes, pt. 2, Famille des Crinoïdes*: in Barrande, Joachim, *Système silurien du centre de la Bohême*, Prague, p. 1-215, 33 fig., pl. 40-79.
- WACHSMUTH, CHARLES, 1896, *Class Crinoidea*: in Zittel, K. A., *Text-book of palaeontology*, Eastman, C. R. (ed.), Macmillan, London, New York, v. 1, p. 124-177, p. 219-291.
- , & SPRINGER, FRANK, 1879, [1880]-85, *Revision of the Palaeocrinoidea*: Acad. Nat. Sci. Philadelphia, Proc., pt. 1 (1879) [1880], p. 226-378, pl. 15-17; pt. 2 (1881), p. 177-411, pl. 17-19; pt. 3 (1885), p. 225-364, pl. 4-9.
- , & ———, 1886 [1887], *Revision of the Palaeocrinoidea, pt. 3, Discussion of the classification and relations of the brachiate crinoids, and conclusion of the generic descriptions. Sec. 2*: Acad. Nat. Sci. Philadelphia, Proc., p. 64-226.
- , & ———, 1897, *North American Crinoidea Camerata*: Harvard Coll., Mus. Comp. Zoology, Mem. 20, v. 21, 22, 897 p., 21 fig., 83 pl.
- WANNER, JOHANN, 1916, *Die permischen Echinodermen von Timor, Teil I*: Paläontologie von Timor, no. 6, pt. 11, p. 1-329, 88 fig., pl. 96-114 (1-19).
- , 1924, *Die permischen Krinoiden von Timor*: Mijnwezen Ned.-Oost Indië, Verh. 1921, pt. 3, p. 1-348, 61 fig., pl. 1-22.
- , 1929, *Neue Beiträge zur Kenntnis der permischen Echinodermen von Timor; I, Allagecrinus, II, Hypocrinites*: Nederlandsch-Indië, Dienst mijnbouw, Wetensch. Mededeel., no. 11, p. 1-116, pl. 1-7.
- , 1930a, *Neue Beiträge zur Kenntnis der permischen Echinodermen von Timor, III, Hypocrinitinae, Paracatillocrinus und Allagecrinus Dux*: Nederlandsch-Indië, Dienst mijnbouw, Wetensch. Mededeel., no. 13, p. 1-30, 9 fig., pl. 1-2.
- , 1930b, *Neue Beiträge zur Kenntnis der permischen Echinodermen von Timor, IV, Flexibilia*: Nederlandsch-Indië, Dienst mijnbouw, Wetensch. Mededeel., no. 14, p. 1-60, 31 fig., pl. 1-4.
- , 1931, *Neue Beiträge zur Kenntnis der permischen Echinodermen von Timor; V, Poteriocrinidae, Teil I*: Nederlandsch-Indië, Dienst mijnbouw, Wetensch. Mededeel., no. 16, p. 1-27, 19 fig. pl. 1-4.
- , 1937, *Neue Beiträge zur Kenntnis der permischen Echinodermen von Timor, VIII-XIII*: Palaeontographica, Suppl.-Bd. 4, Abt. 4, no. 2, p. 59-212, 82 fig., pl. 5-14.
- , 1938, *Beiträge zur Paläontologie des ostindischen Archipels; XV, Balanocrinus sundaicus n. sp. und sein Epöke aus dem Altmiocän der Insel Madura*: Neues Jahrb. Mineralogie, Beil. Bd. 79, Abt. B, no. 3, p. 385-402, 2 pl.
- , 1940, *Neue Beiträge zur Kenntnis der permischen Echinodermen von Timor, XIV, Poteriocrinidae, Teil 3*: Palaeontographica, Suppl.-Bd. 4, p. 215-242.
- , 1949, *Neue Beiträge zur Kenntnis der permischen Echinodermen von Timor, XVI, Poteriocriniden, Teil 4*: Palaeontographica, Suppl.-Bd. 4, 5, 56 p., 15 fig., 3 pl.
- WARREN, J. S., 1962, *Form classification of crinoid stems*: Geol. Soc. America, Spec. Paper 68, p. 81.
- WELLER, J. M., 1930, *On the occurrence of Platycrinus in Pennsylvanian strata of western Indiana*: Illinois Acad. Sci., Trans., v. 22, p. 478-484, 1 pl.
- WELLER, STUART, 1900, *The paleontology of the Niagaran limestone in the Chicago area; the Crinoidea*: Chicago Acad. Sci., Nat. History Survey Bull. 4, pt. 1, 152 p., 57 fig., 15 pl.

- , 1909a, *Kinderhook faunal studies; V, The fauna of the Fern Glen Formation*: Geol. Soc. America, Bull., v. 20, p. 265-332, pl. 10-15.
- , 1909b, *Description of a Permian crinoid fauna from Texas*: Jour. Geology, v. 17, p. 623-635, 1 pl.
- , 1916, *Atactocrinus, a new crinoid genus from the Richmond of Illinois; Description of a Ste. Genevieve Limestone fauna from Monroe County, Illinois*: Contrib. Walker Mus., v. 1, p. 239-265, pl. 15-19.
- WILSON, D. W. R., 1964, *The two-holed ossicle—a practical zone index fossil*: Canadian Inst. Mining and Metall., Petroleum and Nat. Gas Div., Tech. meeting, abstr. Program, p. 17 [Calgary, Alta., May 6-8, 1964].
- WÖHRMANN, S. VON, 1889, *Die Fauna der sogenannten Cardita—und Raibler-Schichten in dem Nordtiroler und bayerischen Alpen*: Geol. Rheichanst., Jahrb., v. 43, p. 181-258, pl. 5-10 (Wien).
- WRIGHT, JAMES, 1950-60, *A monograph of the British Carboniferous Crinoidea*: London, Palaeontographical Soc., v. 1, pt. 1, p. 1-24, 1 fig., pl. 1-7 (1950); pt. 2, p. 25-46, fig. 5-14, pl. 8-12 (1951); pt. 3, p. 47-102, fig. 15-41, pl. 13-31 (1951); pt. 4, p. 103-148, fig. 42-81, pl. 32-40 (1952); pt. 5, p. 149-190, fig. 82-108, pl. 41-47 (1954); v. 2, pt. 1, p. 191-254, fig. 109-124, pl. 48-63 (1955); pt. 2, p. 255-272, fig. 125, 126, pl. 64-67 (1955); pt. 3, p. 273-306, fig. 127, 128, pl. 68-75 (1956); pt. 4, p. 307-328, fig. 129-132, pl. 76-81 (1958); pt. 5, p. 329-347, pl. A-B (1960). [Pt. 5 is a posthumous appendix by W. H. C. Ramsbottom.]
- YAKOVLEV, N. N., 1933, *Deux crinoïdes du permien supérieur de la Transcaucasie*: Akad. Nauk SSSR, Izv., no. 7, p. 975-978, 1 pl.
- , & FAAS, A., 1938, *Novi echinodermi permiani di Sicilia*: Palaeontographia Italica, v. 38, p. 115-125, 1 pl.
- , & IVANOV, A. P., 1956, *Morskie lilii i blastoidei Kamennougolnykh i Permskikh otlozheniy SSR* [Crinoids and blastoids of the Carboniferous and Permian of the USSR]: Vses. Nauchno-Issled. Geol. Inst., Trudy (VSEGEI), new ser., v. 11, p. 1-142, 23 fig., 21 pl.
- YELTSHEVA [ELTYSHEVA], R. S., 1955, *Klass Crinoidea, morskie lilii, stebli morskikh lilii* [Class Crinoidea, sea lilies, crinoid stems]: (p. 40-47, fig. 7-8, pl. 23, 37, 54), in Nikiforova, O. I. (ed.), *Field Atlas of ordoviksnoi i Silurijskoi fauny Sibirskoi platformy*, Vses. Nauchno-Issled. Geol. Inst. [VSEGEI], Trudy, 140 p., 17 fig., 62 pl.
- , 1956, *Stebli morskikh lilii i ikh klassifikatsiya* [Crinoid stems and their classification]: Leningrad. Univ., Vestnik, ser. Geologiya i Geografii, no. 12, vyp. 2, p. 40-46, 3 fig.
- , 1957, *O novom semeistve paleozoiskikh morskikh lilii* [On a new family of Paleozoic crinoids]: Vses. Paleont. Obshch., Ezhegodnik, v. 16, p. 218-235, 9 fig., 3 pl.
- , 1959, *Pprintsipy klassifikatsii, metodika izucheniya i stratigraficheskoe znachenie stebley morskikh lilii* [Principles of classification and methods of study for stratigraphic use of crinoid stems]: in Voprosy paleobiologii i biostratigrafii [Problems of paleobiology and biostratigraphy], Vses. Paleont. Obshch., Ezhegodnik, Trudy, v. 2, p. 230-235, 1 pl.
- , 1960, *Ordovikske i Silurijske krinoidei Sibirskoy platformy* [Ordovician and Silurian crinoids of the Siberian platform]: in Biostratigrafiya Paleozoya Sibirskoy Platformy, Ordovik i Silur. Vses. Nauchno-Issled. Geol. Inst. Trudy (VSEGEI), n. ser., v. 3, p. 1-39, 6 pl.
- , 1964a, *Stebli Ordoviskikh morskikh lilii Prebaltiki (Nizhniy Ordovik)* [Prebaltic Ordovician crinoid stems (Lower Ordovician)]: Leningradskiy Ordena Lenina Gosudarstvennyy Universitet, Nauchno-Issled. Inst. Zemnoy Kory Paleont. Lab., p. 59-82, pl. 1-4.
- , 1964b [1965], *O klassifikatsii stebley morskikh lilii* [On classification of crinoid stems]: in Yakovlev, N. N., *Klass Crinoidea, obshchaya chasty, in Orlov, Yu. A., Osnovy paleologii, Iglokozhi, Gemikhordovy, Pogonofory, i Shchetinkochelyustnye*, p. 74, 75, 80, fig. 114, pl. 115, fig. 8-23.
- , 1966, *Stebli Ordoviskikh morskikh lilii Pribaltiki (Sredniy Ordovik)* [Prebaltic Ordovician crinoid stems (Middle Ordovician)]: Leningradskiy Ordena Lenina Gosudarstvennyy Universitet, Voprosy paleologii, v. 5, p. 53-70, pl. 1-3.
- , & DUBATOLOVA, YU. A., 1960 (1961), *Novye vidy Devoniskikh crinoidey Verkhnego Amura* [New species of Devonian crinoids from the Upper Amour]: in *Novye vidy drevnikh rasteniy i bespozvonochnykh*, Sbornik, pt. 2, Vses. Nauchno-Issled. Geol. Inst., Trudy [VSEGEI], p. 367-372, pl. 70.
- , & STUKALINA, G. A., 1963, *Stebli Ordoviskikh i nizhesiluriyskikh krinoidey tsentralnogo Taymyra, Novoy Zemli i Vaygacha* [Ordovician and Lower Silurian crinoid stems from Central Taimyr, Novaya Zemlya, and Vaygachal]: Nauchno-Issled. Inst. Geologii Arktiki, Uchennye zapiski, ser. paleont. i biostrat., Bull. 2, p. 23-62, 22 fig., 4 pl.
- ZITTEL, K. A. VON, 1879, *Handbuch der Paläontologie, v. 1, Paläozoologie*: R. Oldenbourg, Munchen & Leipzig, Abt. 1, 765 p., 557 fig.
- , 1900, *Text-book of paleontology*: transl. & ed. by Eastman, C. R.: Macmillan & Co., London & New York, v. 1, 706 p., 1476 fig. [The 1st edit.; p. 1-372 published in 1896, p. 373-706 published in 1900.]
- , 1913, *Text-book of paleontology*: transl. & ed. by Eastman, C. R.: Macmillan & Co., London, 2nd edit., 839 p., 1594 fig.

INDEX

This index is chiefly concerned with taxonomic units described or discussed in the Articles, in connection with which an asterisk (*) identifies type species of genera. Morphological terms are explained in the alphabetical glossary given in Article 8, and therefore mostly are omitted in the general index. Pages cited refer to Article 9 unless indicated otherwise by numbers in bold face type (e.g., **8** for Article 8, and **10** for Article 10).

- Acrochordocrinus*, 23
Actinocrinites, 23
Adelocrinus, 23
Ageneracrinus, 25
Ampholenium, 75
 " **apolegma*, 75
Amsdenanteris, 60
 " **tennesseensis*, 60
Ancyrocrinus, 9, 23
Angulocrinus, 23
Anthinocrinus, 23
Antunia, 23
Apiastrum, 58
 " **candidum*, 58
Apiocrinites, 23
 articularia (-um), **8**, 14
 Articulata, 48
Ascarum, 23
Aspidocrinus, 23
Asterocrinus, 23
Asteromischus, 54
 " **stellatus*, 54
 " **stellatus?*, 54
Astroporites, 23
Atactocrinus, 37
 " ? sp., 37
Austinocrinus, 23, 49
 " *mexicanus*, 28, 49
Avicantus, 70
 " **dunbari*, 70
 axial canal, **8**, 16

Balanocrinus, 23
Baryschr, 64
 " **anostus*, **8**, 17; 64
Blothronagma, 63
 " **cinctutum*, **8**, 17; 63
Bornium, 23
 Bourgueticrinidae, 49
 Bourgueticrinina, 49
Bourgueticrinus, 23, 50
 " *alabamensis*, 50
Brachiocrinus, 13, 23
Bystrouicrinus, 23, 26

Camarocrinus, 23
 Camerata, 36
Camptocrinus, 48
 " *beaveri*, 28, 48
Catholichorhachis, 63
 " **multifaria*, 63
Cionerisma, 76
 " **exile*, 76
 cirral (-s), **8**, 10
 cirrus fragment (-s), **8**, 26
 Cladida, 38
Clematidiscus, 74
 " **denotatus*, 74
Coenocrinus, 23
 columnal articulation, types, **8**, 15
 " facet-part indices, **8**, 21
 " indices, **8**, 21
 " shape indices, **8**, 24
 columnal (-s), **8**, 10
Concretum, 23
Cophinus, 23
Crenatames, 71
 " **amicabilis*, 71
Crisantum, 23
Crotalocrinites, 17, 23
Ctenocrinus, 23
Cupressocrinites, 23
Cupressocrinus, 13
Cyathocrinites, 23
 Cyclici, **8**, 26; 37, 46, 57
Cyclocaudex, 65
 " *aptus*, 66
 " *congregalis*, 65
 " *costatus*, 66
 " *insaturatus*, 66
 " *jucundus*, 66
 " *plenus*, **8**, 17, 19, 24; 66; **10**, 11
 " **typicus*, 65
Cyclocaudiculus, 83
 " **regularis*, 83
 Cyclocharacidae, 76
Cyclocharax, 76
 " **fasciatus*, 76
 " *modestus*, 76
Cyclocion, 78
 " **distinctus*, 78
Cyclocrinus, 23
Cyclocrista, 79
 " *cheneyi*, 80
 " **lineolata*, 80
 " *martini*, 83; **10**, 13
Cyclocyclicus, 23, 26
 Cyclomischidae, 58
Cyclomischus, 58
 " *alternatus*, 59
 " **shelbyensis*, 59
 " *tennesseensis*, 59
Cyclomonile, 57
 " **monile*, 58
Cyclopagoda, 57
 " **alternata*, 57
 " *costata*, 57
 Cyclopagodidae, 57
Cyclopentagonalis, 23, 26
Cycloscapus, 83
 " **laevis*, 83
Cylostelechus, 75
 " **turratus*, 76
Cylindrocantiscus, 62
 " **fiski*, 62
Cyphostelechus, 62
 " **claudus*, 63
Cystocrinus, 23

Decacrinus, 23, 26, 35
 " **pennatus*, 35

 " *tennicrenulatus*, 35
 Dendrocrinina, 33
Desidiamphtidia, 70
 " **frondea*, 70
Desmidocrinus, 23
Dianthicoeloma, 50
 " **insuetum*, 50
 " *monile*, 51
 " *pentaporus*, 51
 Dianthicoelomatidae, 50
 Dichocrinidae, 48
Dierocalipter, 73
 " **doter*, 73
Dilanteris, 72
 " **trestes*, 72
 Diplobathrida, 36
 Disparida, 31
 Dolatocrinidae, 47
Dolatocrinus, 47
 " *avis*, 47
 " *exculptus*, 28, 47
Dorekicrinus, 23
Dunnicrinus, 50
 " **mississippiensis*, 50

Edriocrinus, 12
 Elliptici, **8**, 25; 40, 49
Elytroclon, 40
 " **elimatus*, 40
Encrinites, 23, 25
Entrochites, 23, 25
Eomyelodactylus, 23, 25
Eucalyptocrinites, 17, 46
 " *caelatus*, 28
 " ? *shelbyensis*, 26, 46
 Eucalyptocrinitidae, 46
Eucladocrinus, 9, 42
 " ? *kentuckyensis*, 42
 " ? *springeri*, 28, 42
Eugeniocrinites, 23
Euloncherostigma, 58
 " **impunitum*, 58
 Euracidae, 68
Eurax, 68
 " **ethas*, 68
 " *eugenes*, 68
 Exaesiiodiscidae, 73
Exaesiiodiscus, 73
 " **acutus*, 73
 " *minutus*, 74
 " *truncatus*, 73
Exedrodiscus, 74
 " **excussus*, 74

Fabaliium, 71
 " **fabale*, 71
 Floricyclidae, 76
Floricyclus, 76
 " *angustimargo*, 77
 " *granulosus*, **8**, 15; 77
 " **hebes*, 77

- " *kansasensis*, 77
 " *pulcher*, 77
 " *welleri*, 77
Floripila, 56
 " **florealis*, 57
Flucticharacidae, 70
Flucticharax, 70
 " **undatus*, 71

Gilbertsocrinus, 17, 38
 " *aequalis*, 39
 " *cassiope*, 39
 " *concinus*, 39
 " *vetulus*, 28, 38
 Glossary of morphological terms, 8, 27
Goniocion, 75
 " **gonimus*, 75
 " *turgidus*, 75
Goniostathmus, 75
 " **annexus*, 75
Grammocrinus, 23
Graphosterigma, 61
 " *grammodes*, 62
 " **scriptum*, 62
 " *synthetes*, 62

Haplocrinites, 23
Hattinanteris, 59
 " **indianensis*, 59
 " *regularis*, 59
Herpetocrinus, 13
 " *fletcheri*, 28
 heteromorphic pluricolumnals, 8, 11
Heterostaurus, 83
 " **belknapensis*, 83
Heterosteichus, 81
 " *jeffordsi*, 82; 10, 12
 " *keithi*, 8, 15, 24; 82
 " **texanus*, 8, 15; 81
Hexacrinites, 23
 holdfasts, 8, 26
Holocrinus, 23
 homeomorphic pluricolumnals, 8, 11
Hybochylocrinus, 12
Hyperexochus, 33
 " **immodicus*, 33

Idromecrinus, 23
Ilematerisma, 40
 " **enamma*, 40
Imperatoria, 23
Inadunata, 31
 indices (of columnals and pluricolumnals), 8, 20
 internodal (-s), 8, 14
Iocrinidae, 31
Iocrinus, 32
 " **subcrassus*, 32
Isocrinida, 48
Isocrinidae, 48
Isocrinus, 23
Isselicrinus, 23, 48
 " *bermudezi*, 28, 49

Kasachstanocrinus, 23
Kstutocrinus, 23
Kuzbassocrinus, 23, 26, 35
 " **bystrowi*, 36

Lamprosterigma, 78
 " *erathense*, 79
 " **mirificum*, 79
Lampteroocrinidae, 36
Lampteroocrinus, 36
 " sp. A, 37
 " sp. B, 37
 " **tennesseensis*, 36
latera (-us), 8, 11
Laudonomphalus, 71
 " *ornatus*, 72
 " **regularis*, 72
 " *tuberosus*, 72
Leptocarphiidae, 79
Leptocarphum, 79
 " **gracile*, 79
 " *regulare*, 79
Leseus, 23
Lichenocrinus, 23
 " *crateriformis*, 28
Lomalegnum, 61
 " **hormidium*, 61

Mariacrinus, 23
Medineocrinus, 23
Mediocrinus, 23
Melocrinites, 23
Mespilocrinus, 9
 " *konincki*, 8, 25
Millericrinida, 49
Millericrinina, 49
Millericrinus, 23
Monobathrida, 40
Mooreanteris, 66
 " *perforatus*, 71
 " sp., 71
 " **waylandensis*, 8, 15; 67
Musivocrinus, 23
Myelodactylidae, 32
Myelodactylus, 9, 23, 32
 " *ammonis*, 28, 32
Myrtillocrinus, 23

*Nevadacrinus *geniculatus*, 8, 25
Nielsenocrinus, 23
 nodal (-s), 8, 14
 noditaxes (-is), 8, 14
 noditaxial indices, 8, 21
Nothrosterigma, 83
 " **merum*, 84

Obutierinus, 23, 26
 ontogeny, 10, 6
Onychocrinus, 9
 " *diversus*, 8, 25
 " *pulaskiensis*, 8, 25
Orocrinus, 23

Pachycrinites, 23
Palermocrinus, 23
Pandocrinus, 23, 69
 " *stoloniferus*, 69
Pentacauliscidae, 51
Pentacauliscus, 52
 " **nodosus*, 52
 " **nodosus?*, 52
Pentacrinites, 23
Pentacrinus asteriscus, 28

Pentagonocyclicus, 23
Pentagonomischus, 53
 " **plebeius*, 53
 " **plebeius?*, 54
Pentagonopentagonalis, 23, 26
Pentagonopternix, 55
 " **insculptus*, 55
Pentagonostaurus, 54
 " **leptus*, 54
Pentagonostipes, 52
 " **petaloides*, 52
 Pentameri, 8, 25; 31, 33, 36, 48, 50
Pentamerostela, 56
 " **delicatula*, 56
 " *minuta*, 56
Pentamerostelidae, 55
Pentardica, 54
 " *pentagonalis*, 55
 " **rothi*, 55
 " *simplicis*, 55
Petalerisma, 53
 " **eriense*, 53
Petalocrinus, 17
 " **mirabilis*, 28
Phialocrinus, 23
Pisocrinus, 12
Platycion, 46
 " **mingusensis*, 46
Platyclonus, 43
 " **dispar*, 44
Platycrinites, 23, 41
 " *?irroratus*, 28, 41
Platycrinitidae, 40
Platyparallelus, 43
 " **parilis*, 43
Platyplateium, 44
 " *providencense*, 45
 " sp., 45
 " **texanum*, 44
Platystela, 41
 " **proiecta*, 28, 42
Plummeranteris, 78
 " **sansaba*, 78
 pluricirral (-s), 8, 10
 pluricolumnal (-s), 8, 10
Plussacrinus, 23
Podolocrinus, 23
Podolithus, 23
Poteriocrinites, 23
Preptopremnum, 80
 " *laeve*, 81
 " **rugosum*, 8, 19, 24, 25; 81; 10, 12
Proctothylacocrinidae, 34
Proctothylacocrinus, 34
 " *esseri*, 35
 " **longus?*, 34
Pterotocrinus bifurcatus, 28

Rhenocrinus, 12
Rhizocrinus, 23
Rhodocrinites, 23
Rhodocrinitidae, 37
Rhysocamax, 60
 " **cristata*, 60
 " *grandis*, 60
 " *tuberculata*, 60

- Ristnacrinus*, 23
Salagastiana, 23
Scelidiopternix, 74
 " **norops*, 74
Schizocrinus, 23
Scillus, 23
Scyphocrinites, 12
Seiocrinus, 23
Sphenocrinus, 23
Stiberostaurus, 61
 " **aestimatus*, 61
- sutures, 8, 15
Syndetocrinus, 23
Taxocrinus colleti, 8, 25
 " **communis*, 8, 25
techniques, 8, 8
Teliocrinus, 13
Tetragonocrinus, 23
Tetragonocyclicus, 23
Tetragonotetragonalis, 23, 26
Thiolliericrinus, 23
- Traumatocrinus*, 23
Trigonocyclicus, 23
Trigonotrigonalis, 23
Varii, 8, 26; 47
Vasocrinus, 23
Wanakastaurus, 53
 " **delicatus*, 53
xenomorphic crinoid columns, 8, 19
Zeravshanocrinus, 23