

# A New Species of *Dialictus* from Sombrero Island, Anguilla (Hymenoptera, Halictidae)

Michael S. Engel

*Division of Entomology, Natural History Museum, and Department of Ecology & Evolutionary Biology, 1501 Crestline Drive – Suite 140, University of Kansas, Lawrence, Kansas 66049–2811, USA*

*urn:lsid:zoobank.org:author:3714A7FF-E19E-495A-AAF9-98D2F597B757*

Corresponding author: *Michael S. Engel* (msengel@ku.edu)

---

Academic editor: *Michael Ohl* | Received 15 January 2011 | Accepted 9 March 2011 | Published 19 March 2011

---

*urn:lsid:zoobank.org:pub:48BE1134-8166-4E1D-A95A-CFD0167475FD*

---

**Citation:** Engel MS (2011) A New Species of *Dialictus* from Sombrero Island, Anguilla (Hymenoptera, Halictidae). ZooKeys 86: 61–68. doi: 10.3897/zookeys.86.909

---

## Abstract

A new species of *Lasioglossum* Curtis subgenus *Dialictus* Robertson (Halictinae, Halictini) is described and figured from a series of female and males collected on Sombrero Island, Anguilla; the northernmost island of the Lesser Antilles. *Lasioglossum* (*Dialictus*) *sombrerense* **sp. n.** is distinguished from its congeners and the name made available for a forthcoming work on the arthropod diversity of Sombrero Island.

## Keywords

Apoidea, Anthophila, Halictidae, Halictinae, Halictini, taxonomy, Anguilla, West Indies

## Introduction

Since the overview by Eickwort (1988), work has steadily increased on the West Indian halictine fauna (e.g., Snelling 2005; Genaro 2006, 2007, 2008; Genaro and Franz 2008; Smith-Pardo 2009) and it is greatly hoped that a new synthesis of this entire region is undertaken. The purpose of the present contribution is to provide a name for a new species of halictine bee on Sombrero Island, Anguilla. The name is established here so that it may be used in a forthcoming study of

Sombrero Island arthropod diversity by Mike Ivie and Justin Runyon (Ivie and Runyon, in prep.).

Institutional acronyms used in the sections on the type material are: SEMC, Snow Entomological Collection, Division of Entomology, University of Kansas Natural History Museum, Lawrence; NHML, Department of Entomology, The Natural History Museum, London; FSCA, Museum of Entomology, Florida State Collection of Arthropods, Gainesville; and MTEC, Montana Entomology Collection, Montana State University, Bozeman. Morphological terminology generally follows that of Engel (2001, 2009) and Michener (2007), while the format for the description loosely follows that of Engel (2000) and Engel et al. (2007). Measurements were taken using an ocular micrometer on an Olympus SZX-12 stereomicroscope while photomicrographs were prepared with a Nikon D1x digital camera attached to an Infinity K-2 long-distance microscopic lens. Measurements are of the holotype, with ranges for paratypes provided in parentheses.

## Systematics

### Genus *Lasioglossum* Curtis

#### Subgenus *Dialictus* Robertson

#### *Lasioglossum (Dialictus) sombreroense* Engel, sp. n.

urn:lsid:zoobank.org:act:380A073D-2F45-4933-9B64-697690DF6D03

Figs 1–9

**Holotype.** ♀, Sombrero, West Indies: 18°35.19'N, 63°25.63'W, 12 November 1999, M.A. Ivie & J.B. Runyon (SEMC).

**Paratypes.** 5 ♀♀ Sombrero, West Indies: 18°35.19'N, 63°25.63'W, 12 November 1999, M.A. Ivie & J.B. Runyon (SEMC, MTEC, NHML); 5 ♀♀ 1 ♂, Sombrero, West Indies: 18°35.171'N, 63°25.631'W, 12–13 November 1999, M.A. Ivie & J.B. Runyon (SEMC, MTEC, FSCA).

**Diagnosis.** Small bees, around 5 mm or less in total length; integument of head and mesosoma dark metallic green with bluish and coppery tints in places (Figs 1–5); apical half of clypeus of male and female dark brown to brown (Figs 3, 6); flagellum two-toned, brown to dark brown above, honey brown below (Figs 1, 3, 4, 6); female metasoma with abundant appressed tomentum on metasomal terga III–V and basally and laterally on tergum II (Fig. 1); male face densely covered with plumose appressed to subappressed setae (Fig. 6); and male genitalia as in figures 7–9.

**Description.** *Female:* Total body length 4.93 mm (4.03–5.00 mm); forewing length 2.83 mm (2.53–2.86 mm). Head longer than wide, length 1.32 mm (1.23–1.34 mm), width 1.22 mm (1.13–1.22 mm); upper interorbital distance 0.77 mm (0.73–0.77 mm); lower interorbital distance 0.66 mm (0.60–0.66 mm). Mandible with weak subapical tooth; labrum with weak apical callosity; majority of clypeus




**Figures 1–3.** Female of *Lasioglossum (Dialictus) sombreroense* Engel, sp. n. **1** Lateral habitus **2** Dorsal habitus **3** Facial aspect.

below lower tangent of compound eyes; malar space linear. Intertegular distance 0.89 mm (0.81–0.91 mm); mesoscutellum slightly longer than metanotum, about as long as basal area of propodeum. Forewing with distal venation weakened (1rs-m, 2rs-m, and 2m-cu); basal vein 2–2.5 times distad cu-a; combined lengths of second and third submarginal cells slightly less than length of first submarginal cell; second submarginal cell slightly narrowed anteriorly, anterior border along Rs about as long as that of third submarginal cell along Rs; hind wing with distal hamuli arranged 2-1-2. Inner metatibial spur with three distinct branches distributed along rachis of spur and decreasing in size and angle relative to rachis from proximal to apical, sometimes with a minute fourth branch apically and poorly differentiated from rachis.

Mandible outer surface smooth with scattered minute punctures; labrum smooth with scattered minute punctures, punctures more sparse leading up to apical callosity; clypeus with apical half smooth and with sparse shallow coarse punctures, basally faintly imbricate with small punctures separated by 1–2 times a puncture width; supra-clypeal area imbricate with small punctures separated by a puncture width except along border with clypeus integument between punctures smooth to faintly imbricate; face below level of antennal toruli imbricate with small punctures, punctures separated by less than a puncture width to nearly contiguous except bordering compound eye and clypeus integument smooth to faintly imbricate and punctures separated by a puncture width or less; face above level of antennal toruli with small punctures separated by less than a puncture width, integument between punctures, where evident, faintly imbricate; punctures separated by a puncture width or less in parocular area and between lateral ocelli, integument between punctures smooth; vertex with scattered small punctures; gena with small punctures separated by a puncture width or less, integument smooth to faintly imbricate; postgena imbricate. Pronotum imbricate; mesoscutum imbricate with small punctures separated by a puncture width or less, such punctures more faint anteromedially and becoming separated by 1–2 times a puncture width; punctures outside of parapsidal lines separated by a puncture width or less; punctures along posterior border separated by less than a puncture width; mesoscutellum imbricate with small punctures separated by 1–2 times a puncture width on disc, punctures more closely spaced along borders; metanotum with minute punctures separated by less than a puncture width, integument between punctures apparently smooth; pleura with small punctures separated by less than a puncture width in hypoepimeral area, integument between punctures smooth to faintly imbricate; remainder of pleuron with slightly larger punctures separated by less than a puncture width although becoming more widely spaced ventrally and ventroposteriorly, integument between punctures faintly imbricate; metepisternum faintly imbricate to imbricate with punctures separated by less than a puncture width; basal area of propodeum strongly imbricate to nearly granulose, with irregular striae radiating from basal margin, such striae stronger laterally, medially becoming more rugulose apically; lateral surface of propodeum smooth to faintly imbricate with small punctures separated by less than a puncture width; posterior surface of propodeum imbricate. Metasomal terga faintly imbricate with scattered minute shallow punctures, punctures separated by 2–3 times a puncture width, becoming sparser and fainter in apical margins; sterna faintly imbricate.

Integument of head and mesosoma dark metallic green, with some faint bluish and or coppery tints except labiomaxillary complex and labrum dark brown; apical half of clypeus brown to dark brown; mandible dark brown at base, blending to honey brown medially, then to reddish brown at apex; scape, pedicel, and first flagellomere dark brown, remainder of flagellum brown to dark brown above and honey brown ventrally. Pronotal lobe honey brown; tegula honey brown and semi-translucent; legs dark brown except lighter on tarsi and at femorotibial joints. Wing membranes hyaline, veins generally honey brown except C, Sc+R, and R distad pterostigma brown. Metasoma generally dark brown except apical margins of terga semi-translucent and



**Figures 4–6.** Male of *Lasioglossum (Dialictus) sombreroense* Engel, sp. n. **4** Lateral habitus **5** Dorsal habitus. **6** Facial aspect.

lighter brown, pseudopygidial area lighter brown; sterna brown to dark brown with narrow lighter apical margins.

Pubescence generally white (Figs 1–6). Head and mesosoma with scattered short to long, branched, suberect to erect setae, not obscuring integument; such setae more numerous on face around and below level of antennal toruli, and on vertex, gena, pronotal dorsal ridge, around pronotal lobe, on metanotum, pleura, and lateral and posterior surfaces of propodeum. Metasomal terga III–V with abundant plumose, appressed tomentum (Fig. 1), similar tomentum on basally and laterally on second metasomal tergum, first metasomal tergum without such tomentum but with sparse, suberect to erect setae, such setae minute over disc, becoming longer laterally, such setae appressed

and posterolaterally-directed basally on anterior-facing surface; such suberect to erect setae scattered on remaining metasomal terga but most prominent and numerous laterally and on more apical terga; sterna with areas of longer erect setae apically on discs and margins.

*Male:* As described for the female except in usual gender differences and as follows: Total body length 5.17 mm; forewing length 2.63 mm. Head longer than wide, length 1.21 mm, width 1.09 mm; upper interorbital distance 0.68 mm; lower interorbital distance 0.46 mm. Mandible simple; labrum transverse, without callosity; apical half of clypeus below lower tangent of compound eyes. Intertegular distance 0.73 mm. Forewing second submarginal cell with anterior border along Rs much shorter than that of third submarginal cell along Rs. Inner metatibial spur simple. Male genitalia as in figures 7–9.


Face below level of antennal toruli imbricate with small punctures, punctures separated by less than a puncture width to nearly contiguous; face above level of antennal toruli with small punctures separated by less than a puncture width, integument between punctures, where evident, faintly imbricate or smooth; punctures separated by less than a puncture width in parocular area and between lateral ocelli, integument between punctures smooth. Punctuation of mesosoma coarser than in female; mesoscutum imbricate with small punctures separated by a puncture width, such punctures more faint anteromedially and becoming more widely spaced; punctures outside of parapsidal lines and along posterior border separated by less than a puncture width.

Mandible largely honey brown with reddish brown apex; scape and pedicel dark brown, flagellum brown to dark brown above and honey brown ventrally.

Face densely covered in appressed to subappressed, plumose setae, largely obscuring integument (Fig. 6); metasomal terga without abundant plumose, appressed tomentum (Fig. 4) except for thin patches basolaterally on II and III.

**Etymology.** The specific epithet is based on the name of the island to which the species was captured, Sombrero Island (a.k.a. Hat Island), Anguilla, West Indies, northernmost island of the Lesser Antilles.

**Comments.** The new species has some similarities to the widespread *Lasioglossum (Dialictus) parvum* (Cresson). Both are of relatively similar proportions and coloration but the antenna in *L. parvum* is not so strikingly two-toned, the integument tends to have a more bluish cast, the apex of the clypeus is more dark brown, the abdomen lacks the dense covering of tomentum observed in *L. sombrense* (e.g., Fig. 1), the mesoscutal punctures are more widely spaced (separated by a puncture width or more on disc), and the propodeum is much more strongly rugosostriate throughout and reaching to the margins, particularly laterally where such striae extend onto the upper portions of the lateral surfaces. In other small West Indian species such as *L. (D.) buskiellum* (Cockerell), the abdomen is darker with reddish brown apical margins to the terga, the tegulae are reddish brown, the apical abdominal terga are not obscured by tomentum, the head is broader, and the propodeum has a finer file-like striation (Cockerell, 1915), and in *L. (D.) gundlachii* (Baker) the punctuation is finer and more widely spaced (Baker, 1906).



**Figures 7–9.** Male genitalia of *Lasioglossum (Dialictus) sombreroense* Engel, sp. n. **7** Dorsal aspect **8** Ventral aspect **9** Lateral aspect.

## Acknowledgements

I am grateful to Michael Ivie and Justin B. Runyon, Montana State University, for making this material available for study and permitting deposition of material in SEMC, NHML, and FSCA; to Charles D. Michener, University of Kansas, for assistance and advice; to two anonymous reviewers for their thoughtful comments; and to Ismael A. Hinojosa-Díaz, University of Kansas, for taking the photomicrographs reproduced herein. This is a contribution of the Division of Entomology, University of Kansas Natural History Museum.

## References

- Baker CF (1906) Halictinae de Cuba. In: Primer Informe Anual de la Estación Central Agronómica de Cuba. La Universal de Ruiz y Hermano, Havana, 253–274. [total volume pages 430 pp., +52 pls.]
- Cockerell TDA (1915) Descriptions and records of bees—LXVIII. *Annals and Magazine of Natural History*, Series 8 16: 1–9.
- Eickwort GC (1988) Distribution patterns and biology of West Indian sweat bees (Hymenoptera: Halictidae). In: Liebherr JK (Ed) *Zoogeography of Caribbean Insects*. Cornell University Press, Ithaca, 231–253. [total volume pages ix+[i]+285 pp.]

- Engel MS (2000) A new *Lasioglossum* from the Juan Fernandez Islands (Hymenoptera: Halictidae). *Revista Chilena de Entomología* 27: 5–10.
- Engel MS (2001) A monograph of the Baltic amber bees and evolution of the Apoidea (Hymenoptera). *Bulletin of the American Museum of Natural History* 259: 1–192.
- Engel MS (2009) Revision of the bee genus *Chlerogella* (Hymenoptera, Halictidae), Part I: Central American species. *ZooKeys* 23: 47–75.
- Engel MS, Hinojosa-Díaz IA, Yáñez-Ordóñez O (2007) The *Augochlora*-like *Dialictus* from Guatemala and southern México (Hymenoptera: Halictidae). *Acta Zoológica Mexicana* 23(3): 125–134.
- Genaro JA (2006) A history of systematic studies of the bees of Cuba (Insecta: Hymenoptera, Anthophila). *Zootaxa* 1195: 39–60.
- Genaro JA (2007) Las abejas (Hymenoptera: Apoidea: Anthophila) de la Hispaniola, Antillas. *Boletín Sociedad Entomológica Aragonesa* 40: 247–254.
- Genaro JA (2008) Origins, composition and distribution of the bees of Cuba (Hymenoptera: Apoidea: Anthophila). *Insecta Mundi* 52: 1–16.
- Genaro JA, Franz NM (2008) The bees of Greater Puerto Rico (Hymenoptera: Apoidea: Anthophila). *Insecta Mundi* 40: 1–24.
- Michener CD (2007) *The Bees of the World* [2<sup>nd</sup> Edition]. Johns Hopkins University Press; Baltimore, MD; xvi+[i]+953 pp., +20 pls.
- Smith-Pardo AH (2009) A new species of *Habralictus* (Hymenoptera, Halictidae) from the Island of Grenada (Lesser Antilles) with comments on the insular species of the genus. *ZooKeys* 27: 51–58.
- Snelling RR (2005) Wasps, ants, and bees: aculeate Hymenoptera. In: Lazell JD (Ed) *Island: Fact and Theory in Nature*. University of California Press, Berkeley, 283–296. [total volume pages xx+382 pp.]