

## Measurement of the Semileptonic Charge Asymmetry using $B_s^0 \rightarrow D_s \mu X$ Decays

V. M. Abazov,<sup>32</sup> B. Abbott,<sup>69</sup> B. S. Acharya,<sup>26</sup> M. Adams,<sup>46</sup> T. Adams,<sup>44</sup> G. D. Alexeev,<sup>32</sup> G. Alkhazov,<sup>36</sup> A. Alton,<sup>58,\*</sup> G. Alverson,<sup>57</sup> A. Askew,<sup>44</sup> S. Atkins,<sup>55</sup> K. Augsten,<sup>7</sup> C. Avila,<sup>5</sup> F. Badaud,<sup>10</sup> L. Bagby,<sup>45</sup> B. Baldin,<sup>45</sup> D. V. Bandurin,<sup>44</sup> S. Banerjee,<sup>26</sup> E. Barberis,<sup>57</sup> P. Baringer,<sup>53</sup> J. F. Bartlett,<sup>45</sup> U. Bassler,<sup>15</sup> V. Bazterra,<sup>46</sup> A. Bean,<sup>53</sup> M. Begalli,<sup>2</sup> L. Bellantoni,<sup>45</sup> S. B. Beri,<sup>24</sup> G. Bernardi,<sup>14</sup> R. Bernhard,<sup>19</sup> I. Bertram,<sup>39</sup> M. Besançon,<sup>15</sup> R. Beuselinck,<sup>40</sup> P. C. Bhat,<sup>45</sup> S. Bhatia,<sup>60</sup> V. Bhatnagar,<sup>24</sup> G. Blazey,<sup>47</sup> S. Blessing,<sup>44</sup> K. Bloom,<sup>61</sup> A. Boehnlein,<sup>45</sup> D. Boline,<sup>66</sup> E. E. Boos,<sup>34</sup> G. Borissov,<sup>39</sup> T. Bose,<sup>56</sup> A. Brandt,<sup>72</sup> O. Brandt,<sup>20</sup> R. Brock,<sup>59</sup> A. Bross,<sup>45</sup> D. Brown,<sup>14</sup> J. Brown,<sup>14</sup> X. B. Bu,<sup>45</sup> M. Buehler,<sup>45</sup> V. Buescher,<sup>21</sup> V. Bunichev,<sup>34</sup> S. Burdin,<sup>39,†</sup> C. P. Buszello,<sup>38</sup> E. Camacho-Pérez,<sup>29</sup> B. C. K. Casey,<sup>45</sup> H. Castilla-Valdez,<sup>29</sup> S. Caughron,<sup>59</sup> S. Chakrabarti,<sup>66</sup> D. Chakraborty,<sup>47</sup> K. M. Chan,<sup>51</sup> A. Chandra,<sup>74</sup> E. Chapon,<sup>15</sup> G. Chen,<sup>53</sup> S. Chevalier-Théry,<sup>15</sup> D. K. Cho,<sup>71</sup> S. W. Cho,<sup>28</sup> S. Choi,<sup>28</sup> B. Choudhary,<sup>25</sup> S. Cihangir,<sup>45</sup> D. Claes,<sup>61</sup> J. Clutter,<sup>53</sup> M. Cooke,<sup>45</sup> W. E. Cooper,<sup>45</sup> M. Corcoran,<sup>74</sup> F. Couderc,<sup>15</sup> M.-C. Cousinou,<sup>12</sup> A. Croc,<sup>15</sup> D. Cutts,<sup>71</sup> A. Das,<sup>42</sup> G. Davies,<sup>40</sup> S. J. de Jong,<sup>30,31</sup> E. De La Cruz-Burelo,<sup>29</sup> F. Déliot,<sup>15</sup> R. Demina,<sup>65</sup> D. Denisov,<sup>45</sup> S. P. Denisov,<sup>35</sup> S. Desai,<sup>45</sup> C. Deterre,<sup>15</sup> K. DeVaughan,<sup>61</sup> H. T. Diehl,<sup>45</sup> M. Diesburg,<sup>45</sup> P. F. Ding,<sup>41</sup> A. Dominguez,<sup>61</sup> A. Dubey,<sup>25</sup> L. V. Dudko,<sup>34</sup> D. Duggan,<sup>62</sup> A. Duperrin,<sup>12</sup> S. Dutt,<sup>24</sup> A. Dyshkant,<sup>47</sup> M. Eads,<sup>61</sup> D. Edmunds,<sup>59</sup> J. Ellison,<sup>43</sup> V. D. Elvira,<sup>45</sup> Y. Enari,<sup>14</sup> H. Evans,<sup>49</sup> A. Evdokimov,<sup>67</sup> V. N. Evdokimov,<sup>35</sup> G. Facini,<sup>57</sup> L. Feng,<sup>47</sup> T. Ferbel,<sup>65</sup> F. Fiedler,<sup>21</sup> F. Filthaut,<sup>30,31</sup> W. Fisher,<sup>59</sup> H. E. Fisk,<sup>45</sup> M. Fortner,<sup>47</sup> H. Fox,<sup>39</sup> S. Fuess,<sup>45</sup> A. Garcia-Bellido,<sup>65</sup> J. A. García-González,<sup>29</sup> G. A. García-Guerra,<sup>29,‡</sup> V. Gavrilov,<sup>33</sup> P. Gay,<sup>10</sup> W. Geng,<sup>12,59</sup> D. Gerbaudo,<sup>63</sup> C. E. Gerber,<sup>46</sup> Y. Gershtein,<sup>62</sup> G. Ginther,<sup>45,65</sup> G. Golovanov,<sup>32</sup> A. Goussiou,<sup>76</sup> P. D. Grannis,<sup>66</sup> S. Greder,<sup>16</sup> H. Greenlee,<sup>45</sup> G. Grenier,<sup>17</sup> Ph. Gris,<sup>10</sup> J.-F. Grivaz,<sup>13</sup> A. Grohsjean,<sup>15,§</sup> S. Grünendahl,<sup>45</sup> M. W. Grünewald,<sup>27</sup> T. Guillemin,<sup>13</sup> G. Gutierrez,<sup>45</sup> P. Gutierrez,<sup>69</sup> S. Hagopian,<sup>44</sup> J. Haley,<sup>57</sup> L. Han,<sup>4</sup> K. Harder,<sup>41</sup> A. Harel,<sup>65</sup> J. M. Hauptman,<sup>52</sup> J. Hays,<sup>40</sup> T. Head,<sup>41</sup> T. Hebbeker,<sup>18</sup> D. Hedin,<sup>47</sup> H. Hegab,<sup>70</sup> A. P. Heinson,<sup>43</sup> U. Heintz,<sup>71</sup> C. Hensel,<sup>20</sup> I. Heredia-De La Cruz,<sup>29</sup> K. Herner,<sup>58</sup> G. Hesketh,<sup>41,||</sup> M. D. Hildreth,<sup>51</sup> R. Hirosky,<sup>75</sup> T. Hoang,<sup>44</sup> J. D. Hobbs,<sup>66</sup> B. Hoeneisen,<sup>9</sup> J. Hogan,<sup>74</sup> M. Hohlfeld,<sup>21</sup> I. Howley,<sup>72</sup> Z. Hubacek,<sup>7,15</sup> V. Hynek,<sup>7</sup> I. Iashvili,<sup>64</sup> Y. Ilchenko,<sup>73</sup> R. Illingworth,<sup>45</sup> A. S. Ito,<sup>45</sup> S. Jabeen,<sup>71</sup> M. Jaffré,<sup>13</sup> A. Jayasinghe,<sup>69</sup> M. S. Jeong,<sup>28</sup> R. Jesik,<sup>40</sup> K. Johns,<sup>42</sup> E. Johnson,<sup>59</sup> M. Johnson,<sup>45</sup> A. Jonckheere,<sup>45</sup> P. Jonsson,<sup>40</sup> J. Joshi,<sup>43</sup> A. W. Jung,<sup>45</sup> A. Juste,<sup>37</sup> K. Kaadze,<sup>54</sup> E. Kajfasz,<sup>12</sup> D. Karmanov,<sup>34</sup> P. A. Kasper,<sup>45</sup> I. Katsanos,<sup>61</sup> R. Kehoe,<sup>73</sup> S. Kermiche,<sup>12</sup> N. Khalatyan,<sup>45</sup> A. Khanov,<sup>70</sup> A. Kharchilava,<sup>64</sup> Y. N. Kharzhev,<sup>32</sup> I. Kiselevich,<sup>33</sup> J. M. Kohli,<sup>24</sup> A. V. Kozelov,<sup>35</sup> J. Kraus,<sup>60</sup> S. Kulikov,<sup>35</sup> A. Kumar,<sup>64</sup> A. Kupco,<sup>8</sup> T. Kurča,<sup>17</sup> V. A. Kuzmin,<sup>34</sup> S. Lammers,<sup>49</sup> G. Landsberg,<sup>71</sup> P. Lebrun,<sup>17</sup> H. S. Lee,<sup>28</sup> S. W. Lee,<sup>52</sup> W. M. Lee,<sup>45</sup> X. Lei,<sup>42</sup> J. Lellouch,<sup>14</sup> H. Li,<sup>11</sup> L. Li,<sup>43</sup> Q. Z. Li,<sup>45</sup> J. K. Lim,<sup>28</sup> D. Lincoln,<sup>45</sup> J. Linnemann,<sup>59</sup> V. V. Lipaev,<sup>35</sup> R. Lipton,<sup>45</sup> H. Liu,<sup>73</sup> Y. Liu,<sup>4</sup> A. Lobodenko,<sup>36</sup> M. Lokajicek,<sup>8</sup> R. Lopes de Sa,<sup>66</sup> H. J. Lubatti,<sup>76</sup> R. Luna-Garcia,<sup>29,\*\*</sup> A. L. Lyon,<sup>45</sup> A. K. A. Maciel,<sup>1</sup> R. Madar,<sup>15</sup> R. Magaña-Villalba,<sup>29</sup> S. Malik,<sup>61</sup> V. L. Malyshev,<sup>32</sup> Y. Maravin,<sup>54</sup> J. Martínez-Ortega,<sup>29</sup> R. McCarthy,<sup>66</sup> C. L. McGivern,<sup>41</sup> M. M. Meijer,<sup>30,31</sup> A. Melnitchouk,<sup>60</sup> D. Menezes,<sup>47</sup> P. G. Mercadante,<sup>3</sup> M. Merkin,<sup>34</sup> A. Meyer,<sup>18</sup> J. Meyer,<sup>20</sup> F. Miconi,<sup>16</sup> N. K. Mondal,<sup>26</sup> M. Mulhearn,<sup>75</sup> E. Nagy,<sup>12</sup> M. Naimuddin,<sup>25</sup> M. Narain,<sup>71</sup> R. Nayyar,<sup>42</sup> H. A. Neal,<sup>58</sup> J. P. Negret,<sup>5</sup> P. Neustroev,<sup>36</sup> T. Nunnemann,<sup>22</sup> J. Orduna,<sup>74</sup> N. Osman,<sup>12</sup> J. Osta,<sup>51</sup> M. Padilla,<sup>43</sup> A. Pal,<sup>72</sup> N. Parashar,<sup>50</sup> V. Parihar,<sup>71</sup> S. K. Park,<sup>28</sup> R. Partridge,<sup>71,||</sup> N. Parua,<sup>49</sup> A. Patwa,<sup>67</sup> B. Penning,<sup>45</sup> M. Perfilov,<sup>34</sup> Y. Peters,<sup>41</sup> K. Petridis,<sup>41</sup> G. Petrillo,<sup>65</sup> P. Pétrouff,<sup>13</sup> M.-A. Pleier,<sup>67</sup> P. L. M. Podesta-Lerma,<sup>29,††</sup> V. M. Podstavkov,<sup>45</sup> A. V. Popov,<sup>35</sup> M. Prewitt,<sup>74</sup> D. Price,<sup>49</sup> N. Prokopenko,<sup>35</sup> J. Qian,<sup>58</sup> A. Quadt,<sup>20</sup> B. Quinn,<sup>60</sup> M. S. Rangel,<sup>1</sup> K. Ranjan,<sup>25</sup> P. N. Ratoff,<sup>39</sup> I. Razumov,<sup>35</sup> P. Renkel,<sup>73</sup> I. Ripp-Baudot,<sup>16</sup> F. Rizatdinova,<sup>70</sup> M. Rominsky,<sup>45</sup> A. Ross,<sup>39</sup> C. Royon,<sup>15</sup> P. Rubinov,<sup>45</sup> R. Ruchti,<sup>51</sup> G. Sajot,<sup>11</sup> P. Salcido,<sup>47</sup> A. Sánchez-Hernández,<sup>29</sup> M. P. Sanders,<sup>22</sup> A. S. Santos,<sup>1,‡‡</sup> G. Savage,<sup>45</sup> L. Sawyer,<sup>55</sup> T. Scanlon,<sup>40</sup> R. D. Schamberger,<sup>66</sup> Y. Scheglov,<sup>36</sup> H. Schellman,<sup>48</sup> S. Schlobohm,<sup>76</sup> C. Schwanenberger,<sup>41</sup> R. Schwienhorst,<sup>59</sup> J. Sekaric,<sup>53</sup> H. Severini,<sup>69</sup> E. Shabalina,<sup>20</sup> V. Shary,<sup>15</sup> S. Shaw,<sup>59</sup> A. A. Shchukin,<sup>35</sup> R. K. Shivpuri,<sup>25</sup> V. Simak,<sup>7</sup> P. Skubic,<sup>69</sup> P. Slattery,<sup>65</sup> D. Smirnov,<sup>51</sup> K. J. Smith,<sup>64</sup> G. R. Snow,<sup>61</sup> J. Snow,<sup>68</sup> S. Snyder,<sup>67</sup> S. Söldner-Rembold,<sup>41</sup> L. Sonnenschein,<sup>18</sup> K. Soustruznik,<sup>6</sup> J. Stark,<sup>11</sup> D. A. Stoyanova,<sup>35</sup> M. Strauss,<sup>69</sup> L. Suter,<sup>41</sup> P. Svoisky,<sup>69</sup> M. Takahashi,<sup>41</sup> M. Titov,<sup>15</sup> V. V. Tokmenin,<sup>32</sup> Y.-T. Tsai,<sup>65</sup> K. Tschann-Grimm,<sup>66</sup> D. Tsybychev,<sup>66</sup> B. Tuchming,<sup>15</sup> C. Tully,<sup>63</sup> L. Uvarov,<sup>36</sup> S. Uvarov,<sup>36</sup> S. Uzunyan,<sup>47</sup> R. Van Kooten,<sup>49</sup> W. M. van Leeuwen,<sup>30</sup> N. Varelas,<sup>46</sup> E. W. Varnes,<sup>42</sup> I. A. Vasilyev,<sup>35</sup> P. Verdier,<sup>17</sup> A. Y. Verkheev,<sup>32</sup> L. S. Vertogradov,<sup>32</sup> M. Verzocchi,<sup>45</sup> M. Vesterinen,<sup>41</sup> D. Vilanova,<sup>15</sup> P. Vokac,<sup>7</sup> H. D. Wahl,<sup>44</sup> M. H. L. S. Wang,<sup>45</sup> J. Warchol,<sup>51</sup> G. Watts,<sup>76</sup> M. Wayne,<sup>51</sup> J. Weichert,<sup>21</sup> L. Welty-Rieger,<sup>48</sup> A. White,<sup>72</sup> D. Wicke,<sup>23</sup> M. R. J. Williams,<sup>39</sup> G. W. Wilson,<sup>53</sup> M. Wobisch,<sup>55</sup> D. R. Wood,<sup>57</sup> T. R. Wyatt,<sup>41</sup> Y. Xie,<sup>45</sup> R. Yamada,<sup>45</sup> S. Yang,<sup>4</sup> W.-C. Yang,<sup>41</sup> T. Yasuda,<sup>45</sup> Y. A. Yatsunenko,<sup>32</sup> W. Ye,<sup>66</sup> Z. Ye,<sup>45</sup> H. Yin,<sup>45</sup>

K. Yip,<sup>67</sup> S. W. Youn,<sup>45</sup> J. M. Yu,<sup>58</sup> J. Zennamo,<sup>64</sup> T. Zhao,<sup>76</sup> T. G. Zhao,<sup>41</sup> B. Zhou,<sup>58</sup> J. Zhu,<sup>58</sup>  
M. Zielinski,<sup>65</sup> D. Zieminska,<sup>49</sup> and L. Zivkovic<sup>71</sup>

(D0 Collaboration)

<sup>1</sup>LAFEX, Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro, Brazil

<sup>2</sup>Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil

<sup>3</sup>Universidade Federal do ABC, Santo André, Brazil

<sup>4</sup>University of Science and Technology of China, Hefei, People's Republic of China

<sup>5</sup>Universidad de los Andes, Bogotá, Colombia

<sup>6</sup>Charles University, Faculty of Mathematics and Physics, Center for Particle Physics, Prague, Czech Republic

<sup>7</sup>Czech Technical University in Prague, Prague, Czech Republic

<sup>8</sup>Center for Particle Physics, Institute of Physics, Academy of Sciences of the Czech Republic, Prague, Czech Republic

<sup>9</sup>Universidad San Francisco de Quito, Quito, Ecuador

<sup>10</sup>LPC, Université Blaise Pascal, CNRS/IN2P3, Clermont, France

<sup>11</sup>LPSC, Université Joseph Fourier Grenoble 1, CNRS/IN2P3, Institut National Polytechnique de Grenoble, Grenoble, France

<sup>12</sup>CPPM, Aix-Marseille Université, CNRS/IN2P3, Marseille, France

<sup>13</sup>LAL, Université Paris-Sud, CNRS/IN2P3, Orsay, France

<sup>14</sup>LPNHE, Universités Paris VI and VII, CNRS/IN2P3, Paris, France

<sup>15</sup>CEA, Irfu, SPP, Saclay, France

<sup>16</sup>IPHC, Université de Strasbourg, CNRS/IN2P3, Strasbourg, France

<sup>17</sup>IPNL, Université Lyon 1, CNRS/IN2P3, Villeurbanne, France and Université de Lyon, Lyon, France

<sup>18</sup>III. Physikalisches Institut A, RWTH Aachen University, Aachen, Germany

<sup>19</sup>Physikalisches Institut, Universität Freiburg, Freiburg, Germany

<sup>20</sup>II. Physikalisches Institut, Georg-August-Universität Göttingen, Göttingen, Germany

<sup>21</sup>Institut für Physik, Universität Mainz, Mainz, Germany

<sup>22</sup>Ludwig-Maximilians-Universität München, München, Germany

<sup>23</sup>Fachbereich Physik, Bergische Universität Wuppertal, Wuppertal, Germany

<sup>24</sup>Panjab University, Chandigarh, India

<sup>25</sup>Delhi University, Delhi, India

<sup>26</sup>Tata Institute of Fundamental Research, Mumbai, India

<sup>27</sup>University College Dublin, Dublin, Ireland

<sup>28</sup>Korea Detector Laboratory, Korea University, Seoul, Korea

<sup>29</sup>CINVESTAV, Mexico City, Mexico

<sup>30</sup>Nikhef Science Park, Amsterdam, The Netherlands

<sup>31</sup>Radboud University Nijmegen, Nijmegen, The Netherlands

<sup>32</sup>Joint Institute for Nuclear Research, Dubna, Russia

<sup>33</sup>Institute for Theoretical and Experimental Physics, Moscow, Russia

<sup>34</sup>Moscow State University, Moscow, Russia

<sup>35</sup>Institute for High Energy Physics, Protvino, Russia

<sup>36</sup>Petersburg Nuclear Physics Institute, Saint Petersburg, Russia

<sup>37</sup>Institució Catalana de Recerca i Estudis Avançats (ICREA) and Institut de Física d'Altes Energies (IFAE), Barcelona, Spain

<sup>38</sup>Uppsala University, Uppsala, Sweden

<sup>39</sup>Lancaster University, Lancaster LA1 4YB, United Kingdom

<sup>40</sup>Imperial College London, London SW7 2AZ, United Kingdom

<sup>41</sup>The University of Manchester, Manchester M13 9PL, United Kingdom

<sup>42</sup>University of Arizona, Tucson, Arizona 85721, USA

<sup>43</sup>University of California Riverside, Riverside, California 92521, USA

<sup>44</sup>Florida State University, Tallahassee, Florida 32306, USA

<sup>45</sup>Fermi National Accelerator Laboratory, Batavia, Illinois 60510, USA

<sup>46</sup>University of Illinois at Chicago, Chicago, Illinois 60607, USA

<sup>47</sup>Northern Illinois University, DeKalb, Illinois 60115, USA

<sup>48</sup>Northwestern University, Evanston, Illinois 60208, USA

<sup>49</sup>Indiana University, Bloomington, Indiana 47405, USA

<sup>50</sup>Purdue University Calumet, Hammond, Indiana 46323, USA

<sup>51</sup>University of Notre Dame, Notre Dame, Indiana 46556, USA

<sup>52</sup>Iowa State University, Ames, Iowa 50011, USA

<sup>53</sup>University of Kansas, Lawrence, Kansas 66045, USA

<sup>54</sup>Kansas State University, Manhattan, Kansas 66506, USA

<sup>55</sup>Louisiana Tech University, Ruston, Louisiana 71272, USA

- <sup>56</sup>*Boston University, Boston, Massachusetts 02215, USA*  
<sup>57</sup>*Northeastern University, Boston, Massachusetts 02115, USA*  
<sup>58</sup>*University of Michigan, Ann Arbor, Michigan 48109, USA*  
<sup>59</sup>*Michigan State University, East Lansing, Michigan 48824, USA*  
<sup>60</sup>*University of Mississippi, University, Mississippi 38677, USA*  
<sup>61</sup>*University of Nebraska, Lincoln, Nebraska 68588, USA*  
<sup>62</sup>*Rutgers University, Piscataway, New Jersey 08855, USA*  
<sup>63</sup>*Princeton University, Princeton, New Jersey 08544, USA*  
<sup>64</sup>*State University of New York, Buffalo, New York 14260, USA*  
<sup>65</sup>*University of Rochester, Rochester, New York 14627, USA*  
<sup>66</sup>*State University of New York, Stony Brook, New York 11794, USA*  
<sup>67</sup>*Brookhaven National Laboratory, Upton, New York 11973, USA*  
<sup>68</sup>*Langston University, Langston, Oklahoma 73050, USA*  
<sup>69</sup>*University of Oklahoma, Norman, Oklahoma 73019, USA*  
<sup>70</sup>*Oklahoma State University, Stillwater, Oklahoma 74078, USA*  
<sup>71</sup>*Brown University, Providence, Rhode Island 02912, USA*  
<sup>72</sup>*University of Texas, Arlington, Texas 76019, USA*  
<sup>73</sup>*Southern Methodist University, Dallas, Texas 75275, USA*  
<sup>74</sup>*Rice University, Houston, Texas 77005, USA*  
<sup>75</sup>*University of Virginia, Charlottesville, Virginia 22901, USA*  
<sup>76</sup>*University of Washington, Seattle, Washington 98195, USA*

(Received 9 July 2012; revised manuscript received 14 September 2012; published 3 January 2013)

We present a measurement of the time-integrated flavor-specific semileptonic charge asymmetry in the decays of  $B_s^0$  mesons that have undergone flavor mixing,  $a_{\text{sl}}^s$ , using  $B_s^0(\bar{B}_s^0) \rightarrow D_s^\mp \mu^\pm X$  decays, with  $D_s^\mp \rightarrow \phi \pi^\mp$  and  $\phi \rightarrow K^+ K^-$ , using  $10.4 \text{ fb}^{-1}$  of proton-antiproton collisions collected by the D0 detector during Run II at the Fermilab Tevatron Collider. A fit to the difference between the time-integrated  $D_s^-$  and  $D_s^+$  mass distributions of the  $B_s^0$  and  $\bar{B}_s^0$  candidates yields the flavor-specific asymmetry  $a_{\text{sl}}^s = [-1.12 \pm 0.74(\text{stat}) \pm 0.17(\text{syst})]\%$ , which is the most precise measurement and in agreement with the standard model prediction.

DOI: [10.1103/PhysRevLett.110.011801](https://doi.org/10.1103/PhysRevLett.110.011801)

PACS numbers: 13.20.He, 11.30.Er, 14.40.Nd

$CP$  violation has been observed in the decay and mixing of neutral mesons containing strange, charm, and bottom quarks. Currently, all measurements of  $CP$  violation, either in decay, mixing, or the interference between the two, have been consistent with the presence of a single phase in the Cabibbo-Kobayashi-Maskawa matrix. An observation of anomalously large  $CP$  violation in  $B_s^0$  oscillations can indicate the existence of physics beyond the standard model (SM) [1]. Measurements of the like-sign dimuon asymmetry by the D0 Collaboration [2,3] show evidence of anomalously large  $CP$ -violating effects using data corresponding to  $9 \text{ fb}^{-1}$  of integrated luminosity. Assuming that this asymmetry originates from mixed neutral  $B$  mesons, the measured value is  $A_{\text{sl}}^b = C_d a_{\text{sl}}^d + C_s a_{\text{sl}}^s = [-0.787 \pm 0.172(\text{stat}) \pm 0.021(\text{syst})]\%$ , where  $a_{\text{sl}}^{s(d)}$  is the time-integrated flavor-specific semileptonic charge asymmetry in  $B_s^0(B_d^0)$  decays that have undergone flavor mixing and  $C_{d(s)}$  is the fraction of  $B_d^0(B_s^0)$  events. The value of  $a_{\text{sl}}^s$  is extracted from this measurement and found to be  $a_{\text{sl}}^s = (-1.81 \pm 1.06)\%$  [3]. This Letter presents an independent measurement of  $a_{\text{sl}}^s$  using the decay  $B_s^0 \rightarrow D_s^- \mu^+ X$ , where  $D_s^- \rightarrow \phi \pi^-$  and  $\phi \rightarrow K^+ K^-$  (charge conjugate states are assumed in this Letter).

The asymmetry  $a_{\text{sl}}^s$  is defined as

$$a_{\text{sl}}^s = \frac{\Gamma(\bar{B}_s^0 \rightarrow B_s^0 \rightarrow \ell^+ \nu X) - \Gamma(B_s^0 \rightarrow \bar{B}_s^0 \rightarrow \ell^- \bar{\nu} \bar{X})}{\Gamma(\bar{B}_s^0 \rightarrow B_s^0 \rightarrow \ell^+ \nu X) + \Gamma(B_s^0 \rightarrow \bar{B}_s^0 \rightarrow \ell^- \bar{\nu} \bar{X})}, \quad (1)$$

where in this analysis  $\ell = \mu$  and  $X = D_s^{(*)-}$ . This includes all decay processes of  $B_s^0$  mesons that result in a  $D_s^-$  meson and an oppositely charged muon in the final state. To study  $CP$  violation, we identify events with the decay  $B_s^0 \rightarrow D_s^- \mu^+ X$ . The flavor of the  $B_s^0$  meson at the time of decay is identified using the charge of the associated muon, and this analysis does not make use of initial-state tagging. The fraction of mixed events integrated over time is extracted using Monte Carlo (MC) simulations. We assume there is no production asymmetry between  $B_s^0$  and  $\bar{B}_s^0$  mesons, that there is no direct  $CP$  violation in the decay of  $D_s$  mesons to the indicated states or in the semileptonic decay of  $B_s^0$  mesons, and that any  $CP$  violation in  $B_s^0$  mesons only occurs in mixing. We also assume that any direct  $CP$  violation in the decay of  $b$  baryons and charged  $B$  mesons is negligible. This analysis does not make use of the decay  $D_s^- \rightarrow K^{*0} K^-$ , as used in Ref. [4], as the expected statistical uncertainty in this channel is 2.5 times worse than the decay  $D_s^- \rightarrow \phi \pi^-$ .

The value of the SM prediction for  $a_{\text{sl}}^s = (1.9 \pm 0.3) \times 10^{-5}$  [1] is negligible compared with current experimental precision. The best direct measurement of  $a_{\text{sl}}^s$  was performed by the D0 Collaboration using data corresponding to  $5 \text{ fb}^{-1}$  of integrated luminosity, giving  $a_{\text{sl}}^s = [-0.17 \pm 0.91(\text{stat})_{-0.15}^{+0.14}(\text{syst})]\%$  [4]. This Letter presents a new and improved measurement of  $a_{\text{sl}}^s$  using the full Tevatron data sample with an integrated luminosity of  $10.4 \text{ fb}^{-1}$ .

The measurement is performed using the raw asymmetry

$$A = \frac{N_{\mu^+ D_s^-} - N_{\mu^- D_s^+}}{N_{\mu^+ D_s^-} + N_{\mu^- D_s^+}}, \quad (2)$$

where  $N_{\mu^+ D_s^-}$  ( $N_{\mu^- D_s^+}$ ) is the number of reconstructed  $B_s^0 \rightarrow \mu^+ D_s^- X$  ( $\bar{B}_s^0 \rightarrow \mu^- D_s^+ X$ ) decays. The time-integrated flavor-specific semileptonic charge asymmetry in  $B_s^0$  decays that have undergone flavor mixing,  $a_{\text{sl}}^s$ , is then given by

$$a_{\text{sl}}^s F_{B_s^0}^{\text{osc}} = A - A_\mu - A_{\text{track}} - A_{KK}, \quad (3)$$

where  $A_\mu$  is the reconstruction asymmetry between positive and negatively charged muons in the detector [5],  $A_{\text{track}}$  is the asymmetry between positive and negative tracks,  $A_{KK}$  is the residual kaon asymmetry from the decay of the  $\phi$  meson, and  $F_{B_s^0}^{\text{osc}}$  is the fraction of  $D_s^- \rightarrow \phi \pi^-$  decays that originate from the decay of a  $B_s^0$  meson after a  $\bar{B}_s^0 \rightarrow B_s^0$  oscillation. The  $F_{B_s^0}^{\text{osc}}$  factor corrects the measured asymmetry for the fraction of events in which the  $B_s^0$  meson is mixed under the assumptions outlined earlier that no other physics asymmetries are present in the other  $b$ -hadron backgrounds. While the data selection, fitting models,  $A_\mu$ ,  $A_{\text{track}}$ , and  $A_{KK}$  were studied, the value of the raw asymmetry was offset by an unknown arbitrary value and any distribution that gave an indication of the value of the asymmetry was not examined.

The D0 detector has a central tracking system, consisting of a silicon microstrip tracker and a central fiber tracker, both located within a 2 T superconducting solenoidal magnet [5,6]. An outer muon system, at  $|\eta| < 2$  [7], consists of a layer of tracking detectors and scintillation trigger counters in front of 1.8 T toroidal magnets, followed by two similar layers after the toroids [8].

The data are collected with a suite of single and dimuon triggers. The selection and reconstruction of  $\mu^+ D_s^- X$  decays requires tracks with at least two hits in both the central fiber tracker and the silicon microstrip tracker. Muons are required to have hits in at least two layers of the muon system, with segments reconstructed both inside and outside the toroid. The muon track segment has to be matched to a particle found in the central tracking system that has momentum  $p > 3 \text{ GeV}/c$  and transverse momentum  $2 < p_T < 25 \text{ GeV}/c$ .

The  $D_s^- \rightarrow \phi \pi^-$ ;  $\phi \rightarrow K^+ K^-$  decay is reconstructed as follows. The two particles from the  $\phi$  decay are assumed

to be kaons and are required to have  $p_T > 0.7 \text{ GeV}/c$ , opposite charge, and a mass  $M(K^+ K^-) < 1.07 \text{ GeV}/c^2$ . The charge of the third particle, assumed to be the charged pion, has to be opposite to that of the muon with  $0.5 < p_T < 25 \text{ GeV}/c$ . The three tracks are combined to create a common  $D_s^-$  decay vertex using the algorithm described in Ref. [9]. To reduce combinatorial background, the  $D_s^-$  vertex is required to have a displacement from the  $p\bar{p}$  interaction vertex (PV) in the transverse plane with a significance of at least 4 standard deviations. The cosine of the angle between the  $D_s^-$  momentum and the vector from the PV to the  $D_s^-$  decay vertex is required to be greater than 0.9. The trajectories of the muon and  $D_s^-$  candidates are required to be consistent with originating from a common vertex (assumed to be the  $B_s^0$  decay vertex) and to have an effective mass of  $2.6 < M(\mu^+ D_s^-) < 5.4 \text{ GeV}/c^2$ , consistent with coming from a  $B_s^0$  semileptonic decay. The cosine of the angle between the combined  $\mu^+ D_s^-$  direction, an approximation of the  $B_s^0$  direction in the direction from the PV to the  $B_s^0$  decay vertex, has to be greater than 0.95. The  $B_s^0$  decay vertex has to be displaced from the PV in the transverse plane with a significance of at least 4 standard deviations. These angular criteria ensure that the  $D_s^-$  and  $\mu^+$  momenta are correlated with that of their  $B_s^0$  parent and that the  $D_s^-$  is not mistakenly associated with a random muon. If more than one  $B_s^0$  candidate passes the selection criteria in an event, then all candidates are included in the final sample.

To improve the significance of the  $B_s^0$  selection, we use a likelihood ratio taken from Refs. [10,11]. It combines several discriminating variables: the helicity angle between the  $D_s^-$  and  $K^+$  momenta in the center-of-mass frame of the  $\phi$  meson; the isolation of the  $\mu^+ D_s^-$  system, defined as  $I = p(\mu^+ D_s^-) / [p(\mu^+ D_s^-) + \sum p_i]$ , where  $p(\mu^+ D_s^-)$  is the sum of the momenta of the three tracks that make up the  $D_s^-$  meson and  $\sum p_i$  is the sum of momenta for all tracks not associated with the  $\mu^+ D_s^-$  in a cone of  $\sqrt{(\Delta\phi)^2 + (\Delta\eta)^2} < 0.5$  around the  $\mu^+ D_s^-$  direction [7]; the  $\chi^2$  of the  $D_s^-$  vertex fit; the invariant masses  $M(\mu^+ D_s^-)$ ,  $M(K^+ K^-)$ ; and  $p_T(K^+ K^-)$ .

The final requirement on the likelihood ratio variable,  $y_{\text{sel}}$ , is chosen to maximize the predicted ratio  $N_S / \sqrt{N_S + N_B}$  in a data subsample corresponding to 20% of the full data sample, where  $N_S$  is the number of signal events and  $N_B$  is the number of background events determined from the signal and sideband regions of the  $M(K^+ K^- \pi^-)$  distributions.

The  $M(K^+ K^- \pi^-)$  distribution is analyzed in bins of  $6 \text{ MeV}/c^2$ , over a mass range of  $1.7 < M(K^+ K^- \pi^-) < 2.3 \text{ GeV}/c^2$ . The number of events is extracted by fitting the data to a model using a  $\chi^2$  fit. The  $D_s^-$  meson mass distribution is well modeled by two Gaussian functions constrained to have the same mean but with different widths and relative normalizations. A second peak in the  $M(K^+ K^- \pi^-)$  distribution corresponding to the

Cabibbo-suppressed decay of the  $D^-$  meson is also similarly modeled by two Gaussian functions, and the combinatoric background is modeled by a third-order polynomial function. The number of  $D_s^\pm$  signal decays determined from the fit is  $N(\mu^\pm D_s^\mp) = 215763 \pm 1467$ , where the uncertainty is statistical.

The polarities of the toroidal and solenoidal magnetic fields are reversed on average every two weeks so that the four solenoid-toroid polarity combinations are exposed to approximately the same integrated luminosity. This allows for a cancelation of first-order effects related to instrumental asymmetries. To ensure full cancelation, the events are weighted according to the number of  $\mu^+ D_s^-$  decays for each data sample corresponding to a different configuration of the magnets' polarities. The data are then fitted to obtain the number of weighted events,  $N(\mu^\pm D_s^\mp) = 203513 \pm 1337$ . This is shown in Fig. 1, where the weighted  $M(K^+ K^- \pi^-)$  invariant mass distributions in data are compared to the signal and background fits.

The raw asymmetry [Eq. (2)] is extracted by fitting the  $M(\phi \pi^\mp)$  distribution of the  $D_s^\mp$  candidates using a  $\chi^2$  minimization. The fit is performed simultaneously, using the same models, on the sum (Fig. 1) and the difference (Fig. 2) of the  $M(\phi \pi^-)$  distribution associated with a positively charged muon and  $M(\phi \pi^+)$  distribution associated with a negatively charged muon. The functions  $W$  used to model the two distributions are

$$W_{\text{sum}} = W^{\text{sig}}(D_s) + W^{\text{sig}}(D) + W_{\text{sum}}^{\text{bg}}, \quad (4)$$

$$W_{\text{diff}} = A W^{\text{sig}}(D_s) + A_D W^{\text{sig}}(D) + A_{\text{bg}} W_{\text{sum}}^{\text{bg}}, \quad (5)$$

where  $W^{\text{sig}}(D_s)$ ,  $W^{\text{sig}}(D)$  and  $W_{\text{sum}}^{\text{bg}}$  describe the  $D_s^-$ ,  $D^-$  mass peaks and the combinatorial background, respectively. The asymmetry of the  $D^-$  mass peak is  $A_D$ , and  $A_{\text{bg}}$  is the asymmetry of the combinatorial background.




FIG. 1 (color online). The weighted  $K^+ K^- \pi^-$  invariant mass distribution for the  $\mu^+ \phi \pi^-$  sample with the solid line representing the signal fit and the dashed line showing the background fit. The lower mass peak is due to the decay  $D^\mp \rightarrow \phi \pi^\mp$ , and the second peak is due to the  $D_s^\mp$  meson decay. Note the zero suppression on the vertical axis.

The result of the fit is shown in Fig. 2 with fitted asymmetry parameters  $A = (-0.40 \pm 0.33)\%$ ,  $A_D = (-1.21 \pm 1.00)\%$ , and  $A_{\text{bg}} = (0.00 \pm 0.11)\%$ .

The  $\chi^2$  of the fit model with respect to the difference histogram is 129.7/97 degrees of freedom over the whole mass range and 34.7 for 25 bins in the mass range  $1.90 < M(\mu^+ D_s^-) < 2.05 \text{ GeV}/c^2$ , which corresponds to a  $p$  value of 9.7%. The value of the extracted raw asymmetry,  $A$ , is checked by calculating the difference between the number of  $\mu^+ D_s^-$  and  $\mu^- D_s^+$  events in the mass range  $1.92 < M(\mu^+ D_s^-) < 2.00 \text{ GeV}/c^2$  without using a fit. In this region, we observe an asymmetry of  $(-0.5 \pm 0.3)\%$ , which is consistent with the value of  $A$  extracted by the fitting procedure.

To test the sensitivity of the fitting procedure, the charge of the muon is randomized to introduce an asymmetry signal. We use a range of raw signals from  $-2.0\%$  to  $+2.0\%$  in  $0.2\%$  steps with 1000 trials performed for each step, and the result of these pseudoexperiments, each with the same statistics as the measurement, is found. In each case, the central value of the asymmetry distribution is consistent with the input value, with a fitted width of 0.33% and no observable bias. The uncertainty found in data agrees with this expected statistical sensitivity.

Systematic uncertainties in the fitting method are evaluated by making reasonable variations to the fitting procedure. The mass range of the fit is shifted from  $1.700 < M(K^+ K^- \pi^-) < 2.300 \text{ GeV}/c^2$  to  $1.724 < M(K^+ K^- \pi^-) < 2.270 \text{ GeV}/c^2$ . The functions modeling the signal,  $W^{\text{sig}}$ , are modified so that the  $D^-$  and  $D_s^-$  peaks are fitted by single Gaussian functions. The background function,  $W_{\text{sum}}^{\text{bg}}$ , is varied from a second-order polynomial function to a fifth-order polynomial function, and the asymmetry is extracted. Instead of setting the background of  $W_{\text{diff}}$  to  $A_{\text{bg}} W_{\text{sum}}^{\text{bg}}$ , the background is either set to zero, a constant, or a polynomial function of up to degree 3. The width of the mass bins is varied between 2 and 12  $\text{MeV}/c^2$ . Instead of using the fitted number of  $B_s^0$  decays per magnet




FIG. 2. The fit to the difference distribution for the data (for clarity, the data have been rebinned).

polarity to weight the events, the total number of candidates in the mass range  $1.7 < M(K^+K^-\pi^-) < 2.3 \text{ GeV}/c^2$  is used. The systematic uncertainty is assigned to be half the maximal variation in the asymmetry for each of these sources, added in quadrature. The total effect of all of these systematic sources of uncertainty is a systematic uncertainty of 0.051% on the raw asymmetry  $A$ , giving

$$A = [-0.40 \pm 0.33(\text{stat}) \pm 0.05(\text{syst})]\%. \quad (6)$$

To extract  $a_{\text{sl}}^s$  from the raw asymmetry, corrections to the charge asymmetries in the reconstruction have to be made. These corrections are described in detail in Ref. [12]. The residual detector tracking asymmetry,  $A_{\text{track}}$ , has been studied in Ref. [2] and by using  $K_S^0 \rightarrow \pi^+\pi^-$  and  $K^{*\pm} \rightarrow K_S^0\pi^\pm$  decays. No significant residual track reconstruction asymmetries are found, and no corrections for tracking asymmetries need to be applied. The tracking asymmetry of charged pions has been studied using MC simulations of the detector. The asymmetry is found to be less than 0.05%, which is assigned as a systematic uncertainty. The muon and the pion have opposite charge, so any remaining track asymmetries will cancel to first order.

Any asymmetry between the reconstruction of  $K^+$  and  $K^-$  mesons cancels as we require that the two kaons form a  $\phi$  meson. However, there is a small residual asymmetry in the momentum of the kaons produced by the decay of the  $\phi$  meson due to  $\phi$ - $f_0(980)$  interference [13]. The kaon asymmetry is measured using the decay  $K^{*0} \rightarrow K^+\pi^-$  [12] and is used to determine the residual asymmetry due to this interference,  $A_{KK} = [0.020 \pm 0.002(\text{syst})]\%$ .

The residual reconstruction asymmetry of the muon system,  $A_\mu$ , has been measured using  $J/\psi \rightarrow \mu^+\mu^-$  decays as described in Refs. [2,3,12]. This asymmetry is determined as a function of  $p_T$  and  $|\eta|$  of the muons, and the correction is obtained by a weighted average over the normalized yields, as determined from fits to the  $M(\phi\pi^-)$  distribution. The resulting correction is  $A_\mu = (0.11 \pm 0.03)\%$ , and the combined corrections are  $A_\mu + A_{\text{track}} + A_{KK} = [0.13 \pm 0.06(\text{syst})]\%$ , including the statistical uncertainties combined in quadrature.

The remaining variable required is  $F_{B_s^0}^{\text{osc}}$  (Eq. (3)), which is the only correction extracted from a MC simulation. The  $D_s^-$  signal decays can also be produced via the decay of  $B_d^0$  mesons,  $B^\pm$  mesons, and from prompt  $c\bar{c}$  production. The  $B_s^0$  ( $B_d^0$ ) mesons can oscillate to  $\bar{B}_s^0$  ( $\bar{B}_d^0$ ) states before decaying. We split these MC samples into mixed and unmixed decays. This classification is inclusive and includes most intermediate excited states of both  $B$  and  $D$  meson decays.

The MC sample is created using the PYTHIA event generator [14] modified to use EVTGEN [15] for the decay of hadrons containing  $b$  and  $c$  quarks. Events recorded in random beam crossings are overlaid over the simulated

events to quantify the effect of additional collisions in the same or nearby bunch crossings. The PYTHIA inclusive jet production model is used, and events are selected that contain at least one muon and a  $D_s^- \rightarrow \phi\pi^-$ ;  $\phi \rightarrow K^+K^-$  decay. The generated events are processed by the full simulation chain and then by the same reconstruction and selection algorithms as used to select events from data. Each event is classified based on the decay chain that is matched to the reconstructed particles.

The mean proper decay lengths of the  $b$  hadrons are fixed in the simulation to values close to the current world-average values [16]. To correct for these differences, a correction is applied to all nonprompt events in simulation, based on the generated lifetime of the  $B$  candidate, to give the appropriate world-average  $B$  meson lifetimes and measured value of the width difference  $\Delta\Gamma_s$  [17].

To estimate the effects of trigger selection and track reconstruction, we weight each event as a function of  $p_T$  of the reconstructed muon so that it matches the distribution in the data and as a function of the lifetime to ensure that the  $B$ -meson lifetimes and  $\Delta\Gamma_s$  match the world average [16].

In the case of the  $B_s^0$  meson, the time-integrated oscillation probability is essentially 50% and is insensitive to the exact value of  $\Delta M_s$ . Combining the fraction of  $B_s^0$  decays in the sample and the time-integrated oscillation probability, we find  $F_{B_s^0}^{\text{osc}} = 0.465$ .

To determine the systematic uncertainty on  $F_{B_s^0}^{\text{osc}}$ , the branching ratios and production fractions of  $B$  mesons are varied by their uncertainties. We also vary the  $B$ -meson lifetimes and  $\Delta\Gamma_s$  and use a coarser  $p_T$  binning in the  $p_T$  event weighting. The total resulting systematic uncertainty on  $F_{B_s^0}^{\text{osc}}$  is determined to be 0.017, which includes the statistical uncertainty from the MC simulation. An asymmetry of  $B_d^0$  decays of 1% would contribute 0.005% to the total asymmetry, which is negligible compared to the statistical uncertainties and therefore neglected.

The uncertainty due to the fitting procedure (0.05%) and the asymmetry corrections (0.06%) are added in quadrature and scaled by the dilution factor,  $F_{B_s^0}^{\text{osc}}$ . The effect of the uncertainty on the dilution factor is then added in quadrature, giving a total systematic uncertainty of 0.17%.

The resulting time-integrated flavor-specific semileptonic charge asymmetry is found to be

$$a_{\text{sl}}^s = [-1.12 \pm 0.74(\text{stat}) \pm 0.17(\text{syst})]\%, \quad (7)$$

superseding the previous measurement of  $a_{\text{sl}}^s$  by the D0 Collaboration [4,18] and in agreement with the SM prediction. This result can be combined with the two  $A_{\text{sl}}^b$  measurements that depend on the impact parameter of the muons [3] and the average of  $a_{\text{sl}}^d$  measurements from the  $B$  factories,  $a_{\text{sl}}^d = (-0.05 \pm 0.56)\%$  [16] (Fig. 3). As a




FIG. 3 (color online). A combination of this result with two measurements of  $A_{sl}^b$  with different muon impact parameter selections made using like-sign dimuons [3] and the average of  $a_{sl}^d$  measurements from  $B$  factories [16]. The error bands represent the  $\pm 1$  standard deviation uncertainties on each individual measurement. The ellipses represent the 1, 2, 3, and 4 standard deviation two-dimensional C.L. regions, respectively, in the  $a_{sl}^s$  and  $a_{sl}^d$  plane.

result of this combination, we obtain  $a_{sl}^s = (-1.42 \pm 0.57)\%$  and  $a_{sl}^d = (-0.21 \pm 0.32)\%$  with a correlation of  $-0.53$ , which is a significant improvement on the precision of the measurement of  $a_{sl}^d$  and  $a_{sl}^s$  obtained in Ref. [3]. These results have a probability of agreement with the SM of  $0.28 \times 10^{-2}$ , which corresponds to 3.0 standard deviations from the SM prediction.

In summary, we have presented the most precise measurement to date of the time-integrated flavor-specific semileptonic charge asymmetry,  $a_{sl}^s = [-1.12 \pm 0.74(\text{stat}) \pm 0.17(\text{syst})]\%$ , which is in agreement with the standard model prediction and the D0 like-sign dimuon result [3].

We thank the staffs at Fermilab and collaborating institutions and acknowledge support from the DOE and NSF (U.S.A.); CEA and CNRS/IN2P3 (France); MON, NRC KI, and RFBR (Russia); CNPq, FAPERJ, FAPESP, and FUNDUNESP (Brazil); DAE and DST (India); Colciencias (Colombia); CONACyT (Mexico); NRF (Korea); FOM (The Netherlands); STFC and the Royal Society (United Kingdom); MSMT and GACR (Czech Republic); BMBF and DFG (Germany); SFI (Ireland); The Swedish Research Council (Sweden); and CAS and CNSF (China).

- \*Visitor from Augustana College, Sioux Falls, SD, USA.
- †Visitor from The University of Liverpool, Liverpool, United Kingdom.
- ‡Visitor from UPIITA-IPN, Mexico City, Mexico.
- §Visitor from DESY, Hamburg, Germany.
- ||Visitor from SLAC, Menlo Park, CA, USA.
- ¶Visitor from University College London, London, United Kingdom.
- \*\*Visitor from Centro de Investigacion en Computacion-IPN, Mexico City, Mexico.
- ††Visitor from ECFM, Universidad Autonoma de Sinaloa, Culiacán, Mexico.
- ‡‡Visitor from Universidade Estadual Paulista, São Paulo, Brazil.

- [1] A. Lenz and U. Nierste, [arXiv:1102.4274](https://arxiv.org/abs/1102.4274); A. Lenz and U. Nierste, *J. High Energy Phys.* **06** (2007) 072.
- [2] V.M. Abazov *et al.* (D0 Collaboration), *Phys. Rev. D* **82**, 032001 (2010); V.M. Abazov *et al.* (D0 Collaboration), *Phys. Rev. Lett.* **105**, 081801 (2010).
- [3] V.M. Abazov *et al.* (D0 Collaboration), *Phys. Rev. D* **84**, 052007 (2011).
- [4] V.M. Abazov *et al.* (D0 Collaboration), *Phys. Rev. D* **82**, 012003 (2010).
- [5] V.M. Abazov *et al.* (D0 Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **565**, 463 (2006).
- [6] R. Angstadt *et al.* (D0 Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **622**, 298 (2010).
- [7]  $\eta = -\ln[\tan(\theta/2)]$  is the pseudorapidity and  $\theta$  is the polar angle between the track momentum and the proton beam direction.  $\phi$  is the azimuthal angle of the track.
- [8] V.M. Abazov *et al.* (D0 Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **552**, 372 (2005).
- [9] J. Abdallah *et al.* (DELPHI Collaboration), *Eur. Phys. J. C* **32**, 185 (2004).
- [10] V.M. Abazov *et al.* (D0 Collaboration), *Phys. Rev. Lett.* **97**, 021802 (2006).
- [11] G. Borisov, *Nucl. Instrum. Methods Phys. Res., Sect. A* **417**, 384 (1998).
- [12] V.M. Abazov *et al.* (D0 Collaboration), *Phys. Rev. D* **86**, 072009 (2012).
- [13] M. Starič *et al.* (Belle Collaboration), *Phys. Rev. Lett.* **108**, 071801 (2012).
- [14] T. Sjöstrand, S. Mrenna, and P.Z. Skands, *J. High Energy Phys.* **05** (2006) 026.
- [15] D.G. Lange, *Nucl. Instrum. Methods Phys. Res., Sect. A* **462**, 152 (2001); For details, see <http://www.slac.stanford.edu/~lange/EvtGen>.
- [16] D. Asner *et al.* (Heavy Flavor Averaging Group (HFAG)), [arXiv:1010.1589](https://arxiv.org/abs/1010.1589); [http://www.slac.stanford.edu/xorg/hfag/osc/PDG\\_2012/](http://www.slac.stanford.edu/xorg/hfag/osc/PDG_2012/).
- [17] R. Aaij *et al.* (LHCb Collaboration), *Phys. Rev. Lett.* **108**, 241801 (2012); R. Aaij *et al.* (LHCb Collaboration), *Phys. Rev. Lett.* **108**, 101803 (2012).
- [18] The analysis presented in this Letter has the same statistical uncertainty as the analysis presented in Ref. [4] when performed on the same data sample.