

# Phylogenetics of Hydroidolina (Hydrozoa: Cnidaria)

PAULYN CARTWRIGHT<sup>1</sup>, NATHANIEL M. EVANS<sup>1</sup>, CASEY W. DUNN<sup>2</sup>, ANTONIO C. MARQUES<sup>3</sup>,  
MARIA PIA MIGLIETTA<sup>4</sup>, PETER SCHUCHERT<sup>5</sup> AND ALLEN G. COLLINS<sup>6</sup>

<sup>1</sup>Department of Ecology and Evolutionary Biology, University of Kansas, Lawrence, KS 66049, USA, <sup>2</sup>Department of Ecology and Evolutionary Biology, Brown University, Providence RI 02912, USA, <sup>3</sup>Departamento de Zoologia, Instituto de Biociências, Universidade de São Paulo, São Paulo, SP, Brazil, <sup>4</sup>Department of Biology, Pennsylvania State University, University Park, PA 16802, USA, <sup>5</sup>Muséum d'Histoire Naturelle, CH-1211, Genève, Switzerland, <sup>6</sup>National Systematics Laboratory of NOAA Fisheries Service, NMNH, Smithsonian Institution, Washington, DC 20013, USA

*Hydroidolina* is a group of hydrozoans that includes Anthoathecata, Leptothecata and Siphonophorae. Previous phylogenetic analyses show strong support for *Hydroidolina* monophyly, but the relationships between and within its subgroups remain uncertain. In an effort to further clarify hydroidolinan relationships, we performed phylogenetic analyses on 97 hydroidolinan taxa, using DNA sequences from partial mitochondrial 16S rDNA, nearly complete nuclear 18S rDNA and nearly complete nuclear 28S rDNA. Our findings are consistent with previous analyses that support monophyly of Siphonophorae and Leptothecata and do not support monophyly of Anthoathecata nor its component subgroups, Filifera and Capitata. Instead, within Anthoathecata, we find support for four separate filiferan clades and two separate capitate clades (*Aplanulata* and *Capitata* sensu stricto). Our data however, lack any substantive support for discerning relationships between these eight distinct hydroidolinan clades.

**Keywords:** phylogenetics, Hydroidolina, Hydrozoa, Cnidaria

Submitted 30 November 2007; accepted 12 May 2008; first published online 29 July 2008

## INTRODUCTION

Hydroidolina (=Leptolina) is a clade of hydrozoans comprising Leptothecata (=Leptomedusae, Thecata), Anthoathecata (=Anthomedusae, Athecata) and Siphonophorae (Collins, 2002; Marques & Collins, 2004; Collins *et al.*, 2006). Amongst the approximately 3220 valid species of Hydroidolina (Bouillon *et al.*, 2006), there exist vast amounts of diversity in the morphologies of hydroids and medusae as well as in life cycles. Uncovering a robust phylogeny for Hydroidolina would shed insight into the patterns underlying this diversity and provide a framework for generating hypotheses concerning processes responsible for their evolution. In addition, molecular phylogenies of Hydroidolina could help serve as a guide to taxonomic classification, which has been somewhat problematic, in large part due to inconsistencies in classifications of hydroids and medusae (e.g. Bouillon, 1985, 1994).

Hydrozoan phylogenetics has seen much progress in recent years, particularly in revealing major hydrozoan lineages and questioning others. For example, phylogenetic analyses have shown that Hydrozoa comprises two well-supported, reciprocally monophyletic clades, Trachylina and Hydroidolina (Marques & Collins, 2004; Marques, 2001a; Collins, 2002; Collins *et al.*, 2006; Van Iten *et al.*, 2006). Siphonophorae is a clade (Collins, 2002; Dunn *et al.*, 2005), but its phylogenetic

position within Hydroidolina is uncertain (Collins, 2002; Collins *et al.*, 2006). Similarly, there is strong support for the monophyly of Leptothecata (Collins *et al.*, 2006; Leclère *et al.*, 2007), but no well-supported hypotheses have emerged regarding its relationship with other hydroidolinans. Molecular phylogenetic studies do not support the monophyly of Anthoathecata and instead suggest that it is a paraphyletic assemblage that has given rise to one or more hydroidolinan groups (Marques & Collins, 2000; Marques, 2001a; Collins, 2002; Collins *et al.*, 2006; Van Iten *et al.*, 2006).

Although these studies illuminated the phyletic status of the three main groups of Hydroidolina, the relationships within and between these groups remain uncertain (Collins *et al.*, 2006). In addition, with exception to studies on Kirchenpaueriidae (Peña Cantero & Marques, 1999), Corynidae (Collins *et al.*, 2005), Siphonophorae (Dunn *et al.*, 2005), Tubulariidae (Marques & Migotto, 2001), Campanulariidae (Govindarajan *et al.*, 2006), Hebellidae–Lafoeidae (Marques *et al.*, 2006) and Plumularioidea (Leclère *et al.*, 2006), relationships within component hydroidolinan groups have not been studied within a detailed phylogenetic framework. In an effort to further clarify relationships within Hydroidolina, we greatly augmented the published molecular dataset of hydroidolinan taxa using three molecular markers, the nuclear large (28S) and small (18S) subunit rDNAs and the mitochondrial large subunit rDNA (16S). We present combined phylogenetic analyses of 97 hydroidolinan taxa (plus 13 trachylina taxa as outgroups) under maximum likelihood (ML) and parsimony (MP) criteria. The augmented dataset reveals new evolutionary

**Corresponding author:**

P. Cartwright

Email: [pcart@ku.edu](mailto:pcart@ku.edu)

patterns in morphology, although a more thorough sampling is needed to further clarify these patterns. These data suggest that a molecular phylogenetic approach is promising for guiding future taxonomic classifications but further study is needed to elucidate phylogenetic patterns of the deeper nodes within Hydroidolina.

## MATERIALS AND METHODS

### Taxa sampled, DNA isolation, amplification and sequencing

The 110 hydrozoan taxa used in this study are arranged taxonomically in Table 1, including GenBank accession and museum voucher numbers. The sequences in Table 1 comprise both published and new DNA sequences generated for this study. Although most new sequences correspond to museum voucher specimens, some were included that had no associated vouchers, but for which published sequences of other markers were generated from the same DNA pool. For new sequences, genomic DNA was extracted using Qiagen DNeasy kits according to the manufacturer's protocol (QIAGEN Inc., Mississauga, ON) or a standard phenol/chloroform protocol. The latter method involved tissue digestion with proteinase K (20 mg/ml) in a lysis buffer (20 mM Tris-CL pH 8.0, 5 mM EDTA pH 8.0, 400 mM NaCl, 2% SDS), extraction with phenol/chloroform (1:1), precipitation with 2.5 vol. 95% EtOH and elution in TE or H<sub>2</sub>O.

An approximately 600 bp fragment of 16S was amplified using a modified forward primer (F1Mod: TCGACTGTTTACCAAAAACATA) and reverse primer (R2: ACGGAATGA ACTCAAATCATGTAAAG) from Cunningham & Buss (1993). Amplifications of 16S were conducted with the following thermal profile: 5 minutes (min.) at 94°C; 5 cycles of 50 seconds(s) at 94°C, 50 s at 45°C and 1 min. at 72°C; 30 cycles of 5 s at 94°C, 50 s at 50°C and 1 min. at 72°C; 10 min. at 72°C. An approximately 1.8 kb portion of the gene coding for 18S was amplified with universal eukaryotic primers as described by Medlin *et al.* (1988). Nearly complete, an approximately 3 kb portion of the gene coding for 28S was amplified and sequenced according to Evans *et al.* (2008).

All gene fragments were purified and sequenced by Cogenics, Inc. (Houston, TX) and assembled and edited using Sequencher v4.5 (Gene Code Co., 2005). Sequences for each marker were aligned using the program MUSCLE (Edgar, 2004). Regions containing alignment ambiguities were removed using Gblocks v0.91b (Castresana, 2000) with default parameters except the minimum length block was set to 5 and half the taxa were allowed to be gaps for any given position (Table 2). The three datasets were concatenated into one combined dataset.

### Phylogenetic analysis

Phylogenetic analyses were performed on individual markers and on the combined dataset using both maximum likelihood (ML) and parsimony (MP) criteria. ML searches were performed using GARLI v0.951.OsX-GUI (Zwickl, 2006) under an assumed GTR + I + G model with rates estimated from the data. The assumed model of nucleotide substitution was selected by using the Akaike information criterion (AIC)

as implemented in ModelTest (Posada & Crandall, 2000). For the combined dataset the ML analysis was repeated 10 times from random starting trees using default termination conditions. Each run gave identical topologies and similar likelihood scores. 100 bootstrap replications were run in GARLI v0.951.OsX-GUI (Zwickl, 2006) under the same parameters.

MP analyses were performed using PAUP\* 4.0.ob10 (Swofford, 1998). Heuristic analyses were run using 500 random addition sequences and TBR branch swapping. 100 bootstrap replications were run using 10 random addition sequences per replicate and TBR branch swapping. Most parsimonious trees were summarized as a strict consensus.

The concatenated, Gblocked DNA alignment and corresponding trees can be found in TreeBASE (<http://www.treebase.org/treebase/index.html>, accession No. S2066).

## RESULTS AND DISCUSSION

After excluding the ambiguously aligned regions, the combined dataset of nearly complete 28S rDNA, nearly complete 18S rDNA and partial 16S rDNA contained 5046 characters, 1699 of which are parsimony informative. Information about individual markers is shown in Table 2. The markers were analysed separately under a ML optimality criterion and in a combined dataset under ML and MP optimality criteria. There is incongruence in topologies between the individual markers and very little support for most of the nodes in the 16S and 18S datasets (not shown). By contrast, the 28S topology is almost identical to the combined dataset (not shown) but the combined dataset shows a higher frequency of well-supported nodes (bootstrap values >50%), than the 28S topology (not shown). Given that the 28S and combined dataset are congruent but the combined dataset gives better overall support values, we concluded that the combined dataset provides the most robust hypotheses. Thus, all subsequent discussions are confined to the analyses of the combined dataset (Figures 1 & 2).

MP and ML analyses both support the monophyly of Hydroidolina (Figures 1 & 2). Trachyline relationships are treated in detail in this volume (see Collins *et al.*) and are therefore not discussed here. The hydroidolinan taxa included in these analyses sort out into eight different monophyletic clades (Figures 1 & 2; Table 1). The composition of taxa in these clades is identical in the ML and MP analyses (Figures 1 & 2). Both optimality criteria support the monophyly of Leptothecata and Siphonophorae. In the ML and MP topologies, 'Anthoathecata' is a polyphyletic assemblage with leptothecates and siphonophores derived within anthoathecate lineages. Although it should be noted that all of the nodes separating the different anthoathecate lineages are weakly supported, consistent with previous phylogenetic analyses (Collins, 2002; Marques & Collins, 2004; Collins *et al.*, 2005, 2006; Dunn *et al.*, 2005; Van Iken *et al.*, 2006; Leclère *et al.*, 2007). The separate anthoathecate clades that emerge from both the ML and MP analyses are Aplanulata (Collins *et al.*, 2005), Capitata *sensu stricto* and four filiferan clades (Figures 1 & 2; Table 1). The composition and relationships within these major clades are discussed below.

Relationships among these major clades of Hydroidolina are uncertain. There is very little bootstrap support (<50%) in the deeper nodes under both optimality criteria and there is incongruence in the ML and MP topologies between the

**Table 1.** Taxon and sequence list. A complete list of sequences used in the analyses with GenBank accession numbers and museum voucher numbers. Those in bold represent new sequences generated for this study.

| Taxonomic hierarchy   |   | 28S             | 18S             | 16S             | Voucher         |
|-----------------------|---|-----------------|-----------------|-----------------|-----------------|
| <b>Anthoathecata</b>  |   |                 |                 |                 |                 |
| Capitata              |   |                 |                 |                 |                 |
| Cladocorynidae        | <i>Cladocoryne floccosa</i>             | EU272551        | EU272608        | AY512535        |                 |
| Corynidae             | <i>Dipurena ophiogaster</i>             | EU272560        | EU272615        | <b>EU305473</b> | KUNHM 2803      |
| Corynidae             | <i>Sarsia nipponica</i>                 | <b>EU305530</b> | <b>EU448096</b> | <b>EU448100</b> | KUNHM 2627      |
| Moerisiidae           | <i>Moerisia</i> sp.                     | AY920801        | AF358083        | AY512534        |                 |
| Pennariidae           | <i>Pennaria disticha</i>                | EU272581        | AY920762        | AM088481        |                 |
| Polyorchidae          | <i>Scrippsia pacifica</i>               | AY920804        | AF358091        | AY512551        |                 |
| Porpitidae            | <i>Porpita</i> sp.                      | AY920803        | AF358086        | AY512529        |                 |
| Porpitidae            | <i>Verella</i> sp./ <i>V. vellela</i>   | EU272597        | AF358087        | <b>EU305487</b> |                 |
| Solanderiidae         | <i>Solanderia ericopsis</i>             | EU272593        | EU272636        | AY78788         | MHNG INVE 29593 |
| Solanderiidae         | <i>Solanderia secunda</i>               | <b>EU305533</b> | <b>EU305502</b> | <b>EU305484</b> | KUNHM 2611      |
| Zanclidae             | <i>Zanclaea prolifera</i>               | EU272598        | EU272639        | <b>EU305488</b> | KUNHM 2793      |
| Aplanulata            |   |                 |                 |                 |                 |
| Candelabridae         | <i>Candelabrum cocksii</i>              | AY920796        | AY920758        | AY512520        | MHNG INVE 29591 |
| Corymorphidae         | <i>Corymorpha pendula</i>               | <b>EU305510</b> | <b>EU305494</b> | <b>EU448098</b> | KUNHM 2962      |
| Corymorphidae         | <i>Euphysora bigelowi</i>               | EU272563        | EU272618        | <b>EU448099</b> | KUNHM 2829      |
| Hydridae              | <i>Hydra circumcincta</i>               | AY026371        | AF358080        | AY512521        |                 |
| Tubulariidae          | <i>Ectopleura dumortieri</i>            | EU272561        | EU272616        | <b>EU305474</b> |                 |
| Tubulariidae          | <i>Ralpharia gorgoniae</i>              | EU272590        | EU272633        | <b>EU305482</b> | KUNHM 2778      |
| Tubulariidae          | <i>Zyzyzus calderi</i>                  | EU272599        | EU272640        | <b>EU305489</b> | KUNHM 2777      |
| Filifera I            |   |                 |                 |                 |                 |
| Eudendriidae          | <i>Eudendrium californicum</i>          | <b>EU305513</b> | <b>EU305492</b> | <b>EU305475</b> | KUNHM 2850      |
| Eudendriidae          | <i>Eudendrium capillare</i>             | <b>EU305514</b> |                 | <b>EU305476</b> | KUNHM 2625      |
| Eudendriidae          | <i>Eudendrium racemosum</i>             | EU272562        | EU272617        | AY787896        |                 |
| Filifera II           |   |                 |                 |                 |                 |
| <i>incertae sedis</i> | <i>Brinckmannia hexactinellidophila</i> | EU272550        | EU272607        | AM183123        | MHNG INVE 38148 |
| Laingiidae            | <i>Fabienna sphaerica</i>               | AY920797        | AY920767        | AM183133        | MHNG INVE 33453 |
| Proboscidiactylidae   | <i>Proboscidiactyla flavicirrata</i>    | <b>EU305527</b> | <b>EU305500</b> | <b>EU305480</b> | USNM 1074994    |
| Proboscidiactylidae   | <i>Proboscidiactyla ornata</i>          | EU272587        | EU272631        | <b>EU305481</b> | KUNHM 2767      |
| Ptilocodiidae         | <i>Hydrichthella epigorgia</i>          | EU272569        | EU272622        | <b>EU305478</b> | KUNHM 2665      |
| Filifera III          |   |                 |                 |                 |                 |
| Hydractiniidae        | <i>Clava multicornis</i>                | EU272552        | EU272609        | <b>EU305471</b> |                 |
| Hydractiniidae        | <i>Clavactinia gallensis</i>            | EU272553        | EU272610        | <b>EU448101</b> | MHNG INVE 33470 |
| Hydractiniidae        | <i>Hydractinia</i> sp.                  | <b>EU305518</b> | <b>EU305495</b> | <b>EU305477</b> | KUNHM 2876      |
| Hydractiniidae        | <i>Hydractinia symbiolongicarpus</i>    | EU272568        | EU272621        |                 |                 |
| Hydractiniidae        | <i>Podocoryne carnea</i>                | AY920802        | AF358092        | AY512513        |                 |
| Stylasteridae         | <i>Adelopora crassilabrum</i>           | EU272541        | EU272642        |                 | USNM 1027760    |
| Stylasteridae         | <i>Conopora anthohelia</i>              | <b>EU305509</b> |                 |                 |                 |
| Stylasteridae         | <i>Crypthelia cryptotrema</i>           | EU272558        | EU272641        |                 | USNM 1027758    |
| Stylasteridae         | <i>Lepidopora microstylus</i>           | EU272572        | EU272644        |                 | USNM 1027724    |
| Stylasteridae         | <i>Pseudocrypthelia pachypoma</i>       | EU272589        | EU272643        |                 | USNM 1027728    |
| Filifera IV           |   |                 |                 |                 |                 |
| Bougainvilliidae      | <i>Dicoryne conybeari</i>               | EU272559        | EU272614        | AM183141        | MHNG INVE 32949 |
| Bougainvilliidae      | <i>Bimeria vestita</i>                  | EU272548        | EU272605        | AM183130        |                 |
| Bougainvilliidae      | <i>Bougainvillia carolinensis</i>       | EU272549        | EU272606        |                 |                 |
| Bougainvilliidae      | <i>Bougainvillia fulva</i>              | <b>EU305507</b> | <b>EU305490</b> | <b>EU305470</b> | KUNHM 2816      |
| Bougainvilliidae      | <i>Garveia annulata</i> /Garveia sp.    | EU272564        | AY920766        |                 | KUNHM 2860      |
| Bougainvilliidae      | <i>Koellikerina fasciculata</i>         | EU272571        | EU272623        | AM183129        |                 |
| Bougainvilliidae      | <i>Pachycordyle pusilla</i>             | EU272579        | EU272627        | AM183132        | MHNG INVE 32953 |
| Bougainvilliidae      | <i>Pruvotella (Garvia) grisea</i>       | EU272588        | EU272632        | AM183131        | MHNG INVE 34436 |
| Oceaniidae            | <i>Cordylophora caspia</i>              | EU272556        | EU272612        | <b>EU305472</b> |                 |
| Oceaniidae            | <i>Corydendrium</i> sp.                 | EU272557        | EU272613        |                 | KUNHM 2764      |
| Oceaniidae            | <i>Rhizogeton nudus</i>                 | EU272592        | EU272635        | AY787883        | MHNG INVE 35757 |
| Oceaniidae            | <i>Turritopsis dohrnii</i>              | EU272596        | EU272638        | AY787889        | MHNG INVE 29753 |
| Oceaniidae            | <i>Turritopsis nutricula</i>            | <b>EU305538</b> | <b>EU305504</b> | <b>EU305486</b> | KUNHM 2817      |
| Pandeidae             | <i>Hydrichthys boycei</i>               | EU272570        | <b>EU305496</b> | <b>EU448102</b> | MHNG INVE 37417 |
| Pandeidae             | <i>Leuckartiara octona</i>              | EU272573        | EU272624        | AM411421        |                 |
| Pandeidae             | <i>Neoturtis brevicornis</i>            | <b>EU305524</b> | <b>EU448097</b> | <b>EU448103</b> | KUNHM 002961    |
| Pandeidae             | <i>Pandea</i> sp.                       | EU272580        | AY920765        |                 |                 |
| Rathkeidae            | <i>Lizzia blondina</i>                  | EU272574        | EU272625        | AM411417        |                 |
| Rathkeidae            | <i>Rathkea octopunctata</i>             | EU272591        | EU272634        | <b>EU305483</b> | KUMIP 314321    |
| <b>Leptothecata</b>   |   |                 |                 |                 |                 |
| Conica                |   |                 |                 |                 |                 |
| Aequoreidae           | <i>Aequorea aequorea</i>                | <b>EU305505</b> | AF358076        | AY512518        |                 |
| Aequoreidae           | <i>Aequorea florida</i>                 | <b>EU305506</b> |                 |                 | USNHM PENDING   |

Continued

Table 1. Continued

| Taxonomic hierarchy  |   | 28S      | 18S      | 16S      | Voucher         |
|----------------------|---|----------|----------|----------|-----------------|
| Aequoreidae          | <i>Aequorea victoria</i>                  | AY920799 | AF358077 | EU305469 | KUNHM 2867      |
| Aequoreidae          | <i>Rhacostoma atlantica</i>               | EU305528 | EU305501 |          |                 |
| Aglaopheniidae       | <i>Aglaophenia tubiformis</i>             | EU272543 | EU272601 | AY787914 | MHNG INVE 29967 |
| Blackfordiidae       | <i>Blackfordia virginica</i>              | AY920800 | AF358078 | AY512516 |                 |
| Eirenidae            | <i>Eutima sapinhao</i>                    | EU305515 | EU305493 |          |                 |
| Haleciidae           | <i>Halecium muricatum</i>                 | EU272565 | EU272619 | AY787915 | MHNG INVE 29028 |
| Halopterididae       | <i>Halopterus minuta</i>                  | EU272567 | EU272620 | AY787912 | MHNG INVE 25073 |
| Halopterididae       | <i>Monostaechas quadridens</i>            | EU305521 | EU305497 | DQ855941 |                 |
| Hebellidae           | <i>Anthohebella parasitica</i>            | EU272545 | EU272603 | AY787918 | MHNG INVE 29762 |
| Lafoeidae            | <i>Lafoea dumosa</i>                      | EU305520 |          | AY787917 | MHNG INVE 29952 |
| Laodiceidae          | <i>Melicertissa</i> sp.                   | AY920798 | AF358075 | AY512515 |                 |
| Malagazziidae        | <i>Octophialucium indicum</i>             | EU272577 | EU272626 | AY787897 | MHNG INVE 29970 |
| Melicertidae         | <i>Melicertum octocostatum</i>            | EU272575 | AY920757 | EU305479 | USNM 1073342    |
| Mitrocomidae         | <i>Tiaropsidium kelseyi</i>               | EU305537 | AF358079 | EU305485 |                 |
| Plumulariidae        | <i>Nemertesia antennina</i>               | EU305523 | EU305498 | AY787910 | MHNG INVE 29954 |
| Plumulariidae        | <i>Plumularia hyalina</i>                 | EU305525 | EU305499 | AY787913 | MHNG INVE 25333 |
| Sertulariidae        | <i>Abietinaria filicula</i>               | EU272540 | EU272600 | AY787899 | MHNG INVE 29947 |
| Sertulariidae        | <i>Diphasia fallax</i>                    | EU305511 | EU305491 | AY787901 | MHNG INVE 29950 |
| Sertulariidae        | <i>Hydrallmania falcata</i>               | EU305519 |          | AY787900 | MHNG INVE 29948 |
| Sertulariidae        | <i>Sertularia cupressina</i>              | EU305531 |          | AY787905 | MHNG INVE 29949 |
| Sertulariidae        | <i>Sertularia perpusilla</i>              | EU305532 |          | AY787894 | MHNG INVE 29765 |
| Sertulariidae        | <i>Thuiaria thuja</i>                     | EU305536 | EU305503 | AY787908 | MHNG INVE 29951 |
| Proboscidoidea       |   |          |          |          |                 |
| Campanulariidae      | <i>Clytia noliformis</i>                  | EU272554 | EU272611 | DQ064792 |                 |
| <b>Siphonophorae</b> |   |          |          |          |                 |
| Calycophorae         |   |          |          |          |                 |
| Clausophyidae        | <i>Kephyes ovata</i>                      | EU305508 | AY937336 | AY935294 | YPM 35349       |
| Diphyidae            | <i>Sulculeolaria quadrivalvis</i>         | EU272594 | AY937353 | AY935311 | YPM 35357       |
| Hippopodiidae        | <i>Hippopodius hippopus</i>               | EU305517 | AY937341 | AY935314 | YPM 35045       |
| Prayidae             | <i>Nectadamas diomedea</i>                | EU305522 | AY937348 | AY935306 | YPM 35352       |
| Prayidae             | <i>Nectopyramis</i> sp./ <i>N. natans</i> | AY026377 | AF358068 | AY935307 |                 |
| Prayidae             | <i>Praya dubia</i>                        | EU305526 | AY937326 | AY935285 | YPM 35346       |
| Prayidae             | <i>Rosacea flaccida</i>                   | EU305529 | AY937328 |          | YPM 35041       |
| Physonectae          |   |          |          |          |                 |
| Agalmatidae          | <i>Agalma elegans</i>                     | EU272542 | AY937313 | AY935271 | YPM 35029       |
| Agalmatidae          | <i>Cordagalma cordiforme</i>              | EU272555 | AY937317 | AY935275 | YPM 35032       |
| Agalmatidae          | <i>Halistemma rubrum</i>                  | EU272566 | AY937358 | AY935316 | YPM 35359       |
| Agalmatidae          | <i>Nanomia bijuga</i>                     | EU272576 | AY937338 | AY935296 | YPM 35043       |
| Agalmatidae          | <i>Stephanomia amphitridis</i>            | EU305535 | AY937322 | AY935280 | YPM 35076       |
| Apolemiidae          | <i>Apolemia</i> sp.                       | EU272546 | AY937331 | AY935290 | YPM 35090       |
| Erennidae            | <i>Erenna</i> sp.                         | EU305512 | AY937361 | AY935319 | YPM 35362       |
| Forskaliidae         | <i>Forskalia edwardsi</i>                 | EU305516 | AY937320 | AY935278 | YPM 35036       |
| Physophoridae        | <i>Physophora hydrostatica</i>            | EU272582 | AY937342 | AY935300 | YPM 35046       |
| Rhodaliidae          | <i>Stephalia dilata</i>                   | EU305534 | AY937357 | AY935315 | YPM 35358       |
| Cystonectae          |   |          |          |          |                 |
| Physaliidae          | <i>Physalia physalis</i>                  | EU448095 | AY358065 | AY935284 | YPM 35345       |
| <b>Trachylina</b>    |   |          |          |          |                 |
| Aeginidae            | <i>Aegina citrea</i>                      | AY920789 | AF358058 | EU293997 |                 |
| Cuninidae            | <i>Solmissus marshalli</i>                | AY920790 | AF358060 |          |                 |
| Cuninidae            | <i>Solmundella bitentaculata</i>          | EU247797 | EU247812 |          | MHNG 31746      |
| Cuninidae            | <i>Solmundella bitentaculata</i>          |          |          | EU293998 | USNM 1107456    |
| Halicreatidae        | <i>Haliscera conica</i>                   | EU247797 | AF358064 | EU293981 |                 |
| Oliandiasidae        | <i>Limnocnida tanganyicae</i>             | AY920795 | AY920755 | EU293972 | USNM 1075114    |
| Oliandiasidae        | <i>Aglauroopsis aeora</i>                 | AY920793 | AY920754 | EU293973 | USNM 1073327    |
| Oliandiasidae        | <i>Astrohydra japonica</i>                | AY920794 |          | EU293975 |                 |
| Oliandiasidae        | <i>Olindias sambaquiensis</i>             | EU247809 | EU247814 |          |                 |
| Rhopalonematidae     | <i>Aglantha digitale</i>                  | AY920791 | EU247821 | EU293985 | USNM 1073329    |
| Rhopalonematidae     | <i>Aglaura hemistoma</i>                  | EU247803 | EU247818 |          | MHNG 31745      |
| Rhopalonematidae     | <i>Aglaura hemistoma</i>                  |          |          | EU293984 | KUMIP 314322    |
| Rhopalonematidae     | <i>Pantachogon haeckeli</i>               | AY920792 | AF358062 |          |                 |
| Rhopalonematidae     | <i>Pantachogon haeckeli</i>               |          |          | EU293988 | USNM 111078     |
| Rhopalonematidae     | <i>Rhopalonema velatum</i>                | EU247804 | EU247819 | EU293992 |                 |
| Tetraplatiidae       | <i>Tetraplatia volitans</i>               | DQ002502 | DQ002501 | EU293999 | KUMIP 314322    |

KUMIP, University of Kansas Museum of Invertebrate Paleontology; KUNHM, University of Kansas Natural History Museum; MHNG, Muséum d'Histoire Naturelle de Genève; YPM, Yale–Peabody Museum; USNM, US National Museum of Natural History.



**Table 2.** Summary of genetic markers used in this study.

| Marker | Primer source               | Length after gblocks (bp) (% retained) | No. of parsimony informative characters (% informative) |
|--------|-----------------------------|--|---|
| 28S    | Evans <i>et al.</i> , 2008  | 2959 (81%)                             | 969 (33%)   |
| 18S    | Medina <i>et al.</i> , 2001 | 1648 (82%)                             | 407 (25%)   |
| 16S    | Cunningham & Buss, 1993     | 439 (55%)                              | 323 (74%)   |

major clades. Given the inconclusiveness of these results, any discussion of relationships between major hydroidolinan clades would be premature. By contrast, within each of the major clades, the topologies in the ML and MP analyses are largely congruent and most of the nodes within these clades display high bootstrap support (Figures 1 & 2). Thus we focus our discussion below on the composition and relationships within these clades.

### Capitata *sensu stricto*

Capitata is traditionally defined by the presence of capitate tentacles at some stage in its life cycle (Reese, 1957; Petersen, 1990). Recent molecular phylogenetic analyses have questioned the monophyly of Capitata and instead suggest that there are two clades, Aplanulata (*sensu* Collins *et al.*, 2005) and non-Aplanulata capitates (Collins, 2002; Collins *et al.*, 2005, 2006). Our ML and MP analyses provide strong support (bootstrap values = 100 and 96 respectively) for a clade of capitates to the exclusion of aplanulata taxa. We refer to this clade as Capitata *sensu stricto* herein. Within Capitata *sensu stricto* the topologies between the ML and MP analyses are nearly identical (Figures 1 & 2). Both optimality criteria indicate support for the suborder Zancleida including Cladocorynidae, Porpitiidae and Zancleidae (*sensu* Peterson, 1990), but also including Solanderiidae. *Moerisia* and *Pennaria* together form a sister taxon to the Zancleida clade (MP; Figure 2) or as successive sister taxa (ML; Figure 1). A Corynidae + Polyorchidae clade is strongly supported under both optimality criteria (Figures 1 & 2). These topologies are largely consistent with that of Collins *et al.* (2005, 2006).

### Aplanulata

Aplanulata (Collins *et al.*, 2005) is a clade supported by previous molecular phylogenetic analyses (Collins *et al.*, 2006) and is united by the lack of a ciliated planula stage (Petersen, 1990). Our analyses of Corymorphidae, Hydridae, Candelabridae and Tubulariidae representatives provide strong support for the monophyly of Aplanulata (bootstrap values = 100 for ML and MP) (Figures 1 & 2). Although our sampling is limited, within Aplanulata, there is strong support and nearly complete congruence between ML and MP topologies and these relationships are largely consistent with that recovered from Collins *et al.* (2005) that used partial 16S data. Corymorphidae and Tubulariidae are both monophyletic and there is strong support for a Corymorphidae + Tubulariidae clade (bootstrap values = 100 for ML and MP). The *Hydra* + *Candelabrum* clade is the sister group to the rest of Aplanulata in the MP analysis (Figure 2) but are successive sister taxa in the ML analysis (Figure 1). As discussed in Collins *et al.* (2006), there are other putative Aplanulata families that await future sampling and analyses.

### Filifera I: Eudendriidae


Our MP and ML analyses provide strong support for a Eudendriidae clade (bootstrap values = 98 for ML and 85 for MP), apart from other filiferan clades (Figures 1 & 2). Eudendriidae as a clade distinct from other filiferans is supported by many synapomorphies including the absence of desmoneme nematocysts, a styloid-shaped gonophore and a trumpet-shaped hypostome (Marques, 1996). Because of these unique traits, a possible sister-group relationship of the Eudendriidae with other filiferans remains dubious (Marques, 1996, 2001b).

### Filifera II: *Fabienna/Proboscidactyla/Brinckmannia/Hydrichthella*

The monophyly of *Fabienna* + *Proboscidactyla* + *Brinckmannia* is well supported (bootstrap values = 94 for ML and 79 for MP) but the node that includes *Hydrichthella* as its sister taxon has relatively low support (bootstrap values = 57 for ML and MP). An association between the Laingiomedusae *Fabienna* and Proboscidactylidae is supported by morphological evidence including a solid ring canal and macrobasic euryteles (Schuchert, 1996). A previous molecular analysis supported this relationship (Collins *et al.*, 2006). In addition, there are a number of morphological features that support the association of *Fabienna/Proboscidactyla* with *Brinckmannia* (*Filifera incertae sedis*) and the ptilocodiid *Hydrichthella*. Schuchert & Reising (2006) argued for a close relationship between *Brinckmannia* and Proboscidactylidae based on the shape of their hydranths and 16S sequence similarity. Although the polyp stage in *Fabienna* is unknown, the other taxa share a synapomorphy of hydranths with reduced tentacles: in *Brinckmannia*, hydranths have no tentacles (Schuchert & Reising, 2006), *Hydrichthella* has no tentacles on its gastrozooids (dactylozooids have many tentacles) and *Proboscidactyla* has only two tentacles on its hydranths. Interestingly, many of the species in this group are closely associated with another invertebrate as a substrate: *Hydrichthella* is found on an octocoral, *Brinckmannia* within the tissues of a hexactinellid sponge and *Proboscidactyla* on tubes of sabellid polychaetes. In addition, although the ptilocodiid *Hydrichthella* does not have a medusa, the medusae of *Fabienna* are strikingly similar to that of another ptilocodiid species, *Thecocardium quadratum* (Collins *et al.*, 2006).

### Filifera III: Hydractiniidae/Stylasteridae

Our ML and MP analyses are congruent in identifying a clade that includes Hydractiniidae and Stylasteridae (bootstrap values = 86 for ML and <50 for MP). There is strong support for monophyly of Stylasteridae (bootstrap values = 100 for ML and 99 for MP) (Figures 1 & 2).



**Fig. 1.** Phylogenetic hypothesis among 110 hydrozoan taxa, based on a maximum likelihood criterion of a combined dataset of nearly complete 28S, nearly complete 18S and partial 16S rDNA sequences. Bootstrap values greater than 50 are indicated above nodes. The assumed model (GTR + I + G) with six substitution rates estimated from the data (A-C, 0.8735; A-G, 2.9730; A-T, 1.6586; C-G, 0.8463; C-T, 5.2641; G-T, 1.0000), an assumed proportion of invariant sites (0.5740) and a gamma shaped parameter of (0.6021).

Although the hydractiniids are monophyletic in the ML analysis (bootstrap value = 60) (Figure 1) the MP analysis places them as paraphyletic relative to the stylasterids (Figure 2). Our analyses show *Clava multicornis* as the sister taxon to the hydractiniid *Podocoryne carnea* with strong support in both ML and MP trees (bootstrap values = 94 for ML and 89 for MP). *Clava* has traditionally been placed in the family Clavidae, although Schuchert

(2001) argued, based on the similarities of *Clava* to other hydractiniids (Bouillon *et al.*, 1997), that the genus *Clava* should be moved to the hydractiniids and the other Clavidae genera moved to the nominal family Oceaniidae (Schuchert, 2004). Our analysis supports the interpretation that *Clava* is a hydractiniid.

The close relationship between the Hydractiniidae and Stylasteridae families has previously been suggested based




Fig. 2. Phylogenetic hypothesis among 110 hydrozoan taxa, based on parsimony criterion of a combined dataset of nearly complete 28S, nearly complete 18S and partial 16S rDNA sequences. Bootstrap values greater than 50 are indicated above nodes. Topology is a strict consensus of the 10 most parsimonious trees (5046 characters). Length: 13246 steps, CI=0.26; RI=0.59.

on a number of synapomorphies including polymorphic polyps and the perisarc or skeleton covered stolons (Bouillon, 1978; Petersen, 1979). Bouillon (1978) placed these families in the superfamily Hydractinoidea, which also includes Ptilocodiidae, Rathkeidae and Rhysiidae. Though we did not sample any members of Rhysiidae, our analyses do not support Hydractinoidea, as sampled Ptilocodiidae and Rathkeidae members are placed outside this clade.

**Filifera IV: Gonoproxima + Dicoryne—  
Bougainvilliidae/Oceaniidae/Pandeidae/  
Rathkeidae**

Our ML and MP analyses support the monophyly of a clade that includes representatives of Bougainvilliidae, Oceaniidae, Pandeidae and Rathkeidae. This clade has relatively weak support (bootstrap values <50 for ML and MP) and must be viewed as tentative. Within the clade the topologies are

congruent between the ML and MP analyses. Of the four families, Pandaeidae and Rathkeidae are monophyletic, whereas Bougainvilliidae and Oceaniidae are polyphyletic. Although association of these four families is somewhat surprising, they all share a striking synapomorphy. The species in all four families bear gonophores on hydrocauli, pedicels, or stolons and not on the hydranth body. The shift of the gonophores from the hydranth body to the region below is an apomorphy (Schuchert, 2001). Backed by this synapomorphy, we name this clade Gonoproxima. The 'bougainvilliid' *Dicoryne*, which is distinct from other bougainvilliid species in that it produces gonophores on blastostyles, is placed as the sister taxon to Gonoproxima in both the ML and MP analyses (Figures 1 & 2). Interestingly, many taxa included in this group have perisarc extending over the hydranth body, either as a gelatinous structure or pseudohydrotheca.

Two species within Gonoproxima, *Cordylophora caspia* (sampled here) and *Pachycordyle kubotai* (not sampled) live in fresh water. In our analyses, *Cordylophora* and *Pachycordyle*, which also contains brackish and marine species, are indicated to be close relatives under both optimality criteria, forming a clade with the bougainvilliid *Bimeria*. With denser taxon sampling within Gonoproxima and more targeted phylogenetic analyses, it should be possible to ascertain whether the fresh water habit was evolved one or more times in this clade and potentially whether freshwater species are descended from ancestors that lived in brackish environments.

## Siphonophorae

The siphonophores have historically been split into three major groups, Cystonectae, Physonectae and Calyphorae. Collins (2002) placed the cystonect *Physalia* as sister to the other included siphonophores and suggested that Physonectae may be paraphyletic with respect to the Calyphorae. A later study (Dunn *et al.*, 2005), that considered additional taxa and two genes (18S and 16S) found that cystonects form a monophyletic group that is sister to the remaining siphonophores and the paraphyly of Physonectae was recovered with significant support. Dunn *et al.* (2005) erected the name Codonophora to refer to the clade comprising taxa assigned to Physonectae and Calyphorae (i.e. the clade that is sister to Cystonectae).

Our ML and MP analyses are consistent with the findings of Collins (2002) and Dunn *et al.* (2005) in that Siphonophorae is a strongly supported monophyletic group (bootstrap values = 100 for ML and MP). Our ML and MP analysis also recovered Physonectae as paraphyletic and Calyphorae derived within this clade (Figures 1 & 2). We did not however find support for Codonophora (*sensu* Dunn *et al.*, 2005). Our ML analysis placed the cystonect *Physalia* as the sister taxon to Calyphorae (Figure 1), not as the earliest diverging member of siphonophores (Collins 2002; Dunn *et al.*, 2005). The MP analysis also recovered a probable paraphyletic Physonectae, but unlike the ML analyses, *Physalia* was nested within Calyphorae (Figure 2). Under both optimality criteria, the physonect, *Apolemia*, was the earliest diverging siphonophore. Given that we have only one cystonect representative (*Physalia*) and that its placement is dependent on optimality criteria, we view the placement of Cystonectae relative to other siphonophores as equivocal and await further study.

## Leptothecata

Our ML and MP analyses found strong support for the Leptothecata clade (bootstrap values = 100 for ML and MP) (Figures 1 & 2). Sampling was concentrated amongst the Conica subgroup, with the inclusion of only one Proboscoida representative, *Clytia noliformis*. This sampling is therefore insufficient to address the question of monophyly of its subgroups, Conica and Proboscoida.

The ML and MP topologies within Leptothecata are nearly congruent except for the placement of *Lafoea* relative to *Melicertum* (discussed below). The traditional taxonomy of Leptothecata, including the relationships of its higher groups, is largely based on similarities in the morphology of the hydrotheca and nematotheca (e.g. Bouillon, 1985, 1994). Many groups found in our analyses corroborate Bouillon's hypotheses, including the monophyly of the Plumularioidea taxa, Plumulariidae and Halopterididae, Sertulariidae and the affinities of these with Haleciidae. Recent molecular and morphological analyses also have corroborated or are consistent with these hypotheses (Leclère *et al.*, 2007).

The affinities of the Hebellidae and Lafoeidae, based on morphological characters, were investigated by Marques *et al.* (2006). Although the authors hypothesized the exclusive monophyly of each family, they considered the possibility that the families are distantly related, a finding consistent with our analyses.

Campanulinida is a group of leptothecates including many diverse families: Aequoreidae, Blackfordiidae, Eirenidae, Laodiceidae, Malagazziidae, Melicertidae and Mitrocomidae. The Campanulinida taxa belonging to Aequoreidae, Blackfordiidae, Eirenidae, Malagazziidae and Mitrocomidae are a strongly supported clade that also includes *Clytia noliformis* (bootstrap values = 100 for ML and MP). The Campanulinida belonging to Melicertidae, *Melicertum octocostatum* is the sister taxon to the rest of the Leptothecata in the MP analysis (Figure 2). This analysis corroborates the hypothesis of an early divergence of *M. octocostatum* (Collins *et al.*, 2006), a species that lacks a theca but has typical leptothecate medusae. The ML analysis places *Melicertum* + *Lafoea dumosa* as the sister taxon to the rest of Leptothecata (Figure 1).

## CONCLUSIONS

Anthoathecata represents a diverse order of hydroidolins that traditionally comprises two suborders, Filifera and Capitata (reviewed in Daly *et al.*, 2007). Although our analyses and previous molecular phylogenetic analyses (Marques & Collins, 2000; Marques, 2001a; Collins, 2002; Collins *et al.*, 2006; Van Iken *et al.*, 2006) do not support the monophyly of Anthoathecata, the dissolution or re-definition of Anthoathecata is premature and should await clarification of relationships between major hydroidolinan clades. Capitata in the traditional sense comprises two clades, the Aplanulata, recognized by the lack of a free-swimming planula (Petersen, 1990) and Capitata *sensu stricto*. Given that there is strong support for these two groups and that there is no support for the monophyly of traditional 'Capitata' in these analyses and in previous phylogenetic analyses (Collins, 2002; Collins *et al.*, 2005, 2006), the validity of Capitata in the traditional sense is questioned. If these clades



are indeed separate, then Aplanulata should be referred to as its own order, separate from Capitata *sensu stricto*. Re-defining Capitata however, should await further clarification of Hydroidolina phylogeny. Our analyses do not support the monophyly of Filifera but this too is preliminary as the nodes separating the filiferan subgroups are weakly supported (Figures 1 & 2).

The new augmented dataset used in our analyses provide support for four distinct filiferan clades. Notably, all of these clades possess compelling morphological synapomorphies; Gonoproxima is characterized by gonophores on regions of the colony proximal to the hydranth; Eudendriidae displays distinct polyp and hypostome morphology; the *Fabienna/Proboscidactyla/Brinckmannia/Hydrichthella* clade displays polyps with a reduced number of tentacles and the Hydractiniidae/Sylasteridae clade displays polymorphism. A more comprehensive sampling of hydroidolinan families should provide greater insight into these emerging patterns.

Despite increased sampling, relationships between major hydroidolinan clades remain elusive. The lack of resolution suggests that the initial radiation of Hydroidolina may have been rapid, leaving little clues regarding the sequence of hydroidolinan diversification. New molecular markers, especially if combined with other types of data, may prove helpful in resolving these deep nodes.

## ACKNOWLEDGEMENTS

We would like to thank A. Lindner for DNA samples and anonymous referees for helpful comments. This study was supported from grants from NSF ATOL EF-0531779 (to P.C. and A.G.C.). A.C.M. has financial support from CNPq (55.7333/2005-9, 490348/2006-8, 305735/2006-3) and FAPESP (2004/09961-4).

## REFERENCES

- Bouillon J.** (1978) Sur un nouveau genre et une nouvelle espèce de Ptilocodidae, *Hydrichthelloides reticulata* et la super-famille des Hydractinoidea (Hydroida – Athecata). *Steenstrupia* 5, 53–67.
- Bouillon J.** (1985) Essai de classification des hydropolypes–hydroméduses (Hydrozoa – Cnidaria). *Indo-Malayan Zoology* 2, 29–243.
- Bouillon J.** (1994) Classe des hydrozoaires (Hydrozoa Owen, 1843). In P.P. Grassé (ed.) *Traité de Zoologie T III fasc. 2 Cnidaires*. Paris: Masson, pp. 29–416.
- Bouillon J., Gravili C., Pagès F., Gili J.M. and Boero F.** (2006) *An introduction to Hydrozoa*. Paris: Publications Scientifiques du Muséum, Paris.
- Bouillon J., Medel D. and Peña Cantero A.L.** (1997) The taxonomic status of the genus *Stylactaria* Stechow, 1921 (Hydroidomedusae, Anthomedusae, Hydractiniidae), with the description of a new species. *Scientia Marina* 61, 471–486.
- Castresana J.** (2000) Selection of conserved blocks from multiple alignments for their use in phylogenetic analysis. *Molecular Biology and Evolution* 17, 540–552.
- Collins A.G.** (2002) Phylogeny of Medusozoa and the evolution of cnidarian life cycles. *Journal of Evolutionary Biology* 15, 418–432.
- Collins A.G., Schuchert P., Marques A.C., Jankowski T., Medina M. and Schierwater B.** (2006) Medusozoan phylogeny and character evolution clarified by new large and small subunit rDNA data and an assessment of the utility of phylogenetic mixture models. *Systematic Biology* 55, 97–115.
- Collins A.G., Winkelmann S., Hadrys H. and Schierwater B.** (2005) Phylogeny of Capitata and Corynidae (Cnidarian, Hydrozoa) in light of mitochondrial 16S rDNA data. *Zoologica Scripta* 34, 91–99.
- Cunningham C.W. and Buss L.W.** (1993) Molecular evidence for multiple episodes of paedomorphosis in the family Hydractiniidae. *Biochemical Systematics and Ecology* 21, 57–69.
- Daly M., Brugler M.R., Cartwright P., Collins A.G., Dawson M.N., Fautin D.G., France S.C., McFadden C.S., Opreko D.M., Rodrigues E., Romano S.L. and Stake J.L.** (2007) The phylum Cnidaria: a review of phylogenetic patterns and diversity 300 years after Linnaeus. *Zootaxa* 1668, 127–182.
- Dunn C.W., Pugh P.R. and Haddock S.H.D.** (2005) Molecular phylogenetics of siphonophora (Cnidaria), with implications for the evolution of functional specialization. *Systematic Biology* 54, 916–935.
- Evans N.M., Lindner A., Raikova E.V., Collins A.G. and Cartwright P.** (2008) Phylogenetic placement of the enigmatic parasite, *Polypodium hydriforme*, within the Phylum Cnidaria. *BMC Evolutionary Biology* 8, 139.
- Edgar R.C.** (2004) MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Research* 32, 1792–1797.
- Govindarajan A.F., Boero F. and Halanich K.M.** (2006) Phylogenetic analysis with multiple markers indicates repeated loss of the adult medusa stage in Campanulariidae (Hydrozoa, Cnidaria). *Molecular Phylogenetics and Evolution* 38, 820–834.
- Leclère L., Schuchert P. and Manuel M.** (2007) Phylogeny of the Plumularioidea (Hydrozoa, Leptothecata): evolution of colonial organisation and life cycle. *Zoologica Scripta* 36, 371–394.
- Marques A.C.** (1996) A critical analysis of a cladistic study of the genus *Eudendrium* (Cnidaria: Hydrozoa), with some comments on the family Eudendriidae. *Journal of Comparative Biology* 1, 153–162.
- Marques A.C.** (2001a) Simplifying hydrozoan classification: inappropriateness of the group Hydroidomedusae in a phylogenetic context. *Contributions to Zoology* 70, 175–179.
- Marques A.C.** (2001b) O gênero *Eudendrium* (Cnidaria, Hydrozoa, Anthomedusae) no Brasil. *Papéis Avulsos de Zoologia* 41, 329–405.
- Marques A.C. and Collins A.G.** (2004) Cladistic analysis of Medusozoa and cnidarian evolution. *Invertebrate Biology* 123, 23–42.
- Marques A.C. and Migotto A.E.** (2001) Cladistic analysis and new classification of the family Tubulariidae (Hydrozoa, Anthomedusae). *Papéis Avulsos de Zoologia* 41, 465–488.
- Marques A.C., Peña Cantero A.L. and Migotto A.E.** (2006) An overview of the phylogeny of the families Lafoeidae and Hebellidae (Hydrozoa: Leptothecata): their composition and classification. *Invertebrate Systematics* 20, 43–58.
- Medina M., Collins A.G., Silberman J.D. and Sogin M.L.** (2001) Evaluating hypotheses of basal animal phylogeny using complete sequences of large and small subunit rRNA. *Proceedings of the National Academy of Science of the USA* 98, 9707–9712.
- Medlin L.H., Elwood H.J., Stickel S. and Sogin M.L.** (1988) The characterization of enzymatically amplified eukaryotic 16S-like rRNA-coding regions. *Gene* 71, 491–499.
- Peña Cantero A.L. and Marques A.C.** (1999) Phylogenetic analysis of the Antarctic genus *Oswaldella* Stechow, 1919 (Hydrozoa, Leptomedusae, Kirchenpaueriidae). *Contributions to Zoology* 68, 83–93.
- Petersen K.W.** (1979) Development of coloniality in Hydrozoa. In G. Larwood and B.R. Rosen (eds) *Biology and systematics of colonial organisms*. London: Academic Press, pp. 105–139.

- Petersen K.W.** (1990) Evolution and taxonomy in capitate hydroids and medusae (Cnidaria: Hydrozoa). *Zoological Journal of the Linnean Society* 100, 1–231.
- Posada D. and Crandall K.A.** (2000) Modeltest: testing the model of DNA substitution. *Bioinformatics* 14, 817–818.
- Rees W.J.** (1957) Evolutionary trends in the classification of capitate hydroids and medusae. *Bulletin of the British Museum (Natural History) Zoology* 4, 455–534.
- Schuchert P.** (1996) The marine fauna of New Zealand: athecate hydroids and their medusae (Cnidaria: Hydrozoa). *New Zealand Oceanographic Institute Memoir* 106, 1–159.
- Schuchert P.** (2001) Hydroids of Greenland and Iceland (Cnidaria, Hydrozoa). *Meddelelser om Grønland, Bioscience* 53, 1–184.
- Schuchert P.** (2004) Revision of the European athecate hydroids and their medusae (Hydrozoa, Cnidaria): families of Oceanidae and Pachycordylidae. *Revue Suisse de Zoologie* 111, 315–369.
- Schuchert P. and Reiswig H.M.** (2006) *Brinckmannia hexactinellidophila*, n. gen., n. sp.: a hydroid living in tissues of glass sponges of the reefs, fjords, and seamounts of Pacific Canada and Alaska. *Canadian Journal of Zoology* 84, 564–572.
- Swofford D.L.** (1998) *PAUP\*—Phylogenetic Analysis Using Parsimony (\*and other methods)*. Sunderland, MA: Sinauer.
- Van Iten H., Leme J.M., Simões M.G., Marques A.C. and Collins A.G.** (2006) Reassessment of the phylogenetic position of conulariids (?Ediacaran–Triassic) within the subphylum Medusozoa (Phylum Cnidaria). *Journal of Systematic Paleontology* 4, 109–118.
- and
- Zwickl D.J.** (2006) *Genetic algorithm approaches for the phylogenetic analysis of large biological sequence datasets under the maximum likelihood criterion*. PhD thesis. Austin: The University of Texas.

**Correspondence should be addressed to:**

Paulyn Cartwright  
Department of Ecology and Evolutionary Biology  
University of Kansas, Lawrence, KS 66049, USA  
email: pcart@ku.edu