

AzTEC/ASTE 1.1-mm survey of the AKARI Deep Field South: source catalogue and number counts

B. Hatsukade,^{1*} K. Kohno,^{2,3} I. Aretxaga,⁴ J. E. Austermann,⁵ H. Ezawa,⁶
D. H. Hughes,⁴ S. Ikarashi,² D. Iono,¹ R. Kawabe,¹ S. Khan,⁷ H. Matsuo,⁶
S. Matsuura,⁸ K. Nakanishi,⁶ T. Oshima,¹ T. Perera,⁹ K. S. Scott,¹⁰ M. Shirahata,⁸
T. T. Takeuchi,¹¹ Y. Tamura,¹ K. Tanaka,¹² T. Tosaki,¹³ G. W. Wilson⁹ and M. S. Yun⁹

¹Nobeyama Radio Observatory, Minamimaki, Minamisaku, Nagano 384-1805, Japan

²Institute of Astronomy, The University of Tokyo, 2-21-1 Osawa, Mitaka, Tokyo 181-0015, Japan

³Research Center for the Early Universe, University of Tokyo, 7-3-1 Hongo, Bunkyo, Tokyo 113-0033, Japan

⁴Instituto Nacional de Astrofísica, Óptica y Electrónica (INAOE), Aptdo. Postal 51 y 216, 72000 Puebla, Pue., Mexico

⁵Center for Astrophysics and Space Astronomy, University of Colorado, Boulder, CO 80309, USA

⁶National Astronomical Observatory of Japan, 2-21-1 Osawa, Mitaka, Tokyo 181-8588, Japan

⁷Shanghai Key Lab for Astrophysics, Shanghai Normal University, Shanghai 200234, China

⁸Institute of Space and Astronautical Science, Japan Aerospace Exploration Agency, 3-1-1 Yoshinodai, Sagami-hara, Kanagawa 229-8510, Japan

⁹Department of Astronomy, University of Massachusetts, Amherst, MA 01003, USA

¹⁰Department of Physics and Astronomy, University of Pennsylvania, Philadelphia, PA 19104, USA

¹¹Institute for Advanced Research, Nagoya University, Chikusa-ku, Nagoya 464-8601, Japan

¹²Department of Physics, Faculty of Science and Technology, Keio University, 3-14-1 Hiyoshi, Kohoku-ku, Yokohama, Kanagawa 223-8522, Japan

¹³Joetsu University of Education, Yamayashiki, Joetsu, Niigata 943-8512, Japan

Accepted 2010 September 7. Received 2010 August 18; in original form 2010 June 21

ABSTRACT

We present results of a 1.1-mm deep survey of the AKARI Deep Field South (ADF-S) with AzTEC mounted on the Atacama Submillimetre Telescope Experiment (ASTE). We obtained a map of 0.25-deg² area with an rms noise level of 0.32–0.71 mJy. This is one of the deepest and widest maps thus far at millimetre and submillimetre wavelengths. We uncovered 198 sources with a significance of 3.5 σ –15.6 σ , providing the largest catalogue of 1.1-mm sources in a contiguous region. Most of the sources are not detected in the far-infrared bands of the AKARI satellite, suggesting that they are mostly at $z \geq 1.5$ given the detection limits. We constructed differential and cumulative number counts in the ADF-S, the Subaru/XMM–Newton Deep Field and the SSA 22 field surveyed by AzTEC/ASTE, which provide currently the tightest constraints on the faint end. The integration of the best-fitting number counts in the ADF-S finds that the contribution of 1.1-mm sources with fluxes of ≥ 1 mJy to the cosmic infrared background (CIB) at 1.1 mm is 12–16 per cent, suggesting that the large fraction of the CIB originates from faint sources of which the number counts are not yet constrained. We estimate the cosmic star formation rate density contributed by 1.1-mm sources with ≥ 1 mJy using the best-fitting number counts in the ADF-S and find that it is lower by about a factor of 5–10 compared to those derived from UV/optically selected galaxies at $z \sim 2$ –3. The fraction of stellar mass of the present-day universe produced by 1.1-mm sources with ≥ 1 mJy at $z \geq 1$ is ~ 20 per cent, calculated by the time integration of the star formation rate density. If we consider the recycled fraction of >0.4 , which is the fraction of materials forming stars returned to the interstellar medium, the fraction of stellar mass produced by 1.1-mm sources decreases to $\lesssim 10$ per cent.

Key words: galaxies: evolution – galaxies: formation – galaxies: high-redshift – galaxies: starburst.

1 INTRODUCTION

Over the past decade, millimetre and submillimetre observations have shown that (sub)millimetre-bright galaxies (SMGs) hold

*E-mail: hatsukade@nro.nao.ac.jp

important clues to galaxy evolution and the cosmic star formation history (Blain et al. 2002 for a review). SMGs are highly obscured by dust, and the resulting thermal dust emission dominates the bolometric luminosity. The source of heating energy is dominated by vigorous star formation with star formation rates (SFRs) of several $100\text{--}1000 M_{\odot} \text{yr}^{-1}$. Optical/near-infrared (near-IR) spectroscopy of a sample of SMGs with radio counterparts revealed a median redshift of $z \sim 2$ for the population (Swinbank et al. 2004; Chapman et al. 2005). Recently, SMGs at $z > 4$ have been confirmed (Capak et al. 2008; Coppin et al. 2009; Daddi et al. 2009; Knudsen et al. 2010), and there is now a spectroscopically confirmed source at $z = 5.3$ (Riechers et al. 2010).

Coupled with reports of high dynamical mass and gas mass (e.g. Greve et al. 2005; Tacconi et al. 2006), it is suggested that SMGs are the progenitors of massive spheroidal galaxies observed during their formation phase (e.g. Lilly et al. 1996; Smail et al. 2004).

Mounting evidence shows that the cosmic infrared background (CIB; Puget et al. 1996; Fixsen et al. 1998) at millimetre and submillimetre wavelengths is largely contributed by high-redshift galaxies (Lagache, Puget & Dole 2005). The CIB is the integral of unresolved emission from extragalactic sources and contains information on the evolutionary history of galaxies. While 850- μm surveys have resolved 20–40 per cent of the CIB into point sources in blank fields (e.g. Eales et al. 2000; Borys et al. 2003; Coppin et al. 2006) and 50–100 per cent in lensing cluster fields (e.g. Blain et al. 2002; Cowie, Barger & Kneib 2002; Knudsen et al. 2008), 1-mm blank field surveys have resolved only ~ 10 per cent (e.g. Greve et al. 2004; Laurent et al. 2005; Maloney et al. 2005; Perera et al. 2008; Scott et al. 2008, 2010). A large portion of the CIB at millimetre and submillimetre wavelengths likely arises from galaxies with fainter flux densities.

In conjunction with constraints from the CIB, the number counts of SMGs are sensitive to the history of galaxy evolution at high redshifts. This requires constraining both the faint and bright ends of the number counts, which in turn requires a suitable combination of small, deep surveys along with shallower wide-area surveys. Blank field surveys at millimetre and submillimetre wavelengths have been carried out with large bolometer arrays such as the Submillimetre Common User Bolometer Array (SCUBA; Holland et al. 1999) on the James Clerk Maxwell Telescope (JCMT) (e.g. Smail, Ivison & Blain 1997; Hughes et al. 1998; Coppin et al. 2006), the Max-Planck Millimetre Bolometer Array (MAMBO; Kreysa et al. 1998) on the IRAM 30-m telescope and the Bolocam (Glenn et al. 1998) on the Caltech Submillimetre Observatory (e.g. Greve et al. 2004, 2008; Laurent et al. 2005; Bertoldi et al. 2007), the Large Apex BOLometer CAMera (LABOCA; Siringo et al. 2009) on the Atacama Pathfinder EXperiment (APEX; e.g. Weiß et al. 2009; Swinbank et al. 2010) and AzTEC (Wilson et al. 2008a) on the JCMT (Perera et al. 2008; Scott et al. 2008; Austermann et al. 2009, 2010). However, the total area covered in existing surveys is still small ($\lesssim 1 \text{ deg}^2$) compared to the cosmic large-scale structure, and substantial field-to-field variations can be seen in the published number counts. In addition, because of the limited depth of these surveys, the number counts of SMGs at faint flux densities ($\sim 1 \text{ mJy}$) are still not well constrained.

We performed extensive surveys at 1.1 mm with AzTEC mounted on the Atacama Submillimetre Telescope Experiment (ASTE; Ezawa et al. 2004; Ezawa et al. 2008) in 2007 and 2008 (Wilson et al. 2008b; Tamura et al. 2009; Scott et al. 2010). The ASTE is a 10-m submillimetre telescope located at Pampa la Bola in the Atacama desert in Chile. Some of the AzTEC/ASTE sources

are followed up by submillimetre/millimetre interferometers (Hatsukade et al. 2010; Tamura et al. in preparation; Ikarashi et al. 2010). In this paper, we report on a deep blank field survey of the AKARI Deep Field-South (ADF-S). The ADF-S is a multiwavelength deep survey field near the South Ecliptic Pole. It is known to be one of the lowest cirrus regions in the whole sky ($100\text{-}\mu\text{m}$ flux density of $< 0.5 \text{ MJy sr}^{-1}$ and an H I column density of $\sim 5 \times 10^{19} \text{ cm}^{-2}$; Schlegel, Finkbeiner & Davis 1998), providing a window to the high-redshift dusty universe. AKARI, an IR satellite (Matsuhara et al. 2005), has conducted deep surveys with the InfraRed Camera (IRC; Onaka et al. 2007) at 2.4, 3.2, 4.1, 7, 11, 15 and 24 μm and with the Far-Infrared Surveyor (FIS; Kawada et al. 2007) at 65, 90, 140 and 160 μm , down to the confusion limit (Shirahata et al. 2009). Multiwavelength follow-up observations from the UV to the radio are underway. The full data sets of IR to submillimetre bands with AKARI, *Spitzer* (Clements et al. 2010), the Balloon-borne Large-Aperture Submillimetre Telescope (BLAST), *Herschel Space Observatory* and the AzTEC/ASTE offer a unique opportunity to study the dusty galaxy population that contributes to the cosmic background at IR–mm wavelengths.

This paper presents the 1.1-mm map and source catalogue of the ADF-S. Together with the results from the Subaru/*XMM-Newton* Deep Field (SXDF; Ikarashi et al., in preparation) and the SSA 22 field (Tamura et al. 2009) surveyed by AzTEC/ASTE, we present statistical properties of the SMG population. Currently, these are the deepest wide-area surveys ever made at millimetre wavelengths along with the AzTEC/ASTE GOODS-S survey (Scott et al. 2010), providing the tightest constraints on number counts towards the faint flux density, albeit with lower resolution than has been typically employed to date. Comparisons with multiwavelength data and statistical studies such as clustering analysis of this data set will be presented in future papers.

The arrangement of this paper is as follows. Section 2 summarizes the observations of the ADF-S with AzTEC/ASTE. Section 3 outlines the data reduction and calibration details. In Section 4, we present the 1.1-mm map and the source catalogue. In Section 5, we derive number counts of the ADF-S, the SXDF and the SSA 22 field and compare them with other 1-mm wavelength surveys and luminosity evolution models. In Section 6, we estimate the contribution of 1.1-mm sources to the CIB. In Section 7, we constrain the redshifts of the AzTEC sources using flux ratios of 1.1 mm to 90 μm obtained with AKARI/FIS. In Section 8, we discuss the cosmic star formation history traced by 1.1-mm sources. A summary is presented in Section 9.

Throughout the paper, we adopt a cosmology with $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $h = 0.7$, $\Omega_M = 0.3$ and $\Omega_{\Lambda} = 0.7$.

2 OBSERVATIONS

The centre region of the ADF-S was observed with AzTEC on the ASTE. The observations were made from 2007 September 16 to October 14 and from 2008 August 4 to December 21. The operations were carried out remotely from the ASTE operation rooms in San Pedro de Atacama, Chile, and in Mitaka, Japan, through the network observation system N-COSMOS3 developed by the National Astronomical Observatory of Japan (NAOJ; Kamazaki et al. 2005). The full width at half-maximum (FWHM) of the AzTEC detectors on the ASTE is $\sim 30 \text{ arcsec}$ at 1.1 mm (270 GHz), and the field of view of the array is roughly circular with a diameter of 8 arcmin. During the observing run, 107 and 117 out of the 144 AzTEC detectors were operational in 2007 and 2008, respectively.

In order to maximize the observing efficiency, we used a Lissajous scan pattern to map the field. We chose a high maximum velocity of $300 \text{ arcsec s}^{-1}$ to mitigate low-frequency atmospheric fluctuations. The Lissajous scan pattern provides $\sim 20 \times 20 \text{ arcmin}^2$ coverage on the sky, in which the central $\sim 12 \times 12 \text{ arcmin}^2$ area is nearly uniform, with decreasing integration time from the inside to the outside. We covered the ADF-S with seven different field centres to make the noise level of the entire map uniform. We obtained a total of 319 individual observations under conditions where the atmospheric zenith opacity at 220 GHz as monitored with a radiometer at the ASTE telescope site was $\tau_{220\text{GHz}} = 0.02\text{--}0.1$. The total time spent on-field was $\sim 216 \text{ h}$. Uranus or Neptune was observed at least once a night in raster-scan mode to measure each detector's point spread function and relative position and to determine the flux conversion factor for absolute calibration (Wilson et al. 2008a). Pointing observations with the quasar J0455–462 that lies $\sim 7^\circ$ from the field centre were performed every 2 h, bracketing science observations. A pointing model for each observing run in 2007 and 2008 is constructed from these data, and we make corrections to the telescope astrometry. The resulting pointing accuracy is better than 3 arcsec (Wilson et al. 2008b).

3 DATA REDUCTION

The data were reduced in a manner identical to those described in Scott et al. (2010). We used a principle component analysis (PCA) to remove the low-frequency atmospheric signal from the time-stream data (Laurent et al. 2005; Scott et al. 2008). The PCA method AC couples the bolometer time-stream data, making the entire map and point-source kernel have a mean of zero. The cleaned time series data are projected into map space using $3 \times 3 \text{ arcsec}^2$ pixels in right-ascension–declination, and the individual observations are co-added into a single map by weighted averaging. We also create 100 noise realizations by ‘jackknifing’ the time series data as described in Scott et al. (2008). These noise maps represent realizations of the underlying random noise in the map in the absence of sources (both bright and confused) and are used throughout this paper to characterize the properties of the map and source catalogue. The point source profile is affected by the PCA method since the faint point sources with low spatial frequencies are also attenuated. The PCA method makes the mean of the map zero, causing negative side lobes around the peak of the point source profile. We trace the effects of PCA and other processes in the analysis on the point source profile; this ‘point source kernel’ is used to optimally filter the co-added map and the 100 noise realizations for the detection of point sources. A two-dimensional Gaussian fitting to the point source kernel gives an FWHM of 35.9 arcsec.

4 MAP AND SOURCE CATALOGUE

4.1 Map

The signal map and the corresponding noise map are shown in Figs 1 and 2, respectively. The dashed curves represent regions with ≥ 30 per cent (outer contour) and ≥ 50 per cent (inner contour) of the maximum weights (hereafter called as 30 and 50 per cent coverage regions, respectively). The area and noise level are 709 and 200 arcmin^2 , and $0.32\text{--}0.55$ and $0.55\text{--}0.71 \text{ mJy}$ in the 50 and 30–50 per cent coverage regions, respectively (Table 1). This survey is confusion-limited, where the 5σ -confusion limit estimated by Takeuchi & Ishii (2004) is 4.4 mJy using the point source kernel and the differential number counts in the ADF-S derived in Section 5.1.

Fig. 3 shows the distribution of flux values in the map, compared to that averaged over the 100 noise realizations within the 50 per cent coverage region. The result of a Gaussian fit to the averaged noise map is superimposed in Fig. 3. The presence of real sources in the map makes excess of both positive- and negative-valued pixels over the histogram of the noise map, since the signal map is created to have a mean of zero. This fit deviates from the distribution of pixel values at high positive and negative fluxes because the map is not uniform over the entire region, with the outer region being slightly noisier.

4.2 Source catalogue

Source extraction is performed on the signal-to-noise (SN) map using a criterion of $\geq 3.5\sigma$. The source positions are determined by flux-squared weighting on pixels within a 15-arcsec radius of the nominal peak. We detect 198 and 169 sources with a significance of $3.5\sigma\text{--}15.6\sigma$ in the 30 and 50 per cent coverage regions, respectively. The source catalogue is given in Table 2, where both the observed and the deboosted flux densities (Section 4.6) are listed. 169 sources (ADFS-AzTEC1–169) detected within the 50 per cent coverage region, the deeper and more uniform coverage region, are listed first, followed by the remaining 29 sources (ADFS-AzTEC170–198) detected outside the 50 per cent coverage region.

4.3 False detections

Monte Carlo simulations are carried out to estimate the number of spurious sources due to positive noise fluctuations. We conduct the standard source extraction on the 100 synthesised noise realizations and count the number of ‘sources’ above given signal-to-noise ratio (S/N) thresholds in steps of 0.5σ . Fig. 4 shows the average number of false detections as a function of S/N. The expected number of false detections in our $\geq 3.5\sigma$ source catalogue is $\sim 4\text{--}5$ and $\sim 1\text{--}2$ in the 50 and 30–50 per cent coverage regions, respectively.

4.4 Completeness

The survey completeness is computed by injecting simulated point sources of known flux densities into the real signal map one at a time. The input positions are randomly selected within the 50 per cent coverage region, but are required to be outside a 20-arcsec radius from a real source in the map to avoid blending. When a simulated source is extracted within 20 arcsec of its input position with $S/N \geq 3.5$, the source is considered to be recovered. We repeat this 1000 times for each flux bin and compute the fraction of output sources to input sources. The completeness as a function of intrinsic flux density is shown in Fig. 5. The error bars are the 68 per cent confidence intervals from the binomial distribution. The completeness is about 50 per cent at a flux density of 2.0 mJy .

4.5 Positional uncertainty

The positional error is an important indicator when identifying other-wavelength counterparts. The positional uncertainties for detected sources are calculated in a similar manner for the completeness calculation. Fake sources of known flux densities (ranging from 1 to 10 mJy in steps of 0.25 mJy) are injected into the real signal map one at a time. The input positions are selected randomly outside a 20-arcsec radius from the real sources. Source extractions with $S/N \geq 3.5$ are performed within 20-arcsec search radius from the input positions. These procedures are repeated 1000 times for




Figure 1. Signal map of the ADF-S. White contours represent the 30 per cent (outer contour) and 50 per cent (inner contour) coverage regions (corresponding to rms noise levels of 0.55 and 0.71 mJy beam⁻¹), respectively. The 198 sources detected with $\geq 3.5\sigma$ in the 30 per cent coverage region are indicated by green circles with a 30-arcsec diameter and numbered in order of significance (Table 2). The pixel size is 3×3 arcsec².




Figure 2. Noise map of the AzTEC/ASTE ADF-S. The outer and inner contours indicate the 30 and 50 per cent coverage regions, respectively.

each flux bin and measure the angular distances between the input and output positions. The probability that a source is extracted outside an angular distance from its input position is shown in Fig. 6 for different S/N ranges. The 68.3 per cent confidence interval for

Table 1. Map properties in the 50 and the 30–50 per cent coverage regions.

Coverage (per cent)	Area (arcmin ²)	Noise level (mJy beam ⁻¹)	Source ^a	False detections ^b
50 per cent	709	0.32–0.55	169	4.9 ± 0.22
30–50 per cent	200	0.55–0.71	29	1.4 ± 0.12

^aNumber of sources ($\geq 3.5\sigma$).

^bNumber of false detection ($\geq 3.5\sigma$).

a 3.5σ source is ~ 9 arcsec. We compare the positional uncertainty with theoretical predictions for uncorrelated Gaussian noise derived by Ivison et al. (2007). Shaded regions in Fig. 6 represent the theoretical probability distributions for sources with $3.5 \leq S/N < 4.0$, $4.5 \leq S/N < 5.0$ and $5.5 \leq S/N < 6.0$ from top to bottom. The positional uncertainties of AzTEC sources are broad compared to the theoretical predictions. It is possible that the ADF-S map is confused and the confusion noise affects source positions.

4.6 Flux deboosting

When dealing with a low S/N map, we need to consider the effect that flux densities of low S/N sources are boosted to above detection thresholds, due to the steep slope of number counts in the flux range of our map (Murdoch, Crawford & Jauncey 1973; Hogg & Turner 1998). We correct for the flux boosting effect to estimate intrinsic




Figure 3. Distribution of flux density in the real map (solid histogram) and averaged over the 100 noise realizations (dashed histogram) in the 50 per cent coverage region. The result of a Gaussian fit to the flux distribution of the noise maps is plotted as a dashed curve.

flux densities of sources based on the Bayesian estimation (Coppin et al. 2005; Coppin et al. 2006; Austermann et al. 2009, 2010).

To calculate the probability distribution of its intrinsic flux density [posterior flux distribution (PFD)], we use the best-fitting differential number counts in the ADF-S derived in Section 5.1. We inject fake sources with flux densities ranging from -5 to 20 mJy into a synthesized noiseless sky at random positions. We iterate this process 10 000 times, and the prior distribution is given as the averaged flux distribution of the sources. The deboosted flux densities are given in Table 2.

5 1.1-MM NUMBER COUNTS

5.1 Number counts of the ADF-S

We create number counts for the 50 per cent coverage region, where the noise distribution is more uniform, the survey completeness at faint flux densities is high and the number of false detections is low compared to the 30 per cent coverage region. We employ the Bayesian method, which is now commonly used for deriving number counts in millimetre/submillimetre surveys (e.g. Coppin et al. 2005, 2006; Perera et al. 2008; Austermann et al. 2009, 2010; Scott et al. 2010).

The PFD of each source is calculated in the same manner as used in flux deboosting (Section 4.6). We adopt the best-fitting Schechter function of the SCUBA/SHADES 850- μ m number counts (Coppin et al. 2006) scaled to 1.1 mm as an initial prior distribution function. We create 20 000 sample catalogues by bootstrapping off (i.e. sampling with replacement) the PFDs. We sample only from the PFDs with a criterion of $P(S < 0) \leq 0.05$ as adopted in Scott et al. (2010), where $P(S < 0)$ is the probability that the flux densities of the detected sources deboosted to < 0 mJy. The number of sources in each sample catalogue is chosen randomly as a Poisson deviate from the real number of sources. The survey completeness is calculated by tracing output to input sources from the simulated maps. After correcting for the survey completeness, the mean counts and the 68.3 per cent confidence intervals in each flux bin with a bin size of 1 mJy are calculated from the 20 000 sample catalogues. The deboosted fluxes in the source catalogue range from 1 to 6.4 mJy. Each of the derived number counts for the 20 000 sample catalogues

is fitted to a Schechter function of the form

$$\frac{dN}{dS} = N_{3\text{mJy}} \left(\frac{S}{3\text{mJy}} \right)^{\alpha+1} \exp \left[\frac{-(S - 3\text{mJy})}{S'} \right], \quad (1)$$

where $N_{3\text{mJy}}$ is a differential count at 3 mJy and best-fitting parameters are obtained in $S' - N_{3\text{mJy}}$ parameter space. We adopt the above Schechter functional form with $\alpha = -2$ since it well describes number counts derived in previous deep SMG surveys (e.g. Coppin et al. 2006; Perera et al. 2008; Austermann et al. 2009, 2010; Scott et al. 2010). The derived best-fitting function is then used as a new prior distribution function, and the procedure described above is repeated. The resultant differential and cumulative number counts are presented in Figs 7 and 8 and Table 3. The errors indicate the 68.3 per cent confidence intervals. The best-fitting parameters of the Schechter functional form in equation (1) are $N_{3\text{mJy}} = 169 \pm 19$ and $S' = 1.48 \pm 0.20$.

5.2 Number counts of the SXDF and the SSA 22 fields surveyed by AzTEC/ASTE

We extract number counts for two other deep fields surveyed by AzTEC on the ASTE: the SXDF and the SSA 22 field.

The SXDF is a blank field with deep multiwavelength observations from X-ray to radio. The AzTEC/ASTE observations covered ~ 0.27 deg² of the central part of the SXDF with an rms noise level of ~ 0.5 – 0.9 mJy (Ikarashi et al., in preparation), which is about a factor of 2 deeper than the AzTEC/JCMT survey of this field (Austermann et al. 2010). In total, ~ 200 sources ($\geq 3.5\sigma$) are detected.

The SSA 22 field is thought to be a protocluster region since it has an overdensity of UV/optically selected galaxies such as Lyman α emitters and Lyman-break galaxies at $z \sim 3.1$ (e.g. Steidel et al. 1998; Steidel et al. 2000; Hayashino et al. 2004; Matsuda et al. 2005). The SSA 22 field characterized by the overdensity is a good comparison field with other blank fields to see the relation between SMGs and other galaxy populations and the large-scale structure of the universe. The AzTEC/ASTE observations covered ~ 0.28 deg² with an rms noise level of ~ 0.6 – 1.2 mJy and detected ~ 100 sources ($\geq 3.5\sigma$) (Tamura et al. 2009; Tamura et al., in preparation).

The procedure and parameters in data reductions and extracting number counts of these two fields are the same as used in this paper. The differential and cumulative number counts are presented in Figs 7 and 8 and Table 4. The best-fitting parameters of the Schechter functional form in equation (1) are shown in Table 5.

5.3 Comparison among 1-mm surveys

In Figs 7 and 8, we compare the number counts in the ADF-S, the SXDF and the SSA 22 fields surveyed by AzTEC/ASTE with those of previous 1-mm surveys: the AzTEC/JCMT surveys of GOODS-N (Perera et al. 2008), COSMOS (Austermann et al. 2009) and SHADES (combined counts in the Lockman Hole and SXDF) (Austermann et al. 2010), the AzTEC/ASTE survey of GOODS-S (Scott et al. 2010) and 1.2-mm MAMBO surveys of the Lockman Hole and ELAIS N2 (Greve et al. 2004).

The ADF-S and the SXDF provide the tightest constraints on the faint end of the number counts because of their depth and large survey areas. On the whole, the 1-mm counts of various surveys are consistent within errors. This is interesting since the SSA 22 field has overdensity of UV/optically selected galaxies. It is possible that the overdensity of sources at $z = 3.1$ traced by the

Table 2. AzTEC/ASTE ADF-S 3.5σ source catalogue. AzTEC1–169 are detected in the 50 per cent coverage region and AzTEC170–198 are detected in the region with a coverage of 30–50 per cent. The columns give (1) source name, (2) ID, (3) right ascension, (4) declination, (5) observed flux density and 1σ error, (6) deboosted flux density and 68 per cent confidence level and (7) signal-to-noise ratio.

Name	ID	RA (^h ^m ^s)	Dec. ([°] ['] ^{''})	S_{observed} (mJy)	$S_{\text{deboosted}}$ (mJy)	S/N
AzTEC J044511.52–533807.3	ADFS-AzTEC1	4 45 11.52	–53 38 07.3	5.9 ± 0.4	$5.7^{+0.4}_{-0.4}$	15.6
AzTEC J044621.97–533630.4	ADFS-AzTEC2	4 46 21.97	–53 36 30.4	5.5 ± 0.4	$5.3^{+0.5}_{-0.4}$	14.2
AzTEC J044730.07–531928.1	ADFS-AzTEC3	4 47 30.07	–53 19 28.1	5.1 ± 0.4	$4.9^{+0.5}_{-0.4}$	13.8
AzTEC J044441.16–534543.4	ADFS-AzTEC4	4 44 41.16	–53 45 43.4	6.5 ± 0.5	$6.2^{+0.5}_{-0.6}$	12.8
AzTEC J044419.58–533422.9	ADFS-AzTEC5	4 44 19.58	–53 34 22.9	6.7 ± 0.5	$6.4^{+0.6}_{-0.6}$	12.2
AzTEC J044702.37–532049.8	ADFS-AzTEC6	4 47 02.37	–53 20 49.8	4.9 ± 0.4	$4.7^{+0.4}_{-0.5}$	12.1
AzTEC J044706.89–531827.3	ADFS-AzTEC7	4 47 06.89	–53 18 27.3	4.9 ± 0.4	$4.7^{+0.4}_{-0.5}$	12.0
AzTEC J044711.18–532154.4	ADFS-AzTEC8	4 47 11.18	–53 21 54.4	4.9 ± 0.4	$4.7^{+0.5}_{-0.5}$	12.0
AzTEC J044544.27–533124.0	ADFS-AzTEC9	4 45 44.27	–53 31 24.0	4.0 ± 0.3	$3.8^{+0.4}_{-0.4}$	11.9
AzTEC J044441.47–534222.3	ADFS-AzTEC10	4 44 41.47	–53 42 22.3	5.3 ± 0.5	$5.1^{+0.5}_{-0.5}$	11.8
AzTEC J044452.99–533810.5	ADFS-AzTEC11	4 44 52.99	–53 38 10.5	4.8 ± 0.4	$4.5^{+0.5}_{-0.5}$	11.4
AzTEC J044733.99–531837.5	ADFS-AzTEC12	4 47 33.99	–53 18 37.5	4.0 ± 0.4	$3.8^{+0.4}_{-0.4}$	11.0
AzTEC J044736.90–532527.5	ADFS-AzTEC13	4 47 36.90	–53 25 27.5	4.5 ± 0.4	$4.3^{+0.5}_{-0.5}$	10.7
AzTEC J044442.42–534122.9	ADFS-AzTEC14	4 44 42.42	–53 41 22.9	4.8 ± 0.4	$4.5^{+0.5}_{-0.5}$	10.6
AzTEC J044844.20–531443.4	ADFS-AzTEC15	4 48 44.20	–53 14 43.4	5.0 ± 0.5	$4.7^{+0.5}_{-0.5}$	10.5
AzTEC J044802.69–531712.7	ADFS-AzTEC16	4 48 02.69	–53 17 12.7	4.2 ± 0.4	$3.9^{+0.4}_{-0.5}$	10.5
AzTEC J044743.53–531541.8	ADFS-AzTEC17	4 47 43.53	–53 15 41.8	4.2 ± 0.4	$4.0^{+0.5}_{-0.5}$	10.3
AzTEC J044753.52–532331.4	ADFS-AzTEC18	4 47 53.52	–53 23 31.4	4.1 ± 0.4	$3.9^{+0.5}_{-0.4}$	10.3
AzTEC J044543.44–532524.3	ADFS-AzTEC19	4 45 43.44	–53 25 24.3	3.8 ± 0.4	$3.6^{+0.4}_{-0.4}$	10.1
AzTEC J044813.58–531859.6	ADFS-AzTEC20	4 48 13.58	–53 18 59.6	4.1 ± 0.4	$3.9^{+0.5}_{-0.5}$	10.0
AzTEC J044658.41–531534.1	ADFS-AzTEC21	4 46 58.41	–53 15 34.1	5.0 ± 0.5	$4.7^{+0.6}_{-0.5}$	9.9
AzTEC J044455.57–533423.0	ADFS-AzTEC22	4 44 55.57	–53 34 23.0	3.9 ± 0.4	$3.7^{+0.5}_{-0.4}$	9.8
AzTEC J044540.95–533318.2	ADFS-AzTEC23	4 45 40.95	–53 33 18.2	3.2 ± 0.3	$3.0^{+0.4}_{-0.4}$	9.7
AzTEC J044522.00–532700.9	ADFS-AzTEC24	4 45 22.00	–53 27 00.9	3.6 ± 0.4	$3.4^{+0.4}_{-0.4}$	9.7
AzTEC J044659.94–531907.3	ADFS-AzTEC25	4 46 59.94	–53 19 07.3	4.0 ± 0.4	$3.7^{+0.5}_{-0.5}$	9.7
AzTEC J044629.83–533345.0	ADFS-AzTEC26	4 46 29.83	–53 33 45.0	3.3 ± 0.4	$3.1^{+0.4}_{-0.4}$	9.3
AzTEC J044658.48–532327.8	ADFS-AzTEC27	4 46 58.48	–53 23 27.8	3.6 ± 0.4	$3.4^{+0.4}_{-0.5}$	9.1
AzTEC J044612.33–532743.6	ADFS-AzTEC28	4 46 12.33	–53 27 43.6	2.9 ± 0.3	$2.7^{+0.4}_{-0.4}$	9.1
AzTEC J044714.90–531738.9	ADFS-AzTEC29	4 47 14.90	–53 17 38.9	3.7 ± 0.4	$3.4^{+0.4}_{-0.5}$	9.1
AzTEC J044805.34–532433.2	ADFS-AzTEC30	4 48 05.34	–53 24 33.2	4.4 ± 0.5	$4.1^{+0.6}_{-0.5}$	9.0
AzTEC J044351.90–534443.3	ADFS-AzTEC31	4 43 51.90	–53 44 43.3	4.8 ± 0.5	$4.5^{+0.6}_{-0.6}$	8.8
AzTEC J044553.57–533520.6	ADFS-AzTEC32	4 45 53.57	–53 35 20.6	2.8 ± 0.3	$2.6^{+0.4}_{-0.4}$	8.7
AzTEC J044706.39–531612.5	ADFS-AzTEC33	4 47 06.39	–53 16 12.5	4.0 ± 0.5	$3.7^{+0.5}_{-0.5}$	8.7
AzTEC J044538.56–532827.6	ADFS-AzTEC34	4 45 38.56	–53 28 27.6	2.8 ± 0.3	$2.6^{+0.4}_{-0.4}$	8.4
AzTEC J044853.11–531557.9	ADFS-AzTEC35	4 48 53.11	–53 15 57.9	4.1 ± 0.5	$3.7^{+0.5}_{-0.6}$	8.2
AzTEC J044556.43–533929.7	ADFS-AzTEC36	4 45 56.43	–53 39 29.7	3.5 ± 0.4	$3.3^{+0.5}_{-0.5}$	8.1
AzTEC J044348.44–534314.2	ADFS-AzTEC37	4 43 48.44	–53 43 14.2	4.3 ± 0.5	$3.9^{+0.6}_{-0.6}$	8.0
AzTEC J044824.87–531501.3	ADFS-AzTEC38	4 48 24.87	–53 15 01.3	3.5 ± 0.4	$3.2^{+0.5}_{-0.5}$	8.0
AzTEC J044609.02–532710.8	ADFS-AzTEC39	4 46 09.02	–53 27 10.8	2.6 ± 0.3	$2.4^{+0.4}_{-0.4}$	8.0
AzTEC J044534.14–533939.6	ADFS-AzTEC40	4 45 34.14	–53 39 39.6	3.4 ± 0.4	$3.1^{+0.5}_{-0.5}$	7.9
AzTEC J044354.85–533928.8	ADFS-AzTEC41	4 43 54.85	–53 39 28.8	3.9 ± 0.5	$3.6^{+0.6}_{-0.5}$	7.9
AzTEC J044630.39–533154.7	ADFS-AzTEC42	4 46 30.39	–53 31 54.7	2.7 ± 0.3	$2.5^{+0.4}_{-0.4}$	7.8
AzTEC J044607.77–532813.4	ADFS-AzTEC43	4 46 07.77	–53 28 13.4	2.5 ± 0.3	$2.3^{+0.4}_{-0.3}$	7.8
AzTEC J044643.79–533002.4	ADFS-AzTEC44	4 46 43.79	–53 30 02.4	3.0 ± 0.4	$2.7^{+0.4}_{-0.5}$	7.6
AzTEC J044451.57–533544.1	ADFS-AzTEC45	4 44 51.57	–53 35 44.1	3.1 ± 0.4	$2.9^{+0.5}_{-0.5}$	7.5
AzTEC J044420.88–534011.4	ADFS-AzTEC46	4 44 20.88	–53 40 11.4	3.4 ± 0.5	$3.1^{+0.5}_{-0.5}$	7.3
AzTEC J044541.74–533445.2	ADFS-AzTEC47	4 45 41.74	–53 34 45.2	2.3 ± 0.3	$2.1^{+0.4}_{-0.4}$	7.3
AzTEC J044437.74–534637.8	ADFS-AzTEC48	4 44 37.74	–53 46 37.8	3.8 ± 0.5	$3.4^{+0.6}_{-0.6}$	7.2

Table 2 – *continued*

Name	ID	RA (h m s)	Dec. (° ′ ″)	S _{observed} (mJy)	S _{deboosted} (mJy)	S/N
AzTEC J044528.39–533715.8	ADFS-AzTEC49	4 45 28.39	–53 37 15.8	2.4 ± 0.3	2.2 ^{+0.4} _{–0.4}	7.1
AzTEC J044456.63–533616.2	ADFS-AzTEC50	4 44 56.63	–53 36 16.2	2.8 ± 0.4	2.5 ^{+0.4} _{–0.5}	7.1
AzTEC J044623.49–533552.4	ADFS-AzTEC51	4 46 23.49	–53 35 52.4	2.6 ± 0.4	2.4 ^{+0.4} _{–0.4}	6.9
AzTEC J044344.41–534325.9	ADFS-AzTEC52	4 43 44.41	–53 43 25.9	3.8 ± 0.6	3.4 ^{+0.6} _{–0.6}	6.9
AzTEC J044742.49–531951.2	ADFS-AzTEC53	4 47 42.49	–53 19 51.2	2.4 ± 0.4	2.2 ^{+0.4} _{–0.4}	6.9
AzTEC J044721.15–532659.2	ADFS-AzTEC54	4 47 21.15	–53 26 59.2	2.9 ± 0.4	2.6 ^{+0.5} _{–0.5}	6.8
AzTEC J044423.80–533413.5	ADFS-AzTEC55	4 44 23.80	–53 34 13.5	3.7 ± 0.6	3.3 ^{+0.6} _{–0.6}	6.8
AzTEC J044435.35–534346.6	ADFS-AzTEC56	4 44 35.35	–53 43 46.6	3.1 ± 0.5	2.8 ^{+0.5} _{–0.5}	6.7
AzTEC J044611.47–532152.8	ADFS-AzTEC57	4 46 11.47	–53 21 52.8	3.5 ± 0.5	3.1 ^{+0.5} _{–0.6}	6.7
AzTEC J044516.36–533309.9	ADFS-AzTEC58	4 45 16.36	–53 33 09.9	2.2 ± 0.3	2.0 ^{+0.4} _{–0.4}	6.7
AzTEC J044639.12–532518.6	ADFS-AzTEC59	4 46 39.12	–53 25 18.6	2.5 ± 0.4	2.2 ^{+0.4} _{–0.4}	6.7
AzTEC J044527.26–533518.6	ADFS-AzTEC60	4 45 27.26	–53 35 18.6	2.1 ± 0.3	1.9 ^{+0.4} _{–0.4}	6.7
AzTEC J044858.01–531524.3	ADFS-AzTEC61	4 48 58.01	–53 15 24.3	3.5 ± 0.5	3.2 ^{+0.6} _{–0.6}	6.6
AzTEC J044722.55–531825.8	ADFS-AzTEC62	4 47 22.55	–53 18 25.8	2.5 ± 0.4	2.3 ^{+0.5} _{–0.4}	6.6
AzTEC J044400.63–534228.2	ADFS-AzTEC63	4 44 00.63	–53 42 28.2	3.3 ± 0.5	2.9 ^{+0.5} _{–0.6}	6.6
AzTEC J044813.94–532244.7	ADFS-AzTEC64	4 48 13.94	–53 22 44.7	3.0 ± 0.5	2.7 ^{+0.5} _{–0.5}	6.6
AzTEC J044822.25–531420.0	ADFS-AzTEC65	4 48 22.25	–53 14 20.0	2.9 ± 0.4	2.6 ^{+0.5} _{–0.5}	6.5
AzTEC J044503.53–532816.7	ADFS-AzTEC66	4 45 03.53	–53 28 16.7	2.8 ± 0.4	2.5 ^{+0.5} _{–0.5}	6.5
AzTEC J044826.06–531413.7	ADFS-AzTEC67	4 48 26.06	–53 14 13.7	2.9 ± 0.4	2.6 ^{+0.5} _{–0.5}	6.5
AzTEC J044857.68–531827.0	ADFS-AzTEC68	4 48 57.68	–53 18 27.0	3.5 ± 0.5	3.1 ^{+0.6} _{–0.6}	6.5
AzTEC J044556.20–532541.2	ADFS-AzTEC69	4 45 56.20	–53 25 41.2	2.2 ± 0.4	2.0 ^{+0.4} _{–0.4}	6.2
AzTEC J044603.17–533847.5	ADFS-AzTEC70	4 46 03.17	–53 38 47.5	2.6 ± 0.4	2.3 ^{+0.5} _{–0.4}	6.2
AzTEC J044358.65–533731.2	ADFS-AzTEC71	4 43 58.65	–53 37 31.2	3.1 ± 0.5	2.7 ^{+0.5} _{–0.6}	6.2
AzTEC J044717.48–532614.8	ADFS-AzTEC72	4 47 17.48	–53 26 14.8	2.5 ± 0.4	2.2 ^{+0.5} _{–0.4}	6.1
AzTEC J044428.74–534129.4	ADFS-AzTEC73	4 44 28.74	–53 41 29.4	2.8 ± 0.5	2.4 ^{+0.5} _{–0.5}	6.0
AzTEC J044747.96–531305.5	ADFS-AzTEC74	4 47 47.96	–53 13 05.5	3.1 ± 0.5	2.7 ^{+0.6} _{–0.6}	6.0
AzTEC J044545.17–533815.3	ADFS-AzTEC75	4 45 45.17	–53 38 15.3	2.2 ± 0.4	1.9 ^{+0.4} _{–0.4}	6.0
AzTEC J044502.41–533934.4	ADFS-AzTEC76	4 45 02.41	–53 39 34.4	2.5 ± 0.4	2.2 ^{+0.5} _{–0.5}	6.0
AzTEC J044554.73–533829.6	ADFS-AzTEC77	4 45 54.73	–53 38 29.6	2.3 ± 0.4	2.0 ^{+0.5} _{–0.4}	5.9
AzTEC J044744.88–531622.8	ADFS-AzTEC78	4 47 44.88	–53 16 22.8	2.3 ± 0.4	2.0 ^{+0.4} _{–0.5}	5.9
AzTEC J044614.68–532631.6	ADFS-AzTEC79	4 46 14.68	–53 26 31.6	2.0 ± 0.3	1.7 ^{+0.4} _{–0.4}	5.9
AzTEC J044635.07–532950.7	ADFS-AzTEC80	4 46 35.07	–53 29 50.7	2.1 ± 0.4	1.8 ^{+0.4} _{–0.4}	5.8
AzTEC J044409.74–534607.4	ADFS-AzTEC81	4 44 09.74	–53 46 07.4	3.1 ± 0.5	2.7 ^{+0.6} _{–0.6}	5.8
AzTEC J044432.70–534419.3	ADFS-AzTEC82	4 44 32.70	–53 44 19.3	2.7 ± 0.5	2.4 ^{+0.6} _{–0.5}	5.8
AzTEC J044429.65–534659.4	ADFS-AzTEC83	4 44 29.65	–53 46 59.4	3.1 ± 0.5	2.7 ^{+0.6} _{–0.6}	5.8
AzTEC J044531.48–533019.1	ADFS-AzTEC84	4 45 31.48	–53 30 19.1	1.9 ± 0.3	1.6 ^{+0.4} _{–0.4}	5.7
AzTEC J044553.94–532908.4	ADFS-AzTEC85	4 45 53.94	–53 29 08.4	1.8 ± 0.3	1.6 ^{+0.4} _{–0.4}	5.6
AzTEC J044434.71–534125.5	ADFS-AzTEC86	4 44 34.71	–53 41 25.5	2.5 ± 0.4	2.2 ^{+0.5} _{–0.5}	5.6
AzTEC J044738.06–532754.8	ADFS-AzTEC87	4 47 38.06	–53 27 54.8	2.9 ± 0.5	2.5 ^{+0.6} _{–0.6}	5.5
AzTEC J044439.10–533704.0	ADFS-AzTEC88	4 44 39.10	–53 37 04.0	2.5 ± 0.5	2.2 ^{+0.5} _{–0.5}	5.5
AzTEC J044642.77–532926.9	ADFS-AzTEC89	4 46 42.77	–53 29 26.9	2.1 ± 0.4	1.8 ^{+0.4} _{–0.5}	5.5
AzTEC J044732.69–531422.0	ADFS-AzTEC90	4 47 32.69	–53 14 22.0	2.6 ± 0.5	2.2 ^{+0.5} _{–0.5}	5.4
AzTEC J044509.91–532831.2	ADFS-AzTEC91	4 45 09.91	–53 28 31.2	2.1 ± 0.4	1.8 ^{+0.4} _{–0.5}	5.4
AzTEC J044749.46–532244.4	ADFS-AzTEC92	4 47 49.46	–53 22 44.4	2.0 ± 0.4	1.8 ^{+0.5} _{–0.4}	5.4
AzTEC J044438.94–533316.8	ADFS-AzTEC93	4 44 38.94	–53 33 16.8	2.9 ± 0.5	2.4 ^{+0.6} _{–0.6}	5.4
AzTEC J044504.27–533030.9	ADFS-AzTEC94	4 45 04.27	–53 30 30.9	2.0 ± 0.4	1.8 ^{+0.4} _{–0.4}	5.3
AzTEC J044632.65–533457.6	ADFS-AzTEC95	4 46 32.65	–53 34 57.6	2.1 ± 0.4	1.8 ^{+0.4} _{–0.5}	5.3
AzTEC J044751.90–531920.3	ADFS-AzTEC96	4 47 51.90	–53 19 20.3	1.9 ± 0.4	1.6 ^{+0.4} _{–0.4}	5.3
AzTEC J044627.13–533224.4	ADFS-AzTEC97	4 46 27.13	–53 32 24.4	1.8 ± 0.3	1.5 ^{+0.4} _{–0.4}	5.2
AzTEC J044607.98–533416.2	ADFS-AzTEC98	4 46 07.98	–53 34 16.2	1.7 ± 0.3	1.5 ^{+0.4} _{–0.4}	5.2

Table 2 – continued

Name	ID	RA ($^{\circ}$ $^{\prime}$ $^{\prime\prime}$)	Dec. ($^{\circ}$ $^{\prime}$ $^{\prime\prime}$)	S _{observed} (mJy)	S _{deboosted} (mJy)	S/N
AzTEC J044537.32–532311.4	ADFS-AzTEC99	4 45 37.32	–53 23 11.4	2.8 ± 0.5	2.3 ^{+0.6} _{–0.6}	5.2
AzTEC J044608.37–533550.4	ADFS-AzTEC100	4 46 08.37	–53 35 50.4	1.7 ± 0.3	1.5 ^{+0.4} _{–0.4}	5.1
AzTEC J044654.55–532308.3	ADFS-AzTEC101	4 46 54.55	–53 23 08.3	2.0 ± 0.4	1.8 ^{+0.5} _{–0.5}	5.1
AzTEC J044747.12–532011.2	ADFS-AzTEC102	4 47 47.12	–53 20 11.2	1.8 ± 0.4	1.6 ^{+0.4} _{–0.4}	5.1
AzTEC J044526.81–533031.2	ADFS-AzTEC103	4 45 26.81	–53 30 31.2	1.7 ± 0.3	1.4 ^{+0.4} _{–0.4}	5.1
AzTEC J044628.14–533254.4	ADFS-AzTEC104	4 46 28.14	–53 32 54.4	1.7 ± 0.3	1.5 ^{+0.4} _{–0.4}	5.1
AzTEC J044837.01–531924.2	ADFS-AzTEC105	4 48 37.01	–53 19 24.2	2.3 ± 0.5	2.0 ^{+0.5} _{–0.5}	5.0
AzTEC J044558.64–533747.4	ADFS-AzTEC106	4 45 58.64	–53 37 47.4	1.8 ± 0.4	1.6 ^{+0.4} _{–0.4}	5.0
AzTEC J044821.59–531219.6	ADFS-AzTEC107	4 48 21.59	–53 12 19.6	2.5 ± 0.5	2.1 ^{+0.6} _{–0.5}	5.0
AzTEC J044741.65–531839.1	ADFS-AzTEC108	4 47 41.65	–53 18 39.1	1.8 ± 0.4	1.5 ^{+0.4} _{–0.4}	5.0
AzTEC J044442.20–534413.9	ADFS-AzTEC109	4 44 42.20	–53 44 13.9	2.4 ± 0.5	2.0 ^{+0.5} _{–0.6}	5.0
AzTEC J044720.44–532526.5	ADFS-AzTEC110	4 47 20.44	–53 25 26.5	2.0 ± 0.4	1.7 ^{+0.4} _{–0.5}	5.0
AzTEC J044543.78–532612.1	ADFS-AzTEC111	4 45 43.78	–53 26 12.1	1.7 ± 0.3	1.5 ^{+0.4} _{–0.4}	4.9
AzTEC J044403.61–533816.8	ADFS-AzTEC112	4 44 03.61	–53 38 16.8	2.4 ± 0.5	2.0 ^{+0.6} _{–0.5}	4.9
AzTEC J044517.36–532619.7	ADFS-AzTEC113	4 45 17.36	–53 26 19.7	2.0 ± 0.4	1.7 ^{+0.5} _{–0.5}	4.9
AzTEC J044511.45–533328.2	ADFS-AzTEC114	4 45 11.45	–53 33 28.2	1.6 ± 0.3	1.4 ^{+0.4} _{–0.4}	4.9
AzTEC J044834.22–531727.4	ADFS-AzTEC115	4 48 34.22	–53 17 27.4	2.1 ± 0.4	1.8 ^{+0.5} _{–0.5}	4.9
AzTEC J044759.97–531235.0	ADFS-AzTEC116	4 47 59.97	–53 12 35.0	2.5 ± 0.5	2.1 ^{+0.6} _{–0.6}	4.8
AzTEC J044540.52–532927.6	ADFS-AzTEC117	4 45 40.52	–53 29 27.6	1.6 ± 0.3	1.3 ^{+0.4} _{–0.4}	4.8
AzTEC J044529.40–534100.9	ADFS-AzTEC118	4 45 29.40	–53 41 00.9	2.4 ± 0.5	2.0 ^{+0.6} _{–0.6}	4.7
AzTEC J044553.28–533017.6	ADFS-AzTEC119	4 45 53.28	–53 30 17.6	1.5 ± 0.3	1.3 ^{+0.4} _{–0.4}	4.7
AzTEC J044634.10–533721.6	ADFS-AzTEC120	4 46 34.10	–53 37 21.6	2.4 ± 0.5	2.0 ^{+0.6} _{–0.6}	4.7
AzTEC J044515.32–533354.3	ADFS-AzTEC121	4 45 15.32	–53 33 54.3	1.5 ± 0.3	1.3 ^{+0.4} _{–0.4}	4.7
AzTEC J044435.41–533529.4	ADFS-AzTEC122	4 44 35.41	–53 35 29.4	2.2 ± 0.5	1.8 ^{+0.5} _{–0.6}	4.6
AzTEC J044458.94–533504.3	ADFS-AzTEC123	4 44 58.94	–53 35 04.3	1.7 ± 0.4	1.4 ^{+0.4} _{–0.5}	4.5
AzTEC J044636.36–532915.4	ADFS-AzTEC124	4 46 36.36	–53 29 15.4	1.6 ± 0.4	1.3 ^{+0.4} _{–0.4}	4.5
AzTEC J044605.18–533137.8	ADFS-AzTEC125	4 46 05.18	–53 31 37.8	1.5 ± 0.3	1.2 ^{+0.4} _{–0.4}	4.5
AzTEC J044640.93–532608.2	ADFS-AzTEC126	4 46 40.93	–53 26 08.2	1.6 ± 0.4	1.3 ^{+0.4} _{–0.4}	4.4
AzTEC J044634.09–532612.2	ADFS-AzTEC127	4 46 34.09	–53 26 12.2	1.6 ± 0.4	1.3 ^{+0.4} _{–0.5}	4.4
AzTEC J044756.14–531727.7	ADFS-AzTEC128	4 47 56.14	–53 17 27.7	1.6 ± 0.4	1.4 ^{+0.5} _{–0.4}	4.3
AzTEC J044609.04–532913.4	ADFS-AzTEC129	4 46 09.04	–53 29 13.4	1.4 ± 0.3	1.1 ^{+0.4} _{–0.4}	4.3
AzTEC J044756.92–531637.4	ADFS-AzTEC130	4 47 56.92	–53 16 37.4	1.7 ± 0.4	1.4 ^{+0.5} _{–0.4}	4.3
AzTEC J044528.27–533551.3	ADFS-AzTEC131	4 45 28.27	–53 35 51.3	1.4 ± 0.3	1.1 ^{+0.4} _{–0.4}	4.3
AzTEC J044532.76–532921.6	ADFS-AzTEC132	4 45 32.76	–53 29 21.6	1.4 ± 0.3	1.2 ^{+0.4} _{–0.4}	4.3
AzTEC J044551.09–532505.4	ADFS-AzTEC133	4 45 51.09	–53 25 05.4	1.6 ± 0.4	1.3 ^{+0.4} _{–0.5}	4.3
AzTEC J044438.82–534047.6	ADFS-AzTEC134	4 44 38.82	–53 40 47.6	1.9 ± 0.4	1.5 ^{+0.5} _{–0.5}	4.3
AzTEC J044449.52–533723.0	ADFS-AzTEC135	4 44 49.52	–53 37 23.0	1.8 ± 0.4	1.5 ^{+0.5} _{–0.5}	4.3
AzTEC J044723.01–531414.1	ADFS-AzTEC136	4 47 23.01	–53 14 14.1	2.2 ± 0.5	1.7 ^{+0.6} _{–0.6}	4.2
AzTEC J044746.74–532129.6	ADFS-AzTEC137	4 47 46.74	–53 21 29.6	1.5 ± 0.4	1.3 ^{+0.4} _{–0.4}	4.2
AzTEC J044522.17–533345.0	ADFS-AzTEC138	4 45 22.17	–53 33 45.0	1.3 ± 0.3	1.1 ^{+0.4} _{–0.4}	4.2
AzTEC J044351.18–534228.7	ADFS-AzTEC139	4 43 51.18	–53 42 28.7	2.2 ± 0.5	1.7 ^{+0.6} _{–0.6}	4.2
AzTEC J044758.09–531825.1	ADFS-AzTEC140	4 47 58.09	–53 18 25.1	1.6 ± 0.4	1.3 ^{+0.5} _{–0.4}	4.2
AzTEC J044733.34–532222.1	ADFS-AzTEC141	4 47 33.34	–53 22 22.1	1.6 ± 0.4	1.3 ^{+0.5} _{–0.4}	4.2
AzTEC J044746.06–532623.4	ADFS-AzTEC142	4 47 46.06	–53 26 23.4	2.0 ± 0.5	1.6 ^{+0.5} _{–0.6}	4.1
AzTEC J044715.12–532500.9	ADFS-AzTEC143	4 47 15.12	–53 25 00.9	1.6 ± 0.4	1.3 ^{+0.5} _{–0.5}	4.1
AzTEC J044458.75–534319.5	ADFS-AzTEC144	4 44 58.75	–53 43 19.5	2.1 ± 0.5	1.6 ^{+0.6} _{–0.6}	4.1
AzTEC J044625.74–531931.6	ADFS-AzTEC145	4 46 25.74	–53 19 31.6	2.1 ± 0.5	1.7 ^{+0.6} _{–0.6}	4.1
AzTEC J044637.83–531851.1	ADFS-AzTEC146	4 46 37.83	–53 18 51.1	1.8 ± 0.5	1.4 ^{+0.5} _{–0.6}	4.0
AzTEC J044524.83–533134.0	ADFS-AzTEC147	4 45 24.83	–53 31 34.0	1.3 ± 0.3	1.0 ^{+0.4} _{–0.4}	3.9
AzTEC J044608.77–533716.8	ADFS-AzTEC148	4 46 08.77	–53 37 16.8	1.4 ± 0.4	1.1 ^{+0.4} _{–0.4}	3.9

Table 2 – *continued*

Name	ID	RA (^h ^m ^s)	Dec. ([°] ['] ^{''})	S _{observed} (mJy)	S _{deboosted} (mJy)	S/N
AzTEC J044515.33–532955.2	ADFS-AzTEC149	4 45 15.33	–53 29 55.2	1.3 ± 0.3	1.1 ^{+0.4} _{–0.4}	3.9
AzTEC J044654.28–533116.4	ADFS-AzTEC150	4 46 54.28	–53 31 16.4	1.9 ± 0.5	1.5 ^{+0.6} _{–0.6}	3.9
AzTEC J044634.68–532251.6	ADFS-AzTEC151	4 46 34.68	–53 22 51.6	1.5 ± 0.4	1.2 ^{+0.5} _{–0.5}	3.8
AzTEC J044508.10–534204.2	ADFS-AzTEC152	4 45 08.10	–53 42 04.2	2.0 ± 0.5	1.5 ^{+0.6} _{–0.6}	3.8
AzTEC J044456.83–533216.7	ADFS-AzTEC153	4 44 56.83	–53 32 16.7	1.5 ± 0.4	1.2 ^{+0.5} _{–0.5}	3.8
AzTEC J044629.37–532922.3	ADFS-AzTEC154	4 46 29.37	–53 29 22.3	1.3 ± 0.3	1.0 ^{+0.4} _{–0.4}	3.8
AzTEC J044655.70–532707.5	ADFS-AzTEC155	4 46 55.70	–53 27 07.5	1.5 ± 0.4	1.2 ^{+0.5} _{–0.5}	3.8
AzTEC J044834.45–531642.0	ADFS-AzTEC156	4 48 34.45	–53 16 42.0	1.6 ± 0.4	1.3 ^{+0.5} _{–0.5}	3.8
AzTEC J044346.80–534131.8	ADFS-AzTEC157	4 43 46.80	–53 41 31.8	2.0 ± 0.5	1.5 ^{+0.7} _{–0.6}	3.8
AzTEC J044822.88–532151.5	ADFS-AzTEC158	4 48 22.88	–53 21 51.5	1.7 ± 0.5	1.3 ^{+0.5} _{–0.6}	3.8
AzTEC J044533.34–532423.6	ADFS-AzTEC159	4 45 33.34	–53 24 23.6	1.7 ± 0.4	1.3 ^{+0.6} _{–0.5}	3.7
AzTEC J044522.66–532518.8	ADFS-AzTEC160	4 45 22.66	–53 25 18.8	1.6 ± 0.4	1.2 ^{+0.5} _{–0.5}	3.7
AzTEC J044407.06–533950.0	ADFS-AzTEC161	4 44 07.06	–53 39 50.0	1.8 ± 0.5	1.3 ^{+0.6} _{–0.6}	3.7
AzTEC J044646.61–531632.6	ADFS-AzTEC162	4 46 46.61	–53 16 32.6	1.8 ± 0.5	1.3 ^{+0.6} _{–0.6}	3.7
AzTEC J044516.25–534025.3	ADFS-AzTEC163	4 45 16.25	–53 40 25.3	1.7 ± 0.5	1.2 ^{+0.5} _{–0.5}	3.7
AzTEC J044512.72–533644.8	ADFS-AzTEC164	4 45 12.72	–53 36 44.8	1.3 ± 0.4	1.0 ^{+0.4} _{–0.5}	3.7
AzTEC J044738.40–532327.9	ADFS-AzTEC165	4 47 38.40	–53 23 27.9	1.4 ± 0.4	1.1 ^{+0.5} _{–0.5}	3.6
AzTEC J044719.99–532902.4	ADFS-AzTEC166	4 47 19.99	–53 29 02.4	1.8 ± 0.5	1.3 ^{+0.6} _{–0.6}	3.6
AzTEC J044422.24–534544.5	ADFS-AzTEC167	4 44 22.24	–53 45 44.5	1.8 ± 0.5	1.3 ^{+0.6} _{–0.6}	3.6
AzTEC J044410.42–533838.0	ADFS-AzTEC168	4 44 10.42	–53 38 38.0	1.7 ± 0.5	1.2 ^{+0.5} _{–0.6}	3.5
AzTEC J044701.87–532915.7	ADFS-AzTEC169	4 47 01.87	–53 29 15.7	1.7 ± 0.5	1.2 ^{+0.5} _{–0.6}	3.5
AzTEC J044327.57–534158.07	ADFS-AzTEC170	4 43 27.57	–53 41 58.07	6.6 ± 0.7	6.1 ^{+0.8} _{–0.7}	9.5
AzTEC J044435.40–533316.47	ADFS-AzTEC171	4 44 35.40	–53 33 16.47	4.4 ± 0.6	4.0 ^{+0.6} _{–0.6}	7.8
AzTEC J044803.16–532615.99	ADFS-AzTEC172	4 48 03.16	–53 26 15.99	4.6 ± 0.6	4.2 ^{+0.7} _{–0.6}	7.7
AzTEC J044819.80–532444.49	ADFS-AzTEC173	4 48 19.80	–53 24 44.49	4.4 ± 0.6	3.9 ^{+0.6} _{–0.7}	7.3
AzTEC J044421.55–534823.46	ADFS-AzTEC174	4 44 21.55	–53 48 23.46	4.4 ± 0.6	3.9 ^{+0.7} _{–0.7}	7.2
AzTEC J044502.44–534452.07	ADFS-AzTEC175	4 45 02.44	–53 44 52.07	4.2 ± 0.6	3.8 ^{+0.7} _{–0.7}	7.0
AzTEC J044511.21–532400.67	ADFS-AzTEC176	4 45 11.21	–53 24 00.67	4.0 ± 0.7	3.4 ^{+0.7} _{–0.7}	6.0
AzTEC J044636.65–531618.03	ADFS-AzTEC177	4 46 36.65	–53 16 18.03	3.3 ± 0.6	2.9 ^{+0.7} _{–0.6}	5.7
AzTEC J044652.58–531505.07	ADFS-AzTEC178	4 46 52.58	–53 15 05.07	3.1 ± 0.6	2.6 ^{+0.6} _{–0.6}	5.5
AzTEC J044352.49–534647.20	ADFS-AzTEC179	4 43 52.49	–53 46 47.20	3.4 ± 0.6	2.9 ^{+0.7} _{–0.7}	5.4
AzTEC J044335.26–534340.94	ADFS-AzTEC180	4 43 35.26	–53 43 40.94	3.4 ± 0.6	2.8 ^{+0.7} _{–0.7}	5.4
AzTEC J044331.59–533937.66	ADFS-AzTEC181	4 43 31.59	–53 39 37.66	3.6 ± 0.7	3.0 ^{+0.7} _{–0.7}	5.3
AzTEC J044446.10–532828.82	ADFS-AzTEC182	4 44 46.10	–53 28 28.82	3.6 ± 0.7	2.9 ^{+0.8} _{–0.8}	5.1
AzTEC J044813.09–531023.30	ADFS-AzTEC183	4 48 13.09	–53 10 23.30	3.1 ± 0.6	2.5 ^{+0.7} _{–0.7}	4.9
AzTEC J044559.84–534117.11	ADFS-AzTEC184	4 45 59.84	–53 41 17.11	3.0 ± 0.6	2.4 ^{+0.7} _{–0.7}	4.7
AzTEC J044657.28–533204.37	ADFS-AzTEC185	4 46 57.28	–53 32 04.37	2.6 ± 0.6	2.1 ^{+0.7} _{–0.6}	4.5
AzTEC J044853.36–531151.36	ADFS-AzTEC186	4 48 53.36	–53 11 51.36	2.8 ± 0.6	2.2 ^{+0.7} _{–0.7}	4.5
AzTEC J044845.15–531020.30	ADFS-AzTEC187	4 48 45.15	–53 10 20.30	2.9 ± 0.7	2.3 ^{+0.8} _{–0.7}	4.4
AzTEC J044911.57–531409.24	ADFS-AzTEC188	4 49 11.57	–53 14 09.24	3.0 ± 0.7	2.3 ^{+0.8} _{–0.7}	4.4
AzTEC J044739.25–531234.57	ADFS-AzTEC189	4 47 39.25	–53 12 34.57	2.5 ± 0.6	1.9 ^{+0.7} _{–0.7}	4.1
AzTEC J044410.78–534755.51	ADFS-AzTEC190	4 44 10.78	–53 47 55.51	2.4 ± 0.6	1.8 ^{+0.7} _{–0.7}	4.0
AzTEC J044722.60–533044.49	ADFS-AzTEC191	4 47 22.60	–53 30 44.49	2.7 ± 0.7	2.0 ^{+0.8} _{–0.8}	3.9
AzTEC J044639.24–531515.80	ADFS-AzTEC192	4 46 39.24	–53 15 15.80	2.4 ± 0.7	1.7 ^{+0.8} _{–0.8}	3.8
AzTEC J044822.21–532519.16	ADFS-AzTEC193	4 48 22.21	–53 25 19.16	2.6 ± 0.7	1.8 ^{+0.8} _{–0.9}	3.7
AzTEC J044638.14–533806.11	ADFS-AzTEC194	4 46 38.14	–53 38 06.11	2.4 ± 0.6	1.7 ^{+0.7} _{–0.8}	3.7
AzTEC J044607.81–532028.67	ADFS-AzTEC195	4 46 07.81	–53 20 28.67	2.6 ± 0.7	1.8 ^{+0.8} _{–0.9}	3.7
AzTEC J044553.73–532144.72	ADFS-AzTEC196	4 45 53.73	–53 21 44.72	2.4 ± 0.7	1.6 ^{+0.8} _{–0.8}	3.6
AzTEC J044648.62–531447.09	ADFS-AzTEC197	4 46 48.62	–53 14 47.09	2.2 ± 0.6	1.5 ^{+0.8} _{–0.7}	3.6
AzTEC J044831.02–531003.43	ADFS-AzTEC198	4 48 31.02	–53 10 03.43	2.3 ± 0.6	1.5 ^{+0.8} _{–0.8}	3.5




Figure 4. Number of false detections expected above given signal-to-noise ratio thresholds calculated in the 50 and 30–50 per cent coverage regions. The error bars are 1σ Poisson uncertainties.




Figure 5. Survey completeness for the 50 per cent coverage region. The error bars are 1σ from the binomial distribution.

UV/optical galaxies does not significantly change the SMG number counts given the large volume and redshift space sampled by the millimetre-wavelength observations. Compared to the ADF-S, the counts from GOODS-N and COSMOS are higher, while the counts from SHADES are lower. The overdensity of bright SMGs in AzTEC/COSMOS compared to other blank fields has been shown to be correlated with foreground structure at $z \lesssim 1$ (Austermann et al. 2009). Since the GOODS and the SHADES fields have no known biases, the diversity in the number counts likely arises from cosmic variance given the small areas of these surveys.

5.4 Comparison with models

We compare the cumulative number counts with the models of Takeuchi et al. (2001a,b), Franceschini et al. (2010) and Rowan-Robinson (2009) which successfully reproduce the observed CIB and number counts at IR and submillimetre wavelengths. The models of Franceschini et al. (2010) and Rowan-Robinson (2009) are constructed to match observed 1.1-mm number counts of AzTEC




Figure 6. Probability that a source is detected outside an angular distance from its true position. The probability is calculated for sources with $3.5 \leq S/N < 4.0$, $4.5 \leq S/N < 5.0$ and $5.5 \leq S/N < 6.0$. The horizontal dashed lines indicate 68.3 and 99.5 per cent confidence intervals. The shaded regions represent theoretical predictions derived in Ivison et al. (2007) for the S/N ranges.




Figure 7. Differential number counts in the ADF-S, the SXDF and the SSA 22 fields surveyed by AzTEC/ASTE. The counts are calculated for the 50 per cent coverage regions. The error bars represent the 68.3 per cent confidence intervals. The red curve represents the best-fitting Schechter functional form of the ADF-S: $dN/dS = N_{3\text{mJy}}(S/3\text{mJy})^{-1} \exp(-(S - 3\text{mJy})/S')$ with $N_{3\text{mJy}} = 169 \pm 19$ and $S' = 1.48 \pm 0.20$. The number counts of previous 1.1-mm surveys are also shown: GOODS-N (Perera et al. 2008), COSMOS (Austermann et al. 2009), SHADES (Austermann et al. 2010) and GOODS-S (Scott et al. 2010). The horizontal dashed line represents the survey limit of the ADF-S, which Poisson deviations to zero sources per bin 32.7 per cent. The bin centres of some number counts are shifted by ± 3 per cent for easier comparison.

surveys of the COSMOS field (Austermann et al. 2009) and GOODS-N field (Perera et al. 2008), respectively.

The model of Takeuchi et al. (2001a,b) consists of three components: (i) the far-IR (FIR) spectral energy distribution (SED) based on the *IRAS* colour–luminosity relation at 60 and 100 μm , (ii) the local 60- μm luminosity function adopted from the *IRAS* data and (iii) galaxy evolution with redshift. The 60- μm luminosity of a galaxy is




Figure 8. Cumulative number counts in the ADF-S, the SXDF and the SSA 22 fields surveyed by AzTEC/ASTE. The counts are calculated for the 50 per cent coverage regions. The error bars indicate 68.3 per cent confidence intervals. The red curve represents the best-fitting Schechter functional form of the ADF-S. Number counts of previous 1-mm surveys are also shown: the AzTEC/JCMT surveys of GOODS-N (Perera et al. 2008), COSMOS (Austermann et al. 2009), SHADES (Austermann et al. 2010), the AzTEC/ASTE GOODS-S (Scott et al. 2010), 1.2-mm MAMBO surveys of the Lockman Hole and ELAIS N2 regions (Greve et al. 2004). The horizontal dashed line represents the survey limit of the ADF-S, which Poisson deviations to zero sources per bin 32.7 per cent. The bin centres of some number counts are shifted by ± 3 per cent for easier comparison.

Table 3. Differential and cumulative number counts in the ADF-S. The flux bin centres for the differential counts (first column) are weighted by the assumed prior. The errors are 68.3 per cent confidence intervals.

Flux density (mJy)	Differential counts ($\text{mJy}^{-1} \text{deg}^{-2}$)	Flux density (mJy)	Cumulative counts (deg^{-2})
1.39	1074^{+123}_{-137}	1.0	1590^{+134}_{-151}
2.41	312^{+44}_{-51}	2.0	515^{+54}_{-64}
3.42	109^{+24}_{-28}	3.0	204^{+31}_{-38}
4.42	63^{+17}_{-21}	4.0	95^{+20}_{-26}
5.43	25^{+10}_{-14}	5.0	31^{+11}_{-15}
6.43	$5.7^{+3.4}_{-5.7}$	6.0	$5.8^{+3.4}_{-5.7}$

thus described as a function of redshift, assuming pure luminosity evolution:

$$L_{60}(z) = L_{60}(z=0)f(z), \quad (2)$$

where L_{60} is the luminosity at $60 \mu\text{m}$ and $L_{60}(z=0)$ represents the local $60\text{-}\mu\text{m}$ luminosity function. The form of $f(z)$ is a step-wise non-parametric function. Takeuchi et al. (2001b) assume three evolutionary scenarios (see fig. 2 of Takeuchi et al. 2001b) within the permitted range derived from the observed CIB and number counts at 15, 60, 90, 170, 450 and $850 \mu\text{m}$ as follows. (i) Evolution 1: $f(z)$ rises steeply between $z=0$ and 0.5 , reaches $f(z)=10$, is constant from $z=0.5$ to 2 and decreases slowly between $z=2$ and 5 . (ii) Evolution 2: $f(z)$ quickly rises between $z=0$ and 0.5 , peaking with $f(z)=30$ between $z=0.5$ and 0.75 and having a long plateau with $f(z)=10$ between $z=0.75$ and 5 . (iii) Evolution 3: $f(z)$ rises

Table 4. Differential and cumulative number counts in the SXDF and the SSA 22 field. The flux bin centres for the differential counts (first column) are weighted by the prior. The errors are 68.3 per cent confidence intervals.

Flux density (mJy)	Differential counts ($\text{mJy}^{-1} \text{deg}^{-2}$)	Flux density (mJy)	Cumulative counts (deg^{-2})
SXDF			
1.38	1110^{+148}_{-148}	1.0	1578^{+156}_{-158}
2.40	314^{+44}_{-48}	2.0	468^{+50}_{-56}
3.41	95^{+20}_{-23}	3.0	154^{+24}_{-20}
4.41	32^{+11}_{-14}	4.0	59^{+13}_{-19}
5.42	$13^{+5.7}_{-9.0}$	5.0	$27^{+7.6}_{-13}$
6.42	$7.4^{+3.5}_{-6.9}$	6.0	$14^{+5.1}_{-8.9}$
SSA 22			
1.40	572^{+152}_{-178}	1.0	1006^{+163}_{-191}
2.42	251^{+52}_{-60}	2.0	434^{+59}_{-69}
3.43	103^{+24}_{-27}	3.0	182^{+28}_{-34}
4.44	46^{+13}_{-17}	4.0	80^{+16}_{-21}
5.44	$19^{+7.7}_{-11}$	5.0	$34^{+8.8}_{-13}$
6.44	$7.0^{+3.3}_{-6.6}$	6.0	$15^{+4.2}_{-7.0}$

Table 5. Best-fitting parameters of parametric fits to differential number counts in the ADF-S, SXDF and SSA 22 fields using the Schechter functional form in equation (1). The errors are 1σ uncertainty.

Field	$N_{3\text{mJy}}$ ($\text{mJy}^{-1} \text{deg}^{-2}$)	S' (mJy)
ADF-S	169 ± 19	1.48 ± 0.20
SXDF	144 ± 17	1.24 ± 0.15
SSA 22	132 ± 18	1.85 ± 0.38

between $z=0$ and 1 , peaks with $f(z)=30$ between $z=1$ and 2 and has a plateau with $f(z)=20$ between $z=2$ and 5 . They adopt two additional evolution models which are made by modifying evolution 1 to rise to $f(z)=10^{1.5}$ and $f(z)=10^{2.0}$ between $z=1$ and 2 , which we will refer to as ‘evolution 4’ and ‘evolution 5’, respectively. Fig. 9 compares 1.1-mm observed number counts to the five models, along with a no-evolution model. The no-evolution, evolution 1 and evolution 2 models are lower than the observed counts, suggesting that significant luminosity evolution with $f(z) > 10$ is needed. Given that these five models fail to explain the observed counts, the models need to be modified. Possible ways to solve this discrepancy are to (i) change the functional form of the evolutionary scenarios and (ii) make the luminosity evolution dependent on luminosity. These issues are discussed in more detail in Takeuchi et al. (in preparation).

The model of Franceschini et al. (2010) is constructed to reproduce the latest observed counts at 15, 24, 70, 350, 850 and $1100 \mu\text{m}$; the redshift-dependent luminosity functions at $15 \mu\text{m}$ and the CIB. They assume both luminosity and number density evolution and create number counts starting from the *IRAS* $12\text{-}\mu\text{m}$ luminosity function. The model population consists of four galaxy classes: non-evolving normal spirals, Type-I active galactic nuclei (AGNs), starburst galaxies of moderate luminosities (or luminous IR galaxies) and very luminous starburst galaxies [or ultra-luminous IR galaxies (ULIRGs)]. The four populations follow different evolution in luminosity and number density. The model of Franceschini et al. (2010) overestimates the ADF-S counts, while it is consistent




Figure 9. Comparison of 1.1-mm observed number counts with the models of Takeuchi et al. (2001b). Descriptions of the models are in Section 5.4.




Figure 10. Comparison of 1.1-mm observed number counts with the models of Rowan-Robinson (2009) and Franceschini et al. (2010). Descriptions of the models are in Section 5.4.

with the counts of the AzTEC/COSMOS survey. This is not surprising, since the model is created to reproduce the 1.1-mm counts of the AzTEC/COSMOS, where a significant overdensity of sources has been reported (Austermann et al. 2009).

The model of Rowan-Robinson (2009) is created by modifying the model of Rowan-Robinson (2001) to reproduce the latest observed counts, particularly at 24 μm . The model assumes pure luminosity evolution. The model consists of four spectral components: IR cirrus, M82-like starburst, Arp 220-like starburst and AGN dust torus. Rowan-Robinson (2009) creates three models with formation redshifts of $z_f = 4, 5$ and 10. Fig. 10 shows that the model with $z_f = 4$ well describes the ADF-S counts down to the faint end, but at the bright end it overestimates the ADF-S counts.

All of the models presented in this section do not match simultaneously the faint and bright ends of observed counts, requiring the models to be modified.

6 CONTRIBUTION TO THE COSMIC INFRARED BACKGROUND

We estimate the fraction of the CIB resolved by the ADF-S survey. The total deboosted flux density of the $\geq 3.5\sigma$ sources in the 50 per cent coverage region is $1.9 \pm 0.03 \text{ Jy deg}^{-2}$. The expected 1.1-mm background as measured by the *Cosmic Background Explorer* satellite is 18–24 Jy deg^{-2} (Puget et al. 1996; Fixsen et al. 1998); therefore, we have resolved about 7–10 per cent of the CIB into discrete sources. This is similar to the resolved fraction of the CIB reported by other 1-mm blank field surveys (Greve et al. 2004; Laurent et al. 2005; Maloney et al. 2005; Scott et al. 2008, 2010). This could be caused by the survey incompleteness due to the confusion noise and the fewer bright sources in the ADF-S, despite the deeper sensitivity compared to the other surveys.

To estimate the total integrated flux density corrected for the survey incompleteness, and to include fainter sources below the detection threshold, we integrate the best-fitting Schechter function of the ADF-S obtained in Section 5. The integration of the best-fitting function at $\geq 1 \text{ mJy}$, where the number counts are tightly constrained, is 2.9 Jy deg^{-2} , which corresponds to 12–16 per cent of the CIB at 1.1 mm. This suggests that a large fraction of the CIB originates from submillimetre-faint sources for which the number counts have not yet been constrained. Integration of the best-fitting number counts extrapolating to lower fluxes (down to 0 mJy) results in a total flux density of 5.7 Jy deg^{-2} , which is only 24–32 per cent of the CIB at 1.1 mm, suggesting that the faint-end slope of the actual number counts should be steeper than that of the present best-fitting model. It is possible that a Schechter functional form is not appropriate for representing 1.1-mm number counts.

7 REDSHIFT CONSTRAINT

In order to constrain redshifts of AzTEC sources, we compare AzTEC sources in the 30 per cent coverage region with the far-IR images obtained by *AKARI/FIS* (Shirahata et al., in preparation). The ADF-S is observed at four bands: 65, 90, 140 and 160 μm , with FWHMs of 37, 39, 58 and 61 arcsec, respectively. The detection limits of the *AKARI* data are 46.5, 15.7, 183 and 608 mJy (3σ) at 65, 90, 140 and 160 μm , respectively. We compare the AzTEC sources with the 90- μm source catalogue, which is most sensitive and reliable among the four bands, and found that only 11 AzTEC sources are within a 20-arcsec radius from the 90- μm sources. A detailed multiwavelength study of these sources will be made in a future paper. We constrain the redshifts of the AzTEC sources using their flux ratios of 1.1 mm to 90 μm . Fig. 11 shows the expected flux ratio as a function of redshift for two different SED models: Arp 220 (Silva et al. 1998), a typical ULIRG, and the average SED of 76 SMGs with spectroscopic redshifts (Michałowski, Hjorth & Watson 2010). The horizontal dotted lines represent the flux ratios of AzTEC sources with 90- μm counterpart candidates. The shaded region represents upper limits on the flux ratios of the remaining AzTEC sources from the 3σ detection limit at 90 μm . The figure suggests that most of the AzTEC sources are likely to be at $z \gtrsim 1.5$.

At $z > 1$, the flux density of SMGs is nearly redshift independent and thus is proportional to the IR luminosity. By scaling the IR luminosities of the SED models, we estimate IR luminosities of the AzTEC sources to be $\sim 3\text{--}14 \times 10^{12} L_{\odot}$. If the emission is




Figure 11. Observed flux ratios of 1.1 mm to 90 μm as a function of redshift. The solid and dashed curves represent the SED models of Arp 220 (Silva et al. 1998) and the average SED of 76 SMGs with spectroscopic redshifts (Michałowski et al. 2010), respectively. The horizontal dotted lines indicate the flux ratios of AzTEC sources with 90- μm counterpart candidates. The shaded region represents the 3σ upper limits of 90 μm for AzTEC sources without 90- μm detections.

powered solely by star formation activity, the inferred SFRs are $\sim 500\text{--}2400 M_{\odot} \text{ yr}^{-1}$ (Kennicutt 1998).

8 COSMIC STAR FORMATION HISTORY TRACED BY 1.1-MM SOURCES

8.1 Star formation rate density

From UV/optical surveys, the total SFR per unit comoving volume (SFR density) is observed to increase with redshift from $z \sim 7$ to 3, peak at $z \sim 3\text{--}1$ and decline steadily from $z = 1$ to 0 (e.g. Lilly et al. 1996; Madau et al. 1996; Steidel et al. 1999; Giavalisco et al. 2004; Bouwens et al. 2010). It is suggested that SFRs per unit comoving volume (SFR density) increase with redshift from $z \sim 7$ to 3, peaks at $z \sim 1\text{--}3$ and decreases from $z \sim 1$. However, the SFR density derived by UV/optically selected galaxies has large uncertainty due to the extinction by dust. It is also possible that dusty galaxies are missed entirely by these previous studies. Comparatively speaking, millimetre and submillimetre wavelengths have a great advantage in tracing dusty starburst galaxies at high redshifts. Previous submillimetre surveys suggested that SMGs contribute significantly ($\gtrsim 10\text{--}20$ per cent) to the cosmic SFR density at $z \sim 2\text{--}3$ (e.g. Hughes et al. 1998; Chapman et al. 2005; Aretxaga et al. 2007; Dye et al. 2008; Wardlow et al. 2010).

We estimate the SFR density contributed by 1.1-mm sources using the best-fitting number counts in the ADF-S derived in Section 5. FIR luminosities of the 1.1-mm sources are calculated by assuming the SED models of Arp 220 (Silva et al. 1998) and the average SED of SMGs (Michałowski et al. 2010), and SFRs are derived from FIR luminosity using the equation of Kennicutt (1998). The largest uncertainty comes from the lack of redshift information. Since the redshifts of the 1.1-mm sources are not known, we assume redshift distributions based on previous studies. The largest spectroscopic sample of SMGs obtained by Chapman et al. (2005) has a median redshift of $z = 2.2$ with an interquartile range of 1.7–2.8, and the redshift distribution is well fitted by Gaussian. Pope et al. (2006) found a median redshift of $z = 2.2$ with an interquartile range of 1.4–2.6 using spectroscopic and photometric redshifts of SMGs

in the *Hubble Deep Field-North*. Aretxaga et al. (2007) estimated photometric redshifts of SHADES sources and found a median of $z = 2.4$ with an interquartile range of $z = 1.8\text{--}3.1$, and that redshift distribution has a near-Gaussian form. Chapin et al. (2009) derive a higher median redshift of $z = 2.7$ using spectroscopic and photometric redshifts of 1.1-mm sources detected in the AzTEC/JCMT GOODS-N survey.

Based on these measurements, we adopt Gaussian functional forms of redshift distributions with central redshifts of $z_c = 2.2, 2.4$ and 2.7. We assume $\sigma_z = 0.5$ and $\sigma_z = 1.0$ for narrow and broad redshift distributions, respectively. These redshift distributions are consistent with the fact that the most of the AzTEC sources are at $z \gtrsim 1.5$ (see Section 7). We calculate the total flux by integrating the differential number counts at ≥ 1 mJy and distribute the total flux following the assumed redshift distributions. If we integrate the number counts down to 0.1 mJy, the total flux density would increase by about a factor of 2.

The derived SFR densities using the Arp 220 SED model are shown in Fig. 12 and the average values in redshift bins are presented in Table 6. The results derived from the two assumed SED models are consistent within 30 per cent. Although the derived SFR densities largely depend on the assumed redshift distribution, they are within the range of those derived in previous studies with SMGs (Hughes et al. 1998; Chapman et al. 2005; Aretxaga et al. 2007; Dye et al. 2008; Wardlow et al. 2010). In Fig. 12, extinction-corrected




Figure 12. SFR density estimated from the best-fitting number counts in the ADF-S. Four redshift distributions are assumed. The SFR densities are calculated by integrating the number counts at ≥ 1 mJy and using the SED model of Arp 220. The results from previous UV/optical observations are plotted for comparison (compiled data are taken from Hopkins 2004, and references therein).

Table 6. Comoving SFR density averaged in redshift bins estimated from the best-fitting number counts in the ADF-S. Four redshift distributions are assumed. The SFR densities are calculated by integrating the number counts at ≥ 1 mJy and using the SED model of Arp 220.

(z_c, σ_z)	Comoving SFR density ($M_{\odot} \text{ yr}^{-1} \text{ Mpc}^{-3}$)			
	$z = 1\text{--}2$	$z = 2\text{--}3$	$z = 3\text{--}4$	$z = 4\text{--}5$
(2.2, 0.5)	3.0×10^{-2}	4.6×10^{-2}	4.1×10^{-3}	1.2×10^{-5}
(2.4, 0.5)	1.9×10^{-2}	5.1×10^{-2}	8.5×10^{-3}	5.2×10^{-5}
(2.4, 1.0)	2.3×10^{-2}	2.9×10^{-2}	1.6×10^{-2}	3.8×10^{-3}
(2.7, 1.0)	1.7×10^{-2}	2.9×10^{-2}	2.1×10^{-2}	6.5×10^{-3}

SFR densities derived from previous UV/optical observations are also shown for comparison (Hopkins 2004). The SFR densities of 1.1-mm sources are lower by about a factor of 5–10 at $z \sim 2$ –3 compared to those of the UV/optically selected galaxies.

In Section 6, we found that integrating the AzTEC/ADF-S number counts at ≥ 1 mJy accounts for 12–16 per cent of the CIB at 1.1 mm. If we assume that the rest of the CIB comes from fainter sources, the total SFR density contributed by 1.1-mm sources including <1 -mJy sources would increase by about a factor of 6–8, which is comparable to or higher than that of the UV/optically selected galaxies at $z \sim 2$ –3. We note that in this case, the faint 1.1-mm sources and UV/optical sources can overlap. The large contribution of dusty galaxies to the SFR density is suggested by Goto et al. (2010) based on 8- and 12- μ m observations. They found that the dust-obscured SFR density at $z \sim 2$ is $\sim 0.5 M_{\odot} \text{ yr}^{-1} \text{ Mpc}^{-3}$, which is consistent with the SFR density of 1.1-mm sources including the fainter (<1 -mJy) population sources.

8.2 Stellar mass density

We estimate the fraction of the stellar mass in the present-day universe produced by 1.1-mm sources with ≥ 1 mJy by integrating the SFR density derived in the previous section. The present-day stellar mass density is estimated from local luminosity functions (e.g. Cole et al. 2001; Bell et al. 2003; Kajisawa et al. 2009). Cole et al. (2001) derived the present-day stellar mass density of $(5.6 \pm 0.8) \times 10^8 M_{\odot} \text{ Mpc}^{-3}$ assuming a Salpeter (1955) initial mass function (IMF). The time integration of the SFR densities at $z \geq 1$ for the four assumed redshift distributions [$(z_c, \sigma_z) = (2.2, 0.5), (2.4, 0.5), (2.4, 1.0)$ and $(2.7, 1.0)$] yields $1.2 \times 10^8, 1.0 \times 10^8, 1.0 \times 10^8$ and $0.90 \times 10^8 M_{\odot} \text{ Mpc}^{-3}$, respectively. This corresponds to ~ 20 per cent of the present-day stellar mass density. This is an upper limit since materials forming massive stars returned to the interstellar medium (ISM) via stellar winds and supernovae explosions. The fraction of stellar mass returned to the ISM, called as the recycled fraction, is estimated in semi-analytical models (e.g. Cole et al. 2000; Baugh et al. 2005; Gonzalez et al. 2010; Lacey et al. 2010), and it depends on the IMF: 0.41 for the Kennicutt (1983) IMF and 0.91 for a top-heavy IMF (Cole et al. 2000; Lacey et al. 2010). If we assume the recycled fractions of 0.41 and 0.91, the fraction of the stellar mass in the present-day universe produced by 1.1-mm sources with ≥ 1 mJy decreases to ~ 10 and a few per cent, respectively.

9 SUMMARY

We performed a 1.1-mm deep survey of the ADF-S with AzTEC mounted on the ASTE, obtaining one of the deepest and widest maps at millimetre wavelengths. The 30 and 50 per cent coverage regions have areas of 909 and 709 arcmin², and noise levels of 0.32–0.71 and 0.32–0.55 mJy, respectively. We detected 198 previously unknown millimetre-bright sources with 3.5σ – 15.6σ in the 30 per cent coverage region, providing the largest 1.1-mm source catalogue from a contiguous region.

We constructed differential and cumulative number counts in the ADF-S, the SXDF and the SSA 22 field which probe fainter flux densities (down to 1 mJy) compared to previous surveys except for the AzTEC/ASTE GOODS-S survey. On the whole, the 1-mm counts of various surveys are consistent within errors. We compare the number counts with the luminosity evolution models of Takeuchi et al. (2001b), Franceschini et al. (2010) and Rowan-Robinson (2009). Comparison with the Takeuchi et al. (2001b) model suggests that

a luminosity evolution with a factor of >10 is needed to explain the observed number counts. The observed number counts favour the model of Rowan-Robinson (2009) with $z_f = 4$, but none of these models simultaneously match both the bright and faint ends of the number counts from 1 to 10 mJy.

In the ADF-S survey, we resolve about 7–10 per cent of the CIB at 1.1 mm into discrete sources. The integration of the best-fitting number counts in the ADF-S down to 1 mJy reaches 12–16 per cent of the CIB. This suggests that the large fraction of the CIB at 1.1 mm originates from faint sources ($S_{1.1\text{mm}} < 1$ mJy) for which the number counts have not yet been constrained. The integration of the best-fitting number counts extrapolating to 0 mJy accounts for only 24–32 per cent of the CIB, suggesting that the faint-end slope of the number counts is steeper than that given by our best-fitting model.

The redshifts of the AzTEC sources are constrained from their flux ratios of 1.1 mm to 90 μ m from the AKARI/FIS. Most of the AzTEC sources are not detected at 90 μ m, suggesting that they are likely to be at $z \gtrsim 1.5$. Assuming $z \gtrsim 1$, the inferred IR luminosities of the AzTEC sources are $\sim (3\text{--}14) \times 10^{12} L_{\odot}$, and their SFRs inferred from the IR luminosities are $\sim 500\text{--}2400 M_{\odot} \text{ yr}^{-1}$.

We derived the cosmic SFR density contributed by 1.1-mm sources using the best-fitting model to the differential number counts. Although the derived SFR density largely depends on the assumed redshift distribution, our estimates are within the range of those derived in previous studies with SMGs. The SFR density of 1.1-mm sources with ≥ 1 mJy at $z \sim 2$ –3 is lower by about a factor of 5–10 than that of UV/optically selected galaxies. If we consider the fact that the contribution of 1.1-mm sources with ≥ 1 mJy to the CIB at 1.1 mm is 12–16 per cent, the SFR density of 1.1-mm sources including those fainter than $S_{1.1\text{mm}} < 1$ mJy would become comparable to or higher than that of UV/optically selected galaxies. The fraction of the present-day stellar mass of the universe produced by 1.1-mm sources with ≥ 1 mJy at $z \geq 1$ is ~ 20 per cent, calculated by the time integration of the SFR density. If we consider the recycled fractions of 0.41 and 0.91, the fraction of stellar mass produced by 1.1-mm sources becomes ~ 10 and a few per cent, respectively.

ACKNOWLEDGMENTS

We are grateful to K. Shimasaku, H. Murakami, S. Okumura, N. Yoshida, T. Kodama, and the anonymous referee for helpful comments and suggestions. We would like to thank Alberto Franceschini for providing the data of his number counts models. BH and YT were financially supported by a Research Fellowship from the JSPS for Young Scientists. TTT was supported by Program for Improvement of Research Environment for Young Researchers from Special Coordination Funds for Promoting Science and Technology, and the Grant-in-Aid for the Scientific Research Fund (20740105) commissioned by the Ministry of Education, Culture, Sports, Science and Technology (MEXT) of Japan. TTT was partially supported from the Grant-in-Aid for the Global COE Program ‘Quest for Fundamental Principles in the Universe: from Particles to the Solar System and the Cosmos’ from the MEXT. This study was supported by the MEXT Grant-in-Aid for Specially Promoted Research (no. 20001003). AzTEC/ASTE observations were partly supported by the Grant-in-Aid for the Scientific Research from the Japan Society for the promotion of Science (no. 19403005) The ASTE project is driven by Nobeyama Radio Observatory (NRO), a branch of NAOJ, in collaboration with University of Chile, and Japanese institutes including University of Tokyo, Nagoya University, Osaka Prefecture University, Ibaraki University and Hokkaido University.

Observations with ASTE were in part carried out remotely from Japan by using NTT's GEMnet2 and its partner R&E (Research and Education) networks, which are based on AccessNova collaboration of University of Chile, NTT Laboratories and NAOJ. AzTEC analysis performed at UMass is supported in part by NSF grant 0907952.

REFERENCES

- Aretxaga I. et al., 2007, *MNRAS*, 379, 1571
 Austermann J. E. et al., 2009, *MNRAS*, 393, 1573
 Austermann J. E. et al., 2010, *MNRAS*, 401, 160
 Baugh C. M., Lacey C. G., Frenk C. S., Granato G. L., Silva L., Bressan A., Benson A. J., Cole S., 2005, *MNRAS*, 356, 1191
 Bell E. F., McIntosh D. H., Katz N., Weinberg M. D., 2003, *ApJS*, 149, 289
 Bertoldi F. et al., 2007, *ApJS*, 172, 132
 Blain A. W., Smail I., Ivison R. J., Kneib J.-P., Frayer D. T., 2002, *Phys. Rep.*, 369, 111
 Borys C., Chapman S., Halpern M., Scott D., 2003, *MNRAS*, 344, 385
 Bouwens R. J. et al., 2010, *ApJ*, 709, L133
 Capak P. et al., 2008, *ApJ*, 681, L53
 Chapin E. L. et al., 2009, *MNRAS*, 398, 1793
 Chapman S. C., Blain A. W., Smail I., Ivison R. J., 2005, *ApJ*, 622, 772
 Clements D. L., Bendo G., Pearson C., Khan S. A., Matsuura S., Shirahata M., 2010, *MNRAS*, in press (arXiv:1009.3388)
 Cole S., Lacey C. G., Baugh C. M., Frenk C. S., 2000, *MNRAS*, 319, 168
 Cole S. et al., 2001, *MNRAS*, 326, 255
 Coppin K., Halpern M., Scott D., Borys C., Chapman S., 2005, *MNRAS*, 357, 1022
 Coppin K. et al., 2006, *MNRAS*, 372, 1621
 Coppin K. E. K. et al., 2009, *MNRAS*, 395, 1905
 Cowie L. L., Barger A. J., Kneib J.-P., 2002, *AJ*, 123, 2197
 Daddi E. et al., 2009, *ApJ*, 694, 1517
 Dye S. et al., 2008, *MNRAS*, 386, 1107
 Eales S., Lilly S., Webb T., Dunne L., Gear W., Clements D., Yun M., 2000, *AJ*, 120, 2244
 Ezawa H., Kawabe R., Kohno K., Yamamoto S., 2004, in Oschmann J. M., ed., *Proc. SPIE Vol. 5489, Ground-Based Telescopes*. SPIE, Bellingham, p. 763
 Ezawa H. et al., 2008, in Stepp L. M., Gilmozzi R., eds, *Proc. SPIE Vol. 7012, Ground-Based and Airborne Telescopes II*. SPIE, Bellingham, p. 6
 Fixsen D. J., Dwek E., Mather J. C., Bennett C. L., Shafer R. A., 1998, *ApJ*, 508, 123
 Franceschini A., Rodighiero G., Vaccari M., Berta S., Marchetti L., Mainetti G., 2010, *A&A*, 517, A74
 Giavalisco M. et al., 2004, *ApJ*, 600, L103
 Glenn J. et al., 1998, in Phillips T. G., ed., *Proc. SPIE Vol. 3357, Advanced Technology MMW, Radio, and Terahertz Telescopes*. SPIE, Bellingham, p. 326
 Gonzalez J. E., Lacey C. G., Baugh C. M., Frenk C. S., 2010, preprint (arXiv:1006.0230)
 Goto T. et al., 2010, *A&A*, 514, A6
 Greve T. R., Ivison R. J., Bertoldi F., Stevens J. A., Dunlop J. S., Lutz D., Carilli C. L., 2004, *MNRAS*, 354, 779
 Greve T. R. et al., 2005, *MNRAS*, 359, 1165
 Greve T. R., Pope A., Scott D., Ivison R. J., Borys C., Conselice C. J., Bertoldi F., 2008, *MNRAS*, 389, 1489
 Hatsukade B. et al., 2010, *ApJ*, 711, 974
 Hayashino T. et al., 2004, *AJ*, 128, 2073
 Hogg D. W., Turner E. L., 1998, *PASP*, 110, 727
 Holland W. S. et al., 1999, *MNRAS*, 303, 659
 Hopkins A. M., 2004, *ApJ*, 615, 209
 Hughes D. H. et al., 1998, *Nat*, 394, 241
 Ikarashi S. et al., 2010, *MNRAS*, submitted (arXiv:1009.1455)
 Ivison R. J. et al., 2007, *MNRAS*, 380, 199
 Kajisawa M. et al., 2009, *ApJ*, 702, 1393
 Kamazaki T. et al., 2005, *ASP Conf. Ser. Vol. 347, Astronomical Data Analysis Software and Systems XIV*. Astron. Soc. Pac., San Francisco, p. 533
 Kawada M. et al., 2007, *PASJ*, 59, 389
 Kennicutt R. C., Jr, 1983, *ApJ*, 272, 54
 Kennicutt R. C., Jr, 1998, *ARA&A*, 36, 189
 Knudsen K. K., van der Werf P. P., Kneib J.-P., 2008, *MNRAS*, 384, 1611
 Knudsen K. K., Kneib J.-P., Richard J., Petitpas G., Egami E., 2010, *ApJ*, 709, 210
 Kreysa E. et al., 1998, in Phillips T. G., ed., *Proc. SPIE Vol. 3357, Advanced Technology MMW, Radio, and Terahertz Telescopes*. SPIE, Bellingham, p. 319
 Lacey C. G., Baugh C. M., Frenk C. S., Benson A. J., Orsi A., Silva L., Granato G. L., Bressan A., 2010, *MNRAS*, 405, 2
 Lagache G., Puget J.-L., Dole H., 2005, *ARA&A*, 43, 727
 Laurent G. T. et al., 2005, *ApJ*, 623, 742
 Lilly S. J., Le Fevre O., Hammer F., Crampton D., 1996, *ApJ*, 460, L1
 Madau P., Ferguson H. C., Dickinson M. E., Giavalisco M., Steidel C. C., Fruchter A., 1996, *MNRAS*, 283, 1388
 Maloney P. R. et al., 2005, *ApJ*, 635, 1044
 Matsuda Y. et al., 2005, *ApJ*, 634, L125
 Matsuhara H., Shibai H., Onaka T., Usui F., 2005, *Advances Space Res.*, 36, 1091
 Michałowski M., Hjorth J., Watson D., 2010, *A&A*, 514, A67
 Murdoch H. S., Crawford D. F., Jauncey D. L., 1973, *ApJ*, 183, 1
 Onaka T. et al., 2007, *PASJ*, 59, 401
 Perera T. A. et al., 2008, *MNRAS*, 391, 1227
 Pope A. et al., 2006, *MNRAS*, 370, 1185
 Puget J.-L., Abergel A., Bernard J.-P., Boulanger F., Burton W. B., Desert F.-X., Hartmann D., 1996, *A&A*, 308, L5
 Riechers D. A. et al., 2010, *ApJ*, 720, L131
 Rowan-Robinson M., 2001, *ApJ*, 549, 745
 Rowan-Robinson M., 2009, *MNRAS*, 394, 117
 Salpeter E. E., 1955, *ApJ*, 121, 161
 Schlegel D. J., Finkbeiner D. P., Davis M., 1998, *ApJ*, 500, 525
 Scott K. S. et al., 2008, *MNRAS*, 385, 2225
 Scott K. S. et al., 2010, *MNRAS*, 405, 2260
 Shirahata M. et al., 2009, *PASJ*, 61, 737
 Silva L., Granato G. L., Bressan A., Danese L., 1998, *ApJ*, 509, 103
 Siringo G. et al., 2009, *A&A*, 497, 945
 Smail I., Ivison R. J., Blain A. W., 1997, *ApJ*, 490, L5
 Smail I., Chapman S. C., Blain A. W., Ivison R. J., 2004, *ApJ*, 616, 71
 Steidel C. C., Adelberger K. L., Dickinson M., Giavalisco M., Pettini M., Kellogg M., 1998, *ApJ*, 492, 428
 Steidel C. C., Adelberger K. L., Giavalisco M., Dickinson M., Pettini M., 1999, *ApJ*, 519, 1
 Steidel C. C., Adelberger K. L., Shapley A. E., Pettini M., Dickinson M., Giavalisco M., 2000, *ApJ*, 532, 170
 Swinbank A. M., Smail I., Chapman S. C., Blain A. W., Ivison R. J., Keel W. C., 2004, *ApJ*, 617, 64
 Swinbank A. M. et al., 2010, *Nat*, 464, 733
 Tacconi L. J. et al., 2006, *ApJ*, 640, 228
 Takeuchi T. T., Ishii T. T., 2004, *ApJ*, 604, 40
 Takeuchi T. T., Ishii T. T., Hirashita H., Yoshikawa K., Matsuhara H., Kawara K., Okuda H., 2001a, *PASJ*, 53, 37
 Takeuchi T. T., Kawabe R., Kohno K., Nakanishi K., Ishii T. T., Hirashita H., Yoshikawa K., 2001b, *PASP*, 113, 586
 Tamura Y. et al., 2009, *Nat*, 459, 61
 Wardlow J. L. et al., 2010, preprint (arXiv:1006.2137)
 Weiß A. et al., 2009, *ApJ*, 707, 1201
 Wilson G. W. et al., 2008a, *MNRAS*, 386, 807
 Wilson G. W. et al., 2008b, *MNRAS*, 390, 1061

This paper has been typeset from a $\text{\TeX}/\text{\LaTeX}$ file prepared by the author.