



American Journal
of Speech-Language
Pathology

Investigating a Multimodal Intervention for Children with Limited Expressive Vocabularies Associated with Autism

Journal:	<i>American Journal of Speech-Language Pathology</i>
Manuscript ID:	AJSLP-14-0093.R2
Manuscript Type:	Research Article
Date Submitted by the Author:	13-Mar-2015
Complete List of Authors:	Brady, Nancy; University of Kansas, Scheifelhush Institute for Life Span Studies; Speech Language Hearing Sciences and Disorders Storkel, Holly; University of Kansas, Speech-Language-Hearing: Sciences and Disorders Bushnell, Paige; University of Kansas, Speech-Language-Hearing: Sciences and Disorders Barker, Robert; University of Kansas, Speech-Language-Hearing: Sciences and Disorders Saunders, Kathryn; University of Kansas, Life Span Institute Daniels, Debby; University of Kansas Medical Center, Hearing and Speech Fleming, Kandace; University of Kansas, Life Span Institute
Keywords:	Autism, Intervention, Augmentative and alternative communication, Speech

SCHOLARONE™
Manuscripts

1 INVESTIGATING A MULTIMODAL INTERVENTION

1

2
3 Running head: INVESTIGATING A MULTIMODAL INTERVENTION

4
5
6
7
8 Investigating a Multimodal Intervention for Children with Limited Expressive Vocabularies Associated
9 with Autism

10
11
12 Nancy C. Brady, Holly L. Storkel, Paige Bushnell, R. Michael Barker, Kate Saunders, Debby Daniels and

13
14
15 Kandace Fleming

16
17
18 University of Kansas

19
20
21
22
23
24
25
26 Author Note

27
28
29 Nancy C. Brady, Department of Speech Language Hearing Sciences and Disorders, University of
30 Kansas; Holly L. Storkel, Department of Speech Language Hearing Sciences and Disorders, University of
31 Kansas; Paige Bushnell, Department of Speech Language Hearing Sciences and Disorders, University of
32 Kansas; R. Michael Barker, Schiefelbusch Institute of Lifespan Studies, University of Kansas; Kate
33 Saunders, Schiefelbusch Institute of Lifespan Studies, University of Kansas; Debby Daniels, Department
34 of Hearing and Speech, University of Kansas Medical Center; Kandace Fleming, Schiefelbusch Institute
35 of Lifespan Studies, University of Kansas

36
37
38 R. Michael Barker is now in the Department of Communication Sciences and Disorders,
39 University of South Florida

40
41
42 This research was supported by a Pilot Research grant from Autism Speaks and NIH grant NICHD
43 P30 HD002528; NICHD 0070511; and T32 HD057844

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

INVESTIGATING A MULTIMODAL INTERVENTION

2

Abstract

Purpose: To investigate a new intervention package aimed at increasing expressive word learning by school-age children with autism who have limited expressive vocabularies. This pilot investigation was intended to show proof of concept.

Method: Ten children between the ages of 6-10 years with educational diagnoses of autism and limited expressive vocabularies at the outset of the study participated. A multimodal intervention composed of speech sound practice and AAC was used to teach individualized vocabulary words that were selected based on initial speech sound repertoires and principles of phonotactic probability and neighborhood density. A multiple-probe design was used to evaluate learning outcomes.

Results: Five children showed gains in spoken- word learning across successive word sets (High Responders). Five children did not meet learning criteria (Low Responders). Comparisons of behaviors measured prior to intervention indicated that High Responders had relatively higher skills in receptive language, prelinguistic communication, vocal/verbal imitation, adaptive behavior and consonant productions.

Conclusions: The intervention package holds promise for improving spoken word productions for some children with autism who have limited expressive vocabularies. Further research is needed to better describe who may most benefit from this approach as well as investigate generalized benefits to untaught contexts and targets.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

INVESTIGATING A MULTIMODAL INTERVENTION

3

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29

Despite promising results of intensive early interventions, approximately one third to one half of school-age children with autism do not use speech as a primary communication mode (National Research Council, 2001). Remaining nonverbal past the age of 5 years has been considered a poor prognostic indicator for future language development (Billstedt, Carina Gillberg, & Gillberg, 2007; Pickett, Pullara, O'Grady, & Gordon, 2009). Although there have been reports of individuals older than 5 acquiring speech (e.g., Pickett, et. al., 2009), the characteristics of the successful individuals and the interventions employed are not fully understood. The present study is a pilot intervention aimed at teaching speech in combination with Augmentative or Alternative Communication (AAC) to a group of school-age children with autism diagnoses, and to describe characteristics that appeared to be associated with differential outcomes. The target population for this intervention is individuals who have very limited or minimal verbal skills-- for example, less than 20 words or stereotyped phrases produced in functional contexts (c.f., Kasari, Brady, Lord, & Tager-Flusberg, 2013).

Implications of Preschool Intervention Research for School-Aged Children

30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Most intervention studies aimed at teaching beginning speech and language skills to children with autism have focused on preschool ages or younger (Goods, Ishijima, Chang, & Kasari, 2013; Rogers et al., 2012; Vismara, Comlombi, & Rogers, 2009). The focus on ages 18 months – 5 years is logical given that this is the age range during which the need for direct language intervention becomes apparent. In addition, intensive interventions are implemented early in hopes of preventing further gaps in language development. However, results from early intensive interventions have been mixed. For example, the Early Start Denver Model (ESDM) is a comprehensive intervention that targets language skills as well as other cognitive and adaptive skills in young children with autism. In a randomized clinical trial study of toddlers with Autism Syndrome Disorders (ASDs), Dawson and colleagues (2010) found that ESDM delivered in the home by trained therapists to children between the ages of 18-30 months was significantly better than standard care in improving composite developmental scores on the Mullen scales of early learning (Mullen, 1995) and adaptive-behavior raw scores measured with the Vineland Adaptive

INVESTIGATING A MULTIMODAL INTERVENTION

4

1
2
3 behavior scale-2 (Sparrow, Cicchetti, & Balla, 2005). However, in a follow-up study that taught parents
4 how to provide ESDM to children between 14-20 months with ASD, no significant child learning
5 outcomes were reported after 12 weeks of intervention (Rogers et al., 2012). The authors speculated that
6 more gains may have been detected if proximal measures directly linked to target behaviors had been
7 used, in addition to the standardized test outcomes. In addition, post-hoc analyses found that children who
8 didn't receive the Parent-Implemented ESDM received more services from community providers, which
9 may have affected results.
10
11
12
13
14
15
16
17

18
19 Other studies have focused more specifically on communication outcomes. For example, Goods
20 and colleagues (2013) demonstrated that a pilot intervention based on Joint Attention Symbolic Play
21 Engagement and Regulation (JASPER) (Kasari, Gulsrud, Wong, Kwon, & Locke, 2010) led to gains in
22 play skills and communicative gestures in children between the ages of 3-5 with ASD. However, this
23 limited intervention (24 sessions) did not show changes in word productions.
24
25
26
27
28

29
30 In each of these studies, results were presented in terms of group comparisons, therefore it is
31 difficult to determine individual responsiveness to the intervention. Warren et al (Warren et al., 2011)
32 completed a meta-analysis with a wide age range of children with Autism Spectrum Disorders (ASD) and
33 concluded that there is a need to better characterize subgroups of children who respond differentially to
34 various intervention approaches. This type of characterization is difficult to complete when only group-
35 level analyses are performed.
36
37
38
39
40
41
42

43 Thus, there is growing evidence for interventions directed to children between the ages of 18
44 months and 5 years. However, many of these strategies may not be appropriate for older, school-age
45 children who remain nonverbal. Several of the approaches mentioned above rely on parent
46 implementation and these approaches may not be feasible when children are attending full-day school
47 programs. In addition, some intervention components, such as play, may need to be modified to be age--
48 as well as developmentally-- appropriate. Lastly, school age children who have not yet started talking
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 may need more intensive interventions that target speech deficits while also teaching augmentative means
4
5 of communication.
6

7 8 **The Need for a Multimodal Approach** 9

10 Most intervention research for nonverbal children with autism has focused on either AAC (Ganz
11 et al., 2012; Mirenda & Bopp, 2003; Schlosser & Wendt, 2008) or speech (Rogers et al., 2006), but not
12 both. Brady and colleagues followed the development of 42 children with autism between the ages of 3-6
13 and noted that children who remained nonverbal were unlikely to receive speech intervention when they
14 reached elementary school (Brady, Thiemann-Bourque, Fleming, & Matthews, 2013). The fact that many
15 children with autism remain nonverbal despite considerable efforts to improve communication suggests
16 the need for an intensive intervention that combines components into one multimodal intervention
17 package. Multimodal refers to the combined use of speech and AAC such as a speech generating device
18 during intervention.
19
20
21
22
23
24
25
26
27
28
29

30 There is good reason that previous efforts have focused primarily on AAC. Speech is likely to
31 remain difficult for some children with autism, whereas they may have more immediate communicative
32 success with AAC. The main goal for AAC interventions is to improve expressive communication using
33 AAC. However, some research studies have reported collateral improvements in speech following AAC.
34 For example, collateral gains in speech were reported for some students following intervention with the
35 Picture Exchange Communication System (PECS; Bondy & Frost, 1994) (Carr & Felce, 2007; Flippin,
36 Reszka, & Watson, 2010). Note that a wide age range of children have participated in the PECS studies.
37 For example, the ages of the children in the Carr and Felce (2007) were between 3-7 years. Kasari and
38 colleagues (2014) found that an intervention that added an SGD to a joint attention and play intervention
39 resulted in significantly better communication outcomes, including speech outcomes, compared to the
40 same intervention without the SGD. Children in the Kasari et al. (2014) study were between the ages of 5-
41 8 and had minimal expressive verbal skills at the outset of intervention. Results were reported in terms of
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

INVESTIGATING A MULTIMODAL INTERVENTION

6

1
2
3 group differences, however, limiting the ability to determine speech gains by individual children, or skill
4
5 sets that may be associated with relative differences in communication and speech outcomes.
6
7

8 The PECS studies and the Kasari et al. (2014) study referenced above measured speech outcomes
9
10 but did not specifically target speech as a part of intervention. Studies directly targeting speech in school-
11
12 age children with autism and minimal expressive vocabularies are difficult to find. For example, Rogers
13
14 and colleagues directly taught participants speech skills using the PROMPT method (Rogers et al., 2006),
15
16 but children were all below the age of 65 months. One study directly taught speech along with SGD use
17
18 in a multimodal approach to children between the ages of 4-8, but participants had severe speech-sound
19
20 disorders not associated with autism (King, Hengst, & DeThorne, 2013). Similar research is needed to
21
22 investigate potential gains made by combining speech and AAC interventions for children with autism.
23
24

25 Multimodal approaches such as that used in the King et al. (2013) study have several potential
26
27 advantages over approaches that focus on either AAC or speech alone. A combined multi-modal approach
28
29 is designed to quickly improve communication success through AAC. AAC responses are learned more
30
31 quickly than speech because they are easier to teach through physical prompts. Learning visually based
32
33 responses is (typically) facilitated by presenting a fixed array of choices. In addition, participants obtain
34
35 linguistic input from both AAC and speech models (Binger & Light, 2007; Harris & Reichle, 2004;
36
37 Sevcik, Ronski, Watkins, & Deffebach, 1995). In terms of speech gains, instead of waiting for collateral
38
39 gains, some participants may benefit from directly teaching speech sounds that comprise targeted
40
41 vocabulary within a multimodal approach. Speech gains are likely to come more slowly than AAC gains,
42
43 but practicing the motor movements required for speech-sound productions may improve speech learning
44
45 while also reinforcing word learning (Vihman, DePaolis, & Keren-Portnoy, 2014).
46
47
48

Teaching Contexts

49
50
51 The current study incorporated teaching contexts and strategies that have been shown to be
52
53 effective in previous research—speech-sound practice using massed trials, joint book reading, interactive
54
55 routines incorporating AAC, and receptive vocabulary trials. Massed-trial practice provides multiple
56
57
58
59
60

1
2
3 opportunities, or trials, in succession, within a short period of time. This strategy can be particularly
4
5 helpful for initial phases of learning (Heflin & Alberto, 2001). Repeated opportunities to practice and
6
7 receive feedback for articulatory movements have been found to improve productions (Pomaville &
8
9 Kladopoulos, 2013). Scripted communication routines are teaching contexts that provide multiple
10
11 opportunities for communication within motivating activities such as playing a social game, or making
12
13 and eating a snack, or more advanced thematic play routines (Goldstein, Wickstrom, Hoyson, Jamieson,
14
15 & Odom, 1988; Kashinath, Woods, & Goldstein, 2006; Rollins, Wambacq, Dowell, Mathews, & Reese,
16
17 1998).

20
21 An additional teaching context that has been investigated in recent studies is *Joint book reading*.
22
23 In joint book reading, an interventionist “reads” from a storybook, providing repeated exposures to
24
25 targeted vocabulary paired with pictures (referents) (Bellon, Ogletree, & Harn, 2000; Fleury, Miramontez,
26
27 Hudson, & Schwartz, 2013). Communication opportunities can be created by pausing during joint book
28
29 reading to allow the child to respond with a specific vocabulary item or re-tell part of the book. Although
30
31 studies have focused primarily on receptive vocabulary gains associated with joint book reading (Bellon
32
33 et al., 2000; Hargrave & Sénéchal, 2000; Whitehurst, Falco, Lonigan, & Fischel, 1988), gains in
34
35 vocabulary production have also been noted (King et al., 2013; Soto & Dukhovny, 2008).

37 38 **Target Vocabulary Identification.**

39
40 Target vocabulary may be selected based on numerous criteria including frequency of use across
41
42 contexts (Snodgrass, Stoner, & Angell, 2013), reinforcement value (Charlop-Christy, Carpenter, LeBlanc,
43
44 & Kellet, 2002) or developmental appropriateness (Solomon-Rice & Soto, 2014). In the current study we
45
46 took a different approach and selected vocabulary based on principles of phonotactic probability and word
47
48 neighborhood (Storkel, 2001; Storkel, Maekawa, & Hoover, 2010), and in consideration of each child’s
49
50 speech sound repertoire. According to these principles, words that have higher frequency phonological
51
52 forms (i.e., high probability) that are phonologically similar to many other words (i.e., high density) are
53
54 easier to learn because these characteristics facilitate holding the word in working memory (Gathercole,
55
56
57
58
59
60

INVESTIGATING A MULTIMODAL INTERVENTION

8

1
2
3 Frankish, Pickering, & Peaker, 1999; Thomson, Richardson, & Goswami, 2005) as well as retrieving the
4 word from long-term memory for production (German & Newman, 2004; Newman & German, 2005).
5
6 Further, in accordance with these principles, children are more likely to learn and use words comprised of
7 higher frequency phonological forms that are phonologically similar to many other words (e.g., cat) than
8 low frequency forms that are phonologically similar to few other words (e.g., juice).
9

10
11
12
13
14 Research has also shown that children are more likely to learn new words if they contain
15 phonemes that they consistently produce in spontaneous vocalizations (MacRoy-Higgins, Schwartz,
16 Shafer, & Marton, 2013; Schwartz, Leonard, Messick, & Chapman, 1987). For example, if a child
17 produces sounds /s/ and /d/ but not the sounds /l/ or /k/ during their vocalizations, learning to say /sad/
18 should be easier than learning to say /luck/. As described below under methods, in the current study we
19 identified target vocabulary based on children's existing speech sound repertoires and principles of
20 phonotactic probability and neighborhood density.
21
22
23
24
25
26
27
28

Predicting Response to the Intervention

29
30
31
32 This study was the first attempt to apply our multimodal intervention for word learning. Based on
33 previous research with children with limited expressive vocabularies, we anticipated that not all
34 participants would respond favorably (Odom et al., 2007). In addition to piloting procedures and
35 determining if children learned to produce new words as a result of the intervention, we wanted to
36 identify learner characteristics that may be associated with different outcomes. Numerous variables have
37 been identified as predictive of language outcomes including developmental level, play, and level of joint
38 attention (Kasari, Paparella, Freeman, & Jahromi, 2008; Rogers et al., 2006; Rogers, Hepburn,
39 Stackhouse, & Wehner, 2003; Toth, Munson, N. Meltzoff, & Dawson, 2006); object interest (Sherer &
40 Schreibman, 2005; Vivanti, Dissanayake, Zierhut, & Rogers, 2013); and nonverbal IQ (Thurm, Lord, Lee,
41 & Newschaffer, 2007). Particularly relevant to the current study is research showing strong associations
42 between expressive language and the following variables: imitation, receptive language and early social
43 communication (Bopp & Mirenda, 2011; Poon, Watson, Baranek, & Poe, 2012; Rogers et al., 2006;
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

INVESTIGATING A MULTIMODAL INTERVENTION

9

Rogers et al., 2003). Past studies on predictive relationships have focused more on non-experimental studies demonstrating a longitudinal relationship between predictors and language outcomes. However, there are clinical implications of these studies, including identifying children who may respond better to one type of intervention compared to another.

Purpose of the Current Study.

The current study was a pilot investigation aimed at demonstrating proof of concept for a new multi-modal intervention that combines AAC and speech sound practice for children with autism and minimal expressive vocabularies. Our purposes were (1) to determine if participants showed gains in spoken word production, (2) determine if gains were also made in receptive word learning, and (3) supposing that some but not all participants would show gains during intervention, compare the profiles of children who responded favorably to those who responded less favorably in an effort to identify possible predictors of response to this specific multi-modal intervention. As with any pilot study, an overarching goal was to determine if the results are promising enough to follow the research with a larger study.

Methods**Overview**

Ten children with autism participated in a multi-modal intervention aimed at teaching new word production, and measuring comprehension of these new words. A single-subject design (multiple-probes across-sets-of-vocabulary words) was used to evaluate the success of the intervention package for each child. Target words were selected individually for each child based on the principles of phonotactic probability and neighborhood density. Specifically, words with high probability and high density were selected for each child. In addition, selected words had to include only sounds that were in the child's phonetic repertoire. Intervention sessions were between 45 min. to an hour each and occurred, on average, 4 days per week. Although the total length of intervention depended on the number of sessions to reach learning criteria, the range of the number of intervention sessions was 17 (for a child who did not learn

any words) to 76. After all 10 participants finished participating in the intervention, we evaluated the relative success of individual participants in light of pre-intervention skills in adaptive behavior, receptive language, early communication, and imitation.

Participants

Ten children, nine of whom were boys, participated. The chronological ages of the children were between 7 years 5 months and 11 years 3 months at the time each child began participating in baseline sessions. A brief description of demographic information and communication status at the time of our initial assessment is summarized in Table 1.

All of the children were attending special programs for children with autism in a local school district and had educational diagnoses of autism. Each child had a confirmed diagnosis of autism, according to parent and school reports. Diagnoses were made by professionals not associated with the current research study, based on DSM IV criteria. Parents reported that their children were diagnosed by pediatricians when the children were between 1 and 5 years of age. Two children were diagnosed based on both the CARS and the Autism Diagnostic Observation Schedule (ADOS; Lord, Rutter, DiLavore, & Risi, 1999) while 1 participant was diagnosed based only on the ADOS. Parents could not recall the instrument used to diagnose the remaining children. Participant 1 also had Down syndrome in addition to autism. Consensus clinical diagnoses such as this have been used to describe participants' autism status in past studies (e.g., Gray, Tonge, & Sweeney, 2008; Smith, Barker, Seltzer, Abbeduto, & Greenberg, 2012; Thiemann-Bourque, Brady, & Fleming, 2011), however, we are unable to describe autism severity across participants because we didn't apply a consistent autism measure to all participants.

Our goal was to recruit participants who met the following criteria for minimal expressive vocabulary: less than 20 spoken words produced spontaneously according to (1) teacher report; (2) parent report and (3) a language sample collected during our assessment process. All of the participants met the criteria for teacher report and for productions recorded during our language sample. However, mothers of Participants 1 and 2 indicated on the Vineland Adaptive Behavior Scales (VABS-2) that their child was

1
2
3 able to say 50 different words. We included both of these children in the study because teachers indicated
4
5 the children had vocabularies of less than 20 words, and no words were produced during a 30-minute
6
7 language sample completed by our research staff. Given that our research design was based on data from
8
9 individual subjects, with each child acting as their own control, we were able to evaluate children's
10
11 progress relative to their own baselines without confounding results from other participants.
12
13

14 **Measures**

15
16
17 **Assessment measures.** The following measures were used to assess each child prior to baseline.

18
19 *Vineland Adaptive Behavior Scale-Second Edition (VABS-2)* (Sparrow et al., 2005). We chose to
20
21 use adaptive behavior instead of a measure of nonverbal cognitive development because we were unable
22
23 to identify a standardized assessment that was appropriate for the ages and behaviors of our participants
24
25 (see Kasari, et al., 2013, for a discussion of testing difficulties). The VABS-2 is a measure of adaptive
26
27 behavior for birth through age 90. It is completed through caregiver and/or teacher interview, covering
28
29 four broad domains including communication, daily living skills, socialization, and motor skills. The
30
31 Vineland also includes a maladaptive behavior domain that assesses problem behavior. Each subdomain
32
33 and maladaptive behavior raw score has a v-scale score that corresponds to a standard score. Within the
34
35 communication subdomain, questions are directed to assess development of receptive, expressive, and
36
37 written language skills. The maladaptive behavior index assesses the presence of internalizing and
38
39 externalizing behaviors. The VABS-2 was normed on a national sample of 3,695 individuals and
40
41 represents population demographics (i.e., gender, ethnicity, region, and SES) matching the 2001 U.S.
42
43 Census. The normative sample also included a representative number of individuals with disabilities,
44
45 including relevant groups of individuals with speech-language impairments and intellectual disabilities.
46
47 Reported test-retest reliability and content validity measures are high.
48
49

50
51
52 *Peabody Picture Vocabulary Test-4.* The Peabody Picture Vocabulary Test, Fourth Edition
53
54 (PPVT-4) (Dunn & Dunn, 2007) was administered to all participants as a measure of single word
55
56 vocabulary comprehension. The PPVT-4 norms are based on a representative sample of 3,540 people
57
58
59
60

INVESTIGATING A MULTIMODAL INTERVENTION

12

1
2
3 aged 2 years 6 months through 90 years and older and matching the U.S. Census for gender, ethnicity,
4 region, and SES. Test-retest reliability coefficients and content validity measures for the PPVT-4 are
5 high. Additionally, the demographic characteristics of the PPVT-4 special-population sample included
6 relevant groups of children with identified language delays and intellectual disabilities. The PPVT-4 is
7 administered by presenting test plates with 4 pictures on a grid. The individual being tested is asked to
8 point to, or otherwise indicate, the picture that corresponds to one of the pictures, named by the tester.
9

10
11
12
13
14
15
16 *Communication Complexity Scale (CCS)*. The CCS is a scale describing levels of early
17 communication development (Brady et al., 2012). Each child participated in a scripted assessment
18 protocol consisting of 12 opportunities to initiate communication acts. Responses to each of the 12
19 opportunities were assigned a scaled score between 0 and 11 with 1 representing alerting behavior and 11
20 representing a multi-word utterance. Scores between 0 and 6 are pre-intentional; 7-9 are intentional but
21 nonsymbolic and 10 -11 are symbolic. For each child, the scores of the three opportunities with the
22 highest scores were then averaged to obtain the CCS scores reported in Table 1. Procedures for
23 administering and scoring the CCS are provided in Brady et al. (2012).
24
25
26
27
28
29
30
31
32

33
34 *Imitation*. Imitation was assessed with an instrument developed by Rogers and colleagues
35 (Rogers et al., 2003). This screening instrument contains 8 motor imitation tasks (e.g., clap hands) and 8
36 vocal/verbal items (e.g., say /ba/). Responses to the vocal/verbal items are presented in Table 1 because
37 these responses are most closely related to our word-learning outcomes and because all of the children
38 were able to imitate the motor imitation tasks.
39
40
41
42
43
44

45 **Dependent measures.** The following measures served as the primary measures of word learning.

46
47 *Expressive probes*. Pictures representing each target word were presented to the child, who was
48 asked to name the picture, (e.g., “What’s this?”). No prompts were provided during these probes. Each
49 word was presented five different times for a total of 25 trials. The order of words was randomized by
50 shuffling the pictures before each session. A correct production was recorded if the child produced at
51 least the initial consonant and vowel for a target word. For example, if the child said /mI/ when the
52
53
54
55
56
57
58
59
60

1
2
3 picture of mitt was displayed, this production was scored “correct.” Our rationale for this definition was
4
5 that deleting the final consonant is a common error in early word production (Dollaghan, 1985) and we
6
7 were more concerned with the participant’s attempts to correctly produce a target word, than their correct
8
9 articulation of each sound in the word (c.f.,Storkel, 2001). Research assistants were able to reliably
10
11 determine if the word was correct or not based on this definition.
12
13

14 *Receptive probes.* Computerized matching-to-sample procedures were used to assess receptive
15
16 word learning prior to each intervention session. Sessions were presented on a laptop computer, and
17
18 contained 20 trials, four for each word. For each test trial, a spoken vocabulary word (digitized speech
19
20 recorded with the Proloquo2go app on an iPad) (Sennott & Bowker, 2009) was presented along
21
22 with pictures on the touch-sensitive screen. Child 1 and 3 were provided with a three-choice array. Child
23
24 6 had a great deal of difficulty with the 3-choice array and we moved to a 2-choice array after 10 sessions.
25
26 All the remaining children were presented with a 2-choice array. The spoken word was presented
27
28 every 2 seconds until the child chose one of the pictures by touching it. The software automatically
29
30 recorded the responses. No feedback was given as to the correctness of the response. The next trial began
31
32 after a 1-second blank screen. The experimenter offered praise for participating in the task approximately
33
34 once every 2 minutes on a random schedule.
35
36
37

38 **Procedures**

39
40 **Speech sound identification.** The multimodal intervention described below targeted vocabulary
41
42 selected for each child, based on their extant repertoire of speech sounds. We used a digitized recording
43
44 device (DLP; LENA™) to help us identify their speech sounds because participants could not participate
45
46 in typical speech sound production assessment activities, such as a standardized articulation test. Using
47
48 the DLP, we obtained a recording of 12 hours of spontaneous vocalizations, across 2 days. The recorder
49
50 weighs approximately 2.5 ounces and fits into the front pocket of children’s clothing specially designed to
51
52 hold the device. It records the child’s vocalizations and adult talk near the child within an approximate 6-
53
54 10 foot radius (Thiemann-Bourque, Warren, & Brady, 2010). Participants received a recording packet at
55
56
57
58
59
60

INVESTIGATING A MULTIMODAL INTERVENTION

14

1
2
3 least one day prior to each scheduled recording date. The recording packet contained the DLP, clothing
4 adapted with a pocket for the DLP, and instructional documents. Parents were instructed to turn on the
5 DLP when the child woke up in the morning, insert it into the pocket of the clothing, and dress the child.
6
7 Thereafter, they were to go about their normal daily activities. During the recording day, the recorder
8 stayed on the child at all times except during baths and sleep periods. If the recorder was not on the child,
9
10 parents were instructed to place it nearby within 6-feet and continue to record. Parents were asked not to
11
12 turn off the recorder, but to leave it running continuously until the end of the day, at which time it would
13
14 shut off automatically. The DLPs were picked up by a research assistant and brought to our research lab.
15
16
17
18
19

20
21 Audio recording data from the DLP were then uploaded to a computer and automatically
22 processed using LENA software. The acoustical analysis software separates speech-related sounds from
23 environmental sounds, and segments are identified as adult male, adult female, or child. Using the client
24 manager software, child vocalizations were selected. LENA allows the acoustic information to be binned
25 in specified amounts of time. We chose 5-minute increments and then selected the 10, 5- minute intervals
26 with the highest rates of vocalizations recorded over the 12 hours. Next, a trained graduate student
27 listened to these identified segments and transcribed each sound produced. Because the children were, for
28 the most part, nonverbal, these vocalizations most often occurred during non-speech babbling
29 productions. Thus, the DLP and LENA software increased our efficiency by facilitating our identification
30 of periods of high frequency vocalizations that were later transcribed. The total number of different
31 consonants transcribed for each child is listed in Table 1.
32
33
34
35
36
37
38
39
40
41
42
43
44

45 **Target word selection.** A set of target words was identified for each child using an existing set
46 of consonant vowel consonant real words (Storkel, 2013). This existing corpus contained 1,396 real word
47 CVCs. In the corpus, a real word was any CVC that appeared in an adult or a child corpus. For the current
48 study, we pruned these 1,396 real words to only those that appeared in both the adult and child corpora,
49 yielding a pool of 720 real words. Storkel (2013) provides measures of phonotactic probability and
50 neighborhood density for each real word based on a child corpora. Words with high probability and high
51
52
53
54
55
56
57
58
59
60

INVESTIGATING A MULTIMODAL INTERVENTION

15

1
2
3 density sequences appear to be learned more rapidly than low probability and low density sequences,
4
5 potentially facilitating word learning for the participants in the current study. Thus, these high probability
6
7 and high-density words were targeted for treatment in the current study.
8
9

10 Phonotactic probability refers to the likelihood of occurrence of a sound sequence and can be
11
12 represented by two measures: positional segment sum and biphone sum. Positional segment sum is the
13
14 sum of the positional segment frequencies for each of the three sounds in the CVC. Positional segment
15
16 frequency is computed by summing the log frequency of all the words in the corpus containing the target
17
18 sound in the target word position and dividing by the sum of the log frequency of all the words in the
19
20 corpus containing any sound in the target word position. Biphone sum is computed in the same way
21
22 except that the focus is on an adjacent pair of sounds (i.e., CV or VC) rather than an individual sound.
23
24 Neighborhood density refers to the number of words in the corpus that differ from a given word by a
25
26 single sound substitution, deletion, or addition in any word position. For the 720 real words remaining in
27
28 the pool, the two measures of phonotactic probability and one measure of neighborhood density were
29
30 converted to a z score based on the mean and standard deviation of the items in the pool. Items that had
31
32 negative z scores for any of the measures were removed from the pool. The remaining pool of potential
33
34 targets consisted of 182 real word CVCs with high probability and high density.
35
36
37

38 For each individual child, the pool of 182 high probability and high density real words was
39
40 further reduced based on the child's individual speech sound repertoire. That is, words in the pool that
41
42 contained sounds that were not in the child's phonetic repertoire were removed from the pool for that
43
44 specific child. Thus, the remaining pool for each child contained high probability and high density real
45
46 words composed of sounds that the child produced as part of his/her phonetic repertoire. Thirty target
47
48 words were identified for each child. The thirty selected words were then placed into six sets of five
49
50 words each. Words were assigned to sets to provide a mixture of low and high frequency words (Storkel
51
52 & Hoover, 2010) to ensure that there would be a range of items for each child to learn (e.g., words they
53
54 may have heard before as well as words that they had never encountered). In addition, an attempt was
55
56
57
58
59
60

INVESTIGATING A MULTIMODAL INTERVENTION

16

1
2
3 made to create sets that were phonologically diverse with few words in the set sharing the same initial,
4
5 medial, or final sound.
6

7
8 Sets were then randomly assigned to a treatment order. In terms of the number of sets actually
9
10 treated, 4 children received treatment on 4 word sets; 2 children received treatment on 3 word sets; 3
11
12 children received treatment on 2 word sets and one child received treatment on only 1 set. This yields 145
13
14 treated words across children and sets, although note that some treated words were repeated across
15
16 children. The appendix shows the treated word sets for each child. Characteristics of the treated words
17
18 (i.e., word frequency, positional segment sum, biphone sum, and number of neighbors) was submitted to a
19
20 10 Participant x 4 Set ANOVA. Results showed no significant difference in word characteristics across
21
22 participants, sets, or the interaction of participant and set, all $F < 1.50$, all $p > 0.20$, all $\eta_p^2 < 0.04$. Across
23
24 all participants and sets, the mean log frequency was 2.82 ($SD = 0.86$, range 1.00-4.69); the mean
25
26 segment sum was 0.21 ($SD = 0.03$, range 0.17-0.29); the mean biphone sum was 0.013 ($SD = 0.004$, range
27
28 0.009-0.023); the mean number of neighbors was 20 ($SD = 4$, range 14-34).
29
30
31

32 **Story creation.** Once the set of words was identified for a child, a “story” that could be
33
34 represented in a book and interactive routine was constructed. For example, based on the words: mitt, cap,
35
36 bud, ten, and pin, a story was created about a game between two *buds*, where each child wore a special
37
38 *cap* and a *mitt*, and after the buds scored a *ten* they earned a special *pin*.
39

40 A picture book depicting the story was created. Each target word was presented in the story 5
41
42 different times. Story books were illustrated with clip art and printed words were presented along the
43
44 bottom of each page. Up to three short sentences appeared on each page. Symbols representing each
45
46 target word were presented on an iPad equipped with Proloquo2go™ software. The iPad was made
47
48 available during an interactive teaching routine, described below under Intervention.
49
50

Research Design.

51
52
53
54 In order to determine if participants showed gains in targeted spoken word productions, a
55
56 multiple probe-across-word-sets design was used (Byiers, Reichle, & Symons, 2012; Kazdin, 1982). Six
57
58
59
60

INVESTIGATING A MULTIMODAL INTERVENTION

17

1
2
3 word sets (each set consisting of five real words) were identified for each child based on their phonetic
4 repertoire and word characteristics (described above). Word sets were taught one at a time. Expressive
5 probes for words in the set currently being taught were administered immediately prior to (i.e., on the
6 same day as) each intervention session. These probes constituted the main outcome measure. In addition,
7 probes for the next set to be taught were presented prior to initiating instruction on that set. Decisions
8 regarding changes in experimental conditions were based on the results of these probes. Once a stable
9 baseline was established for set 1, intervention began on set 1. A baseline was considered stable if at least
10 two successive data points showed consistent (flat) performance, or if the performance declined.
11 However, this rule for stable baselines was violated three times during the study due to experimenter
12 error. Participant 1 only had 1 day of baseline for the first word set, and a rising baseline before the
13 introduction of Set 2. Child 2 also showed a small increase immediately before intervention was
14 introduced on Set 2.

15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

The pass criterion for a word set was $\geq 70\%$ correct word approximations over three consecutive expressive probes. When students met this criterion, and also demonstrated a stable baseline on production probes for the next word set, instruction began on the next word set. If a child did not reach criterion after 14 or more intervention sessions for a word set, intervention was discontinued for that set. The baseline-treatment sequence was completed for 4 sets for Child 1, 2, 3, and 4. Due to lack of progress and time constraints, teaching was discontinued after 3 completed sets for Children 6 and 8, and 2 sets for Children 7 and 9, and 1 set for Child 10. Child 5 moved during the study, and had only two sets.

In addition, beginning with Child 2, we included maintenance probes for previously learned words and periodic probes of words that were never targeted for intervention (“control words”). Maintenance probes were identical to expressive word probes and were administered between 2 and 40 sessions following the final instruction session for a successfully learned word set. The purpose of control words was to provide additional evidence that increases in word learning were specific to the words

1
2
3 targeted in intervention. Control words were selected for each child, in the same way as taught words,
4
5 such that there were no differences in the word characteristics (i.e. phonotactic probability or
6
7 neighborhood density) for control vs. taught words. Control words were probed in sets of five, in a
8
9 manner identical to the expressive probes.
10

11 **Multi-Modal Intervention Sessions.**

12
13
14 Following the Receptive and Expressive probes (described above under *Measures*), participants
15
16 received the multimodal intervention for the word set they were currently learning. Multimodal refers to
17
18 combined use of speech and a speech generating device during intervention. All components of the
19
20 intervention were provided in the order listed below, within each intervention session.
21
22

23 **Speech sound practice.** Although children produced target phonemes during their babbling
24
25 recorded with the digitized recording device, we wanted to provide additional practice for the sounds in
26
27 the target words immediately prior to opportunities to use the target vocabulary during joint book reading
28
29 and the interactive routine. Our theory was that this additional practice might increase the likelihood of
30
31 correct productions during these activities and facilitate word learning. For example, for the word set
32
33 mitt, cap, bud, ten and pin, children practiced the sounds /m/, /k/, /t/, /p/, and /n/. The interventionist, a
34
35 speech-language pathologist, provided models, physical prompts and corrective feedback to assist in
36
37 sound production. Each sound was practiced 5 times. Positive feedback (e.g., “good job”) was provided
38
39 for correct sound approximations.
40
41
42

43 **Joint book reading.** The interventionist first read the story book that was created for the
44
45 individualized set of words, emphasizing the target words and pointing to corresponding pictures. Next,
46
47 she read the book with pauses before each target word (i.e, a cloze procedure as in Petersen, Gillam,
48
49 Spencer, & Gillam, 2010). For example, one story begins with, “Bob and Joe are best buds!” On the
50
51 second reading, the interventionist said, “Bob and Joe are best _____” and waited up to 5 seconds for the
52
53 child to fill in the blank. If the child correctly filled in the blank the interventionist provided verbal praise
54
55
56
57
58
59
60

1
2
3 and moved on to the next sentence in the story. If the child said a wrong word or didn't say any word, the
4
5 interventionist provided the correct response and moved on to the next sentence in the story.
6
7

8 **Interactive routines with iPad.** Teaching routines followed a predictable structure with a script
9
10 for each step in the routine (Kashinath et al., 2006; Snyder-McLean, Solomonson, McLean, & Sack,
11
12 1984). Routines also provided opportunities for children to produce target words at least 5 different times.
13
14 In accordance with the script, the interventionist would first model selecting a target symbol on the iPad
15
16 equipped with Proloquo2go software. The arrays contained the five target words. After the child
17
18 participated in the routine several times, the interventionist began pausing before modeling the response
19
20 in order to allow the child an opportunity to initiate the symbol selection. If the child did not select the
21
22 symbol after 5 seconds, the interventionist selected it. Each symbol selection was associated with a
23
24 tangible consequence that also led to the next step in the routine. For example, in a routine about a special
25
26 game between *buds*, children needed to request the *cap* that was worn to play the game. The
27
28 interventionist and child would walk to the shelf where the caps were located and say "Before we play the
29
30 game we need to wear our ____." After the child (or interventionist) selected the *cap* symbol and/or said
31
32 "cap", the child and interventionist put on their caps and went to the location where the game was played.
33
34
35

36 **Receptive matching trials.** At the end of each session, additional practice in associating the
37
38 spoken word with the symbol was provided with another set of receptive matching trials on the laptop
39
40 computer. These trials were identical to the receptive probe trials described above with the following
41
42 exception—correct responses received a visual and auditory consequence on the computer and verbal
43
44 praise by the interventionist.
45
46

47 **Reliability.**

48
49 A second observer (a trained undergraduate student) was present for 22% of the baseline and
50
51 intervention sessions (111 of 500, distributed across participants) to gather inter-observer agreement data
52
53 for the dependent and independent variables. The second observer recorded responses (correct or
54
55 incorrect) to the expressive probes. The second observer also recorded responses to the speech sound
56
57
58
59
60

1
2
3 practice trials. Percent agreement was calculated as the number of agreements / agreements plus
4 disagreements x 100. The percent agreement scores were very high for these variables—97% for the
5 expressive probes and 97% for speech-sound practice. Receptive probe data were recorded directly from
6 the computer and we did not perform reliability checks with the computer.
7
8
9
10

11
12 Procedural reliability for the independent variable was determined by asking the second observer
13 to record whether or not the following intervention components were provided during intervention: 1) did
14 the interventionist pause 5 seconds before each opportunity for an initiated symbol selection during joint
15 book reading and interactive routines? 2) Did the interventionist model the correct response the
16 designated number of times within joint book reading and the interactive routine? The percentage of
17 teaching opportunities for which the interventionist included these components was high -- 99% for
18 pauses, and 87% for modeling.
19
20
21
22
23
24
25
26

27 **Results**

28
29
30 Results will be presented according to our primary purposes: To determine if participants showed
31 gains in (1) spoken word production and (2) receptive word learning, and (3) to compare profiles of
32 children who responded favorably to those who did not respond as favorably to the intervention. The
33 following section presents results pertaining to word learning outcomes.
34
35
36
37

38 **Expressive Word Production**

39
40 Results from the expressive word learning probes are presented in Figures 1-10. Each data point
41 reflects the percent correct out of 25 expressive word production probe trials, collapsed across all 5 words
42 in a set. Child 1 was the first child that participated. It is clear from Figure 1 that correct expressive word
43 productions increased as each new set was introduced to Child 1. Unfortunately, only one baseline session
44 was presented for the first word set and only two baseline sessions were presented for set 4. However,
45 despite these limitations the pattern of responding is strong and consistent. Similarly, the data for child 2,
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 compared to baseline. In addition, data from the control words showed that the gains recorded for
4
5 Participants 1-5 were specific to the words targeted during intervention. Maintenance probes showed that
6
7 word productions maintained at levels similar to those of the final intervention sessions.
8
9

10 Child 6's data is shown in Figure 6. Although child 6 did not meet criterion for any of the word
11
12 sets, he did show substantial progress over baseline production levels for each of three different word sets.
13
14 Child 7 learned to say 1/5 words in each of 2 sets. Child 8 increased word production in set 1, although he
15
16 did not meet criterion. Child 9 and child 10 showed essentially no progress, and hence we discontinued
17
18 intervention after 2 sets for child 9 and 1 set for child 10.
19

20
21 In addition to the graphic data display, we measured each child's changes from baseline to post
22
23 treatment. We calculated confidence intervals for the effect sizes (ds) based on the differences between
24
25 the mean percent correct over the last three days of intervention compared to baseline means. We used
26
27 the method of Odgaard and Fowler (2010) to compute the noncentrality parameter for the upper and lower
28
29 limits of the confidence interval based on dependent samples t-test values. These noncentrality parameters
30
31 were then used to calculate the upper and lower effect size d for the confidence interval. This effect size
32
33 confidence interval, along with the number of different words produced at a criterion of at least 80%
34
35 correct over the last three days of intervention and the total number of treatment sessions are presented in
36
37 Table 2. Effect sizes reflected large to very large effects for participants 1-6.
38
39

40 **Receptive Word Learning.**

41
42
43 Although the focus of this study was on expressive word production, we also measured receptive
44
45 word learning as recorded with the computerized matching software. Words were considered "learned"
46
47 on the receptive probes if children scored more than 80% correct over three consecutive sessions of the
48
49 receptive matching probes. Table 3 shows changes for each child's receptive word learning. For the most
50
51 part, the receptive data were similar to the expressive word data. That is, Participants 1-5 showed positive
52
53 gains and met receptive criteria for most words learned expressively but Participants 6-10 did not.
54
55

56 **Profiles of High and Low Responders.**

57
58
59
60

INVESTIGATING A MULTIMODAL INTERVENTION

22

1
2
3 Participants 1-5 responded to the intervention with significant increases in expressive production
4 of targeted words. We refer to these children as “High Responders.” As shown in Tables 2 and 3, Effect
5 sizes for the “High Responders” were large in the full range of the confidence interval for both receptive
6 and expressive probe data with the exception of Child 5 whose receptive effect size included zero in the
7 confidence interval. Children in the Low Responders group did not meet expressive word learning
8 criteria, although 3 children did show gains in producing some words. Effect size values in the Low
9 Responders group contained zero or were small/moderate within the confidence interval. Child 6 was an
10 exception to this with large effect sizes contained in the full 95% confidence interval for expressive
11 language.
12
13
14
15
16
17
18
19
20
21
22

23 We reviewed the scores for each participant that were collected pre-baseline (see Table 1) to
24 determine if there were differences in scores for children who were High vs. Low Responders to our
25 intervention package. The children who responded best to our intervention (children 1-5) had higher
26 scores on the PPVT-4, indicating better receptive vocabularies. In addition, they also had higher scores on
27 the CCS with High Responders averaging 9.75 and Low Responders averaging 8.0, indicating more
28 advanced prelinguistic communication in the high Responder group. This higher CCS score reflects more
29 frequent use of communicative gestures and/or vocalizations during the scripted interaction. The overall
30 imitation scores were not obviously different across children, a finding similar to Rogers and colleagues
31 (2012). However, when we looked only at imitation for oral, vocal and verbal skills, we saw higher scores
32 for the High-Responders with a mean of 7.6 while the Low Responders had a mean of 3.0. We also
33 reviewed the number of different consonants recorded with the LENA recording system prior to
34 intervention as a possible differentiating variable between high and low responders. The mean number of
35 consonants was 14.0 for High Responders and 11.8 for Low Responders suggesting a possible
36 relationship. Thus, high-responding children had relatively better scores on receptive vocabulary,
37 prelinguistic communication, vocal/verbal imitation and consonant repertoires. However, these
38 observations are merely suggestive due to small sample sizes.
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Discussion

Results of this pilot investigation indicated that children with limited expressive vocabularies learned to say new words with a multimodal intervention package that combined AAC and speech sound practice for individually determined sets of words. Maintenance and control probe data indicated that the responses maintained and were specific to the targeted vocabulary for the four children for whom these data were collected. Given that all the participants had limited expressive vocabularies at the start of intervention, and were between the ages of 6 and 11 years, the gains made by the 5 High Responders are impressive. Currently, the predominant clinical practice for children with this profile is to abandon efforts to increase speech production. Moreover, there is little in the research literature to contraindicate this practice. Our results indicate that an intervention that focuses on speech plus AAC can have positive outcomes on speech for school-age children with autism and limited expressive vocabularies. Given that this was a relatively short-term intervention, and the duration of intervention sessions was only 45 minutes per day, these results are extremely promising.

To our knowledge, this study was also the first reported attempt to apply principles of phonotactic probability and neighborhood density to a word learning intervention for children with intellectual disabilities and limited expressive vocabularies. The vocabulary targeted by this approach may have been easier to learn to produce than other vocabularies, because of these word-learning principles. The approach we used may facilitate initial word learning that can help build a foundation for later functional spoken vocabulary.

Effect sizes for the high responders in our study were for the most part in the high to very high range and compared favorably to other interventions measuring gains in beginning word use by younger children with autism. For example, Rogers and colleagues reported a d of .84 for vocabulary comprehension and a d of .57 for vocabulary production for children who participated in their ESDM intervention. Kasari et al. (2014), reported effect sizes between .21 and .62 across different expressive communication measures for their group who experienced a combined JASPER and SGD treatment,

1
2
3 compared to a treatment group of only JASPER (without the SGD). Thus, our multimodal intervention
4 that includes intensive speech sound practice does seem to be effective for at least some children past the
5 preschool age range. The specific emphasis on speech sound production differs from most interventions
6 that have been used with preschool age children (e.g. Dawson et al., 2010; Goods et al., 2013). It is not
7 known if the additional speech sound practice was a key ingredient of the intervention because all of the
8 children experienced this component (and all of the components). Future research is needed to examine
9 how different components, including speech sound practice contribute to overall outcomes.

10
11
12 Our results were not as positive for all participants, however. Five of the 10 participating children
13 failed to meet our production criterion indicating that they failed to meet our learning criterion for any
14 word sets (Low Responders). However, it is worth noting that three of these children did show increases
15 in word productions over baseline levels, as indicated in Figures 6-8 and Table 2. Perhaps longer or more
16 intense interventions would have improved responding by these three participants, and modifications such
17 as these would be appropriate for clinical applications (outside of a research study). In addition, further
18 analysis could focus on comparing the specific words that children learned to those they did not learn in
19 an effort to identify commonalities in terms of the phonemes included and/or motivations associated with
20 particular vocabulary.

21
22
23 Perhaps the most important aspect of the current study is that we provided results for both
24 children who did and children who did not respond well to our intervention, along with data about some
25 important characteristics that appeared associated with differential outcomes. This type of information is
26 critical for designing future research studies, and ultimately for clinical decision making (Yoder &
27 Compton, 2004). Children with autism or autism symptoms and minimal verbal repertoires present
28 clinicians with various skill repertoires. One goal of intervention research is to identify relative strengths
29 and weaknesses in areas that may facilitate learning with a particular intervention approach.

30
31
32 We retrospectively compared High Responders' and Low Responders' initial skills in areas that
33 have been linked to verbal outcomes (receptive language, prelinguistic communication, imitation, speech
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

INVESTIGATING A MULTIMODAL INTERVENTION

25

1
2
3 sound production and adaptive behavior) because the focus of the current study was on spoken word
4 productions. Our review of differences in entry-level skills of High vs Low responders suggested the
5 intervention may be most helpful for children who have at least some measureable receptive vocabulary,
6 frequent use of intentional communication acts, imitate vocal/verbal sounds and oral movements, and
7 have larger consonant repertoires (note we did not count or compare the size of vowel repertoires in this
8 study). As discussed in the Introduction of this study, each of these skills has been linked to language
9 learning outcomes and hence it is not surprising that they were also associated with different outcomes in
10 our study. It is also noteworthy that Participants 1, 2, 3 and 5 had some correct responses (1-2 words)
11 during baselines, which was not the case for Low Responders. Therefore some initial evidence of correct
12 speech production appears to be predictive of early progress in a multimodal intervention. However, these
13 observations are based on very small sample sizes and replication with larger samples who vary across
14 these skills are needed to make more robust determinations of characteristics of High vs. low Responders.
15 It may be useful to examine additional areas, such as play skills, that have been linked to communication
16 outcomes in future studies.

17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34 Identifying children who respond best to different types of interventions could help maximize
35 positive outcomes by selecting appropriate targets based on individual needs and entry behaviors. For
36 example, in the current study, children who we ultimately described as Low Responders to our
37 intervention package may have made more communication gains in an intervention that emphasized AAC
38 along with prelinguistic social communication, without specific focus on speech sound production. In
39 addition, it may prove beneficial to directly teach skills that are associated with better outcomes (e.g.,
40 receptive language, imitation) as part of a comprehensive communication intervention. However, speech
41 gains made by our High Responding children provide evidence for continued speech-based interventions
42 for some children with autism and limited expressive vocabularies.

43
44
45
46
47
48
49
50
51
52
53
54 The current study used a single subject research design which allowed review of the effectiveness
55 across each child and evaluation of how individual skill sets related to different outcomes. This was very
56
57
58
59
60

1
2
3 important because, if all the children's data had been combined, as in a between-groups design, the
4
5 differences in relative effectiveness across children would have been obscured. Future research that
6
7 further delineates characteristics of responders will increase the generality to participants who are similar
8
9 in terms of the dimensions associated with positive outcomes. However, as in all single subject design
10
11 studies, the outcomes are only generalizable to participants of a given study or individuals highly similar
12
13 to those participants. The detailed information about individual skills provides a basis for comparison that
14
15 clinicians may refer to when evaluating the potential of a given intervention for use with their clients.
16
17

18
19 **Limitations to the current study.** Children in our study were taught a very limited set of
20
21 vocabulary in contrived teaching environments, without any focus on generalized use of targeted
22
23 vocabulary across contexts. While this is a limitation in terms of evaluating the clinical effectiveness of
24
25 intervention outcomes, we propose that our findings be viewed in terms of "proof of concept" for our
26
27 unique multimodal intervention package (Smith et al., 2007). Based on our results, further research is
28
29 warranted that would include measures of generalization to functional communication targets, including
30
31 generalized communication improvements to non-target vocabulary across communication modes. For
32
33 experimental-control purposes, it was important in the current study to show that learning was limited to
34
35 targeted vocabulary. A desirable outcome for future studies would be generalization to non-targeted
36
37 words in addition to learning the words directly taught in intervention.
38
39

40
41 Another area for future research is determining the value of individual components of the
42
43 intervention package. For example, one novel approach we used was to select vocabulary based on both a
44
45 child's current speech sound repertoire and principles of phonotactic probability. This was based on our
46
47 hypothesis that children would learn speech more readily under these conditions. However, we did not
48
49 directly test this hypothesis and future investigations may want to compare word learning using different
50
51 criteria for initial vocabulary selection. In the current study we only measured speech sound productions
52
53 during baseline, and future investigations may also re-measure speech sound productions after
54
55 intervention to determine if there is a change in this variable.
56
57
58
59
60

INVESTIGATING A MULTIMODAL INTERVENTION

27

1
2
3 The lack of a consistent autism measure was another limitation to the current study. We relied on
4 diagnoses performed by outside agencies using a variety of instruments. Therefore, we were not able to
5 quantify the severity of autism in our participants. In future research it will be important to administer an
6 assessment that could reflect autism severity across all participants, in order to determine the possible role
7 of severity in predicting differential outcomes. Finally, the experimenter errors described under the
8 methods section were limitations in the current study. Specifically, experimenter error resulted in baseline
9 errors for one word set each for Child 1 and Child 2.
10
11
12
13
14
15
16
17

18 **Conclusions.** Results from this pilot investigation indicate that our multimodal intervention leads
19 to increased spoken word productions for some school-age children with minimal verbal skills who attend
20 educational programs for children with autism. Future studies are needed to more accurately describe the
21 variables that are associated with positive outcomes, target use in naturalistic contexts and determine the
22 relative contributions of different intervention components. Ultimately, results from this line of research
23 may lead to increased emphasis on speech intervention in addition to AAC instruction for school-age
24 children with autism and minimal verbal skills. Such a combined approach would seem optimal for
25 promoting effective communication with the widest possible set of communication partners and across
26 multiple environments.
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

References

- Bellon, M. L., Ogletree, B. T., & Harn, W. E. (2000). Repeated storybook reading as a language intervention for children with autism. *Focus on Autism & Other Developmental Disabilities*, 15(1), 52.
- Billstedt, E., Gillberg, I., & Gillberg, C. (2007). Autism in adults: symptom patterns and early childhood predictors. Use of the DISCO in a community sample followed from childhood. *Journal of Child Psychology and Psychiatry*, 48(11), 1102-1110. doi: 10.1111/j.1469-7610.2007.01774.x
- Binger, C., & Light, J. (2007). The effect of aided AAC modeling on the expression of multi-symbol messages by preschoolers who use AAC. *Augmentative and Alternative Communication*, 23(1), 30-43. doi: doi:10.1080/07434610600807470
- Bondy, A. S., & Frost, L. A. (1994). The picture exchange communication system. *Focus on Autistic Behavior*, 9, 1-19.
- Bopp, K. D., & Mirenda, P. (2011). Prelinguistic predictors of language development in children with autism spectrum disorders over four–five years. *Journal of Child Language*, 38(03), 485-503. doi: doi:10.1017/S0305000910000140
- Brady, N., Fleming, K., Thiemann-Bourque, K., Olswang, L., Dowden, P., Saunders, M., & Marquis, J. (2012). Development of the Communication Complexity Scale. *American Journal of Speech-Language Pathology*, 21(1), 16-28.
- Brady, N., Thiemann-Bourque, K., Fleming, K., & Matthews, K. (2013). Predicting language outcomes for children learning AAC: Child and environmental factors. *Journal of Speech, Language & Hearing Research*, 56(5), 1595-1612.

INVESTIGATING A MULTIMODAL INTERVENTION

29

- 1
2
3 Byiers, B. J., Reichle, J., & Symons, F. J. (2012). Single-subject experimental design for evidence-based
4
5 practice. *American Journal of Speech-Language Pathology*, 21(4), 397-414. doi: 10.1044/1058-
6
7 0360(2012/11-0036)
8
9
10 Carr, D., & Felce, J. (2007). Brief report: Increase in production of spoken words in some children with
11
12 autism after PECS: Teaching to phase III. *Journal of Autism and Developmental Disorders*,
13
14 37(4), 780-787. doi: 10.1007/s10803-006-0204-0
15
16 Charlop-Christy, M., Carpenter, M., LeBlanc, L., & Kellet, K. (2002). Using the picture exchange
17
18 communication system (PECS) with children with autism: Assessment of PECS acquisition,
19
20 speech, social-communicative behavior and problem behavior. *Journal of Applied Behavior*
21
22 *Analysis*, 35(3), 213-231.
23
24
25 Dawson, G., Rogers, S., Munson, J., Smith, M., Winter, J., Greenson, J., . . . Varley, J. (2010).
26
27 Randomized, controlled trial of an intervention for toddlers with autism: The early start Denver
28
29 model. *Pediatrics*, 125(1).
30
31
32 Dollaghan, C. (1985). Child meets word: Fast mapping in preschool children. *Journal of Speech and*
33
34 *Hearing Research*, 28, 449-454.
35
36
37 Dunn, L., & Dunn, D. (2007). *The Peabody Picture vocabulary Test-Fourth Edition (PPVT-4)*. San
38
39 Antonio, TX: Psychological Corporation.
40
41
42 Fleury, V. P., Miramontez, S. H., Hudson, R. F., & Schwartz, I. S. (2013). Promoting active participation
43
44 in book reading for preschoolers with Autism Spectrum Disorder: A preliminary study. *Child*
45
46 *Language Teaching and Therapy*. doi: 10.1177/0265659013514069
47
48
49 Flippin, M., Reszka, S., & Watson, L. (2010). Effectiveness of the Picture Exchange Communication
50
51 System (PECS) on communication & speech for children with autism spectrum disorders: A
52
53 Metanalysis. *American Journal of Speech-Language Pathology*, 19, 178-195.
54
55
56 Ganz, J. B., Earles-Vollrath, T. L., Heath, A. K., Parker, R. I., Rispoli, M. J., & Duran, J. B. (2012). A
57
58 meta-analysis of single case research studies on aided augmentative and alternative
59
60

- 1
2
3 communication systems with individuals with autism spectrum disorders. *Journal of Autism &*
4
5 *Developmental Disorders*, 42, 60-74.
6
7 Gathercole, S. E., Frankish, C. R., Pickering, S. J., & Peaker, S. (1999). Phonotactic influences of short-
8
9 term memory. *Journal of Experimental Psychology*, 25(1), 84-95.
10
11 German, D. J., & Newman, R. S. (2004). The impact of lexical factors on children's word-finding errors.
12
13 *Journal of Speech, Language & Hearing Research*, 47(3), 624-636. doi: 10.1044/1092-
14
15 4388(2004/048)
16
17 Goldstein, H., Wickstrom, S., Hoyson, M., Jamieson, B., & Odom, S. L. (1988). Effects of sociodramatic
18
19 script training on social and communicative interaction. *Education and Treatment of Children*,
20
21 11, 97-117.
22
23 Goods, K., Ishijima, E., Chang, Y.-C., & Kasari, C. (2013). Preschool based JASPER intervention in
24
25 minimally verbal children with autism: Pilot RCT. *Journal of Autism and Developmental*
26
27 *Disorders*, 43(5), 1050-1056. doi: 10.1007/s10803-012-1644-3
28
29
30
31 Gray, K. M., Tonge, B. J., & Sweeney, D. J. (2008). Using the Autism Diagnostic Interview-Revised and
32
33 the Autism Diagnostic Observation Schedule with young children with developmental delay:
34
35 Evaluating diagnostic validity. *Journal of Autism and Developmental Disorders*, 38(4), 657-667.
36
37 doi: <http://dx.doi.org/10.1007/s10803-007-0432-y>
38
39
40 Hargrave, A. C., & Sénéchal, M. (2000). A book reading intervention with preschool children who have
41
42 limited vocabularies: the benefits of regular reading and dialogic reading. *Early Childhood*
43
44 *Research Quarterly*, 15(1), 75-90. doi: [http://dx.doi.org/10.1016/S0885-2006\(99\)00038-1](http://dx.doi.org/10.1016/S0885-2006(99)00038-1)
45
46
47 Harris, M., & Reichle, J. (2004). The impact of aided language stimulation on symbol comprehension and
48
49 production in learners with moderate cognitive disabilities. *American Journal of Speech-*
50
51 *Language Pathology*, 13(2), 155-167.
52
53 Heflin, L. J., & Alberto, P. A. (2001). Establishing a behavioral context for learning for students with
54
55 autism. *Focus on Autism & Other Developmental Disabilities*, 16(2), 93.
56
57
58
59
60

INVESTIGATING A MULTIMODAL INTERVENTION

31

- 1
2
3 Kasari, C., Brady, N., Lord, C., & Tager-Flusberg, H. (2013). Assessing the minimally verbal school-
4 aged child with autism spectrum disorder. *Autism Research*, 6(6), 479-493.
5
6
7 Kasari, C., Gulsrud, A., Wong, C., Kwon, S., & Locke, J. (2010). Randomized controlled caregiver
8 mediated joint engagement intervention for toddlers with autism. *Journal of Autism &*
9 *Developmental Disorders*, 40(9), 1045-1056.
10
11
12 Kasari, C., Kaiser, A., Goods, K., Nietfeld, J., Mathy, P., Landa, R., . . . Almirall, D. (2014).
13
14 Communication interventions for minimally verbal children with autism: A sequential multiple
15 assignment randomized trial. *Journal of the American Academy of Child and Adolescent*
16 *Psychiatry*, 53(6), 635-646.
17
18
19 Kasari, C., Paparella, T., Freeman, S., & Jahromi, L. (2008). Language outcome in autism: Randomized
20 comparison of joint attention and play interventions. *Journal of Consulting and Clinical*
21 *Psychology*, 76(1), 125-137.
22
23
24 Kashinath, S., Woods, J., & Goldstein, H. (2006). Enhancing generalized teaching strategy use in daily
25 routines by parents of children with autism. *Journal of Speech, Language and Hearing Research*,
26 49(3), 466-485.
27
28
29 Kazdin, A. E. (1982). *Single-Case Research Designs*. New York: Oxford University Press.
30
31
32 King, A., Hengst, J., & DeThorne, L. (2013). Severe speech sound disorders: An integrated multimodal
33 intervention. *Language Speech and hearing Services in the Schools*, 44, 195-210.
34
35
36 Lord, C., Rutter, M., DiLavore, P. C., & Risi, S. (1999). *Autism Diagnostic Observation Schedule-WPS*
37 *Edition (ADOS-WPS)*. Los Angeles: Western Psychological Services.
38
39
40 MacRoy-Higgins, M., Schwartz, R. G., Shafer, V. L., & Marton, K. (2013). Influence of phonotactic
41 probability/neighbourhood density on lexical learning in late talkers. *International Journal of*
42 *Language & Communication Disorders*, 48(2), 188-199. doi: 10.1111/j.1460-6984.2012.00198.x
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 Mullen, E. (1995). *Mullen scales of early learning: AGS edition*. Circle Pines, MN: AGS.
- 4
5 Newman, R. S., & German, D. J. (2005). Life span effects of lexical factors on oral naming. *Language &*
6
7 *Speech, 48*(2), 123-156.
- 8
9
10 National Research Council. (2001). *Educating children with autism: Committee on educational*
11
12 *interventions for children with autism. Division of behavioral and social sciences and education.*
13
14 Washington, DC: National Academy Press.
- 15
16 Odgaard, E. C., & Fowler, R. L. (2010). Confidence intervals for effect sizes: Compliance and clinical
17
18 significance in the Journal of Consulting and Clinical Psychology. *Journal of Consulting and*
19
20 *Clinical Psychology, 78*(3), 287-297. doi: <http://dx.doi.org/10.1037/a0019294>
- 21
22
23 Odom, S. L., Rogers, S., McDougle, C. J., Hume, K., McGee, G., Odom, S., . . . Blacher, J. (2007). Early
24
25 intervention for children with autism spectrum disorder. *Handbook of developmental disabilities,*
26
27 199-223.
- 28
29 Petersen, D. B., Gillam, S. L., Spencer, T., & Gillam, R. B. (2010). The effects of literate narrative
30
31 intervention on children with neurologically based language impairments: An early stage study.
32
33 *Journal of Speech, Language & Hearing Research, 53*(4), 961-981. doi: 10.1044/1092-
34
35 4388(2009/09-0001)
- 36
37
38 Picket, E., Pullara, O., O'Grady, J., & Gordon, B. (2009). Speech acquisition in older nonverbal
39
40 individuals with autism. *Cognitive and Behavioral Neurology, 22*(1), 1-21.
- 41
42 Pomaville, F. M., & Kladopoulos, C. N. (2013). The effects of behavioral Sseech Tterapy on Sseech
43
44 sound Ppduction with adults who have cochlear implants. *Journal of Speech, Language, and*
45
46 *Hearing Research, 56*(2), 531-541. doi: 10.1044/1092-4388(2012/12-0017)
- 47
48
49 Poon, K., Watson, L., Baranek, G., & Poe, M. (2012). To what extent do joint attention, imitation, and
50
51 object play behaviors in infancy predict later communication and intellectual functioning in
52
53 ASD? *Journal of Autism and Developmental Disorders, 42*(6), 1064-1074. doi: 10.1007/s10803-
54
55 011-1349-z
56
57
58
59
60

- 1
2
3 Rogers, S., Estes, A., Lord, C., Vismara, L., Winter, J., Fitzpatrick, A., . . . Dawson, G. (2012). Effects of
4 a brief early start denver model (ESDM)- based parent intervention on toddlers at risk for autism
5 spectrum disorders: A randomized controlled trial. *Journal of the American Academy of Child*
6 *and Adolescent Psychiatry, 5*(10), 1052-1065.
- 7
8 Rogers, S., Hayden, D., Hepburn, S., Charlifue-Smith, R., Hall, T., & Hayes, A. (2006). Teaching young
9 nonverbal children with autism useful speech: A pilot study of the Denver model and PROMPT
10 interventions. *Journal of Autism and Developmental Disorders, 36*, 1007-1024.
- 11
12 Rogers, S., Hepburn, S., Stackhouse, T., & Wehner, E. (2003). Imitation performance in toddlers with
13 autism and those with other developmental disorders. *Journal of Child Psychology and*
14 *Psychiatry, 44*, 763-781.
- 15
16 Rollins, P., Wambacq, I., Dowell, D., Mathews, L., & Reese, P. B. (1998). An intervention technique for
17 children with autistic spectrum disorder: Joint attentional routines. *Journal of Communication*
18 *Disorders, 31*, 181-189.
- 19
20 Schlosser, R., & Wendt, O. (2008). Effects of augmentative and alternative communication intervention
21 on speech production in children with autism: A systematic review. *American Journal of Speech*
22 *Language Pathology, 17*, 212-230.
- 23
24 Schwartz, R. G., Leonard, L. B., Messick, C., & Chapman, K. (1987). The acquisition of object names in
25 children with specific language impairment: Action context and word extension. *8*, 233-244.
- 26
27 Sennott, S., & Bowker, A. (2009). Autism, AAC, and Proloquo2Go. *SIG 12 Perspectives on*
28 *Augmentative and Alternative Communication, 18*, 137-145.
- 29
30 Sevcik, R., Ronski, M., Watkins, R., & Deffenbach, K. (1995). Adult partner-augmented communication
31 input to youth with mental retardation using the system for augmenting language. *Journal of*
32 *Speech, Language and Hearing Research, 38*, 902-912.
- 33
34 Sherer, M. R., & Schreibman, L. (2005). Individual behavioral profiles and predictors of treatment
35 effectiveness for children with autism. *Journal of Consulting and Clinical Psychology, 73*(3),
36 525-538. doi: <http://dx.doi.org/10.1037/0022-006X.73.3.525>
- 37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
- Smith, L., Barker, E. T., Seltzer, M. M., Abbeduto, L., & Greenberg, J. S. (2012). Behavioral phenotype of fragile X syndrome in adolescence and adulthood. *American Journal on Intellectual and Developmental Disabilities, 117*, 1-17.
- Smith, T., Scahill, L., Dawson, G., Guthrie, D., Lord, C., Odom, S., . . . Wagner, A. (2007). Designing research studies on psychosocial interventions in autism. *Journal of Autism and Developmental Disorders, 37*(2), 354-366. doi: 10.1007/s10803-006-0173-3
- Snodgrass, M. R., Stoner, J. B., & Angell, M. E. (2013). Teaching conceptually referenced core vocabulary for initial augmentative and alternative communication. *Augmentative and Alternative Communication, 29*(4), 322-333. doi: doi:10.3109/07434618.2013.848932
- Snyder-McLean, L. K., Solomonson, B., McLean, J., & Sack, S. (1984). Structuring joint action routines: A strategy for facilitating communication and language development in the classroom. *Seminars in Speech and Language, 5*(3), 213-228.
- Solomon-Rice, P. L., & Soto, G. (2014). Facilitating vocabulary in toddlers using AAC: A preliminary study comparing focused stimulation and augmented input. *Communication Disorders Quarterly*. doi: 10.1177/1525740114522856
- Soto, G., & Dukhovny, E. (2008). The effect of shared book reading on the acquisition of expressive vocabulary of a 7 year old who uses AAC. *Seminars in Speech and Language, 29*(2), 133-145.
- Sparrow, S., Cicchetti, D., & Balla, D. (2005). *Vineland-II: Vineland adaptive behavior scales, Second Edition*. Minneapolis: Pearson.
- Storkel, H. L. (2001). Learning new words: Phonotactic probability in language development. *Journal of Speech Language and Hearing Research, 44*, 1321-1337.
- Storkel, H. L. (2013). A corpus of consonant–vowel–consonant real words and nonwords: Comparison of phonotactic probability, neighborhood density, and consonant age of acquisition. *Behavior Research Methods, 45*(4), 1159-1167. doi: 10.3758/s13428-012-0309-7

INVESTIGATING A MULTIMODAL INTERVENTION

35

- 1
2
3 Storkel, H. L., & Hoover, J. R. (2010). An on-line calculator to compute phonotactic probability and
4
5 neighborhood density based on child corpora of spoken American English. *Behavior Research*
6
7 *Methods*, 42(2), 497-506.
8
9
10 Storkel, H. L., Maekawa, J., & Hoover, J. R. (2010). Differentiating the effects of phonotactic probability
11
12 and neighborhood density on vocabulary comprehension and production: A comparison of
13
14 preschool children with versus without phonological delays. *Journal of Speech, Language &*
15
16 *Hearing Research*, 53(4), 933-949. doi: 10.1044/1092-4388(2009/09-0075)
17
18
19 Thiemann-Bourque, K., Brady, N., & Fleming, K. (2011). Symbolic play of preschoolers with severe
20
21 communication impairments with autism and other developmental delays: More similarities than
22
23 differences. *Journal of Autism and Developmental Disorders*, 1-11. doi: 10.1007/s10803-011-
24
25 1317-7
26
27 Thiemann-Bourque, K., Warren, S., & Brady, N. (2010). *What We Can Learn from Counting Words:*
28
29 *Applications and Implications of the Automatic Measurement of Natural Environments*. Paper
30
31 presented at the Society for Research on child Development, Montreal.
32
33
34 Thomson, J., Richardson, U., & Goswami, U. (2005). Phonological similarity neighborhoods and
35
36 children's short-term memory: Typical development and dyslexia. *Memory & Cognition*, 33(7),
37
38 1210-1219. doi: 10.3758/BF03193223
39
40
41 Thurm, A., Lord, C., Lee, L., & Newschaffer, C. (2007). Predictors of language acquisition in preschool
42
43 children with autism spectrum disorders. *Journal of Autism and Developmental Disorders*, 37,
44
45 1721-1734.
46
47 Toth, K., Munson, J., N. Meltzoff, A., & Dawson, G. (2006). Early predictors of communication
48
49 development in young children with autism spectrum disorder: Joint attention, imitation, and toy
50
51 play. *Journal of Autism and Developmental Disorders*, 36(8), 993-1005. doi: 10.1007/s10803-
52
53 006-0137-7
54
55
56
57
58
59
60

INVESTIGATING A MULTIMODAL INTERVENTION

36

- 1
2
3 Vihman, M. M., DePaolis, R. A., & Keren-Portnoy, T. (2014). The role of production in infant word
4 learning. *Language Learning*, 64(s2), 121-140. doi: 10.1111/lang.12058
5
6
7
8 Vismara, L. A., Comlombi, C., & Rogers, S. J. (2009). Can 1 hour per week of therapy lead to lasting
9 changes in young children with autism? *Autism*, 13, 93-115.
10
11
12 Vivanti, G., Dissanayake, C., Zierhut, C., & Rogers, S. (2013). Brief Report: Predictors of outcomes in
13 the Early Start Denver Model delivered in a group setting. *Journal of Autism and Developmental*
14 *Disorders*, 43(7), 1717-1724. doi: 10.1007/s10803-012-1705-7
15
16
17
18 Warren, Z., McPheeters, M. L., Sathe, N., Foss-Feig, J. H., Glasser, A., & Veenstra-VanderWeele, J.
19 (2011). A Systematic review of early intensive intervention for autism spectrum disorders.
20
21
22 *Pediatrics*, 127(5), e1303-e1311. doi: 10.1542/peds.2011-0426
23
24
25 Whitehurst, G., Falco, F., Lonigan, C., & Fischel, J. (1988). Accelerating language development through
26 picture book reading. *Developmental Psychology*, 24(4), 552-559.
27
28
29 Yoder, P., & Compton, D. (2004). Identifying predictors of treatment response. *Mental Retardation and*
30 *Developmental Disabilities Research Reviews*, 10(3), 162-168. doi: 10.1002/mrdd.20013
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

INVESTIGATING A MULTIMODAL INTERVENTION

Table 1.

Participant Information for High and Low Responders

Child	Age	Consonants	VABS-C	VABS-M	PPVT-4	CCS	Imitation-V
High Responders							
1	8;9	12	30 (28)	25	23 (24)	–	8
2	9;8	14	28 (26)	18	24 (20)	10	8
3	7;7	15	14 (28)	20	13 (25)	10	6
4	8;2	16	19 (28)	30	17 (24)	8.67	8
5	6;5	13	–	–	13 (27)	10.33	8
Low Responders							
6	10;11	15	21 (<20)	37	20 (20)	10	2
7	8;2	12	27 (31)	18	7 (20)	8	6
8	7;4	11	24 (33)	35	6 (20)	6.33	5
9	8;11	10	21 (26)	33	3 (20)	7.67	0
10	6;5	11	22 (37)	32	4 (20)	8	2

Note. VABS-C = Vineland Adaptive Behavior Scales Communication Domain; VABS-M = Vineland Adaptive Behavior Scales Maladaptive Behavior Index- high scores indicate increased maladaptive behaviors; PPVT-4 = Peabody Picture Vocabulary Test-4; CCS = Complexity of Communication Scale; Imitation-V = imitation of vocal/verbal/oral tasks. Age is age at start of data collection. Raw scores are reported; standard scores are presented in parentheses. CCS scores are based on a scale of 1-11 with 11 being highest. Imitation-V scores are number correct out of 8 from Early Steps Imitative Sequences Assessment. Child 4 is female.

INVESTIGATING A MULTIMODAL INTERVENTION

38

Table 2.

Summary of Expressive Probe Data

Child	# Words Learned (Speech Only)/words taught	Total Treatment Sessions	Average Percent Correct Production Probes–Baseline	Average Percent Correct Production Probes–Treatment	95% CI d
High Responders					
1	15/20	36	3.06	52.75	2.50 - 5.96
2	17/20	62	12.83	73.25	4.03 - 7.88
3	14/20	63	3.47	65.50	1.81 – 4.80
4	17/20	55	1.08	79.92	3.88 - 9.49
5	7/10	18	29.57	77.67	1.28 - 5.58
Low Responders					
6	4/15	66	0.00	30.67	2.32 – 7.30
7	1/10	29	0.00	10.33	.33 – 3.57
8	3/15	60	0.00	17.00	.52– 2.97
9	0/10	31	0.00	0.00	-
10	0/5	15	0.00	0.00	-

INVESTIGATING A MULTIMODAL INTERVENTION

39

Table 3.

Summary of Receptive Probe data.

Child	# Words learned ¹	Total Treatment Sessions	Average Percent Correct Receptive Probes – Baseline	Average Percent Correct Receptive Probes – Treatment	95% CI d
High responders					
1	15/20	39	38.71	76.00	1.26 – 3.26
2	20/20	62	59.00	94.42	2.40 – 5.13
3	20/20	62	40.68	88.83	2.46 – 4.94
4	18/20	54	58.57	90.50	1.26 – 3.27
5	10/10	17	57.14	64.67	-.26 – 2.70
Low responders					
6	4/15	66	38.60	54.89	.09 – 2.03
7	3/10	29	58.14	41.17	-2.84 - -.29
8	3/15	60	56.36	52.89	-1.25 - .53
9	0/10	31	50.00	56.50	-.52 – 1.82
10	0/5	11	55.67	46.67	-2.47 - .91

Note: Children 1 and 3 had a three-choice array; Child 6 had three choices in baseline for the first set, and a three-choice array thereafter; the remaining children had a two-choice array throughout the study.

¹Words learned indicates the number of words meeting the 80% correct criterion on receptive probes.




Figure 1. Multiple baselines across word sets for Child 1. Phase line indicates the beginning of intervention for the word set.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

INVESTIGATING A MULTIMODAL INTERVENTION




Figure 2. Multiple baselines across word sets for Child 2. Phase line indicates the beginning of intervention for the word set.

INVESTIGATING A MULTIMODAL INTERVENTION




Figure 3. Multiple baselines across word sets for Child 3. Phase line indicates the beginning of intervention for the word set.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

INVESTIGATING A MULTIMODAL INTERVENTION




Figure 4. Multiple baselines across word sets for Child 4. Phase line indicates the beginning of intervention for the word set.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 5. Multiple baselines across word sets for Child 5. Phase line indicates the beginning of intervention for the word set.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

INVESTIGATING A MULTIMODAL INTERVENTION

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 6. Multiple baselines across word sets for Child 6. Phase line indicates the beginning of intervention for the word set.

INVESTIGATING A MULTIMODAL INTERVENTION




Figure 7. Multiple baselines across word sets for Child 7. Phase line indicates the beginning of intervention for the word set.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

INVESTIGATING A MULTIMODAL INTERVENTION




Figure 8. Multiple baselines across word sets for Child 8. Phase line indicates the beginning of intervention for the word set.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 9. Multiple baselines across word sets for Child 9. Phase line indicates the beginning of intervention for the word set.




Figure 10. Multiple baselines across word sets for Child 10. Phase line indicates the beginning of intervention for the word set.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Appendix A. Word lists for each participant**Child 1**

Set 1: mitt, cap, bud, ten, pin
Set 2: pot, men, beat, pick, ton
Set 3: dead, pen, buck, met, bit
Set 4: hot, kick, mitt, tan, hid

Child 2

Set 1: bag, head, mane, tick, sit
Set 2: bud, wet, dock, sing, fin
Set 3: hot, bun, kick, pad, seat
Set 4: ham, pine, mad, sick, cot

Child 3

Set 1: bat, head, mane, tick, sing
Set 2: bun, mad, seat, wine, kick
Set 3: win, dad, king, met, coke
Set 4: bud, wet, cat, dock, tan

Child 4

Set 1: coke, win, dad, bus, met
Set 2: hole, bun, kick, pad, seat
Set 3: pal, wet, dock, sing, fun
Set 4: bag, head, mane, sit, tick

Child 5

Set 1: bag, head, mane, tick, sit
Set 2: pal, wet, dock, sing, fun

Child 6

Set 1: bat, done, head, tick, mane
Set 2: buck, kit, had, ten, man
Set 3: back, wet, coat, dot, tin

Child 7

Set 1: pan, mitt, bud, sing, nine
Set 2: bang, dad, sun, met, Pass

Child 8

Set 1: pen, dad, king, met, coke
Set 2: bat, fun, tick, mane, said
Set 3: pass, sing, tin, coat, fan

Child 9

Set 1: bag, done, mane, gnat, head
Set 2: hot, mitt, bang, dad, win

Child 10

Set 1: bang, dad, cop, pick, win

INVESTIGATING A MULTIMODAL INTERVENTION

50

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60