



HHS PUBLIC ACCESS

Author manuscript

J Org Chem. Author manuscript; available in PMC 2016 October 16.

Published in final edited form as:

J Org Chem. 2015 October 16; 80(20): 9942–9950. doi:10.1021/acs.joc.5b01456.

Silica-Supported Oligomeric Benzyl Phosphate (Si-OBP) and Triazole Phosphate (Si-OTP) Alkylating Reagents


Pradip K. Maity, Saqib Faisal, Alan Rolfe, Diana Stoianova[†], and Paul R. Hanson^{*}

Department of Chemistry, University of Kansas, 1251 Wescoe Hall Drive, Lawrence, KS 66045-7582, United States, and The University of Kansas Center for Chemical Methodologies and Library Development (KU-CMLD), 2034-Becker Drive, Delbert M. Shankel Structural Biology Center, Lawrence, Kansas 66047, United States

Abstract

The syntheses of silica-supported oligomeric benzyl phosphates (Si-OBP_n) and triazole phosphates (Si-OTP_n) using ring-opening metathesis polymerization (ROMP) for use as efficient alkylating reagents is reported. Ease of synthesis and grafting onto the surface of norbornenyl-tagged (Nb-tagged) silica particles has been demonstrated for benzyl phosphate and triazole phosphate monomers. It is shown that these silica polymer hybrid reagents, Si-OBP_n and Si-OTP_n, can be used to carry out alkylation reactions with an array of different nucleophiles to afford the corresponding benzylated and (triazolyl)methylated products in good yield and high purity.

Graphical abstract



INTRODUCTION

The need to rapidly synthesize a wide variety of small molecules in the desired quantities with high purities and yields is an important challenge facing drug discovery and developmental chemistry.¹ To help address this need, immobilized reagents and scavengers have been developed to provide ease of synthesis and to eliminate time-consuming chromatographic separation protocols. These reagents have found frequent use in the arena of facilitated synthesis and high-throughput chemistry,^{2,3} as well as in the scale-up of advanced pharmaceutical intermediates.⁴ Given the large range of chemistries and conditions in which they need to be employed as reagents, scavengers, or catalysts, the field has seen a number of new innovations regarding immobilized polystyrene resins,⁵ silicas,⁶

^{*}Corresponding Author: phanson@ku.edu.**Notes**

The corresponding author (PRH) is a member of the Scientific Advisory Board at Materia, Inc.

[†]Materia, Inc. 60 N. San Gabriel Blvd. Pasadena, CA 91107.

ASSOCIATED CONTENT

Supporting Information. Experimental details and NMR spectral data for new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>

soluble polyethylene glycol (PEG) polymers,⁷ monolith⁸ and fluorine-tagged compounds⁹ to remove impurities or excess reagents from reaction mixtures. Some of the more recent developments reported include silica-supported isocyanide ligands for scavenging ruthenium,¹⁰ mesoporous silica-supported boronic acids as direct amidation catalysts,¹¹ silica-supported rhodium catalysts,¹² *N*-heterocyclic carbenes,¹³ palladium and copper (I) catalysts,¹⁴ phosphines,¹⁵ prolinol and TADDOLs.¹⁶ However, for all of their advantages immobilized reagents are often limited by several factors, including (i) low load levels (ii) heterogeneous reaction kinetics and non-surface diffusion-controlled processes, (iii) immobilized reagent swelling, and (iv) poor solvent tolerance. While some of these issues have been addressed through the development of technologies such as microgels, Janda-gels and micro-porous resins,³ there is still a need for additional materials that can deliver the performance of solution-phase reagents with the ease of removal (via filtration, precipitation, extraction, etc.) afforded by their immobilization.

We have previously examined a variety of soluble, high-load, oligomeric reagents and scavengers derived through the use of ring-opening metathesis polymerization (ROMP) of functionalized norbornene and 7-oxanorbornene monomers.¹⁷ These materials, known as ROMPgels, build on the pioneering efforts of Barrett,¹⁸ Buchmeiser,¹⁹ Bolm²⁰ and others;²¹ and they have effectively been used to mediate a number of chemical reactions. However, while these ROMPgel materials have the advantage of behaving equivalently to traditional homogeneous reagents and catalysts, they are generally removed from solution via precipitation, which can limit their use in pharmaceutical applications.

As a result, inspired by the seminal work of Buchmeister and coworkers,²² efforts were focused on the synthesis and study of a number of hybrid materials that graft several of the ROMPgel materials onto the surface of silica particles.²³ Through this grafting process, the resulting immobilized ROMP reagents can be removed from solution by simple filtration, thus eliminating the precipitation step. In this fashion, silica-supported ROMPgel acid chloride, dichlorotriazine, and triphenylphosphine reagents were generated with properties nearly identical to the equivalent soluble ROMPgel oligomeric reagents.²³

We herein describe our efforts to expand the family oligomeric materials that have been grafted onto silica to include ROMP-derived oligomeric benzyl phosphate (OBP_n) and triazole phosphate (OTP_n), which we previously reported as soluble alkylating reagents that were used in facilitated library generation.²⁴ Benzylation and triazolation are useful diversification reactions in medicinal chemistry, high-throughput chemistry, and diversity-oriented synthesis (DOS).²⁵ The benzylation of amines and alcohols also serves as one of the most utilized protecting group strategies in organic synthesis due to its easy incorporation and removal.²⁶ While these uses have spurred development of a number of alternative approaches to benzylation²⁷ and triazolation, we believe the generation of the analogous silica-oligomeric benzyl phosphate (Si-OBP_n) and triazole phosphate (Si-OTP_n) reagents could find efficient, safe, and cost effective applications in chemical synthesis and library production. Key advantages of these reagents, include: (i) their stability at room temperature (ii) safety in handling when compared to commercially available benzyl bromides or iodides, and (iii) ease of purification via simple filtration through Celite®.

RESULTS AND DISCUSSION

Initial efforts centered on the synthesis of the silica-grafted oligomeric benzyl phosphate reagent, as outlined in Scheme 1, using a modification of the procedure reported by Buchmeiser.²⁸ The activated silica **1** (60 Å, 20 μm) was tagged with commercially available norbornene silyl reagent [(MeO)₃Si-(CH₂)₂-Nb (**2**)] followed by capping with trimethoxymethylsilane and dimethoxydimethylsilane to afford the norbornene-functionalized silica (Si-Nb) **3** (Scheme 1). It was observed that, the use of (MeO)₃Si-(CH₂)₂-Nb (**2**) dramatically increased the norbornene load of Si-Nb (**3**), compare to 5-(bicycloheptenyl)-triethoxysilane [Nb-Si(OEt)₃], which was previously used for silica tagging.²³ By this optimized method, we prepared norbornene-functionalized silica **3** (Si-Nb) on gram scale with 0.4 mmol/g loading (determined by modified bromine titration method).²⁹ With this Si-tagged nanoparticle (**3**) in hand, a metathesis catalyst-armed surface-initiated polymerization was established using the C848 [Grubbs 2nd generation catalyst, **G-II**]³⁰ (20 mol%, based on Si-Nb load) in CH₂Cl₂ and toluene as solvents, followed by addition of the Nb-tagged benzyl phosphate monomer **4** to rapidly generate the desired hybrid material **5a**, silica oligomeric benzyl phosphate (Si-OBP_n). The benzyl phosphate monomer **4** was itself easily synthesized in good yield and purity according to previously reported methods.^{24a} Norbornene *exo*-diol **6** was reacted with POCl₃ and Et₃N in the presence of catalytic DMAP to generate the Nb-tagged monochlorophosphate compound **7** in moderate yields as a white solid (Scheme 1). This material was then reacted with benzyl alcohol in NMI, and CH₂Cl₂ at room temperature to yield **4**. The scale-up synthesis of the monomeric reagent, as well as Si-OBP_n have been carried out on gram scale as stable, free-flowing powders. The SEM images of Si-OBP_n and Si-OTP_n silica hybrid materials demonstrate the grafting of the corresponding monomer on silica surface and inherent morphology of hybrid materials (Figure 1).

The utilization of silica oligomeric benzyl phosphate **5a** (Si-OBP_n) for benzylation with various *N*-, *O*-, *S*- nucleophiles including anilines, amines, phenols, thiols, and sulfonamides (Table 1), were next examined. The reactions were carried out in a sealed pressure tube with different nucleophiles (1 equiv.), Si-OBP_n (1.5 equiv.), Cs₂CO₃ (3.0 equiv.) and NaI (0.2 equiv.) in THF at 80 °C (oil bath temperature) to yield the products after simple filtration via Celite[®]-SPE to remove Si-phosphate by-product. The corresponding benzylated analogs (**8a–8h**) were isolated in excellent purities (>90%, determined by LC-MS) and yields. We have screened wide varieties of nucleophile for benzylation. Initially, a variety of amines and phenols (Table 1, entry 1–4) were utilized and then extended to thiophenols and sulfonamides (Table 1, entry 5–8) to yield (>90%) of the benzylated products.

With these results in hand, an expanded set of Si-OBP_n derivatives was next examined (Figure 2). These high load reagents (**5b–5d**) were synthesized in an analogous fashion to the Si-OBP_n reagent in Scheme 1 and were also obtained on gram scales as free-flowing powders. Utilization of these reagents in substitution reaction with *O*- and *S*-nucleophiles were next carried out to afford benzylated products (**8i–8n**) in good yield (>90%) and excellent purity after Celite[®]-SPE filtration (Table 2). Diversification of the Si-OBP_n reagents via substituting electron withdrawing groups, as well as electron donating groups,

on the aryl moiety were next examined, and in all cases, analogous results to Table 1 were obtained (see Table 2).

Attention was next placed on the gram-scale generation of silica oligomeric triazole phosphate hybrid reagents (Si-OTP_n). Triazoles and their derivatives have demonstrated a wide variety of biological activity, with many reports focusing on antifungal activity.³¹ We previously reported soluble oligomeric triazole phosphates to construct a wide array of triazole-containing compounds. Silica-immobilized oligomeric triazole phosphate **10a** (Si-OTP_n) was synthesized via grafting of the Nb-tagged benzyl triazole phosphate monomer **9** onto the surface of norbornene-functionalized silica (Si-Nb) **3** through the same protocol discussed for the synthesis of Si-OBP_n in Scheme 1. The product, **10a** (Si-OTP_n), was isolated as a free flowing solid in high-load (1.15 mmol/g) (Scheme 2). The triazole phosphate monomer **9** was synthesized in good yield and purity according to previously reported methods.²⁴ Phosphorylation of propargyl alcohol with Nb-tagged phosphonyl chloride **7**, followed by a “Click”-capture event of the corresponding azide, yields the desired monomer **9** in an efficient fashion (Scheme 2). The triazole phosphate monomer, as well as silica oligomeric triazole phosphate reagents (Si-OTP_n) have been synthesized on gram scale as stable, free-flowing solids.

With Si-OTP_n **10a** in hand, the hybrid material was evaluated for (triazolyl)methylation on various nucleophiles including amines, phenols, thiophenols and sulfonamides in good yields and high purity with chromatography-free purification. The reaction was carried out in a pressure tube with different nucleophiles (1 equiv.), Si-OTP_n (1.5 equiv.), Cs₂CO₃ (3.0 equiv.) and NaI (0.2 equiv.) in DMF at 90 °C for overnight. The reaction mixture was diluted with EtOAc and filtered through a pad of Celite[®] to give the corresponding (triazolyl)methylated products. Optimal results were achieved in DMF compared to THF. A variety of *N*-, *O*-, *S*-nucleophiles were utilized for nucleophilic substitution reaction with Si-OTP_n triazolating reagent gave the (triazolyl)methylated products (**11a–11g**) in excellent yield and high purity (>85%, determined by LC-MS) (Table 3). In all cases, similar results (Table 3) were obtained as compared to Si-OBP_n in THF in Table 1.

Building on these results, the project was expanded to the synthesis of additional variants of Si-OTP_n hybrid reagents (**10b–10d**, Figure 3) on gram scales as free-flowing powders in an analogous fashion to the Si-OTP_n reagent in Scheme 2. These Si-OTP_n hybrid reagents were utilized for (triazolyl)methylation of *O*- and *S*-nucleophiles. In all cases, (triazolyl)methylated products (**11h–11m**) were isolated in excellent yield and high purity after passing thru a Celite[®] SPE (Table 4).

In conclusion, the combination of Nb-tagged silica particles and functionalized Nb-tagged monomers efficiently yields high-load, hybrid Si-ROMP benzylating and (triazolyl)methylating reagents. A metathesis catalyst-armed surface (CAS)-initiated polymerization was key to functionalization of units off the silica particle surface. With this technology, we developed Si-immobilized oligomeric benzyl phosphate (Si-OBP_n) as a benzylating reagent and triazole phosphate (Si-OTP_n) as a (triazolyl)methylating reagent on gram scale as stable, free-flowing powders in all cases, utilizing ROMP of Nb-tagged phosphate monomers. SEM imaging was utilized to demonstrate the successful grafting of

the corresponding oligomer and the inherent morphology of the hybrid materials. Utilization of these reagents for representative small molecule synthesis, under purification free protocols, was demonstrated to afford corresponding products in excellent yield and with high purity. Since benzylation is a key transformation in many synthetic processes, we are currently working on expanding the scope of nucleophiles (i.e., alcohols, *bis*-nucleophilic, etc), as well as the scale on which these transformations can be carried out.

EXPERIMENTAL SECTION

General Information

All reactions were carried out under argon atmosphere. Stirring was achieved with oven-dried magnetic stir bars. Et₂O and CH₂Cl₂ were purified by passage through the Solv-Tek purification system employing activated Al₂O₃ (Pangborn, A. B.; Giardello, M. A.; Grubbs, R. H.; Rosen, R. K.; Timmers, F. J. Safe and Convenient Procedure for Solvent Purification *Organometallics* **1996**, *15*, 1518–1520). CHCl₃ was passed through basic alumina and dried over molecular sieves. Et₃N was purified by passage through basic alumina or distilled over CaH₂ and stored over KOH. For Celite-SPE 6 mL empty cartridges were used. All the solvents for routine isolation of products and chromatography were reagent grade. Flash chromatography was performed using silica gel (300–400 mesh) with the indicated solvents. Melting points were recorded on an Electrothermal digital melting point apparatus and were uncorrected. IR spectra were recorded on a spectrophotometer using KBr optics. ¹H NMR and ¹³C NMR spectra were recorded on 400 or 500 MHz (¹H NMR) and 100 or 125 MHz (¹³C NMR) spectrometers using CDCl₃ as the solvent and TMS as the internal standard. The ¹H NMR data are reported as the chemical shift in parts per million, multiplicity (s, singlet; d, doublet; t, triplet; q, quartet; m, multiplet), coupling constant in hertz, and number of protons. High-resolution mass spectra were obtained using a high-resolution ESI-TOF mass spectrometer.

General Procedure for Silica Oligomeric Benzyl Phosphate (Si-OBP_n) and Triazole Phosphate (Si-OTP_n)

Si-Nb (load ~0.4 mmol/g, 1 equiv.) was heated with C848 (**G-II**, 0.2 equiv.) at 45 °C in dichloromethane for 2 hrs under argon. The OBP or OTP monomer was added (50 equiv. w/r to cat. **G-II**) in CH₂Cl₂ and toluene to the reaction mixture and heated at 45 °C for overnight. The reaction mixture was cooled to room temperature and EVE was added, with stirring for an additional 1 hour at room temperature. The reaction mixture was filtered and washed with mixture of toluene:CH₂Cl₂ (1:1), and dried over high *vacuo* pump.

Procedure A: General procedure for different nucleophilic substitution with Si-OBP_n

In a sealed pressure tube was added Si-OBP_n **5a** (1.5 equiv.), followed by addition of sodium iodide (0.2 equiv.), Cs₂CO₃ (3.0 equiv.), and solvent THF (0.2M). The mixture was stirred rapidly and then nucleophiles were added. The reaction was sealed under argon and heated to 80 °C with stirring for 12 h. After such time, the reaction was cooled to rt and the crude mixture was filtered via a Celite[®]-packed SPE and rinsed several times with a mixture of hexanes:EtOAc (1:2). The resulting eluent was concentrated *in vacuo* to yield the benzylation products in good to excellent yields and purities.

Procedure B: General procedure for different nucleophilic substitution with Si-OTP_n

In a pressure tube was added Si-OTP_n **10a** (1.5 equiv.), followed by addition of NaI (0.2 equiv.), Cs₂CO₃ (3.0 equiv.), and solvent DMF (0.2M). The nucleophile was added and the resulting mixture was stirred rapidly. The reaction was sealed under argon and heated to 90 °C w/ stirring for (12–14 hrs) after which time, DMF was removed in *vacuo*. The crude mixture was filtered via celite[®]-SPE and rinsed with EtOAc. The resulting eluent was then concentrated *in vacuo* to yield the products in good to excellent yields and purities.

1-(benzyloxy)-4-(tert-butyl)benzene (8c)

Utilizing general procedure **A**, **8c** (18.5 mg, 0.077 mmol, 97%) was isolated as a white solid. MP: 65 °C; FTIR (neat): 3056, 3024, 2960, 2866, 1610, 1512, 1456, 1242, 1182, 1024 cm⁻¹; ¹H NMR (400 MHz, CDCl₃): δ 7.46–7.43 (m, 2H), 7.42–7.38 (m, 2H), 7.36–7.30 (m, 3H), 6.92–6.96 (m, 2H), 5.06 (s, 2H), 1.33 (s, 9H); ¹³C NMR (126 MHz, CDCl₃): δ 156.6, 143.7, 137.3, 128.5 (2C), 127.9, 127.5 (2C), 126.2 (2C), 114.2 (2C), 70.0, 34.1, 31.4 (3C); GC-MS (EI⁺) calculated for C₁₇H₂₀O 240.15; found 240.1 (M⁺ 7), 91.0 (100).^{32a}

1-(benzyloxy)-2,4-dichlorobenzene (8d)

Utilizing general procedure **A**, **8d** (30 mg, 0.119 mmol, 98%) was isolated as a thick liquid. FTIR (neat): 3031, 2931, 2835, 1597, 1481, 1452, 1382, 1290, 1249, 1060 cm⁻¹; ¹H NMR (400 MHz, CDCl₃): δ 7.47–7.44 (m, 2H), 7.43–7.38 (m, 3H), 7.37–7.33 (m, 1H), 7.18 (dd, *J* = 2.6, 8.7 Hz, 1H), 6.90 (d, *J* = 8.9 Hz, 1H), 5.16 (s, 2H); ¹³C NMR (126 MHz, CDCl₃): δ 153.0, 136.1, 130.1, 128.7 (2C), 128.1, 127.5, 127.1 (2C), 126.0, 124.1, 114.8, 71.2; GC-MS (EI⁺) C₁₃H₁₀Cl₂O calculated 252.01; found 251.9 (M⁺ 4), 91.0 (100).^{32b}

Benzyl(2,4,6-trichlorophenyl)sulfane (8e)

Utilizing general procedure **A**, **8e** (21 mg, 0.07 mmol, 98%) was isolated as a yellow solid. MP: 112 °C; FTIR (neat): 3087, 2935, 2850, 1583, 1454, 1321, 1116, 1054 cm⁻¹. ¹H NMR (400 MHz, CDCl₃): δ 7.47 (s, 1H), 7.37–7.33 (m, 4H), 7.32–7.39 (m, 2H), 4.15 (s, 2H); ¹³C NMR (126 MHz, CDCl₃): δ 136.3, 135.3, 132.0, 131.3, 130.6, 130.0, 129.5, 128.9 (2C), 128.8 (2C), 127.8, 37.6; GC-MS (EI⁺) C₁₃H₉Cl₃S calculated 301.95; found 301.9 (32), 169 (7), 91.0 (100).

Benzyl(3,4-dimethoxyphenyl)sulfane (8f)

Utilizing general procedure **A**, **8f** (22 mg, 0.085 mmol, 98%) was isolated as a thick liquid. FTIR (neat): 2933, 2833, 1595, 1504, 1251, 1228, 1135, 1026 cm⁻¹. ¹H NMR (400 MHz, CDCl₃): δ 7.31–7.19 (m, 5H), 6.95 (dd, *J* = 8.3, 2.2 Hz, 1H), 6.0–6.74 (m, 2H), 4.02 (s, 2H), 3.88 (s, 3H), 3.75 (s, 3H); ¹³C NMR (126 MHz, CDCl₃): δ 148.7 (2C), 138.2, 129.0 (2C), 128.4 (2C), 127.0, 126.2, 125.3, 115.9, 111.3, 55.9, 55.8, 41.2; GC-MS (EI⁺) C₁₅H₁₆O₂S calculated 260.09; found 260.0 (32), 169 (30), 91.0 (100).

***N*-benzyl-4-bromo-*N*-butyl-2-fluorobenzenesulfonamide (8g)**

Utilizing general procedure **A**, **8g** (24 mg, 0.060 mmol, 99%) was isolated as a thick liquid. FTIR (neat): 3089, 3025, 2958, 2931, 1589, 1472, 1456, 1396, 1344, 1161, 1135 cm⁻¹. ¹H NMR (500 MHz, CDCl₃): δ 7.79 (t, *J* = 7.70 Hz, 1H), 7.43–7.38 (m, 2H), 7.34–7.28 (m,

5H), 4.47 (s, 2H), 3.19 (t, $J = 7.60$ Hz, 2H), 1.35–1.29 (m, 2H), 1.16–1.11 (m, 2H), 0.76 (t, $J = 7.45$ Hz, 3H). ^{13}C NMR (126 MHz, CDCl_3): δ 158.44 ($^1J_{\text{C-F}} = 259$ Hz), 136.1, 131.8, 128.6 (2C), 128.2 (2C), 128.0, 127.8 (3C), 120.7 ($^2J = 25.0$ Hz), 51.4, 47.3, 29.7, 19.7, 13.5. GC-MS (EI⁺) $\text{C}_{17}\text{H}_{19}\text{BrFNO}_2\text{S}$ calculated 399.3; found 399.0 (M^+ 3), 401.2 ($\text{M}+2$ 2), 355.8 (60), 357.8 (60), 91 (100).

(S)-2,3-dibenzyl-7-bromo-3,4-dihydro-2H-benzo[b][1,4,5]oxathiazepine1,1-dioxide (8h)

Utilizing general procedure A, **8h** (30 mg, 0.066 mmol, 97%) was isolated as a white solid. MP: 170 °C. FTIR (neat): 3026, 2921, 2852, 1595, 1552, 1325, 1153, 700 cm^{-1} . ^1H NMR (500 MHz, CDCl_3): δ 7.74 (d, $J = 8.5$ Hz, 1H), 7.33–7.28 (m, 4H), 7.25–7.19 (m, 5H), 7.17 (d, $J = 1.8$ Hz, 1H), 6.92 (d, $J = 7.5$ Hz, 2H), 4.85 (t, $J = 12.0$ Hz, 1H), 4.40 (d, $J = 14.3$ Hz, 1H), 4.26 (dd, $J = 4.5, 13.42$ Hz, 1H), 4.09 (d, $J = 14.6$ Hz, 1H), 3.85 (m, 1H), 2.93 (m, 2H). ^{13}C NMR (126 MHz, CDCl_3): δ 156.1, 137.4, 135.2, 129.8, 130.8, 129.0 (2), 128.7 (2), 128.6 (2), 128.1 (2), 127.1 (2), 126.7, 126.4, 124.4, 73.7, 62.4, 55.2, 37.9. HRMS calculated for $\text{C}_{22}\text{H}_{20}\text{BrNO}_3\text{SNa}$ ($\text{M}+\text{Na}$)⁺ 480.0245; found 480.0258 (TOF MS).

1-((4-bromobenzyl)oxy)-2,4-dichlorobenzene (8i)

Utilizing general procedure A, **8i** (29 mg, 0.088 mmol, 98%) was isolated as a yellow solid. MP: 87 °C. FTIR (neat) 3070, 2923, 2866, 1585, 1571, 1480, 1456, 1290, 1262, 1103 cm^{-1} . ^1H NMR (500 MHz, CDCl_3): δ 7.52–7.54 (m, $J = 8.4$ Hz, 2H), 7.40 (d, $J = 2.7$ Hz, 1H), 7.32–7.34 (m, 2H), 7.15–7.18 (dd, $J = 8.6, 2.4$ Hz, 1H), 6.85 (d, $J = 8.8$, 1H), 5.10 (s, 2H). ^{13}C NMR (126 MHz, CDCl_3): δ 152.7, 135.0, 131.8 (2C), 130.1, 128.7 (2C), 127.5, 126.3, 124.1, 122.1, 114.7, 70.3. GC-MS (EI⁺) $\text{C}_{13}\text{H}_9\text{BrCl}_2\text{O}$ calculated 329.92; found 331.9 ($\text{M}+2$, 2), 169 (100), 90 (25).^{32b}

(4-bromobenzyl)(2,4,6-trichlorophenyl)sulfane (8j)

Utilizing general procedure A, **8j** (25 mg, 0.066 mmol, 94%) was isolated as a white solid. MP: 54 °C. FTIR (neat) 2923, 1590, 1487, 1452, 1433, 1323, 1116, 1058, 1012 cm^{-1} . ^1H NMR (500 MHz, CDCl_3): δ 7.44–7.47 (m, 3H), 7.27 (s, 1H), 7.20–7.22 (m, 2H), 4.08 (s, 2H). ^{13}C NMR (126 MHz, CDCl_3): δ 135.6, 134.5, 132.4, 131.9 (2C), 131.4, 130.7, 130.5 (2C), 130.4, 129.9, 121.7, 37.1. GC-MS (EI⁺) $\text{C}_{13}\text{H}_8\text{Cl}_3\text{BrS}$ calculated 379.86; found 379.8 (M^+ 5), 381.8 ($\text{M}+2$, 7), 169 (100), 90 (25).

2,4-dichloro-1-((4-fluorobenzyl)oxy)benzene (8k)

Utilizing general procedure A, **8k** (24 mg, 0.088 mmol, 96%) is isolated as a thick liquid. FTIR (neat) 2931, 1604, 1510, 1483, 1379, 1226, 1060, 823, 730 cm^{-1} . ^1H NMR (500 MHz, CDCl_3): δ 7.41–7.45 (m, 2H), 7.40 (d, $J = 2.5$ Hz, 1H), 7.15–7.18 (dd, $J = 8.6, 2.5$ Hz, 1H), 7.07–7.11 (m, 2H), 6.88 (d, $J = 7.5$, 1H), 5.10 (s, 2H). ^{13}C NMR (126 MHz, CDCl_3): δ 162.5 ($^1J_{\text{C-F}} = 246.5$ Hz) 152.8, 131.8, 130.1, 129.1 ($^3J = 8.4$ Hz, 2), 127.5, 126.6, 124.2, 115.6 ($^2J = 21.6$ Hz, 2) 114.9, 70.5. GC-MS (EI⁺) $\text{C}_{13}\text{H}_9\text{Cl}_2\text{FO}$ calculated 270.00; found 269.9 (M^+ 2), 109 (100).

4-fluorobenzyl(2,4,6-trichlorophenyl)sulfane (8l)

Utilizing general procedure A, **8l** (21 mg, 0.065 mmol, 95%) was isolated as a thick liquid. FTIR (neat) 2933, 1600, 1505, 1433, 1323, 1228, 1116, 1058, 837 cm^{-1} . ^1H NMR (500 MHz, CDCl_3): δ 7.49 (s, 1H), 7.31–7.34 (m, 2H), 7.29 (s, 1H), 7.01–7.05 (m, 2H), 4.13 (s, 2H). ^{13}C NMR (126 MHz, CDCl_3): δ 163.3, ($^1J_{\text{C-F}} = 247.5$ Hz) 135.8, 132.3, 131.3, 131.1 ($^4J_{\text{C-F}} = 3.6$ Hz), 130.7, 130.5 ($^3J = 8.5$ Hz, 2) 130.4, 129.9, 115.7 ($^2J = 22.3$ Hz, 2), 36.9. GC-MS (EI⁺). $\text{C}_{13}\text{H}_8\text{Cl}_3\text{FS}$ calculated 319.94; found 319.9 (M^+ 5), 109 (100).

2,4-dichloro-1-((3,5-dimethoxybenzyl)oxy)benzene (8m)

Utilizing general procedure A, **8m** (28 mg, 0.089 mmol, 98%) was isolated as a yellow solid. MP: 55 °C; FTIR (neat): 2935, 2825, 1598, 1483, 1456, 1292, 1157, 1060 cm^{-1} ; ^1H NMR (500 MHz, CDCl_3): δ 7.38 (d, $J = 2.5$ Hz, 1H), 7.14–7.16 (dd, $J = 8.7, 2.4$ Hz, 1H), 6.88–6.85 (d, $J = 7.8$ Hz, 1H), 6.60 (d, $J = 2.2$ Hz, 2H), 6.42 (t, $J = 2.5$ Hz, 1H), 5.09 (s, 1H), 3.80 (s, 6H); ^{13}C NMR (126 MHz, CDCl_3): δ 161.0 (2C), 152.9, 138.5, 130.0, 127.5, 126.1, 124.0, 114.8, 104.7 (2C), 99.8, 70.9, 55.3 (2C). GC-MS (EI⁺). $\text{C}_{15}\text{H}_{13}\text{Cl}_2\text{O}_2\text{S}$, calculated 312.03; found 312.0 (M^+ , 3), 151.1 (100).

(3,5-dimethoxybenzyl)(2,4,6-trichlorophenyl)sulfane (8n)

Utilizing general procedure A, **8n** (20 mg, 0.055 mmol, 94%) was isolated as a white solid. MP: 84 °C; FTIR (neat): 2956, 2931, 2830, 1610, 1596, 1454, 1431, 1323, 1205, 1157, 1158 cm^{-1} ; ^1H NMR (500 MHz, CDCl_3): δ 7.46 (s, 1H), 7.30 (s, 1H), 6.51 (d, $J = 2.25$ Hz, 2H), 6.38 (t, $J = 2.3$, Hz 1H), 4.08 (s, 2H), 3.79 (s, 6H). ^{13}C NMR (126 MHz, CDCl_3): δ 160.1 (2C), 137.5, 136.3, 131.9, 131.3, 130.6, 130.0, 129.4, 106.9 (2C), 99.8, 55.4 (2C), 37.8. GC-MS (EI⁺). $\text{C}_{15}\text{H}_{13}\text{Cl}_3\text{O}_2\text{S}$, calculated 361.97 found 361.9 (M^+ 3), 151.1 (100).

1-benzyl-4-((3-chlorophenoxy)methyl)-1H-1,2,3-triazole (11a)

Utilizing general procedure B, 1-benzyl-4-((3-chlorophenoxy)methyl)-1H-1,2,3-triazole **11a** (50 mg, 0.167 mmol, 91%) was isolated as a white solid. MP: 96 °C; FTIR (neat): 2960, 2358, 1610, 1511, 1463, 1250, 1184, 1031 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3): δ 7.53 (s, 1H), 7.43 – 7.35 (m, 3H), 7.29 (dd, $J = 4.8, 2.8$ Hz, 2H), 7.20 (t, $J = 8.1$ Hz, 1H), 6.98 – 6.93 (m, 2H), 6.89 – 6.84 (m, 1H), 5.55 (s, 2H), 5.17 (s, 2H). ^{13}C NMR (126 MHz, CDCl_3): δ 158.9, 144.1, 134.9, 134.4, 130.3, 129.2 (2C), 128.9, 128.2 (2C), 122.7, 121.5, 115.4, 113.1, 62.2, 54.3. HRMS calculated for $\text{C}_{16}\text{H}_{15}\text{ClN}_3\text{O}$ ($\text{M}+\text{H}$)⁺ 300.0904; found 300.0906 (TOF MS ES⁺).

1-benzyl-4-((naphthalen-1-yloxy)methyl)-1H-1,2,3-triazole (11b)

Utilizing general procedure B, **11b** (53 mg, 0.168 mmol, 90%) was isolated as a light brown solid. MP: 94 °C; FTIR (neat): 3409, 2918, 1583, 1458, 1390, 1267, 1238, 1155, 1095 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3): δ 8.23 (d, $J = 8.1$ Hz, 1H), 7.82 (d, $J = 7.6$ Hz, 1H), 7.62 (s, 1H), 7.54–7.44 (m, 3H), 7.43–7.36 (m, 4H), 7.34 – 7.25 (m, 2H), 6.98 (d, $J = 7.6$ Hz, 1H), 5.57 (s, 2H), 5.41 (s, 2H). ^{13}C NMR (126 MHz, CDCl_3): δ 153.9, 144.8, 134.5, 129.2 (2C), 128.8, 128.1 (2C), 127.5, 126.5, 125.8, 125.6, 125.3, 122.6, 122.0, 120.9, 105.4, 62.5, 54.3. HRMS calculated for $\text{C}_{20}\text{H}_{18}\text{N}_3\text{O}$ ($\text{M}+\text{H}$)⁺ 316.1450; found 316.1427 (TOF MS ES⁺).

1-benzyl-4-(pyrrolidin-1-ylmethyl)-1H-1,2,3-triazole (11c)

Utilizing general procedure B, **11c** (48 mg, 0.198 mmol, 84%) was isolated as a light brown thick liquid. FTIR (neat): 3307, 2923, 1581, 1448, 1420, 1396, 1238, 1215, 1121, 1008 cm⁻¹; ¹H NMR (400 MHz, CDCl₃): δ 7.59 (s, 1H), 7.41–7.33 (m, 3H), 7.30–7.26 (m, 2H), 5.52 (s, 2H), 3.87 (s, 2H), 2.74–2.67 (m, 4H), 1.89–1.81 (m, 4H). ¹³C NMR (126 MHz, CDCl₃): δ 144.6, 134.6, 129.1 (2C), 128.7, 128.1 (2C), 122.9, 54.2, 53.9 (2C), 50.5, 23.4 (2C). HRMS calculated for C₁₄H₁₉N₄ (M+H)⁺ 243.1610; found 243.1631 (TOF MS ES+).

1-benzyl-4-(((2,4,6-trichlorophenyl)thio)methyl)-1H-1,2,3-triazole (11e)

Utilizing general procedure B, **11e** (63 mg, 0.164 mmol, 92%) was isolated as a white solid. MP: 137 °C; FTIR (neat): 3369, 2923, 1699, 1456, 1433, 1363, 1242, 1116, 1049, 721 cm⁻¹; ¹H NMR (400 MHz, CDCl₃): δ 7.44 (s, 1H), 7.41–7.34 (m, 5H), 7.24–7.20 (m, 2H), 5.52 (s, 2H), 4.25 (s, 2H). ¹³C NMR (126 MHz, CDCl₃): δ 143.8, 135.1, 134.4, 132.3, 131.5, 130.7, 130.6, 130.0, 129.2 (2C), 128.9, 128.0 (2C), 122.1, 54.3, 27.8. HRMS calculated for C₁₆H₁₃Cl₃N₃S (M+H)⁺ 383.9896; found 383.9912 (TOF MS ES+).

1-benzyl-4-(((3,4-dimethoxyphenyl)thio)methyl)-1H-1,2,3-triazole (11f)

Utilizing general procedure B, **11f** (53 mg, 0.155 mmol, 94%) was isolated as a white solid. MP: 101 °C; FTIR (neat): 2952, 1581, 1502, 1438, 1253, 1228, 1135, 1024 cm⁻¹; ¹H NMR (400 MHz, CDCl₃): δ 7.58–7.32 (m, 3H), 7.20 (dd, *J* = 6.7, 2.7 Hz, 2H), 7.17 (s, 1H), 6.89 (dt, *J* = 5.1, 2.1 Hz, 2H), 6.72 (d, *J* = 8.2 Hz, 1H), 5.47 (s, 2H), 4.14 (s, 2H), 3.86 (s, 3H), 3.79 (s, 3H). ¹³C NMR (126 MHz, CDCl₃): δ 149.0, 148.7, 145.5, 134.6, 129.1 (2C), 128.7, 127.9 (2C), 125.7, 124.8, 121.9, 115.2, 111.4, 55.9, 55.9, 54.1, 30.8. HRMS calculated for C₁₈H₁₉N₃O₂S (M+Na)⁺ 364.1096; found 364.1063 (TOF MS ES+).

N-((1-benzyl-1H-1,2,3-triazol-4-yl)methyl)-2-bromo-N-isopropylbenzenesulfonamide (11g)

Utilizing general procedure B, **11g** (47 mg, 0.104 mmol, 89%) was isolated as a colorless thick liquid. FTIR (neat): 3134, 2923, 2846, 1703, 1604, 1487, 1456, 1411, 1328, 1220, 1049 cm⁻¹; ¹H NMR (400 MHz, CDCl₃): δ 8.14 (dd, *J* = 7.8, 1.8 Hz, 1H), 7.70, (dd, *J* = 7.7, 1.4 Hz, 1H), 7.65 (s, 1H), 7.48 – 7.32 (m, 5H), 7.27 – 7.20 (m, 2H), 5.52 (d, *J* = 5.1 Hz, 2H), 4.71 (s, 2H), 4.03 – 3.82 (m, 1H), 1.09 (d, *J* = 6.8 Hz, 6H). ¹³C NMR (126 MHz, CDCl₃): δ 147.1, 139.3, 135.6, 134.7, 133.5, 132.5, 129.1 (2C), 128.7, 127.9 (2C), 127.5, 123.6, 120.3, 54.2, 49.9, 38.6, 21.1 (2C). HRMS calculated for C₁₉H₂₁BrN₄NaO₂S (M+H)⁺ 471.0466; found 471.0478 (TOF MS ES+)

4-((4-bromophenoxy)methyl)-1-(4-methylbenzyl)-1H-1,2,3-triazole (11h)

Utilizing general procedure B, **11h** (52 mg, 0.145 mmol, 85%) was isolated as a white solid. MP: 103 °C; FTIR (neat): 3087, 2920, 1589, 1490, 1384, 1282, 1242, 1022, 825, 757 cm⁻¹. ¹H NMR (400 MHz, CDCl₃): δ 7.49 (s, 1H), 7.37 (d, *J* = 8.8 Hz, 2H), 7.19 (m, 4H), 6.85 (d, *J* = 8.8 Hz, 2H), 5.49 (s, 1H), 5.14 (s, 1H), 2.36 (s, 1H). ¹³C NMR (126 MHz, CDCl₃): δ 157.3, 144.1, 138.8, 132.3 (2C), 131.3, 129.8 (2C), 128.2 (2C), 122.6, 116.6 (2C), 113.5, 62.2, 54.1, 21.2. HRMS calculated for C₁₇H₁₇BrN₃O (M+H)⁺ 358.0555; found 358.0558 (TOF MS ES+).

1-(4-methylbenzyl)-4-(((2,4,6-trichlorophenyl)thio)methyl)-1H-1,2,3-triazole (11i)

Utilizing general procedure B, **11i** (48 mg, 0.120 mmol, 87%) was isolated as a white solid. MP: 136 °C; FTIR (neat): 3083, 2920, 1515, 1434, 1323, 1151, 1116, 1058, 871 cm⁻¹. ¹H NMR (400 MHz, CDCl₃): δ 7.43 (s, 1H), 7.38 (s, 1H), 7.31 (s, 1H), 7.18 (d, *J* = 7.9 Hz, 2H), 7.12 (d, *J* = 8.0 Hz, 2H), 5.45 (s, 2H), 4.23 (s, 2H), 2.36 (s, 3H). ¹³C NMR (126 MHz, CDCl₃): δ 143.7, 138.2, 135.1, 132.3, 131.5, 131.3, 130.7, 130.5, 130.0, 129.8 (2C), 128.0 (2C), 122.0, 54.1, 27.9, 21.2. HRMS calculated for C₁₇H₁₅Cl₃N₃S (M+H)⁺ 398.0052; found 398.0063 (TOF MS ES+).

4-((4-bromophenoxy)methyl)-1-(4-methoxybenzyl)-1H-1,2,3-triazole (11j)

Utilizing general procedure B, **11j** (67 mg, 0.179 mmol, 89%) was isolated as a white solid. MP: 98 °C FTIR (neat): 3137, 2933, 1612, 1514, 1487, 1461, 1247, 1176, 1049, 1031, 821 cm⁻¹. ¹H NMR (400 MHz, CDCl₃): δ 7.48 (s, 1H), 7.36 (dd, *J* = 9.6, 2.5 Hz, 2H), 7.23 (d, *J* = 8.6 Hz, 2H), 6.91 (d, *J* = 8.6 Hz, 2H), 6.85 (d, *J* = 8.9 Hz, 2H), 5.47 (s, 2H), 5.14 (s, 2H), 3.82 (s, 3H). ¹³C NMR (126 MHz, CDCl₃): δ 160.0, 157.3, 144.1, 132.3 (2C), 129.8 (2C), 126.3, 122.4, 116.6 (2C), 114.5 (2C), 113.5, 62.2, 55.4, 53.9. HRMS calculated for C₁₇H₁₆BrN₃NaO₂ (M+Na)⁺ 396.0324; found 396.0331 (TOF MS ES+).

1-(4-methoxybenzyl)-4-(((2,4,6-trichlorophenyl)thio)methyl)-1H-1,2,3-triazole(11k)

Utilizing general procedure B, **11k** (50 mg, 0.120 mmol, 86%) was isolated as a yellow solid. FTIR (neat): 3082, 2931, 1612, 1514, 1434, 1323, 1249, 1176, 1118, 1058 cm⁻¹; MP: 88 °C ¹H NMR (400 MHz, CDCl₃): δ 7.35 (m, 1H), 7.31 (m, 1H), 7.22 (s, 1H), 7.20–7.17 (m, 1H), 7.09 (dd, *J* = 8.5, 1.8 Hz, 2H), 6.82 (dd, *J* = 8.4, 2.1 Hz, 2H), 5.34 (s, 2H), 4.14 (s, 2H), 3.74 (s, 3H). ¹³C NMR (126 MHz, CDCl₃): δ 160.0, 143.6, 135.2, 132.3, 131.5, 130.7, 130.5, 130.0, 129.6 (2C), 126.3, 121.9, 114.5 (2C), 55.4, 53.8, 27.9. HRMS calculated for C₁₇H₁₅Cl₃N₃OS (M+H)⁺ 414.0001; found 414.0007 (TOF MS ES+).

4-((4-bromophenoxy)methyl)-1-(cyclohexylmethyl)-1H-1,2,3-triazole (11l)

Utilizing general procedure B, **11l** (52 mg, 0.148 mmol, 86%) was isolated as a white solid. FTIR (neat): 2921, 2852, 1488, 1446, 1384, 1244, 1224, 1112, 1054 cm⁻¹; MP: 90 °C; ¹H NMR (400 MHz, CDCl₃): 7.54 (s, 1H), 7.41–7.35 (ddd, *J* = 10.2, 3.4, 2.2, 2H), 6.91–6.85 (ddd, *J* = 10.2, 3.4, 2.2, 2H), 5.19 (s, 2H), 4.18 (d, *J* = 7.2 Hz, 2H), 1.94–1.83 (m, 1H), 1.76–1.61 (m, 5H), 1.29–1.14 (m, 3H), 1.04–0.94 (m, 2H). ¹³C NMR (126 MHz, CDCl₃): δ 157.3, 143.5, 132.3, 123.1, 116.7, 113.5, 62.3, 56.6, 38.7, 30.5 (2C), 26.0, 25.5 (2C). HRMS calculated for C₁₆H₂₁BrN₃O (M+H)⁺ 350.0868; found 350.0870 (TOF MS ES+).

1-(cyclohexylmethyl)-4-(((2,4,6-trichlorophenyl)thio)methyl)-1H-1,2,3-triazole (11m)

Utilizing general procedure B, **11m** (48 mg, 0.122 mmol, 88%) was isolated as a white solid. FTIR (neat): 2925, 2852, 1450, 1427, 1215, 1153, 1116 cm⁻¹; MP: 141 °C. ¹H NMR (500 MHz, CDCl₃): δ 7.46 (s, 1H), 7.39 (d, *J* = 5.3 Hz, 2H), 4.27 (s, 2H), 4.14 (d, *J* = 7.2 Hz, 2H), 1.89–1.79 (m, 1H), 1.76–1.61 (m, 4H), 1.58–1.50 (m, 2H), 1.28–1.09 (m, 4H), 0.95 (qd, *J* = 12.2, 3.2 Hz, 2H). ¹³C NMR (126 MHz, CDCl₃): δ 143.1, 135.2, 132.3, 131.5, 130.7, 130.5, 130.0, 122.6, 56.6, 38.7, 30.4 (2C), 27.8, 26.0, 25.5 (2C). HRMS calculated for C₁₆H₁₉Cl₃N₃S (M+H)⁺ 390.0365; found 390.0377 (TOF MS ES+).

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This research was made possible by the National Institute of General Medical Sciences [NIH STTR R42 GM097896 and NIH Center for Chemical Methodologies and Library Development at the University of Kansas (P50 GM069663)]. The authors thank Justin Douglas and Sarah Neuenswander in the University of Kansas NMR Laboratory. We acknowledge Dr. Prem Thapa and Ms. Heather Shinogle for carrying out SEM analysis (KU). We thank Materia Inc. for providing metathesis catalyst. We also thank Dr. Patrick Kearney for helpful discussion and proofing the manuscript.

References

1. (a) Gallop MA, Barrett RW, Dower WJ, Fodor SPA, Gordon EM. *J Med Chem.* 1994; 37:1233–1251. [PubMed: 8176700] (b) Flynn DL, Devraj RV, Naing W, Parlow JJ, Weidner JJ, Yang S. *Med Chem Res.* 1998; 8:219–243. (c) Musonda CC, Chibale K. *Curr Med Chem.* 2004; 11:2519–2533. [PubMed: 15544460] (d) Dolle RE, Bourdonnec BL, Worm K, Morales GA, Thomas CJ, Zhang W. *J Comb Chem.* 2010; 12:765–806. [PubMed: 20923157] (e) Schreiber SL. *Proc Natl Acad Sci U S A.* 2011; 108:6699–6702. [PubMed: 21464328] (f) Chatterjee AK. *J Med Chem.* 2013; 56:7741–7749. [PubMed: 23927720]
2. (a) Kirschning A, Monenschein H, Wittenberg R. *Angew Chem Int Ed.* 2001; 40:650–679. (b) Eames J, Watkinson M. *Eur J Org Chem.* 2001:1213–1224. (c) Booth RJ, Hodges JC. *Acc Chem Res.* 1999; 32:18–26. (d) Ley SV, Baxendale IR, Bream RN, Jackson PS, Leach AG, Longbottom DA, Nesi M, Scott JS, Storer RI, Taylor SJ. *J Chem Soc, Perkin Trans.* 2000; 1:3815–4195. (e) Strohmeier GA, Kappe CO. *Angew Chem Int Ed.* 2004; 43:621–624. (f) Lesch B, Thomson DW, Lindell SD. *Comb Chem High T Scr.* 2008; 11:31–36.
3. (a) Gravert DJ, Janda KD. *Chem Rev.* 1997; 97:489–509. [PubMed: 11848880] (b) Toy PH, Janda KD. *Acc Chem Res.* 2000; 33:546–554. [PubMed: 10955985] (c) Dickerson TJ, Reed NN, Janda KD. *Chem Rev.* 2002; 102:3325–3344. [PubMed: 12371887] (d) Haag R. *Chem Eur J.* 2001; 7:327–335. [PubMed: 11271517] (e) Haag R, Sunder A, Hebel A, Roller S. *J Comb Chem.* 2002; 4:112–119. [PubMed: 11886284] (f) Bergbreiter DE. *Chem Rev.* 2002; 102:3345–3384. [PubMed: 12371888] (g) Bergbreiter DE, Tian J, Hongfa C. *Chem Rev.* 2009; 109:530–582. [PubMed: 19209941]
4. (a) Bhattacharyya S. *Curr Opin Drug Dis Dev.* 2004; 7:752–764. (b) Hudson D. *J Comb Chem.* 1999; 1:333–360. [PubMed: 10748732] (c) Polshettiwar V, Varma RS. *Acc Chem Res.* 2008; 41:629–639. [PubMed: 18419142] (d) Malet-Sanz L, Susanne F. *J Med Chem.* 2012; 55:4062–4098. [PubMed: 22283413]
5. (a) Lu J, Toy PH. *Chem Rev.* 2009; 109:815–838. [PubMed: 19128147] (b) Takashima H, Vigneaud VD, Merrifield RB. *J Am Chem Soc.* 1968; 90:1323–1325. [PubMed: 5636536] (c) Baxendale IR, Brusotti G, Matsuoka M, Ley SV. *J Chem Soc, Perkin Trans 1.* 2002:143–154. (d) Storer RI, Takemoto T, Jackson PS, Brown DS, Baxendale IR, Ley SV. *Chem Eur J.* 2004; 10:2529–2547. [PubMed: 15146525] (e) Baxendale IR, Ley SV, Piutti C. *Angew Chem, Int Ed.* 2002; 41:2194–2197. (f) Roller S, Türk H, Stumbé JF, Rapp W, Haag R. *J Comb Chem.* 2006; 8:350–354. [PubMed: 16677004]
6. (a) Guillier F, Orain D, Bradley M. *Chem Rev.* 2000; 100:2091–2157. [PubMed: 11749285] (b) Polshettiwar V, Len C, Fihri A. *Coord Chem Rev.* 2009; 253:2599–2626. (c) Moreno J, Iglesias J, Melero JA, Sherrington DC. *J Mater Chem.* 2011; 21:6725–6735. (d) Fihri A, Bouhrara M, Patil U, Cha D, Saih Y, Polshettiwar V. *ACS Cat.* 2012; 2:1425–1431. (e) Sharma RK, Yukti M. *App Cat A: General.* 2013; 454:1–10.
7. (a) Crauste C, Périgaud C, Peyrottes S. *J Org Chem.* 2011; 76:997–1000. [PubMed: 21192699] (b) Adams JH, Cook RM, Hudson D, Jammalamadaka V, Lyttle MH, Songster MF. *J Org Chem.* 1998; 63:3706–3716. (c) Gooding OW, Baudart S, Deegan TL, Heisler K, Labadie JW, Newcomb WS, Porco JA Jr, van Eikeren P. *J Comb Chem.* 1999; 1:113–122. (d) Kalinina I, Worsley K, Lugo C,

- Mandal S, Bekyarova E, Haddon RC. *Chem Mater.* 2011; 23:1246–1253.(e) Lee JW, Fuchs PL. *Org Lett.* 1999; 1:179–181. [PubMed: 10822557]
8. (a) Bandari R, Prager-Duschke A, Kühnel C, Decker U, Schlemmer B, Buchmeiser MB. *Macromolecules.* 2007; 28:2090–2094.(b) Mayr M, Mayr B, Buchmeiser MR. *Angew Chem Int Ed.* 2001; 40:3839–3842.(c) Moitra N, Ichii S, Kamei T, Kanamori K, Zhu Y, Takeda K, Nakanishi K, Shimada T. *J Am Chem Soc.* 2014; 136:11570–11573. [PubMed: 25101719]
9. (a) Curran DP. *Aldrichimica Acta.* 2006; 39:3–9.(b) Zhang W. *Chem Rev.* 2004; 104:2531–2556. [PubMed: 15137799] (c) Curran DP. *Med Res Rev.* 2000; 72:1649–1653.(d) Kim J, Lee WS, Koo J, Lee J, Park SB. *ACS Comb Sci.* 2014; 16:24–32. [PubMed: 24215277] (e) Curran DP. *Pure Appl Chem.* 2000; 72:1649–1653.(f) Dandapani S, Curran DP. *J Org Chem.* 2004; 69:8751–8757. [PubMed: 15575753] (g) Fustero S, Garcia SA, Chiva G, Sanz-Cervera JF, Del PC, Acena JL. *J Org Chem.* 2006; 71:3299–3302. [PubMed: 16599635] (h) Leach SG, Cordier CJ, Morton D, McKiernan GJ, Warriner S, Nelson A. *J Org Chem.* 2008; 73:2753–2759. [PubMed: 18327951] (i) Zhang W. *Chem Rev.* 2009; 109:749–795. [PubMed: 19146385] (j) Sugiyama Y, Kurata Y, Kunda Y, Miyazaki A, Matsui J, Nakamura S, Hamamoto H, Shioiri T, Matsugi M. *Tetrahedron.* 2012; 68:3885–3892.(k) Sugiyama Y, Ishihara K, Masuda Y, Kobayashi Y, Hamamoto H, Matsugi M. *Tetrahedron Lett.* 2013; 54:2060–2062.
10. French JM, Caras CA, Diver ST. *Org Lett.* 2013; 15:5416–5419. [PubMed: 24128003]
11. Gu L, Lim J, Cheong JL, Lee SS. *Chem Commun.* 2014; 50:7017–7019.
12. Chepiga KM, Feng Y, Brunelli NA, Jones CW, Davies HML. *Org Lett.* 2013; 15:6136–6139. [PubMed: 24251986]
13. Ghiaci M, Zarghani M, Khojastehnezhad A, Moeinpour F. *RSC Adv.* 2014; 4:15496–15501.
14. (a) Alesi S, Di Maria F, Melucci M, Macquarrie DJ, Luque R, Barbarella G. *Green Chem.* 2008; 10:517–523.(b) Opanasenko M, Stepnicka P, Cejka J. *RSC Adv.* 2014; 4:65137–65162.(c) Miao T, Wang L. *Synthesis.* 2008:363–368.(d) Onozawa S, Fukaya N, Saitou K, Sakakura T, Yasuda H. *Catal Lett.* 2011; 141:866–871.
15. (a) Iwai T, Harada T, Tanaka R, Sawamura M. *Chem Lett.* 2014; 43:548–586.(b) Tsiavalariaris G, Haubrich S, Merckle C, Blümel J. *Synlett.* 2001:391–393.
16. (a) Altava B, Burguete MI, Garcia-Verdugo E, Luis SV, Vicent MJ. *Tetrahedron Lett.* 2001; 42:8459–8462.(b) Heckel A, Seebach D. *Chem Eur J.* 2002; 8:559–572. [PubMed: 11855704]
17. (a) Maity PK, Kainz QM, Faisal S, Rolfe A, Samarakoon TB, Basha FZ, Reiser O, Hanson PR. *Chem Commun.* 2011; 47:12524–12526.(b) Rolfe A, Probst D, Volp K, Omar I, Flynn D, Hanson PR. *J Org Chem.* 2008; 73:8785–8790. [PubMed: 18937412] (c) Stoianova DS, Yao L, Rolfe A, Samarakoon T, Hanson PR. *Tetrahedron Lett.* 2008; 49:4553–4555. [PubMed: 19319202] (d) Herpel RH, Vedantham P, Flynn DL, Hanson PR. *Tetrahedron Lett.* 2006; 47:6429–6432.(e) Harned AM, He Song H, Toy PH, Flynn DL, Hanson PR. *J Am Chem Soc.* 2005; 127:52–53. [PubMed: 15631444] (f) Zhang M, Flynn DL, Hanson PR. *J Org Chem.* 2007; 72:3194–3198. [PubMed: 17407352] (g) Harned AM, Sherrill WM, Flynn DL, Hanson PR. *Tetrahedron.* 2005; 61:12093–12099.(h) Zhang M, Moore JD, Flynn DL, Hanson PR. *Org Lett.* 2004; 6:2657–2660. [PubMed: 15281737] (i) Asad N, Hanson PR, Long TR, Rayabharapu DK, Rolfe A. *Chem Commun.* 2011; 47:9528–9530.(j) Maity PK, Rolfe A, Samarakoon TB, Faisal S, Kurtz RD, Long TR, Schatz A, Flynn DL, Grass RN, Stark WJ, Reiser O, Hanson PR. *Org Lett.* 2011; 13:8–10. [PubMed: 21121636] (k) Kainz QM, Linhardt R, Maity PK, Hanson PR, Reiser O. *ChemSusChem.* 2013; 6:721–729. [PubMed: 23427021] (l) Harned AM, Zhang M, Vedantham P, Mukherjee S, Herpel RH, Flynn DL, Hanson PR. *Aldrichimica Acta.* 2005; 38:3–16.
18. (a) Barrett AGM, Cramp SM, Roberts RS. *Org Lett.* 1999; 1:1083–1086.(b) Ahmed M, Barrett AGM, Braddock DC, Cramp SM, Procopiou PA. *Tetrahedron Lett.* 1999; 40:8657–8662.(c) Fuchter MJ, Hoffman BM, Barrett AGM. *J Org Chem.* 2006; 71:724–729. [PubMed: 16408985] (d) Barrett AGM, Cramp SM, Roberts RS. *Org Lett.* 1999; 1:1083–1086.(e) Barrett AGM, Hopkins BT, Köbberling. *J Chem Rev.* 2002; 102:3301–3324.(f) Barrett AGM, Hopkins BT, Love AC, Tedeschi L. *Org Lett.* 2004; 6:835–837. [PubMed: 14986987] (g) Arstad E, Barrett AGM, Tedeschi L. *Tetrahedron Lett.* 2003; 44:2703–2707.
19. (a) Buchmeiser MR, Atzl N, Bonn GK. *J Am Chem Soc.* 1997; 119:9166–9174.(b) Buchmeiser MR. *Chem Rev.* 2000; 100:1565–1604. [PubMed: 11749276] (c) Buchmeiser MR. *Bioorg Med Chem Lett.* 2002; 12:1837–1840. [PubMed: 12086829] (d) Buchmeiser, MR. In *Handbook of*

- Metathesis. Grubbs, RH., editor. Vol. 3. Wiley-VCH; Weinheim: 2003. p. 226-254.(e) Lubbad SH, Bandari R, Buchmeiser MR. *J Chromatogr A*. 2011; 1218:8897–8902. [PubMed: 21450301] (f) Wang D, Unold J, Bubrin M, Frey W, Kaim W, Buchmeiser MR. *ChemCatChem*. 2012; 4:1808–1812.(g) Buchmeiser, MR. *Synthesis of Polymers*. Schlueter, DA.; Hawker, CJ.; Sakamoto, J., editors. Vol. 2. 2012. p. 547-586.(h) Buchmeiser MR. *Curr Org Chem*. 2013; 17:2764–2775.(i) Naumann S, Schmidt FG, Frey W, Buchmeiser MR. *Polymer Chemistry*. 2013; 4:4172–4181.
20. (a) Bolm C, Dinter CL, Seger A, Höcker H, Brozio J. *J Org Chem*. 1999; 64:5730–5731.(b) Bolm C, Dinter CL, Schiffrers I, Defrere L. *Synlett*. 2001:1875–1877.(c) Bolm C, Tanyeli C, Grenz A, Dinter CL. *Adv Synth Catal*. 2002; 344:649–656.
21. (a) Roberts RS. *J Comb Chem*. 2005; 7:21–32. [PubMed: 15638475] (b) Zhang M, Flynn DL, Hanson PR. *J Org Chem*. 2007; 72:3194–3198. [PubMed: 17407352] (c) Nguyen MH, Smith AB III. *Org Lett*. 2013; 15:4258–4261. [PubMed: 23901881] (d) Nguyen MH, Smith AB III. *Org Lett*. 2013; 15:4872–4875. [PubMed: 24000819] (e) Flynn DL, Hanson PR, Berk SC, Makara GM. *Curr Opin Drug Discovery Dev*. 2002; 5:571–579.(f) Harned, AM.; Probst, DA.; Hanson, PR. In *Handbook of Metathesis*. Grubbs, RH., editor. Wiley-VCH; Weinheim, Germany: 2003. p. 361-402.
22. (a) Buchmeiser MR. *Chem Rev*. 2009; 109:303–321. [PubMed: 18980343] (b) Zou H, Wu S, Shen J. *Chem Rev*. 2008; 108:3893–3957. [PubMed: 18720998] (c) Timofte RS, Woodward S. *Tetrahedron Lett*. 2004; 45:39–42.(d) Butterworth AJ, Clark JH, Walton PH, Barlow SJ. *Chem Commun*. 1996:1859–1860.
23. Rolfe A, Loh JK, Maity PK, Hanson PR. *Org Lett*. 2011; 13:4–7. [PubMed: 21128690]
24. (a) Long T, Maity PK, Samarakoon TB, Hanson PR. *Org Lett*. 2010; 12:2904–2907. [PubMed: 20521800] (b) Long TR, Faisal S, Maity PK, Rolfe A, Kurtz RD, Klimberg SV, Najjar MR, Basha FZ, Hanson PR. *Org Lett*. 2011; 13:2038–2041. [PubMed: 21434675] (c) Faisal S, Farmann U, Maity PK, Rolfe A, Basha FZ, Organ MG, Hanson PR. *ACS Comb Sci*. 2012; 14:268–272. [PubMed: 22384820]
25. (a) Dolle RE, Le Bourdonnec B, Goodman AJ, Morales GA, Thomas CJ, Zhang W. *J Comb Chem*. 2009; 11:739–790. [PubMed: 19715292] (b) Fenster E, Rayabarapu DK, Zhang M, Mukherjee S, Hill D, Neuenswander B, Schoenen F, Hanson PR, Aubé J. *J Comb Chem*. 2008; 10:230–234. [PubMed: 18254600]
26. (a) Paquette, LA. *Encyclopedia of Reagents for Organic Synthesis*. 1. John Wiley and Sons; New York: 1995. p. 316-318.(b) March, J. *Advanced Organic Chemistry*. 4. Wiley; New York: 1991. (c) Greene, TW.; Wuts, PGM. *Protective Groups in Organic Synthesis*. 4. John Wiley and Sons; New York: 2007. p. 102-148.
27. Lopez SS, Dudley GB. *Beilstein J Org Chem*. 2008; 4(44)
28. (a) Buchmeiser MR, Sinner F, Mupa M, Wurst K. *Macromolecules*. 2000; 33:32–39.(b) Eder K, Reichel E, Schottenberger H, Huber CG, Buchmeiser MR. *Macromolecules*. 2001; 34:4334–4341. (c) Buchmeiser MR, Atzl N, Bonn GK. *J Am Chem Soc*. 1997; 119:9166–9174.(d) Ambrose D, Fritz JS, Buchmeiser MR, Altz N, Bonn GK. *J Chromatogr, A*. 1997; 786:259–268.
29. Johnson HL, Clark RA. *Anal Chem*. 1947; 19:869–872.
30. (a) Nguyen ST, Johnson LK, Grubbs RH, Ziller JW. *J Am Chem Soc*. 1992; 114:3974–3975.(b) Schwab P, France MB, Ziller JW, Grubbs RH. *Angew Chem Int Ed*. 1995; 34:2039–2041.(c) Schwab P, Grubbs RH, Ziller JW. *J Am Chem Soc*. 1996; 118:100–110.(d) Huang J-K, Stevens ED, Nolan SP, Petersen JL. *J Am Chem Soc*. 1999; 121:2674–2678.(e) Kingsbury JS, Harrity JPA, Bonitatebus PJ, Hoveyda AH. *J Am Chem Soc*. 1999; 121:791–799.(f) Vougioukalakis GC, Grubbs RH. *Chem Rev*. 2010; 110:1746–1787. [PubMed: 20000700]
31. (a) Xia Y, Qu F, Peng L. *Rev Med Chem*. 2010; 10:806–821.(b) Cronin S, Chandrasekar PHJ. *Antimicrob Chemother*. 2010; 65:410–416.(c) Nivoix Y, Ubeaud-Sequier G, Engel P, Leveque D, Herbrecht R. *Curr Drug Metab*. 2009; 10:395–409. [PubMed: 19519346]
32. (a) Bu X, Jing H, Wang L, Chang T, Jin L, Yongmin L. *J Mol Cat A: Chem*. 2006; 259:121–124.(b) Chakraborti AK, Nayak MK, Sharma L. *J Org Chem*. 1999; 64:8027–8030.




Figure 1.
SEM images of Si-Nb (left), Si-OBP_n (middle) and Si-OTP_n (right).




Figure 2.
Various silica-supported Oligomeric Benzyl Phosphate (Si-OBP_n) analogs.






Figure 3.
Various silica-supported Oligomeric Triazole Phosphate (Si-OTP_n) analogs.



Scheme 1.
Synthesis of silica-supported Oligomeric Benzyl Phosphate (Si-OBP_n).


**Scheme 2.**

Synthesis of silica-supported Oligomeric Triazole Phosphate (Si-OTP_n).


Table 1

Benzylation of *N*-, *O*- and *S*-nucleophiles utilizing Si-OBP_n.

Entry	Nucleophile	Pdt	Yield (%)
1			97
		8a	
2			95
		8b	
3			97
		8c	
4			98
		8d	
5			98
		8e	
6			98
		8f	
7			99
		8g	
8			97
		8h	

Table 2Benzylation of *N*-, *O*- and *S*-nucleophiles utilizing various Si-OBP_n analogs.

Entry	Si-OBP _n	Nucleophile	Pdt	Yield (%)
1	5b	2,4-Cl-PhOH	8i	98
2	5b	2,4,6-Cl-PhSH	8j	94
3	5c	2,4-Cl-PhOH	8k	96
4	5c	2,4,6-Cl-PhSH	8l	95
5	5d	2,4-Cl-PhOH	8m	98
6	5d	2,4,6-Cl-PhSH	8n	94

Table 3Triazolization utilizing Si-OTP_n with *O*-, *N*- and *S*-nucleophiles.
















Entry	Nucleophile	Pdt	yield (%)
1		 11a	91
2		 11b	90
3		 11c	84
4		 11d	83
5		 11e	92
6		 11f	94
7		 11g	89

Table 4Triazolation of *N*-, *O*- and *S*-nucleophiles utilizing various Si-OTP_n Analogs.

Entry	Si-OTP _n	Nucleophile (RB-XH)	Pdt	Yield (%)
1	10b , R ⁴ = 4-MePh	4-Br-PhOH	11h	85
2	10b , R ⁴ = 4-MePh	2,4,6-Cl-PhSH	11i	87
3	10c , R ⁴ = 4-OMePh	4-Br-PhOH	11j	89
4	10c , R ⁴ = 4-OMePh	2,4,6-Cl-PhSH	11k	86
5	10d , R ⁴ = cyclohexyl	4-Br-PhOH	11l	86
6	10d , R ⁴ = cyclohexyl	2,4,6-Cl-PhSH	11m	88