



## NIH PUBLIC ACCESS

## Author Manuscript

*Br J Nutr.* Author manuscript; available in PMC 2013 August 05.

Published in final edited form as:

*Br J Nutr.* 2010 June ; 103(12): 1702–1705. doi:10.1017/S0007114509993771.

## Dietary predictors of visceral adiposity in overweight young adults

Bruce W. Bailey<sup>1</sup>, Debra K. Sullivan<sup>2</sup>, Erik P. Kirk<sup>3</sup>, and Joseph E. Donnelly<sup>4</sup><sup>1</sup>Department of Exercise and Science, Brigham Young University, Provo, Utah 84602, USA<sup>2</sup>Department of Dietetics and Nutrition, University of Kansas Medical Center, 3901 Rainbow Boulevard, Kansas City, KS 66160, USA<sup>3</sup>Department of Kinesiology and Health Education, Southern Illinois University Edwardsville, Campus Box 1126, Edwardsville, IL 62026-1126, USA<sup>4</sup>Energy Balance Laboratory, University of Kansas, 1301 Sunnyside Avenue, Lawrence, KS 66045-7567, USA

### Abstract

The purpose of the present study was to determine the dietary predictors of visceral adipose tissue (VAT) area in overweight young adults. A total of 109 young adults (fifty males and fifty-nine females) ate ad libitum in a university cafeteria for 14 d. All food and beverages consumed in the cafeteria were measured using observer-recorded weighed plate waste. Food consumption outside the cafeteria (i.e. snacks) was assessed by multiple-pass 24 h recall procedures. VAT was determined using computed tomography. Stepwise regression demonstrated that the best predictor of visceral adiposity in women was total dietary fat (P# 0–05). In men, the model for predicting visceral adiposity included Ca and total dietary fat. We concluded that total dietary fat is the best predictor of VAT area in both men and women. While this relationship was independent in women, in men there was a synergistic relationship between dietary fat consumption and Ca consumption in predicting VAT.

### Keywords

Visceral fat; Obesity phenotypes; Dietary fat; Calcium; Waist circumference

---

Excess abdominal fat is associated with increased mortality and risk of diabetes, hyperlipidaemia, hypertension and atherosclerosis(1). Abdominal fat is composed of both subcutaneous and visceral fat. The detrimental influence of abdominal obesity on metabolic processes seems to be mediated by the visceral adipose tissue (VAT) depot, although there is some debate on the role of subcutaneous adipose tissue in these abnormalities, specifically the deep subcutaneous layers(1,2).

---

Correspondence to: Bruce W. Bailey.

The institution where the present study was performed at is the University of Kansas, Energy Balance Laboratory (Lawrence, KS, USA).

B. W. B. was the primary author of the manuscript and participated in all aspects of the study. D. K. S. served as the study dietitian and assisted in the collection and analysis of data, as well as the writing of the manuscript. E. P. K. assisted in data collection and database management, and helped write the methods and proofed the entire paper. J. E. D. was responsible for acquiring the initial funding for this project and also was involved in running the study and writing the manuscript.

There are no conflicts of interest.

Factors that influence VAT have not been clearly defined in young adult populations(3,4). Sex, genetics and sex hormones appear to play a role in the preferential deposition of VAT(5,6). Physical activity may also play a role in VAT accumulation, especially in men(7,8). While diet plays an obvious role in the accumulation of VAT, the complete role of diet on VAT has not been well described. Studies indicate that there is a positive relationship between energy consumption and VAT, but not all studies demonstrate a strong relationship(9,10). Less is known about other components of the diet including macronutrient and micronutrient composition. The purpose of the present study was to examine the relationship between diet and VAT area in overweight young adults.

## Methods

### Design

This is a cross-sectional study using baseline data from participants who were originally recruited for the Midwest Exercise Trial(11). There were four cohorts of participant that took part in the Midwest Exercise Trial. The time between cohorts was roughly 6 months between cohorts 1 and 2, 18 months between cohorts 2 and 3 and 12 months between cohorts 3 and 4. Participants were included in the study once and were not allowed to take part in subsequent cohorts.

### Participants

Participants were aged between 17 and 35 years, sedentary, did not exceed 2093 kJ (500kcal) of physical activity per week, and had a BMI between 25.0 and 34.9kg/m<sup>2</sup>(12). Participants were healthy with no history of chronic disease. Additionally, all participants met or exceeded the 85th percentile for triceps skinfold thickness of the Second National Health and Nutrition Examination Survey. This measurement helped to eliminate subjects who were short and muscular rather than overweight.

The present study was conducted according to the guidelines laid down in the Declaration of Helsinki and all procedures involving human participants were approved by the Institutional Review Board of the University. Written informed consent was obtained from all participants.

### Assessments

**Body weight and composition**—Body weight was measured using a digital scale accurate to within 0.1 kg. Body composition was estimated by hydrostatic weighing using measured residual volume(11). VAT area was determined using computed tomography (CT) at the L4–5 vertebral space. The CT examinations were performed using a CTi scanner (General Electric, Waukesha, WI, USA) at the University of Colorado Hospital (Denver, CO, USA)(11).

**Energy intake and diet composition**—Diet intake was ad libitum and was assessed for 14 d in the university cafeteria. All food and beverages consumed in the cafeteria were measured using observer-recorded weighed plate waste(13). Food consumption outside the cafeteria was assessed by multiple-pass 24 h diet recall procedures(13). This method of evaluating dietary intake was validated against doubly labelled water in fifty-two of the participants in the present study. Results from this validation showed that the participants were in energy balance during the assessment period(13).

The cafeteria featured a food court where different stations offered specialised items such as stir-fry, pasta, salads, sandwiches, pizza and grilled foods. Participants could typically

choose from between eight and ten entrees at each meal. The cafeteria was open for a total of 9–5 h on weekdays and 5–5 h (brunch and dinner) on the weekends.

Results from the plate waste and from diet recalls were entered into a computerised nutrition database for analysis (Food Processor, version 7.1; ESHA Research, Salem, OR, USA).

included in the stepwise regression included total energy intake, dietary fat, dietary protein, dietary carbohydrate, saturated fat, Fe, vitamin A, folic acid, dietary fibre, Ca, alcohol, and Ca:protein ratio. Pearson product moment correlations were calculated to determine the relationship between diet components and VAT or subcutaneous adipose tissue. Independent t tests were used to determine differences between the sexes in descriptive and dietary variables. Significance was set at  $P < 0.05$ . Statistical analysis was performed using SAS (version 8.2; SAS Institute, Inc., Cary, NC, USA).

## Results

Descriptive characteristics of the participants in the study are reported in Table 1. In general, men were heavier than women and had more lean tissue and less fat tissue ( $P < 0.05$ ). Men had a higher VAT area, while women had a higher subcutaneous adipose tissue area ( $P < 0.05$ ). Diet intake of the participants is reported in Table 2. Men consumed more energy and a greater percentage of the energy was from fat (34.6 v. 31.4%) and protein (13.7 v. 12.4%), while women consumed a greater percentage of their energy from carbohydrate (57.6 v. 51%). In general, men consumed more Ca but had a similar Ca:protein ratio compared with women.

Stepwise regression demonstrated that total dietary fat was the best predictor of both VAT and subcutaneous adipose tissue area ( $r = 0.36$  and  $r = 0.48$ ;  $P < 0.05$ ) in women (see Table 3). Saturated fat was similarly associated with both VAT and subcutaneous adipose tissue area in women ( $r = 0.32$  and  $r = 0.40$ ;  $P < 0.05$ ), while total energy intake only predicted subcutaneous adipose tissue area ( $r = 0.36$ ;  $P < 0.05$ ).

There was no single dietary component that predicted VAT area or subcutaneous adipose tissue area in men. Stepwise regression demonstrated that total dietary fat and Ca consumption in combination predicted VAT area ( $r = 0.36$ ;  $P < 0.05$ ) in men (see Table 4).

## Discussion

Total dietary fat predicted VAT in women but did not independently predict VAT in men. While dietary fat did not independently predict VAT in men, a synergistic relationship was observed between dietary fat consumption and Ca intake. If fat intake was similar, men who consumed more Ca had a lower VAT area.

Cross-sectional studies are generally in agreement that dietary fat is associated with excess body weight(14). Less is known about the relationship between dietary fat and VAT. Greenfield et al. demonstrated a relationship between dietary fat and weight in adult women but failed to show a relationship with VAT(15). It is unclear why there is a lack of agreement between these findings and the findings from the present study. It could be the result of methodological differences in assessing diet and VAT between the studies. In the Greenfield study, diet was assessed using the Oxford FFQ and VAT was assessed using dual-energy X-ray absorptiometry. These measurements may lack the precision necessary to determine the relationship between dietary fat and VAT.

In contrast, Dutia et al. in a published abstract showed a positive relationship between MUFA consumption and VAT accumulation in sedentary men and women(16). Diet was

measured using a 3 d food record and VAT was assessed using MRI. Based on the abstract, it is unclear if there were any sex differences or what nutrients were examined in addition to MUFA. The findings from the present study seem to support the role of fat consumption in the accumulation of VAT.

In addition to dietary fat, there is a growing body of literature demonstrating a negative association between increased Ca consumption and adiposity. The impact of Ca on adiposity may be especially important for visceral adiposity, since visceral fat is more metabolically active and sensitive to lipolysis than other adipose tissue(1). It has been reported that increasing dietary Ca accelerates weight and fat loss in energy restricted obese adults, with preferential loss of trunk fat as measured by dual-energy X-ray absorptiometry(17). It has been further shown that elevated Ca consumption alters plasma 1,25-dihydroxyvitamin D<sub>3</sub>, which in turn alters local cortisol levels that may have an impact on the preferential loss of visceral adiposity(18).

The influence of Ca on VAT for men is consistent with the proposed mechanism. It is not clear why this same effect was not seen in women. In general, women have a lower VAT area and it seems to be less metabolically active than in men(1,9). This is specifically true for the population of women in the present study, who were all young adults and less prone to deposit fat in the viscera. It has also been observed in another study in women of a similar age that Ca consumption was related to total body fat in participants who consumed less than 7853 kJ, but not in women who consumed more than 7853kJ(19). Only six women in the present study consumed less than 7853 kJ/d and this could be one of the reasons why no relationship was observed in women. In addition, Ca consumption was 800 (SD 325) mg in women, which is below the recommended intake and was roughly 280 mg lower than men in the study. This level of Ca consumption may be too low to influence fat metabolism and the accumulation of VAT in these women.

There are some limitations that should be considered when interpreting the results of the present study. It should be pointed out that the data available for this investigation did not include MUFA or PUFA. These dietary components may have an impact on VAT and deserve future consideration. In addition, because of the cross-sectional nature of the present investigation, causation cannot be established and caution should be taken in drawing conclusions from the data presented. Finally, because the dietary assessment took place in the cafeteria it is impossible to be sure that dietary patterns remained consistent with patterns established under normal eating conditions. While this is a limitation, the dietary assessment methodology employed in the study directly measured food intake over 14d and results from fifty- two of the participants demonstrated that they were in energy balance during the dietary assessment period when compared with doubly labelled water(13).

## Conclusion

Total dietary fat is the best predictor of VAT area in both men and women. While this relationship was independent in women, in men there was a synergistic relationship between dietary fat consumption and Ca consumption in predicting VAT.

## Acknowledgments

The present study was supported by grant NIHDK49181 from the National Institute of Diabetes and Digestive and Kidney Diseases and by grant M01 RR0051 from the Clinical Research Center of the University of Colorado Health Sciences Center.

## References

1. Wajchenberg BL, Giannella-Neto D, da Silva ME, et al. Depot-specific hormonal characteristics of subcutaneous and visceral adipose tissue and their relation to the metabolic syndrome. *Horm Metab Res.* 2002; 34:616–621. [PubMed: 12660870]
2. Kelley DE, Thaete FL, Troost F, et al. Subdivisions of subcutaneous abdominal adipose tissue and insulin resistance. *Am J Physiol Endocrinol Metab.* 2000; 278:E941–E948. [PubMed: 10780952]
3. Hoffman DJ, Policastro P, Quick V, et al. Changes in body weight and fat mass of men and women in the first year of college: a study of the "freshman 15". *J Am Coll Health.* 2003; 55:41–45. [PubMed: 16889314]
4. Anderson DA, Shapiro JR, Lundgren JD. The freshman year of college as a critical period for weight gain: an initial evaluation. *Eat Behav.* 2003; 4:363–367. [PubMed: 15000962]
5. Katzmarzyk PT, Perusse L, Bouchard C. Genetics of abdominal visceral fat levels. *Am J Hum Biol.* 1999; 11:225–235. [PubMed: 11533946]
6. Lemieux I, Pascot A, Lamarche B, et al. Is the gender difference in LDL size explained by the metabolic complications of visceral obesity? *Eur J Clin Invest.* 2002; 32:909–917. [PubMed: 12534450]
7. Abe T, Sakurai T, Kurata J, et al. Subcutaneous and visceral fat distribution and daily physical activity: comparison between young and middle aged women. *Br J Sports Med.* 1996; 30:297–300. [PubMed: 9015589]
8. Riechman SE, Schoen RE, Weissfeld JL, et al. Association of physical activity and visceral adipose tissue in older women and men. *Obes Res.* 2002; 10:1065–1073. [PubMed: 12376588]
9. Bouchard C, Despres JP, Mauriege P. Genetic and nongenetic determinants of regional fat distribution. *Endocr Rev.* 1993; 14:72–93. [PubMed: 8491156]
10. Bouchard C, Tremblay A, Despres JP, et al. The response to long-term overfeeding in identical twins. *N Engl J Med.* 1990; 322:1477–1482. [PubMed: 2336074]
11. Donnelly JE, Hill JO, Jacobsen DJ, et al. Effects of a 16-month randomized controlled exercise trial on body weight and composition in young, overweight men and women: the Midwest Exercise Trial. *Arch Intern Med.* 2003; 163:1343–1350. [PubMed: 12796071]
12. Blair SN, Haskell WL, Ho P. Assessment of physical activity by a seven-day recall in a community survey and controlled experiments. *Am J Epidemiol.* 1985; 122:794–804. [PubMed: 3876763]
13. Hise ME, Sullivan DK, Jacobsen DJ, et al. Validation of energy intake measurements determined from observer-recorded food records and recall methods compared with the doubly labeled water method in overweight and obese individuals. *Am J Clin Nutr.* 2002; 75:263–267. [PubMed: 11815316]
14. Lissner L, Heitmann BL. Dietary-fat and obesity - evidence from epidemiology. *Eur J Clin Nutr.* 1995; 49:79–90. [PubMed: 7743988]
15. Greenfield JR, Samaras K, Jenkins AB, et al. Moderate alcohol consumption, dietary fat composition, and abdominal obesity in women: evidence for gene-environment interaction. *J Clin Endocrinol Metab.* 2003; 88:5381–5386. [PubMed: 14602777]
16. Dutia R, Wroblewski P, Bacher A, et al. Correlation between dietary fat intake and adipose tissue distribution. *J Am Diet Assoc.* 2005; 105(Suppl. 1):9.
17. Zemel MB, Thompson W, Milstead A, et al. Calcium and dairy acceleration of weight and fat loss during energy restriction in obese adults. *Obes Res.* 2004; 12:582–590. [PubMed: 15090625]
18. Morris KL, Zemel MB. 1,25-Dihydroxyvitamin D3 modulation of adipocyte glucocorticoid function. *Obes Res.* 2005; 13:670–677. [PubMed: 15897475]
19. Lin YC, Lyle RM, McCabe LD, et al. Dairy calcium is related to changes in body composition during a two-year exercise intervention in young women. *J Am Coll Nutr.* 2000; 19:754–760. [PubMed: 11194528]

**Table 1**

Demographic information by sex (Mean values and standard deviations)

	Female (n 57)		Male (n 49)	
	Mean	SD	Mean	SD
Age (years)	22.5	4.7	23.0	4.0
Body weight (kg)	78.7	10.3	96.0*	11.8
Body fat(%)	35.9	4.7	27.6*	4.4
Visceral adipose tissue area (cm <sup>2</sup> )	64.2	24.7	100.3*	38.8
Subcutaneous adipose tissue area (cm <sup>2</sup> )	384.1	104.3	323.2*	100.0
Daily energy consumption (kJ)	10324	1789	13507*	2910
Total dietary fat (g)	86.1	23.3	124.1*	36.4
Total protein (g)	76.7	16.2	108.6*	24.9
Total carbohydrate (g)	355.4	66.2	411.7*	94.2
Total dietary Ca (mg)	800	325	1080*	440

\* Mean value was significantly different from that of the females ( $P$  0.05).

**Table 2**Pearson correlations (*r*) of nutrients and visceral adiposity

	<u>Visceral adipose tissue</u>		<u>Subcutaneous adipose tissue</u>	
	Women	Men	Women	Men
Energy (kJ)	0.25	0.14	0.36*	0.09
Total dietary fat (g)	0.36*	0.21	0.48*	0.20
Total saturated fat (g)	0.32*	0.18	0.40*	0.20
Total protein (g)	0.15	0.13	0.11	0.18
Total carbohydrate (g)	0.11	0.06	0.17	-0.02
Total dietary Ca (mg)	0.02	-0.07	0.03	0.15
Ca:protein ratio	-0.09	-0.22	-0.07	0.11

\* *P* 0.05.

**Table 3**

Stepwise summary table: dietary predictors of visceral and subcutaneous fat in women

	<u>Visceral</u>		<u>Subcutaneous</u>	
	T-statistic	P	T-statistic	P
Total dietary fat	2.89	<0.01	2.68	0.01
Folic acid	-1.25 <sup>†</sup>	0.22	-0.87 <sup>†</sup>	0.39
Total dietary fibre	-1.01 <sup>†</sup>	0.31	-0.02 <sup>†</sup>	0.98
Fe	-0.88 <sup>†</sup>	0.38	-0.96 <sup>†</sup>	0.34
Ca:protein ratio	-0.72 <sup>†</sup>	0.47	-0.08 <sup>†</sup>	0.94
Vitamin A	-0.69 <sup>†</sup>	0.50	-1.38 <sup>†</sup>	0.17
Total energy intake	-0.63 <sup>†</sup>	0.53	-0.54 <sup>†</sup>	0.59
Total dietary carbohydrate	-0.49 <sup>†</sup>	0.62	-0.46 <sup>†</sup>	0.65
Ca	-0.38 <sup>†</sup>	0.70	-0.49 <sup>†</sup>	0.62
Alcohol	0.27 <sup>†</sup>	0.78	0.53 <sup>†</sup>	0.60
Total dietary protein	-0.26 <sup>†</sup>	0.80	-1.27 <sup>†</sup>	0.21
Saturated fat	-0.08 <sup>†</sup>	0.93	-0.70 <sup>†</sup>	0.49

<sup>†</sup>Variables that did not enter the model with dietary fat.



**Table 4**

Stepwise summary table: dietary predictors of visceral and subcutaneous fat in men

	<u>Visceral</u>		<u>Subcutaneous</u>	
	T-statistic	P	T-statistic	P
Total dietary fat	2.68	0.01	1.51 <sup>†</sup>	0.14
Ca	-2.33	0.02	-0.46 <sup>†</sup>	0.65
Total dietary fibre	0.92 <sup>†</sup>	0.36	0.51 <sup>†</sup>	0.61
Fe	-0.79 <sup>†</sup>	0.43	-0.41 <sup>†</sup>	0.69
Vitamin A	0.59 <sup>†</sup>	0.55	0.13 <sup>†</sup>	0.90
Total dietary protein	0.55 <sup>†</sup>	0.59	1.30 <sup>†</sup>	0.20
Folic acid	0.32 <sup>†</sup>	0.75	0.56 <sup>†</sup>	0.58
Saturated fat	0.34 <sup>†</sup>	0.74	1.40 <sup>†</sup>	0.16
Alcohol	-0.21 <sup>†</sup>	0.84	-0.91 <sup>†</sup>	0.37
Total dietary carbohydrate	-0.14 <sup>†</sup>	0.89	-0.12 <sup>†</sup>	0.90
Ca:protein ratio	0.13 <sup>†</sup>	0.90	0.75 <sup>†</sup>	0.46
Total energy intake	-0.00 <sup>†</sup>	0.99	0.63 <sup>†</sup>	0.53

<sup>†</sup>Variables that did not enter the model with total dietary fat and Ca.