



NIH PUBLIC ACCESS


## Author Manuscript

*J Am Chem Soc.* Author manuscript; available in PMC 2011 September 8.

Published in final edited form as:


*J Am Chem Soc.* 2010 September 8; 132(35): 12179–12181. doi:10.1021/ja104196x.**Stereospecific decarboxylative allylation of sulfones**

**Jimmie D. Weaver, Being J. Ka, David K. Morris, Ward Thompson, and Jon A. Tunge**  
 Department of Chemistry, The University of Kansas, 2010 Malott Hall, 1251 Wescoe Hall Drive,  
 Lawrence, KS 66045

Jon A. Tunge: [tunge@ku.edu](mailto:tunge@ku.edu)**Abstract**


Allyl sulfonyl acetic esters undergo highly stereospecific, palladium-catalyzed decarboxylative allylation. The reaction allows the stereospecific formation of tertiary homoallylic sulfones in high yield. In contrast to related reactions that proceed at  $-100\text{ }^{\circ}\text{C}$  and require highly basic preformed organometallics, the decarboxylative coupling described herein occurs under mild non-basic conditions and requires no stoichiometric additives. Allylation of the intermediate  $\alpha$ -sulfonyl anion is more rapid than racemization, leading to a highly enantiospecific process. DFT calculations indicate that the barrier for racemization is 9.9 kcal/mol and thus the barrier of allylation must be  $<9.9$  kcal/mol.

Catalytic decarboxylative coupling is a powerful method for C–C bond formation that avoids the use of strong bases and/or preformed organometallics that are typically required for cross-coupling reactions. In 2004, we reported the first asymmetric decarboxylative allylation of enolates.<sup>1</sup> This was followed by several reports of decarboxylative enolate allylations that allow enantioselective formation of quaternary carbon centers  $\alpha$ -to ketones.<sup>2</sup> Since these reactions proceed through achiral enolate intermediates the reactions are stereoconvergent, allowing racemic starting materials to form highly enantioenriched products (eq. 1). Stereoconvergence is a common reaction profile in carbanion chemistry, where resonance-stabilized carbanions often exist as planar achiral intermediates and unstabilized carbanions rapidly racemize by inversion.<sup>3</sup> Herein we report that optically active sulfonyl acetic esters undergo enantiospecific decarboxylative coupling, where the original stereochemistry is maintained in the product (eq. 2). Furthermore, DFT calculations are used to examine the origin of the enantiospecificity.




(1)

Correspondence to: Jon A. Tunge, [tunge@ku.edu](mailto:tunge@ku.edu).Supporting Information Available: Experimental procedures and characterization data for all new compounds (PDF). This material is available free of charge via the Internet at <http://pubs.acs.org>.



From the elegant work of Corey and Cram in the early 1960's, it was known that the decarboxylative protonation of sulfonyl acetic esters is stereospecific.<sup>4</sup> While there are several examples of stereospecific decarboxylative protonations,<sup>4,5</sup> none has been extended to C–C bond forming reactions. With this in mind, we set out to develop a stereospecific decarboxylative C–C bond forming reaction.

To begin, sulfone **1a** was synthesized and treated with 2 mol% Pd(PPh<sub>3</sub>)<sub>4</sub> in toluene at room temperature (eq. 3).<sup>6</sup> It was gratifying to find that the decarboxylative allylation was highly stereospecific, with a conservation of enantiomeric excess (cee) of 96% (cee = 100 × (product ee)/(reactant ee)).



Next, a variety of different enantioenriched sulfones were similarly subjected to conditions for palladium-catalyzed decarboxylative coupling. As can be seen from the data in table 1, a variety of  $\alpha$ -aryl and  $\alpha,\alpha$ -dialkyl sulfones all undergo decarboxylative allylation with a high degree of stereospecificity.

It is noteworthy that high cee's are obtained even at elevated temperatures (95 °C). Moreover, our reaction proceeds under formally neutral conditions, in contrast to more classical alkylations of sulfones which typically require stoichiometric, highly basic, organolithium reagents.<sup>7</sup> Lastly, an X-ray crystal structure of a derivative of **2k** confirms that the allylation proceeds with overall retention of configuration.<sup>8</sup>

The overall retention of stereochemistry in the decarboxylative allylation can be explained if decarboxylation directly produces either a configurationally stable alkyl palladium complex that undergoes reductive elimination (path 1, Scheme 1) or if decarboxylation produces a configurationally stable  $\alpha$ -sulfonyl anion that attacks the palladium  $\pi$ -allyl complex (path 2, Scheme 1).<sup>9</sup>


To begin addressing whether path 1 or 2 (Scheme 1) is operative, we investigated the effect of palladium on the rates of decarboxylation of several  $\alpha$ -sulfonyl acetates. Control studies show that the rate of decarboxylation is relatively unaffected by the addition of a variety of palladium sources (Table 2).<sup>8</sup> For example, the cesium or triethylammonium salts of  $\alpha$ -sulfonyl acetic acids readily decarboxylate under our reaction conditions. Addition of Pd(OAc)<sub>2</sub> or (Ph<sub>3</sub>P)<sub>2</sub>Pd(allyl)Cl to these reaction mixtures does not substantially affect the rate of decarboxylation.<sup>10</sup> When palladium is directly involved in the decarboxylation, such experiments usually show a dramatic acceleration of the decarboxylation.<sup>11</sup> Ultimately, Corey and Cram have detailed thermal decarboxylations of  $\alpha$ -sulfonyl acetates under similar conditions to those used by us,<sup>4</sup> and we have failed to observe catalysis of decarboxylation

by palladium. Thus, the simplest conclusion is that decarboxylation of the  $\alpha$ -sulfonyl carboxylates is a thermal process that proceeds via path 2.

Reaction via path 1 and path 2 can also be distinguished via the observed diastereoselectivity in nucleophilic attack of a stereochemically labeled allyl ester substrate (Scheme 2).<sup>10</sup> In such a reaction, nucleophiles that are bound to palladium prior to reductive elimination, as in path 1, react with overall inversion of configuration. Alternatively, if the free nucleophile directly attacks the allyl ligand, as in path 2, then one observes retention of stereochemistry by way of the standard double-inversion mechanism.<sup>12</sup> Toward this end, substrate **1n** was prepared and allowed to react under our standard reaction conditions (Scheme 2).<sup>13</sup> The reaction produced the 1,3-*cis* diastereomers as the only C–C bond forming product in 49% yield,<sup>14</sup> suggesting that the decarboxylative coupling proceeds via attack of an  $\alpha$ -sulfonyl anion on the backside of palladium  $\pi$ -allyl complex (path 2, Scheme 2). While we could not generate the  $\alpha$ -chloro- $\alpha$ -sulfonyl ester in enantioenriched form in order to test the enantiospecificity of the coupling, the product is formed as a 1:1 mixture of diastereomers at the sulfonyl stereocenter. Thus, the stereochemistry of the  $\alpha$ -position did not change during the reaction, which is consistent with a stereospecific process.

Our control studies and stereochemical studies both suggest that  $\alpha$ -sulfonyl anions are formed, and react, outside the coordination sphere of palladium. In order to form highly enantioenriched products, these intermediate  $\alpha$ -sulfonyl anions must be configurationally stable. While it is known that  $\alpha$ -sulfonyl anions are more configurationally stable than typical carbanions, Gais has reported the rapid racemization of anion **3a** at  $-80$  °C.<sup>15</sup> Thus, we became curious why our reaction was highly stereospecific, even at elevated temperatures.

To address the reasons for the apparently slower rate of racemization than coupling of the  $\alpha$ -sulfonyl anion and Pd- $\pi$ -allyl, DFT calculations were performed to determine the inversion barrier for the free  $\alpha$ -sulfonyl carbanion. The calculations revealed that the most stable conformation is **3a**, which has the major lobe of the lone-pair oriented anti-periplanar to the S–Ph bond (eq. 4). A definitive number for the inversion barrier could not be determined because all attempts to minimize the energy of conformation **3b** led to convergence to the minimum energy structure (**3a**). While we could not obtain a specific energy for structure **3b**, this result does support the conclusion that the barrier to inversion is small ( $< 2$  kcal/mol). The small inversion barrier is supported by other studies, as is the slightly pyramidal structure of the carbanion.<sup>15–17</sup>



(4)

Taking into account the small barrier to inversion, our mechanistic hypothesis is as follows. Ionization and decarboxylation of **1a** generates sulfonyl anion, **3a** which is likely ion-paired with the Pd- $\pi$ -allyl (Scheme 3). The C–C formation proceeds via path A, where the carbanion can react through the low-energy, staggered, transition state to form the product with retention of stereochemistry. Coupling through the inverted conformer, **3b** (path B) to form the enantiomeric product not only requires the anion to react through a higher energy conformer, but also requires a higher energy transition state to form the eclipsed product.<sup>4</sup> Alternatively, the enantiomer, *ent*-**2a**, could be formed by allylation of the anion that has

undergone inversion and rotation (path C). Thus, we hypothesized that the reaction is stereospecific because rotation about the C–S bond of the sulfone (path C) is slower than reaction of the  $\alpha$ -sulfonyl anion, **3a**, with the palladium  $\pi$ -allyl complex through path A.

DFT calculations carried out at the B3LYP/6-31+G\* level of theory reveal that the lowest barrier to rotation about the  $\alpha$ C–S bond is 9.9 kcal/mol, which is consistent with experimental measurements in related systems (Scheme 4).<sup>15–17</sup> Thus, the barrier for allylation of the  $\alpha$ -sulfonyl anion must be < 9.9 kcal/mol or rotation would lead to racemization. Such a low barrier is consistent with the reaction between an ion-pair of a highly nucleophilic  $\alpha$ -sulfonyl anion and a palladium  $\pi$ -allyl complex.<sup>8,18</sup> Thus, the stark contrast in our observations and that of Gais can be attributed to the fact that we are generating the reactive  $\alpha$ -sulfonyl anion intermediates in the presence of highly electrophilic  $\pi$ -allyl palladium electrophiles, allowing us to effect the C–C bond formation faster than rotation.

In conclusion, we have developed a stereospecific decarboxylative coupling reaction that provides access to enantioenriched homoallylic sulfones. The enantiospecific nature of this reaction is unique among, yet complimentary to, existing decarboxylative allylation methodologies. The stereospecificity is made possible by the relatively high barrier for racemization of axially chiral  $\alpha$ -sulfonyl anions and the low barrier for their reaction with palladium  $\pi$ -allyl complexes.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## Acknowledgments


We thank the National Institute of General Medical Sciences (1R01GM079644) for financial support. We thank Dr. Victor Day for X-ray crystallographic analysis.


## References

1. Burger EC, Tunge JA. *Org Lett.* 2004; 6:4113. [PubMed: 15496112]
2. (a) Trost BM, Xu J. *J Am Chem Soc.* 2005; 127:17180. [PubMed: 16332054] (b) Burger EC, Barron BR, Tunge JA. *Synlett.* 2006:2824. (c) Trost BM, Bream RN, Xu J. *Angew Chem, Int Ed.* 2006; 45:3109. (d) You SL, Dai LX. *Angew Chem, Int Ed.* 2006; 45:5246. (e) Trost BM, Xu J, Schmidt T. *J Am Chem Soc.* 2009; 131:18343. [PubMed: 19928805] (f) Behenna DC, Stoltz BM. *J Am Chem Soc.* 2004; 126:15044. [PubMed: 15547998] (g) Nakamura M, Hajra A, Endo K, Nakamura E. *Angew Chem, Int Ed.* 2005; 44:7248. (h) Mohr JT, Behenna DC, Harned AM, Stoltz BM. *Angew Chem, Int Ed.* 2005; 44:6924.
3. Baechler RD, Andose JD, Stackhouse J, Mislow K. *J Am Chem Soc.* 1972; 94:8060.
4. (a) Corey EJ, Kaiser ET. *J Am Chem Soc.* 1961; 83:490. (b) Corey EJ, Koenig H, Lowry TH. *Tetrahedron Lett.* 1962:515. (c) Corey EJ, Lowry TH. *Tetrahedron Lett.* 1965:803. (d) Corey EJ, Lowry TH. *Tetrahedron Lett.* 1965:793. (e) Cram DJ, Nielsen WD, Rickborn B. *J Am Chem Soc.* 1960; 82:6415. (f) Cram DJ, Rickborn B, Knox GR. *J Am Chem Soc.* 1960; 82:6412. (g) Cram DJ, Wingrove AS. *J Am Chem Soc.* 1963; 85:1100.
5. Rozov LA, Rafalko PW, Evans SM, Brockunier L, Ramig K. *J Org Chem.* 1995; 60:1319.
6. (a) Weaver JD, Morris DK, Tunge JA. *Synlett.* 2010:470. [PubMed: 20526414] (b) Weaver JD, Tunge JA. *Org Lett.* 2008; 10:4657. [PubMed: 18785744]
7. Gais HJ, Hellmann G. *J Am Chem Soc.* 1992; 114:4439.
8. See supporting information for more details.
9. Complete crossover is observed when two sulfonyl esters are allowed to undergo coupling in the same flask. The interpretation of complete crossover is ambiguous since crossover can likely occur

via either the intermediate  $\pi$ -allyl palladium carboxylates or the  $\pi$ -allyl palladium  $\alpha$ -sulfonyl anion complexes.

10. The presence of acidic protons from the carboxylic acid results in the formation of the protonation product rather than the allylation product that may be expected from  $(\text{allyl})\text{PdCl}(\text{PPh}_3)_2$
11. Recio A, Tunge JA. *Org Lett.* 2009; 11:5630. [PubMed: 19921827]
12. (a) Trost BM, Verhoeven TR. *J Am Chem Soc.* 1980; 102:4730. (b) Keinan E, Roth Z. *J Org Chem.* 1983; 48:1770.
13. Unfortunately, the  $\alpha$ -methyl, $\alpha$ -phenyl sulfonyl acetic ester of cyclohexen-3-ol gives 78% elimination product. Thus, it could not be used for the stereochemical test. Elimination is less problematic with an  $\alpha$ -chloro substituent see footnote 6b.
14. Analysis of the crude reaction mixture by  $^1\text{H}$  NMR spectroscopy indicates that the mass balance of the decarboxylative allylation of **1n** is made up of elimination products.
15. Gais HJ, Hellmann G, Gunther H, Lopez F, Lindner HJ, Braun S. *Angew Chem Int Ed.* 1989; 28:1025.
16. (a) Raabe G, Gais HJ, Fleischhauer J. *J Am Chem Soc.* 1996; 118:4622. (b) Reetz MT, Hutte S, Goddard R. *Eur J Org Chem.* 1999:2475.
17. This barrier to rotation is, in part, due to loss of negative hyperconjugation with the Ph-S  $\sigma^*$  orbital. Koch R, Anders E. *J Org Chem.* 1994; 59:4529.
18. (a) Seeliger F, Mayr H. *Org Biomol Chem.* 2008; 6:3052. [PubMed: 18698462] (b) Kuhn O, Mayr H. *Angew Chem Int Ed.* 1999; 38:343.

**Scheme 1.**



Scheme 2.

**Scheme 3.**
















**Scheme 4.**

Table 1

Stereospecificities of decarboxylation allylation

entry	reactant ee (%) <sup>c</sup>	product	procedure <sup>a</sup>	yield (%) <sup>b</sup>	ee (%) <sup>c</sup>	cee (%) <sup>c</sup>
1	46	 <b>2b</b>	A	96	43	91
2	89	 <b>2c</b>	A	99	87	98
3	97	 <b>2d</b>	A	99	96	99
4	61	 <b>2e</b>	A	99	61	99 <sup>d</sup>
5	73	 <b>2f</b>	A	99	69	96 <sup>d</sup>
6	97	 <b>2g</b>	A	99	93	96
7	80	 <b>2h</b>	A	99	80	99

entry	reactant ee(%) <sup>c</sup>	product	procedure <sup>a</sup>	yield (%) <sup>b</sup>	ee(%) <sup>c</sup>	cee (%) <sup>c</sup>
8	94	 <b>2i</b>	B	82	92	99
9	94	 <b>2j</b>	B	95	92	98
10	>99	 <b>2k</b>	B	97	>99	>99
11	98	 <b>2l</b>	B	85	95	97
12	98	 <b>2m</b>	B	93	95	97

<sup>a</sup> Conditions A: 2 mol % Pd(PPh<sub>3</sub>)<sub>4</sub>, 0.2 M toluene, 23 °C, 0.25-2 h. Conditions B: 5 mol % Pd<sub>2</sub>dba<sub>3</sub>, 10 mol % (±)-binap, 0.2 M toluene, 95 °C, 11-15 h.

<sup>b</sup> isolated yields.

<sup>c</sup> determined via chiral stationary phase HPLC analysis

<sup>d</sup> 8: 1 linear:branched

Table 2

<b>R</b>	<b>R'</b>	<b>conditions</b>	<b>Pd source</b>	<b>time</b>	<b>% conv.</b>
H	H	95 °C, Et <sub>3</sub> N toluene-d <sub>6</sub>	none	5 min.	17%
H	H	95 °C, Et <sub>3</sub> N	Pd(allyl)Cl(PPh <sub>3</sub> ) <sub>2</sub> 100 mol%	5 min.	23%
Me	Bn	95 °C, Et <sub>3</sub> N	none	45 min.	25%
Me	Bn	95 °C, Et <sub>3</sub> N	Pd(OAc) <sub>2</sub> 10 mol%	45 min.	27%
Me	H	23 °C, Cs <sub>2</sub> CO <sub>3</sub>	none	36 h	59%
Me	H	23 °C, Cs <sub>2</sub> CO <sub>3</sub>	Pd(OAc) <sub>2</sub> 10 mol%	36 h	59%