



2-2019

Direct Anterior Approach Utilizing a Bikini Incision has Less Wound Related Complications in Patients with High BMI

Taylor Paskey, BS

Thomas Jefferson University, taylor.paskey@jefferson.edu

Jorge Manrique, MD

Thomas Jefferson University, jorge.manrique@jefferson.edu

Majd Tarabichi, MD

Thomas Jefferson University

Camilo Restrepo, MD

Thomas Jefferson University, camilo.restrepo@jefferson.edu

Carol Foltz, PhD

*Thomas Jefferson University, caf012@jefferson.edu**See next page for additional authors*

[Let us know how access to this document benefits you](#)

Follow this and additional works at: https://jdc.jefferson.edu/si_ctr_2021_phase1 Part of the [Surgery Commons](#)

Recommended Citation

Paskey, Taylor; Manrique, Jorge; Tarabichi, Majd; Restrepo, Camilo; Foltz, Carol; Hozack, William J., "Direct Anterior Approach Utilizing a Bikini Incision has Less Wound Related Complications in Patients with High BMI" (2019). SKMC JeffMD Scholarly Inquiry, Phase 1, Project 1.

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Phase 1 by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.

Authors

Taylor Paskey, BS; Jorge Manrique, MD; Majd Tarabichi, MD; Camilo Restrepo, MD; Carol Foltz, PhD; and William J. Hozack, MD

Taylor Paskey
SKMC Class of 2021
SI CTR Abstract
12/15/18

Direct Anterior Approach Utilizing a Bikini Incision has Less Wound Related Complications in Patients with High BMI

*Jorge Manrique MD, Taylor Paskey BS, Majd Tarabichi MD, Camilo Restrepo MD, Carol Foltz
PhD, William J. Hozack MD*

Introduction

Direct anterior approach (DAA) total hip arthroplasty (THA) can be performed through a traditional vertical skin incision, situating the proximal incision at the hip flexion crease, or a horizontal (bikini) skin incision, situating the incision slightly distal and parallel to the hip flexion crease. The dissection beyond the subcutaneous layer is identical for both methods.

Objective

The purpose of this study was to compare these approaches, performed by an experienced single surgeon, in terms of overall wound complications and patient-reported esthetics 6-months post-operatively. It was hypothesized that the bikini incision would result in less wound complications and improved cosmesis due to decreased applied tension from the hip flexion crease.

Methods

A case-control retrospective study was conducted and 86 bikini DAA patients were matched 3:1 to 230 conventional DAA patients for gender, age, body mass index (BMI), and American Society of Anesthesiologists score. The outcomes evaluated included wound

complications, acute periprosthetic joint infection, transfusion, length of surgery, and dysesthesia with an additional subset analysis for obese patients (BMI >30kg/m²). Patients rated incision cosmesis 6 months post-operatively using a Patient Scar Assessment Scale and the Vancouver scar assessment scale.

Results

Bikini patients had lower rates of delayed wound healing compared to conventional incision (2.3% vs. 6.1%; p=0.087). This difference was statistically significant (0% vs. 16.6%; p<0.05) in obese patients with no difference in incision cosmesis in either analysis.

Discussion

The bikini incision could offer safety benefits in selected patients (BMI >30kg/m²) undergoing DAA THA by decreasing wound complications while preserving cosmesis.