



2-2019

Network Influence of the Cerebellum for Predicting DBS Response in Patients with Advanced Parkinson's Disease

Lucy Li

Thomas Jefferson University, lucy.li@jefferson.edu

Jennifer Muller

Thomas Jefferson University, jennifer.muller001@jefferson.edu

Sara Thalheimer

Thomas Jefferson University, sara.thalheimer@jefferson.edu

Mackenzie D. Silverman

Thomas Jefferson University, mackenzie.silverman@jefferson.edu

Mahdi Alizadeh

*Thomas Jefferson University, mahdi.alizadeh.2@jefferson.edu**See next page for additional authors*

[Let us know how access to this document benefits you](#)

Follow this and additional works at: https://jdc.jefferson.edu/si_ctr_2021_phase1 Part of the [Neurology Commons](#), and the [Radiology Commons](#)

Recommended Citation

Li, Lucy; Muller, Jennifer; Thalheimer, Sara; Silverman, Mackenzie D.; Alizadeh, Mahdi; Liang, Tsao-Wei; Layton, Kelly; Kremens, Daniel; Romo, Victor; Mohamed, Feroze B.; Wu, Chengyuan, "Network Influence of the Cerebellum for Predicting DBS Response in Patients with Advanced Parkinson's Disease" (2019). SKMC JeffMD Scholarly Inquiry, Phase 1, Project 1.

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Phase 1 by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.

Authors

Lucy Li, Jennifer Muller, Sara Thalheimer, Mackenzie D. Silverman, Mahdi Alizadeh, Tsao-Wei Liang, Kelly Layton, Daniel Kremens, Victor Romo, Feroze Mohamed, and Chengyuan Wu

Lucy Li
SKMC Class of 2021
SI CTR Abstract
12.15.2018

Network Influence of the Cerebellum for Predicting DBS Response in Patients with Advanced Parkinson's Disease

Jennifer Muller^{1,2}, Lucy Li^{1,2}, Sara Thalheimer^{1,2}, Mackenzie D Silverman^{1,2}, Mahdi Alizadeh^{1,2},
Tsao-Wei Liang⁵, Kelly Layton⁵, Daniel Kremens⁵, Victor Romo⁴, Feroze B. Mohamed¹,
Chengyuan Wu^{1,2}

¹Jefferson Integrated Magnetic Resonance Imaging Center, Department of Radiology, Thomas
Jefferson University, Philadelphia, PA.

²Department of Neurosurgery, Thomas Jefferson University, Philadelphia, PA.

⁴Department of Anesthesiology, Thomas Jefferson University, Philadelphia, PA.

⁵Department of Neurology, Thomas Jefferson University, Philadelphia, PA.

Introduction: Deep brain stimulation (DBS) is a treatment option for reducing motor symptoms in patients with Parkinson's Disease (PD) when first-line medication becomes ineffective.

Existing literature has hypothesized that the clinical outcome of DBS may depend on brain connectivity profiles of the stimulation site to distant brain regions. However, the potential of brain connectivity profiles to predict response to DBS in PD remains unclear.

Objective: This study aimed to investigate how changes in structural and functional connectivity may relate to patient response to DBS, through the examination of brain network changes using graph theory.

Methods: Ten patients with advanced PD were included in this study. Diffusion tensor imaging (DTI) and resting-state fMRI scans were acquired prior to DBS implantation. Pre-DBS and post-DBS UPDRS-III scores were obtained. Network analysis of the DTI and fMRI scans was performed using GRETNA to compute structural and functional graph theory metrics, respectively. Metrics were correlated with UPDRS-III improvement to identify significant correlations to UPDRS improvement due to DBS.

Results: Combined structural and functional graph theory metrics highlighted 32 structures to be significantly correlated with UPDRS-III improvement. Mainly, connections to the cerebellum were found to be significantly correlated with UPDRS-III improvement across several metrics for both structural and functional connectivity.

Discussion: This work combined DTI, fMRI, and graph theory analysis to evaluate improvement with DBS. Several imaging biomarkers were identified that are robust predictors for UPDRS-III improvement. This work warrants investigation into the compensatory effect of the cerebellum and other potential biomarkers for identifying DBS candidates.