

**HHS PUBLIC ACCESS**

Author manuscript

J Child Lang. Author manuscript; available in PMC 2015 December 26.

Published in final edited form as:

J Child Lang. 2015 September ; 42(5): 1036–1072. doi:10.1017/S0305000914000701.

Dense Neighborhoods and Mechanisms of Learning: Evidence from Children with Phonological Delay

Judith A. Gierut and Michele L. Morrisette

Indiana University, Bloomington

Abstract

There is a noted advantage of dense neighborhoods in language acquisition, but the learning mechanism that drives the effect is not well understood. Two hypotheses—long-term auditory word priming and phonological working memory—have been advanced in the literature as viable accounts. These were evaluated in two treatment studies enrolling 12 children with phonological delay. Study 1 exposed children to dense neighbors versus nonneighbors before training sound production in evaluation of the priming hypothesis. Study 2 exposed children to the same stimuli after training sound production as a test of the phonological working memory hypothesis. Results showed that neighbors led to greater phonological generalization than nonneighbors, but only when presented prior to training production. There was little generalization and no differential effect of exposure to neighbors or nonneighbors after training production. Priming was thus supported as a possible mechanism of learning behind the dense neighborhood advantage in phonological acquisition.

INTRODUCTION

Classic research on children's acquisition of phonology has advocated for "the primacy of lexical items in phonological development" (Ferguson & Farwell, 1975: 437). In this context and as used herein, phonology refers to the phones, phonemes, phonotactics, and rules of the sound system of the input language as realized in production. The early thought was that children's mastery of the sound system followed from their acquisition of words. Recently, this view has received renewed attention due, in part, to insights about the developing mental lexicon and its organization into neighborhoods. A neighborhood consists of words that differ by 1-phoneme substitutions, deletions, or additions, e.g., *rat* has neighbors *cat*, *wrap*, *wrote*, *at*, *drat* among others (Luce, 1986). Neighborhoods are described as DENSE when comprised of many words that overlap in phonological form and as SPARSE when comprised of few words that overlap in phonological form. Empirically, the distinction between dense and sparse neighborhoods has been shown to differentially affect phonological acquisition by children with typical (e.g., Stoel-Gammon, 2011) and delayed (e.g., Gierut & Morrisette, 2012b) development. In this paper, we add to this line of investigation by exploring hypotheses about the mechanism of learning behind the observed neighborhood effects specifically for children with phonological delay (PD).

By way of background, we begin with a review of neighborhood effects in typical development and consider two hypotheses advanced in the literature to account for the data: one appeals to long-term auditory word priming and the other, to phonological working memory. We then summarize the findings of Demke, Graham, and Siakaluk (2002) as a specific test of these proposals. The relevance of that research for children with PD is outlined as motivation for the present studies.

TYPICAL PHONOLOGICAL DEVELOPMENT

Stoel-Gammon (2011) published an exhaustive review with commentary on the relationship between phonological acquisition and the lexicon. A key observation to emerge was that children's mastery of phonology is aided by dense neighborhood structure: Similar sounding words in the lexicon are aligned with better phonological skills. To illustrate, Sosa & Stoel-Gammon (2012) reported that toddlers produce sounds with greater accuracy and less variability when these occur in words from dense (as opposed to sparse) neighborhoods. Comparable effects have been reported for nonwords: Toddlers and preschoolers imitate sounds with greater accuracy when nonwords resemble real words from dense neighborhoods (Beckman & Edwards, 2000; Zamuner, Gerken & Hammond, 2004). Dense neighborhoods not only facilitate production accuracy, but presumably, they are the basis from which children extract generalizations about the inherent phonological structure of a language (Pierrehumbert, 2001). Dense neighborhoods support the broad extension, transfer, and representation of sound patterns across lexical items and contexts (Beckman, Munson & Edwards, 2007; Edwards, Beckman & Munson, 2004). Evidence for the dense neighborhood advantage appears to be robust, with convergence of descriptive phonetic (Sosa & Stoel-Gammon, 2012), acoustic phonetic (Munson, 2001), computational (Storkel 2004a), experimental (Storkel, 2002), metalinguistic (De Cara & Goswami, 2003) and cross-linguistic (Stokes, 2012) studies.

In account of the observed effects, Stoel-Gammon (2011), like others (Walley, 1993), suggested that dense neighborhoods trigger the emergence of segmental structure and phonemic contrast in words that start out as holistic or fuzzy in phonological makeup. Because dense neighborhoods consist of many words that overlap in phonological form, this forces a child to build finer-grained phonological representations as reflected behaviorally in their accurate production of new sounds and new contrasts. Notice that this account describes the kinds of phonological gains to result from dense neighborhoods, i.e., new phonemic distinctions; yet, it does not also describe HOW those changes take place. Namely, what is the learning mechanism responsible for enriched phonological structure that arises from dense neighborhoods?

TWO COMPLEMENTARY HYPOTHESES

Church & Fisher (1998) proposed that a child's routine exposure to words of the input acts like a naturalistic case of long-term auditory word priming, which is defined as the presentation of experimental stimuli similar to a set of test stimuli for the purpose of facilitating a behavioral response (Zwitserslood, 1997). Motivation for their proposal came from the observation that priming affords a child many of the same opportunities that are needed in the construction of phonological representations. Priming bolsters the mapping of

sound patterns of words, provides input (contextual) variability, emphasizes similarity of phonological form, and facilitates the differentiation of phonological form among neighbors. The hunch was that priming serves a three-fold purpose: it helps a child encode new words, builds the representation of the sound patterns of those words, and serves as the “link between repeated perception and production of particular sound patterns and subsequent processing of identical or similar items” (Church & Fisher, 1998: 538). In a series of studies, Church & Fisher (1998) demonstrated that children as young as age 2 more accurately identified words that were primed compared to words that were not. The effects were independent of word meaning, attributed instead to the internal representation of sound patterns. Overall, their results were consistent with the general developmental view that priming creates a precise “linguistic experience contributing to learning, changing (strengthening, expanding) linguistic representations” (Savage, Lieven, Theakston & Tomasello, 2006: 29).

Demke and colleagues (2002; also Merriman & Marazita, 1995) carried the hypothesis a step further by employing auditory word priming as a tool for lexical learning. They reported two studies, which are detailed herein as the springboard for the present research. In one study, preschoolers were exposed to dense forms prior to training novel lexical items. This was the pre-exposure condition characteristic of priming. In a second study, preschoolers were exposed to dense forms after training novel items. This was the post-exposure condition because, extending Church & Fisher’s (1998) reasoning, similar-sounding words in the input can occur before or after a child’s exposure to novel items.

Within each study, Demke et al. (2002) manipulated the kind of dense forms that were presented. In one manipulation, dense forms were neighbors of the novel items to be learned in that they shared the same rhyme; e.g., *plane*, *rain*, *train* were affiliated with the novel item *tane*. In a second manipulation, dense forms were phonologically unrelated to the novel items in that there was no rhyme overlap; hence, they were nonneighbors. Thus, neighbor status (neighbor vs. nonneighbor) and timing of exposure (before vs. after training) were the independent variables.

The instructional protocol developed by Demke et al. (2002) consisted of series of story vignettes, with one story for each novel item that was trained. Stories provided a child with multiple exposures to neighbors or nonneighbors depending on experimental assignment. These were visually depicted and a child viewed the pictures while hearing the stories live voice. In training, the examiner labeled visual referents corresponding to the novel items and a child was to imitate. Then, in a delayed test phase, a child recalled the novel items as the dependent variable.

Demke et al. (2002) found no difference in recall of novel items when children were exposed to neighbors versus nonneighbors in the pre-exposure condition; this was at odds with the predicted effect of priming. By comparison, there was greater recall of novel items in the post-exposure condition, but only when neighbors were presented. In account of the results, Demke et al. (2002) appealed to Baddeley’s model of working memory (Baddeley, Gathercole & Papagno, 1998), with attention to the phonological loop. According to the model, the phonological loop is a module of working memory specific to the encoding,

maintenance, and use of auditory linguistic information. The phonological loop is conceived as a temporary buffer for novel words as representations are assembled for eventual storage in long-term memory. To achieve this, the phonological loop must be continually refreshed because the details of novel words decay rapidly when held in the phonological loop. Extending these premises of Baddeley's model, Demke et al. (2002) reasoned that post-exposure to neighbors helped a child retain novel items in the phonological loop. During post-exposure, phonological representations of neighbors were activated in the mental lexicon. Because neighbors were familiar words known by children, their representations were already entrenched in long-term memory. This thereby established a link between novel newly trained items, which were temporarily held in the phonological loop, and neighbors, which were represented and stored in long-term memory. Demke et al. (2002) advanced that post-exposure to neighbors refreshed the phonological loop in this way. They further advanced that post-exposure to neighbors was "used to 'fill in' the representation of the novel words in the phonological loop, leading to enhanced recall" (389). By their account, phonological working memory was introduced as an alternative to auditory word priming as the mechanism behind the dense neighborhood advantage in typical development. Thus, two complementary hypotheses emerged. In the present studies, we aimed to evaluate these by replication and extension to children with PD.

PHONOLOGICAL DELAY

Children with PD present a unique opportunity to assess the impact of dense neighborhoods given their symptomology, etiology, need for treatment, and asymmetries in learning. To begin, the defining characteristic of PD is a severely reduced consonantal inventory relative to age-matched peers. Children with PD who are acquiring English tend to produce nasals, stops, and glides to the exclusion of other sounds. Sounds that are excluded from the repertoire are described linguistically as phonotactic constraints (Dinnsen, 1984), which restrict the occurrence, distribution, and use of sounds at phonemic and phonetic levels. The result is homonymy and a collapse of phonemic distinctiveness among words rendering unintelligible speech.

Beyond this primary deficit, some subsets of children with PD present with co-occurring deficits. The risks do not apply uniformly within or across children, but two are worth mentioning in light of the hypotheses under consideration. Shriberg and Kwiatkowski (1994) found that some children with PD have reduced vocabularies, suggesting a link between phonological and lexical learning. Shriberg and colleagues (2009) also found that some with PD perform poorly on nonword repetition tasks, suggesting a link between phonological learning and phonological working memory.

While there is no known cause of PD, a recurring view is that the problem lies in the way children represent the phonological properties of words (Dinnsen, 1984; Macken, 1980; Stoel-Gammon, 2011). This follows from linguistic accounts of typical and atypical development, which claim that acquisition of the sound system requires change (Dinnsen, 1984), elaboration (Rice & Avery, 1995), and/or reorganization (Gnanadesikan, 1996) of the phonological structure of lexical representations. For children with PD, modifications in phonological structure require explicit treatment. Treatment conventionally targets new

phonemic distinctions and is typically administered as a single-subject experiment (Baker & McLeod, 2011). Sounds excluded from the inventory are trained in production as the independent variable and generalization to treated and untreated (erred) sounds are measured as the dependent variable. The ultimate goal is to induce the broadest system-wide change in the phonologies of these children, thereby optimizing treatment efficacy.

To date, three studies have evaluated dense neighborhoods in treatment of children with PD, yielding mixed results. Two studies found little to no system-wide generalization when sounds were treated in words from dense neighborhoods (Gierut, Morrisette & Champion, 1999; Morrisette & Gierut, 2002). By comparison, a third study showed the expected dense neighborhood advantage (Gierut & Morrisette, 2012b). When sounds were taught in dense words with convergent cues (i.e., dense words that were also frequent early acquired items), children evidenced the greatest system-wide generalization. In the face of asymmetric results, it is noteworthy that the former studies manipulated density as the sole independent variable, whereas the latter co-varied density with other properties of word structure (e.g., frequency, age of word acquisition). It is of further mention that studies of adults and children with typical development (Garlock, Walley & Metsala, 2001; Krull, Choi, Kirk, Prusick & French, 2010; Metsala, 1997) have likewise reported mixed effects when density is manipulated singly versus additively. The literature has appealed to competition models (e.g., MacWhinney, 1987) in account of these observed asymmetries. Competition models claim that a given property of word structure may wax and wane because of cue convergence, collision, or weightings under different scenarios. This account notwithstanding, it remains that the conditions under which dense neighborhoods facilitate (or inhibit) phonological generalization in treatment of PD are not yet fully established or well understood, thus warranting further study.

The purpose was to replicate and extend Demke et al. (2002) by documenting the effects of dense neighborhoods on phonological generalization by children with PD enrolled in treatment. In Study 1, children were exposed to words from dense neighborhoods prior to training sound production, whereas in Study 2, exposure followed training sound production. Dense words presented before or after training were either neighbors or nonneighbors of the words that were taught. Following Stoel-Gammon (2011), one prediction was that neighbors would induce greater system-wide generalization than nonneighbors. Following Church & Fisher (1998), another prediction was that exposure to neighbors prior to production training would result in greater generalization, thereby supporting long-term auditory word priming as a mechanism that drives phonological learning. Following Demke et al. (2002), an alternate prediction was that exposure to neighbors following production training would lead to greater generalization, thereby supporting phonological working memory as responsible for the dense neighborhood advantage.

STUDY 1

METHOD

Participants and their phonologies—Six preschool children with PD ($M = 3;11$; range: 3;5–4;7) were recruited by public announcement. To participate, a child scored 1

standard deviation below the normative mean on the *Goldman-Fristoe Test of Articulation-2* (Goldman & Fristoe, 2000) and produced at least 6 sounds in error on this measure across phonetic contexts. Inclusionary criteria required performance within typical limits on diagnostic tests of hearing acuity, oral-motor structure/function, nonverbal intelligence, expressive/receptive vocabulary, nonword repetition, and working memory. Inclusionary criteria further required that children be preliterate, monolingual speakers of English. Children enrolled in concurrent speech and/or language services were excluded from participation. Table 1 reports the diagnostic profiles of individual participants, along with the mean performance by experimental group.

Children who met these criteria contributed a detailed speech sample. An established probe consisting of 293 words (Gierut, 2008b: 44–48) was used to elicit production of all English consonants in all relevant word positions in multiple exemplars. Each consonant was sampled in approximately 17 unique words: 5 eliciting initial position, 7 intervocalic position, and 5 final position depending on the phonotactics of English. The probe also provided for elicitation of minimal pairs as evidence of the contrastive (i.e., phonemic) status of sounds in a child's inventory. The probe was administered using a spontaneous picture-naming task and a child's responses were digitally recorded. A trained listener, blind to the experiment, phonetically transcribed the data and reliability was established as reported below.

Probe data were used to identify the inventory of sounds that a child used phonemically, along with corresponding exclusions. Sounds excluded from the inventory were of particular interest because these were manipulated in treatment and measured as evidence of phonological generalization. Established criteria were applied to identify sounds excluded from the inventory (Gierut & O'Connor, 2002). Specifically, sounds excluded were produced with near 0% accuracy of probe production and were never used phonemically to mark meaning distinctions in minimal pairs.

Experimental design and variables—A staggered multiple-baseline (MBL) across subjects design was used. The MBL consists of a baseline phase followed by treatment, with the number of baselines increased by 1 as successive children enroll. Baseline performance is expected to remain stable ($\leq 10\%$ accuracy) until the instatement of treatment, thereby establishing cause-effect relationships between instruction and performance.

As applied herein, the baseline was obtained through repeated administrations of the probe. Following baseline, treatment was instated, with children pseudorandomly assigned to 1 of 2 experimental conditions: pre-exposure to dense neighbors versus nonneighbors of the words that were trained in production. Thus, the independent variable was priming neighbors versus nonneighbors.

Children were assigned to an experimental condition in the order they enrolled: A first child was assigned to the pre-exposure neighbor condition, a second to the nonneighbor condition, a third to the neighbor condition, and so on. Assignments were pseudorandom in that treatment was specific to a sound excluded from a given child's phonemic inventory.

The dependent variable was system-wide phonological generalization. GENERALIZATION was operationalized as the percent gain in accuracy of production of sounds excluded from the phonemic inventory relative to baseline performance on the probe. On average, 10 sounds were excluded from children's inventories (Table 1); these were monitored for generalization. Every sound, treated and untreated, that evidenced positive gain over baseline was factored into the evaluation of generalization.

Generalization was measured longitudinally using the aforementioned 293-word probe. The probe was used exclusively as a test measure. Words of the probe were never taught or introduced in treatment. While the probe tests production of all English consonants, only sounds excluded from the phonemic inventory were monitored for generalization; hence, only relevant probe words were examined. There were approximately 170 probe words evaluated for generalization at each longitudinal point in time for each child (i.e., 10 sounds monitored x 17 probe words per sound).

Probes were administered on a variable schedule of two sessions starting at baseline and continuing to the completion of treatment. A variable schedule is a classic way to reduce the predictability of an event because that event is scheduled around an arithmetic average (Hilgard & Bower, 1975). Consequently, a participant's response to the event more closely resembles performance in the natural environment. A variable schedule of probe administration was used herein to best approximate a child's typical productions. Fifteen probe samples were obtained on average for each child over the duration of treatment. These data established generalization as causal to the delivery of treatment. Additional probe samples were collected 2 and 8 weeks after treatment was withdrawn. These data were strictly for descriptive purposes in documenting continued longitudinal gains. Throughout, the procedures outlined above for elicitation, blinded transcription, and reliability of probe data were followed.

Stimuli—Two sets of stimuli were developed: one exclusive to pre-exposure priming and another to training sound production. Words taught in production are described first because their form dictated which words could serve as primes.

Stimuli used in training sound production: Six words were used to teach production of a target sound; Appendix A lists the treated words for /s/. The words taught in production met three criteria: (1) they were from dense neighborhoods comprised of 10 or more neighbors (Luce, 1986), (2) to the extent possible, they were 3 segments in length (CVC), and (3) the treated sound assumed the initial position of the treated words.

Beyond that, there was a general effort to choose words familiar to children based on age-of-word-acquisition norms (Bird, Franklin & Howard, 2001; Gilhooly & Logie, 1980). There were further efforts to equate the words based on log frequency (retrieved from <http://neighborhoodsearch.wustl.edu>) and phonotactic probability (retrieved from <http://www.people.ku.edu/~mvitev/PhonoProbHome.html>). Characteristics of the treated words taught in production are reported in Table 2.

Stimuli used in pre-exposure priming: Consistent with the independent variable, neighbors versus nonneighbors of the treated words were identified. There were 48 words of each type, with examples shown in Appendix A. Following definitions used by Demke et al. (2002), neighbors had the same rhyme structure as corresponding treated words, whereas nonneighbors were phonologically unrelated to treated words.

Neighbors met all of the same criteria outlined above with one exception. They too were dense forms, 3 segments in length to the extent possible, generally familiar to children, and balanced in log frequency and phonotactic probability (Table 2); however neighbors had unique onsets relative to the corresponding words taught in production. Of note, there was no significant difference between neighbors and words taught in production based on age-of-word-acquisition, $t(98) = .43, p = .67$; log frequency, $t(160) = 1.12, p = .26$; sum of segment frequency, $t(160) = 1.87, p = .06$; or sum of biphone frequency $t(160) = .54, p = .59$. Thus, neighbors used in pre-exposure priming were on par with the words taught in production.

Nonneighbors met the same criteria, but were phonologically unrelated to corresponding words taught in production. As in Table 2, density, length, familiarity, log frequency, and phonotactic probability were considered in selection. There was again no significant difference between nonneighbors and words taught in production based on age-of-word-acquisition, $t(98) = 1.13, p = .26$; log frequency, $t(160) = -.16, p = .88$; sum of segment frequency, $t(160) = 1.28, p = .20$; or sum of biphone frequency, $t(160) = -.32, p = .75$. Nonneighbors were thus on par with words taught in production.

For completeness, there was no statistical difference between neighbors and nonneighbors based on age-of-word-acquisition, $F(5, 170) = 1.12, p = .32$; log frequency, $F(5, 282) = 2.05, p = .07$; sum of segment frequency, $F(5, 282) = .60, p = .70$; or sum of biphone frequency $F(5, 282) = 1.93, p = .09$. The stimuli used in pre-exposure priming were comparable across experimental conditions.

Materials—Two sets of materials were developed: one exclusive to pre-exposure priming and another to training sound production. Materials were kept distinct in the experimental sessions.

Materials used in pre-exposure priming: Following Demke et al. (2002), neighbors versus nonneighbors were embedded in stories, as in Appendix A. There were six stories to prime each of the six treated words. Stories averaged 34 words in length. A female talker recorded each story in a speaking style typical of reading to young children. Stories averaged 13.2s in duration with a 2.1s ISI between stories. Visual renditions of the stories were created by Sharp Designs & Illustration Inc. (previously Sharp Designs), measuring 7.5×10 inches. These were assembled into a PowerPoint slide show and synced with the audiofiles for random presentation with automatic advancement during the pre-exposure phase of the treatment protocol.

Materials used in training sound production: Pictures were used to elicit children's responses during production training. There was one picture for each treated word. Pictures

were retrieved from Google images and arranged in PowerPoint for random and repeated presentation during the production phase of the treatment protocol.

Treatment Protocol—Treatment sessions were 1-hour in duration conducted three times weekly. Each session consisted of two sequentially ordered phases: pre-exposure to neighbors (or nonneighbors) followed by training sound production. The treatment protocol was administered by the second author, who is a Ph.D.-level certified speech-language pathologist with 17 years of clinical research experience. Procedures are described below and in Appendix B.

Pre-exposure phase: A child was seated at a small table in a quiet room of an experimental suite. A desktop computer with 17-inch display and 2 desktop speakers were on the table. At the start of each treatment session, a child watched the PowerPoint slide show of either the neighbor or nonneighbor stories, depending on experimental assignment. First, a child was instructed to listen to, and watch the stories; no verbal, physical, or other response was required. Then, the experimenter initiated the PowerPoint slide show. Each of the six pre-exposure stories was presented, with automatic advancement as stated above. Order of the stories was randomized across sessions, also stated above. When the slide show was over, this ended the pre-exposure phase of the session and training sound production immediately commenced.

Production training phase: Each child was taught one sound excluded from, and specific to his/her phonemic inventory. Sounds that were taught were restricted to the late-8 consonants /s/, /l/, or /r/ (Shriberg, Kwiatkowski & Gruber, 1994). These sounds are commonly in error in typical and delayed populations, thereby affording generalizability of results. Sounds were also restricted for practical reasons. Carefully matched experimental stimuli and professional illustrations had to be prepared well in advance of identifying participants. The decision to teach /s/, /l/, or /r/ increased the likelihood that the prepared stimuli would indeed be relevant to the delayed phonologies of eligible children. Within each experimental condition, one child was taught /s/, one /l/, and one /r/. Treating different sounds within and across experimental conditions is conventional and intended to minimize sound-specific learning effects (Rvachew & Nowak, 2001).

The treated sound was taught using the aforementioned treated words. An established treatment protocol was followed (Gierut, 2008a) using drill play (Shriberg & Kwiatkowski, 1982). Specifically, each treated word was introduced as a discrete trial. A child was shown a picture of the treated word. The experimenter modeled the word. Following the model, a child imitated production of the word. 1:1 feedback was provided about accuracy of production, with praise for accurate outputs and placement cues for erred outputs. Production training continued in this way, trial after trial, for the duration of the session: A picture of the treated word was displayed, the experimenter modeled production of that word, a child imitated, and 1:1 feedback was provided. A child was dismissed at the end of the 1-hr session ($M= 71$ trials), only to return the next session for exactly the same instruction, beginning with the pre-exposure slide show followed by training on sound production.

Production training continued in imitation for 7 total sessions or until a child imitated the treated words with 75% accuracy over 2 consecutive sessions, whichever occurred first. When criterion was met, the response mode shifted to spontaneous production. As before, a child produced the treated sound in the treated words, using the same materials, drill play, and 1:1 feedback. The only difference between imitative and spontaneous response modes was that a child produced the treated sound in the treated words without benefit of a preceding model. Treatment continued in the spontaneous response mode for 12 total sessions or until a child independently produced the treated words with 90% accuracy over 3 consecutive sessions, whichever occurred first. When criterion was met in the spontaneous response mode, the protocol was completed and treatment, withdrawn. Thus, each child received treatment for a maximum of 19 sessions. Each treatment session always began with the pre-exposure slide show followed by production training, first in imitative and then spontaneous response modes, until criterion had been met.

Reliability—Transcription reliability was established for 10% of the probe data collected from each child. Two phonetically trained and blinded judges independently transcribed the samples, and these were compared point-to-point for consonant agreement. Reliability was established at 92% mean agreement (range: 85–97%).

Fidelity was assessed for 5% of the experimental sessions using a checklist procedure (Gierut, 2008a). An independent observer monitored randomly selected sessions to ensure that the protocol was administered as directed and probe data sampled as scheduled. The experimenter was blind to the collection of fidelity data. Fidelity was established at 100%.

RESULTS AND DISCUSSION

Two kinds of data were evaluated, performance during treatment and generalization. Performance during treatment established that children learned what had been taught as the foundation from which generalization could occur as the dependent variable.

Performance during treatment—Time in treatment was documented to establish that children received comparable amounts of instruction. Children in the neighbor condition completed the protocol in an average of 15 sessions (range: 6–19). Those in the nonneighbor condition required 17 sessions (range: 13–19) to completion. There was no statistical difference across conditions in the number of sessions in imitative and spontaneous response modes, $\chi^2(1, N=4) = .09, p = .76$.

Accuracy of production of the treated sound in treated words was also considered. In the neighbor condition, children achieved 72% (range: 50–85%) and 84% (range: 73–98%) mean accuracy in imitative and spontaneous response modes, respectively. In the nonneighbor condition, performance was comparable with mean accuracy of 67% (range: 57–83%) and 86% (range: 71–100%) in imitative and spontaneous modes, respectively.

The nonparametric McNemar test for significance of change was computed independently for each condition. The intent was to determine whether improved production accuracy as a consequence of treatment was reliable. Production of the treated sound in treated words was evaluated at baseline and again at completion of treatment. Table 3 reports the number of

treated words produced correctly at each sampling point for the neighbor and nonneighbor conditions. Results showed that both conditions were statistically significant, $p < .001$. As intended, treatment led to improved production of the TREATED SOUND in TREATED WORDS. As will be shown, this result takes on significance when production of the TREATED SOUND in UNTREATED PROBE WORDS is evaluated as a reflection of generalization. Comparable findings across conditions confirmed that treatment provided an equal scaffold from which differential generalization could take place.

Generalization—Generalization was defined previously as system-wide gains in production accuracy of sounds excluded from children’s phonemic inventories relative to baseline performance on the probe. Generalization data were examined in four complementary ways using conventional metrics of description, clinical significance, statistical significance, and practical significance as benchmarks for interpretation (Bain & Dollaghan, 1991; Bothe & Richardson, 2011). Description of generalization from treatment is standard to the evaluation of single-subject research. Clinical significance interprets generalization from treatment in a manner that is recognizable by professionals who work with the population of study (Bothe & Richardson, 2011: 235). Statistical significance establishes that generalization is real and not due to other variables (Bain & Dollaghan, 1991: 266). Practical significance is an adjunct to statistical significance in its use of effect size. Practical significance reveals the importance of generalization by capturing the absolute size of the gain for cross-comparisons and meta-analyses (Bain & Dollaghan, 1991: 267).

Description: Figure 1 plots the mean percent generalization gain over baseline for the neighbor and nonneighbor conditions, with standard error shown. Three key points in time are plotted, showing gains at completion of treatment in imitative and spontaneous response modes, and longitudinally after withdrawal of treatment.

By visual inspection, greater system-wide generalization was associated with the neighbor condition at each point in time. In the neighbor condition, gains ranged from 11.9–21.3% relative to baseline performance on the probe. In the nonneighbor condition, gains were in the range of 3.2–6.2% relative to baseline. This was notable because children in the neighbor condition started with lower levels of baseline accuracy on the probe ($M = 1.9%$) compared to children in the nonneighbor condition ($M = 8.8%$). Despite less accuracy of production at the start of treatment, children pre-exposed to neighbors evidenced greater generalization.

An established criterion cut-off of 10% gain relative to baseline (Elbert, Dinnsen & Powell, 1984) was applied to the generalization data in Figure 1. The 10% cut-off was originally conceptualized as a minimum threshold signaling stable or true generalization gain. The 10% minimum accords with Bain & Dollaghan’s view (1991: 268) that generalization gain, no matter the size, represents a distinct improvement in the daily functioning of a child with linguistic delays/disabilities. The 10% cut-off is conventionally applied in reference to system-wide improvements in production accuracy, not to individual sounds. As such, the criterion cut-off enables the binary coding of experimental conditions as inducing yes/no generalization. Figure 1 shows that the neighbor condition met the 10% criterion cut-off at

each sampling point in time, but the nonneighbor condition did not. By this metric, pre-exposure to neighbors appeared to induce generalization, but nonneighbors did not.

Clinical significance: A conventional index of severity, *Percent Consonants Correct–Revised* (PCC-R; Shriberg, Austin, Lewis, McSweeny & Wilson, 1997), was applied to aid the clinical interpretation of the aforementioned descriptive data. Established procedures for calculating PCC-R were applied to 53-word samples obtained from each child at baseline, completion of treatment, and longitudinally after withdrawal of treatment. PCC-R is computed by tallying the number of accurate consonant productions relative to the total consonants produced in the sample to derive a percentage. Greater PCC-R scores align with less severe phonological delays.

Figure 2 plots the mean gain in PCC-R scores relative to baseline (Table 1) for the neighbor and nonneighbor conditions, with standard error shown. Two points in time are shown: completion of treatment and longitudinally after treatment was withdrawn. Generally, the neighbor condition resulted in greater PCC-R scores than the nonneighbor condition. Children in the neighbor condition improved their PCC-R scores 6.3% at completion of treatment and 14.9% continuing longitudinally. By comparison, those in the nonneighbor condition improved their PCC-R scores 3.4% and 5.7% at completion of treatment and longitudinally, respectively. From a clinical standpoint, pre-exposure to neighbors had a greater (positive) impact on the clinical characterization of severity than nonneighbors.

Statistical significance: Generalization of the treated sound to untreated probe words was evaluated statistically using the McNemar test. This was the direct complement to the aforementioned McNemar test of the treated sound in treated words. The intent was to compare performance during treatment relative to generalization from treatment when the object of learning (i.e., the treated sound) was the same in both cases.

Table 3 shows that, for the neighbor condition, generalization of the treated sound to untreated probe words was statistically reliable and not due to chance, $p = .01$. By comparison, in the nonneighbor condition, generalization of the treated sound was not statistically significant, $p = 1.00$. This finding is of mention because, during treatment, both experimental conditions induced statistically significant change in production of the treated sound in treated stimuli. However, these generalization data show that transfer of the treated sound to other untreated words and contexts was only statistically reliable in the neighbor condition. Pre-exposure to neighbors promoted transfer of the treated sound; nonneighbors did not.

Practical significance: Effect size is a relatively new addition to the single-subject literature that has been introduced with an eye toward meta-analyses of treatment studies (Beeson & Robey, 2006). Effect size captures the absolute magnitude of generalization gain associated with a given experimental condition, and then values obtained across conditions are compared to identify which is relatively more effective in promoting generalization.

Standard mean difference with correction for continuity (d) was applied herein for consistency with other studies of children with PD (Gierut & Morrisette, 2011, 2012a, b).

This statistic is specific to single-subject design (Busk & Serlin, 1992) and not to be confused with effect size for large-N studies (Cohen, 1988). Standard mean difference with correction for continuity is calculated in the following way: Mean accuracies are computed at baseline for each child for all sounds excluded from the phonemic inventory under each experimental condition. Likewise, mean accuracies are computed for each child for each longitudinal sample obtained through completion of treatment for all sounds excluded from the phonemic inventory that generalized. The difference between mean baseline and generalization data is then divided by the mean standard deviation of the baseline for the population of study (i.e., all participants) to yield an effect size, d . The standard deviation of the population takes into account individual variability in baseline performance (Glass, 1977) and accommodates 0% baselines in computation of effect size (Gierut & Morrisette, 2011). The data for each experimental condition are then aggregated for relative comparison.

While benchmarks for interpretation of effect size have been established (arbitrarily) for large-N studies (Cohen, 1988: 532), it is not appropriate to extend these to single-subject research (Beeson & Robey, 2006: 167). Standards that define small, medium, or large effects must be developed empirically for a given population, and to achieve this, effect sizes must first accumulate. To date, treatment studies of PD (Gierut & Morrisette, 2011, 2012a, b), which have used the standard mean difference with correction for continuity, report d values in the range of 2.60 to 16.58 ($M= 7.88$; $Mdn= 5.61$). The present study added to this base as a secondary contribution.

Figure 3 plots the mean effect size associated with neighbor and nonneighbor conditions, with standard error shown. The average d was 7.57 for the neighbor condition compared to 4.00 for the nonneighbor condition. The absolute gain from pre-exposure to neighbors versus nonneighbors was on the order of magnitude of 2:1.

Obtained d values were further aligned with previous reports of effect size for the population of PD cited above. A median split, where $d = 5.61$ for the population of PD, was used to sort the present data into two categories for interpretation (cf. Estes, Edwards & Saffran, 2011). For the neighbor condition, the obtained d of 7.57 was greater than the median d for the population. For the nonneighbor condition, the obtained d of 4.00 was less than the median d . Thus, pre-exposure to neighbors was affiliated with greater magnitude of gain when taken in the context of the population of PD.

Together, the results of description, clinical, statistical and practical significance converged in Study 1. Pre-exposure to dense neighbors led to greater and reliable generalization compared to nonneighbors in treatment of children with PD. This finding is consistent with Stoel-Gammon's (2011) description of the relevance of dense neighborhoods to phonological acquisition. As in typical development, improved production accuracy and generalized use of new phonemes in new contexts and lexical items were associated with dense neighborhoods.

The results were also consistent with Church & Fisher's (1998) hypothesis that long-term auditory word priming bears on the phonological structure of children's representation of

words. Here, change in phonological structure was reflected behaviorally in children's production accuracy and generalized use of sounds excluded from the inventory. Thus, long-term auditory word priming may offer a viable account of the dense neighborhoods effects for children with PD.

Study 1 further demonstrated that the treatment paradigm developed by Demke et al. (2002) for lexical purposes was equally applicable to the phonological domain and appropriate for children with PD. This notwithstanding, the results obtained herein stand apart from the findings of Demke et al. (2002): Pre-exposure to dense neighbors aided phonological learning, but not lexical learning, thus motivating Study 2.

STUDY 2

The purpose was to evaluate the effect of exposing children with PD to neighbors versus nonneighbors after training sound production. The goal was to alter the timing of exposure in evaluation of an alternate hypothesis that phonological working memory guides the dense neighborhood advantage in phonological acquisition.

METHODS

Participants—Six children with PD ($M = 4;4$; range: 3;4–5;5) were recruited following procedures outlined for Study 1. Inclusionary and exclusionary criteria were identical to those described previously, with individual profiles and mean performance by experimental group shown in Table 4.

Experimental design and methods—The design, stimuli, and materials were identical to Study 1. The independent variable was post-exposure to neighbors versus nonneighbors, and the dependent variable was systemwide phonological generalization. The difference between studies was the timing of exposure to neighbors or nonneighbors. Specifically, the order of administration of the treatment protocol was reversed: Study 1 presented neighbors or nonneighbors BEFORE training sound production (Appendix B), whereas Study 2 presented neighbors or nonneighbors AFTER training sound production (Appendix C). This can be seen in the comparison of Appendices B and C, which are identical in all respects except for the timing of exposure relative to production training in a given experimental session.

Reliability—As in Study 1, reliability of transcriptions was computed and established as 92% consonant agreement (range: 89–97%) between independent judges. Fidelity was established as 100% conformity in administration of procedures.

RESULTS AND DISCUSSION

Results were analyzed as in Study 1, with attention to performance during treatment and generalization. A comparison of Studies 1 and 2 was also considered.

Performance during treatment—Time and accuracy of production during treatment were documented. With respect to time, children in the neighbor condition required an average of 11 sessions (range: 5–16) to complete the protocol. The nonneighbor group was

comparable, requiring an average of 12 sessions (range: 10–14) to completion. There was no significant difference across conditions in the number of sessions in imitative and spontaneous response modes, $\chi^2(1, N=4) = .06, p = .81$. With respect to accuracy, children in the neighbor condition produced the treated sound in treated words with 85% mean accuracy (range: 80–93%) in imitative and 97% mean accuracy (range: 95–98%) in spontaneous response modes. Accuracy in the nonneighbor condition was similar: Children achieved 84% (range: 75–95%) and 95% (range: 93–99%) mean accuracy in imitative and spontaneous response modes, respectively.

Independent McNemar tests for the significance of change in accuracy of the treated sound during treatment are reported in Table 5. The neighbor and nonneighbor conditions each resulted in statistically reliable gains in accuracy of the treated sound in treated words as a direct consequence of treatment, $p < .001$. As intended, treatment had comparable effects on children's production of the treated sound across conditions, thereby establishing a foundation for subsequent generalization.

Generalization—Description and clinical, statistical, and practical significance were considered in evaluation of system-wide generalization to sounds excluded from the phonemic inventory relative to baseline performance on the probe. Data analyses paralleled Study 1.

Description: Figure 4 displays the mean generalization gain over baseline performance on the probe for the post-exposure conditions, with three points in time plotted: after completion of treatment in imitative and spontaneous response modes and longitudinally after treatment was withdrawn. Visual inspection revealed little to no difference in generalization gain associated with post-exposure to neighbors versus nonneighbors over time. The neighbor condition resulted in gains in the range of 4.6–6.9% accuracy over baseline. The nonneighbor condition was nearly identical, with gains in the range of 4.2–5.9% accuracy over baseline.

The 10% criterion cut-off (Elbert et al., 1984) was applied to the data in Figure 4 as a metric to binarily code the occurrence of system-wide generalization. Figure 4 shows that neither post-exposure condition met the minimum 10% cut-off. Thus, in descriptive evaluation, post-exposure to dense neighbors and nonneighbors had undifferentiated and minimal effects on system-wide generalization.

Clinical significance: Figure 5 plots the average gains in PCC-R scores relative to baseline (Table 4) for the neighbor and nonneighbor conditions at two points in time. Recall that PCC-R is a clinical index of severity. Figure 5 shows little to no difference in PCC-R scores across conditions. In the neighbor condition, PCC-R scores improved an average of 4.0–8.0% and in the nonneighbor condition, 4.0–6.9%. Thus, post-exposure to neighbors and nonneighbors had comparable modest effects on clinical estimates of severity.

Statistical significance: The McNemar test was used to gauge the statistical reliability of generalization of the treated sound to untreated probe words. Recall that the relevant point of interest was performance during treatment versus transfer from treatment when the object of

learning (i.e., the treated sound) was the same. Table 5 shows that transfer of the treated sound to untreated probe words was not statistically significant for the neighbor, $p = .38$ or nonneighbor, $p = .07$ conditions. Neither post-exposure condition promoted reliable generalization of the treated sound, despite reliable gains in accuracy of that same sound during treatment.

Practical significance: Effect size data are plotted in Figure 6 for the post-exposure neighbor and nonneighbor conditions. The magnitude of system-wide generalization gain was again comparable across conditions, where $d = 2.41$ for neighbors and 2.52 for nonneighbors.

Obtained d values were examined relative to the population of children with PD. Recall that the magnitude of generalization for the population (reported above) ranged from 2.60–16.58 ($M = 7.88$, $Mdn = 5.61$). It is notable that the d values for post-exposure to neighbors (2.41) and nonneighbors (2.52) fell below the median (5.61) obtained for the population. In fact, the d values obtained herein are the lowest reported to date for children with PD. Thus, the post-exposure conditions served to reset the effect size minima for the population.

In sum, Study 2 yielded essentially equivalent effects across experimental conditions based on the convergence of descriptive, clinical, statistical, and practical evidence. Generalization following post-exposure to neighbors versus nonneighbors was undifferentiated and modest at best. These findings are of interest for two reasons. First, they did not support the expected dense neighborhood advantage for phonological learning (Stoel-Gammon, 2011). Second, they are at odds with the results of lexical learning (Demke et al., 2002) where differential effects of neighbors versus nonneighbors were observed post-exposure. The asymmetries are perhaps best understood in a comparison of Studies 1 and 2.

Integration of studies—Thus far, the presentation of results emphasized the differentiation of neighbors versus nonneighbors as the independent variable of the respective studies. Study 1 found differential effects, such that pre-exposure to neighbors > nonneighbors in promoting generalization. In contrast, Study 2 found little to no differential effects, with post-exposure to neighbors = nonneighbors. Visual inspection of the figures associated with Studies 1 and 2 further revealed that the amount of generalization observed in pre-exposure was greater than that of post-exposure. This can be seen in comparisons of descriptive data (cf. Figures 1 and 4), clinical data (cf. Figures 2 and 5) and practical data (cf. Figures 3 and 6). Moreover, the results across studies suggested that neighbor status and timing of exposure were coupled: The greatest generalization occurred when exposure to neighbors took place prior to training production. This is a finding wholly consistent with the predictions of auditory word priming.

It is possible to explore the relative contribution of neighbor status and timing of exposure by aggregating effect sizes across studies (Gierut & Morrisette, 2011). This is achieved by blocking effect sizes for a given variable, while setting aside the other. Specifically, in aggregation of Studies 1 and 2 for the neighbor versus nonneighbor manipulation, timing of exposure was set aside. Based on aggregated d values, neighbors promoted greater generalization than nonneighbors on an order of magnitude of 1.2:1 (i.e., d blocked by

neighbors = 4.5, nonneighbors = 3.3 following from Figures 3 and 6). Likewise, in aggregation of Studies 1 and 2 for the pre-versus post-exposure manipulation, neighbor status was set aside. Based on aggregated d values, pre-exposure induced greater generalization than post-exposure on an order of magnitude of 2.6:1 (i.e., d blocked by pre-exposure = 5.4, post-exposure = 2.5 following from Figures 3 and 6). This hints that timing of exposure to dense neighbors was indeed critical to generalization.

There is one caveat, however, that must be explored in the comparison of studies as it relates to individual differences. While dense neighbors and pre-exposure have emerged as essential variables, the profiles of individual children must be ruled out as possibly contributing to generalization effects. Correlational analyses were completed to establish whether there was a relationship between effect size and the diagnostic data shown in Tables 1 and 4. Results showed no statistically significant correlation between effect size and chronological age, number of sounds excluded from a child's phonemic inventory, or PCC-R scores at enrollment, all r s < .09, all p s > .79. There was also no correlation between effect size and total number of treatment sessions or number of sessions in imitative and spontaneous response modes, all r s < -.39, all p s > .25. There was no correlation between effect size and the results of diagnostic tests of articulation, oral motor structure/function, nonverbal intelligence, expressive vocabulary, receptive vocabulary, nonword repetition, or working memory, all r s < .48, all p s > .14. None of the child-specific variables was predictive of effect size. While it is of course possible that a yet to be determined lurking variable was operative, the data at hand suggest otherwise. Differential generalization within across studies was attributable to the experimental manipulations of neighbor status and timing of exposure.

GENERAL DISCUSSION

Our goal was to evaluate long-term auditory word priming and phonological working memory as possible learning mechanisms behind the dense neighborhood advantage in language acquisition. By replication and extension to the phonological domain for children with PD, results supported the expected benefit of dense neighborhoods, but only under well-defined pre-exposure conditions. In discussion, results are placed in the broader context of phonological acquisition and interpreted in support of priming as a relevant language learning mechanism. Suggestions for future research are offered to better understand the complementary role of phonological working memory in guiding dense neighborhood effects.

DENSE NEIGHBORHOODS AND PHONOLOGICAL ACQUISITION

The present work replicated the dense neighborhood effect in phonological acquisition, but only in part. Neighbors did not uniformly enhance generalization for children with PD, but rather, their beneficial effects were incumbent on timing of exposure.

There are several reasons why the coupling of neighbor status and timing of exposure might have been essential for children with PD. First, previous literature has demonstrated that priming dense neighbors has a magnifying effect on the phonemic distinctions of language (De Cara & Goswami, 2003; Merriman & Marazita, 1995). Recall that a defining trait of

children with PD is a reduced phonemic inventory. Thus, pre-exposure to neighbors may have honed in on the problem at hand. Second, when dense neighbors are repeatedly primed, points of segmental overlap among words habituate and novelty of form is brought forward (Merriman & Marazita, 1995). In the present research, rhymes receded, and onsets were uniquely spotlighted. Importantly, the phoneme to be learned in treatment was in onset position. Third, priming dense neighbors provided children with the opportunity to experience minimal pairs side-by-side. Children heard stories loaded with minimal pairs (e.g., *hail, fail, pail*) and then were taught related neighbors of the same minimal pair set (e.g., *sail*). In typical phonological development, children may not require direct experience with minimal pairs to acquire phonemes (Maye & Gerken, 2000). However, children with PD are highly responsive to minimal pairs, with this being among the efficacious treatments to promote generalization (Baker & McLeod, 2011). It is possible that priming provided a unique way to expose children with PD to minimal pairs. Finally, minimal pairs are central to the differentiation of meaning among words and meaning is at the heart of the lexicon. Priming dense neighbors may have reinforced the critical association between form and meaning for children with PD. Taken together, the pre-exposure neighbor condition may have afforded an ideal scenario for optimizing phonological acquisition in PD.

Given the potential promise of priming for PD, it is especially important to extend this line of investigation to address limitations of the present studies. In particular, the inherent design of single-subject studies restricts the number of participants. Replications enrolling additional children will lend robustness to the data. Likewise, our treatment protocol was designed to affect change in children's phonemic inventories. While gaps in the inventory are a primary trait of PD, it is well known that children often present with a range of production errors. Replications are needed to test the applicability of priming to other error patterns and other contexts. Further, the treated sounds and experimental stimuli need to be expanded. In the present studies, sounds taught in production were restricted to /s/, /l/, or /r/ for reasons associated with generalizability and feasibility. Future studies are needed to establish a broader base of generalization effects in treatment of other early-, mid-, or late-8 segments. Similarly, the words used in priming and production training were chosen based on adult-referenced norms of neighborhood density. There has been considerable debate in the literature (cf. Charles-Luce & Luce, 1990; Dollaghan, 1994), with no clear resolve about the appropriateness of adult estimates of neighborhood density for developmental studies (but see Storkel, 2013). Replications using child-referenced density norms are needed for completeness.

The present work also adds to the treatment literature, which has been inconclusive about the contributions of dense neighborhoods to generalization in PD. Recall that previous studies yielded mixed results about generalization from treatment of dense neighborhoods (cf. Gierut & Morrisette, 2012b; Morrisette & Gierut, 2002). This notwithstanding, there appears to be an emerging pattern about the conditions under which dense neighborhoods help or hinder generalization. In particular, studies that reported dense neighborhoods as facilitating also carried certain qualifications. Consider that dense neighbors aided generalization herein, but ONLY when presented prior to treatment. Similarly, dense neighbors aided generalization (Gierut & Morrisette, 2012b), but ONLY when input cues

converged (i.e., dense words that are frequent and early acquired). In each report of facilitating effects, an added layer of support for learning had been provided. Support came through adjustments in the format of instruction (i.e., priming) and treated stimuli (i.e., convergent input cues). It is possible that these adjustments boosted the utility of dense neighborhoods for children with PD. When such support was absent (Morrisette & Gierut, 2002), dense neighborhoods did not facilitate phonological generalization. Thus, dense neighborhoods seem to be a necessary, but not sufficient condition of phonological treatment.

This observation gives rise to basic questions about the role of dense neighborhoods for children with PD. To address such questions, future research may need to broaden its scope by appealing to, and integrating complementary data from spoken word recognition. This is a recent innovation in the study of SLI (Hoover & Storkel, 2013), where children's recognition of words from dense neighborhoods is explored for predictive value in treatment. New insights may be gleaned from a parallel approach to PD. Perhaps, children's recognition of dense neighbors will inform the success of treatment that employs dense neighbors. Whatever the outcome, the integration of recognition and production is likely to reveal individual differences in the characterization and remediation of PD.

MECHANISMS BEHIND PHONOLOGICAL LEARNING

The present research weighed in on more general hypotheses about the mechanisms of learning behind dense neighborhood effects in language acquisition. For phonological acquisition, in particular, our findings support long-term auditory word priming as a viable account. From this, it is possible to glean a hypothetical course of phonological acquisition by assembly of the literature: A child starts out by building the lexicon (Ferguson & Farwell, 1975), accumulating similar-sounding words from the input (Jusczyk, Luce & Charles-Luce, 1994). Initially, the representation of these words does not fully align with the phonological structure of the input (Dinnsen, 1984; Macken, 1980; Stoel-Gammon, 2011); consequently, a child's early productions are often homophonous. Nevertheless, the lexicon increases in size, as more words that overlap in phonological form are added, rendering dense neighborhoods (Charles-Luce & Luce, 1990). As the lexicon continues to grow, a child repeatedly experiences the same words time and again, akin to a naturalistic case of long-term auditory word priming (Church & Fisher, 1998). With each occurrence, the sound patterns of words from dense neighborhoods are strengthened. The result is change in the representation and use of phonemic distinctions as reflected in production accuracy (Stoel-Gammon, 2011). The contextual variability associated with these repeated occurrences further supports generalization (Beckman et al., 2007), such that a child's experience with similarity of form transfers to new phonemes, features, positions, and lexical items, all in an effort to differentiate meaning. Notice that a long-term auditory word priming account of this sort conforms to classic (Ferguson & Farwell, 1975) and contemporary (Beckman et al., 2007) views that the lexicon gives rise to phonology in acquisition. The account delineates what is learned as a consequence of dense neighborhoods, i.e., phonemic distinctions and how that learning takes place, i.e., priming. While intriguing, empirical validation of the process is essential. Computational studies of longitudinal data might be one way to begin,

with the occurrence of neighbors in the input tallied relative to the emergence of specific phonemes.

Although the explanatory power of priming remains to be determined, it appears that, at the very least, priming offers a ripe situation for advancing change in linguistic structure (Savage et al., 2006). Thus far, priming has been used to trigger various aspects of language learning in children with (Leonard, Miller, Grela, Holland, Gerber & Petucci, 2000) and without (Savage et al., 2003, 2006; Vasilyeva & Waterfall, 2012) language deficits. To our knowledge, the present studies are among the first to apply priming to PD and to phonological treatment (see also Gierut & Morrisette, 2014). As such, the exact conditions of priming that promote generalization warrant elaboration. Rhyme priming was employed herein, but studies of onset priming are needed (Brooks & MacWhinney, 2000; Merriman & Marazita, 1995). This manipulation might involve, for example, exposure to *sun, soap, sit* as primes followed by training production of /s/ in the word *sat*. The optimal mode of priming also needs to be considered (Gierut & Morrisette, 2014). Auditory+visual priming was used herein in keeping with Demke et al. (2002). However, priming might be presented in the auditory mode only as in Church & Fisher (1998), such that children hear neighbor stories in the absence of a corresponding picture. Conversely, pictures might be presented in priming, without corresponding auditory stories. The use of stories in priming is another research need. As priming has been implemented for lexical (Demke et al., 2002) and phonological learning (herein), children heard stories with prime words embedded. Consistent with the SEUSS BOOST (Read, Macauley, Furay, 2014), this may have contributed to word recognition, identification, retention, and learning. In future research, it will be necessary to test the effects of priming using citation lists. Predictably, a list format might facilitate phonological learning because a child would be exposed to minimal pairs without intervening or extraneous content. Yet, it is equally possible that a list format might be detrimental to phonological learning because the contextual support for meaning would be eliminated. Without meaning, the mapping of phonemic distinctions among words might be more challenging. The amount of priming to induce generalization also needs to be established. Children were exposed to primes a maximum of 19 times under the current treatment protocol, but less exposure may be equally effective. Through collective research along these lines, the conditions of priming will be better understood as a necessary foundation for community-based studies of priming in applied settings.

The discussion thus far has centered on priming as a mechanism of learning, but the present studies were designed to also evaluate phonological working memory as an alternative. Results failed to show support for this hypothesis: There was limited and no differential generalization following post-exposure to neighbors or nonneighbors. In contrast to Demke et al. (2002), the post-exposure conditions did not have an apparent effect on refreshing the phonological loop or filling in sound patterns of words for children with PD. At first blush, it might be tempting to reject the phonological working memory hypothesis, but this is premature.

Consider that exposure to dense neighbors before training production affected phonological learning herein, but exposure to dense neighbors after training novel words impacted lexical learning (Demke et al., 2002). One implication is that dense neighbors serve different

purposes depending on the task at hand. For phonological learning, neighbors promote phonological structure; for lexical learning, they refresh phonological structure. The seemingly asymmetric findings may be simply two sides of the same coin.

In future research, it will be important to jointly measure phonological and lexical learning as dependent variables (Ferguson & Farwell, 1975) in manipulations of dense neighborhoods. This idea takes on added interest in the context of PD. Recall that subgroups of children with PD may have co-occurring deficits in lexical learning (Shriberg & Kwiatkowski, 1994) or in phonological working memory (Shriberg et al., 2009). In the present studies, enrollment was restricted to PD in the absence of other deficits. Future studies might broaden the inclusionary criteria to enroll children with co-occurring problems as in PD+lexical deficits or PD+phonological working memory deficits. It is possible that the timing of exposure to dense neighbors will differentially affect these subgroups. Predictably, pre-exposure to neighbors might be optimal in cases of PD+lexical deficits. According to Church & Fisher (1998), pre-exposure assists the encoding of new words (thereby addressing lexical deficits) and the representation of sound patterns in words (thereby addressing PD). Post-exposure to neighbors might be best for those with PD +phonological working memory deficits. According to Demke et al. (2002), post-exposure helps retention of words in the phonological loop (thereby addressing phonological working memory deficits) and refreshes the phonological loop to fill in the details of the representation (thereby addressing PD). Research along these lines has the potential to reveal differential recommendations for treatment, with timing of exposure to dense neighborhoods being tailored specifically to the unique profiles of children with PD.

CONCLUSION

This research added to the body of evidence that supports the contributions of dense neighborhoods to phonological acquisition. The findings disambiguated possible accounts of the effects of dense neighborhoods on phonological structure by pointing to long-term auditory word priming as a mechanism of learning. The finding of a conjunction between neighbor status and timing of exposure sets the stage for continued research to establish the priming conditions that optimize generalization for children with PD. The results further motivate the integration of recognition with production data in the joint evaluation of phonological and lexical learning in children with typical and delayed development.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This research was supported in part by a grant from the National Institute on Deafness and Other Communication Disorders of the National Institutes of Health under Award Number R01DC001694. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. We thank Dan Dinnsen for input on the manuscript, Rachel Dale and Maureen Orawiec for assistance with stimulus development, and members of the Learnability Project for help with phonetic transcription, reliability, data entry and analyses. Paul and Alice Sharp of Sharp Designs & Illustration Inc. supplied the illustrations.

References

- Baddeley A, Gathercole S, Papagno C. The phonological loop as a language learning device. *Psychological Review*. 1998; 105:158–173. [PubMed: 9450375]
- Bain BA, Dollaghan CA. The notion of clinically significant change. *Language, Speech and Hearing Services in Schools*. 1991; 22:264–270.
- Baker E, McLeod S. Evidence-based practice for children with speech sound disorders: Part 1 narrative review. *Language, Speech and Hearing Services in Schools*. 2011; 42:102–139.
- Beckman, ME.; Edwards, J. Lexical frequency effects on young children’s imitative productions. In: Broe, M.; Pierrehumbert, J., editors. *Papers in laboratory phonology V*. Cambridge, UK: Cambridge University Press; 2000. p. 207-217.
- Beckman, ME.; Munson, B.; Edwards, J. Vocabulary growth and the developmental expansion of types of phonological knowledge. In: Cole, JS.; Hualde, JI., editors. *Laboratory Phonology 9*. New York: Mouton de Gruyter; 2007. p. 241-264.
- Bird H, Franklin S, Howard D. Age of acquisition and imageability ratings for a large set of words, including verb and function words. *Behavior Research Methods, Instruments, & Computers*. 2001; 33:73–79.
- Bothe AK, Richardson JD. Statistical, practical, clinical, and personal significance: Definitions and applications in speech-language pathology. *American Journal of Speech-Language Pathology*. 2011; 20:233–242. [PubMed: 21478279]
- Brooks PJ, MacWhinney B. Phonological priming in children’s picture naming. *Journal of Child Language*. 2000; 27:335–366. [PubMed: 10967891]
- Busk, PL.; Serlin, RC. Meta-analysis for single-case research. In: Kratochwill, TR.; Levin, JR., editors. *Single-case research design and analysis*. Hillsdale, NJ: Lawrence Erlbaum; 1992. p. 187-212.
- Charles-Luce J, Luce PA. Similarity neighbourhoods of words in young children’s lexicons. *Journal of Child Language*. 1990; 17:205–215. [PubMed: 2312642]
- Church BA, Fisher C. Long-term auditory word priming in preschoolers: Implicit memory support for language acquisition. *Journal of Memory and Language*. 1998; 39:523–542.
- Cohen, J. *Statistical power analysis for the behavioral sciences*. 2nd. Hillsdale, NJ: Lawrence Erlbaum; 1988.
- De Cara B, Goswami U. Phonological neighborhood density effects in a rhyme awareness task in 5-year old children. *Journal of Child Language*. 2003; 30:695–710. [PubMed: 14513474]
- Demke TL, Graham SA, Siakaluk PD. The influence of exposure to phonological neighbours on preschoolers’ novel word production. *Journal of Child Language*. 2002; 29:379–392. [PubMed: 12109376]
- Dinnsen, DA. Methods and empirical issues in analyzing functional misarticulation. In: Elbert, M.; Dinnsen, DA.; Weismer, G., editors. *Phonological theory and the misarticulating child (ASHA Monographs No. 22)*. Rockville, MD: American Speech-Language-Hearing Association; 1984. p. 5-17.
- Dollaghan CA. Children’s phonological neighbourhoods: Half empty or half full? *Journal of Child Language*. 1994; 21:257–272. [PubMed: 7929681]
- Dollaghan C, Campbell TF. Nonword repetition and child language impairment. *Journal of Speech, Language, and Hearing Research*. 1998; 41:1136–1146.
- Dunn, LM.; Dunn, LM. *Peabody picture vocabulary test*. 3rd. Circle Pines, MN: American Guidance Service; 1997.
- Edwards J, Beckman ME, Munson B. The interaction between vocabulary size and phonotactic probability effects on children’s production accuracy and fluency in nonword repetition. *Journal of Speech, Language, and Hearing Research*. 2004; 47:421–436.
- Elbert M, Dinnsen DA, Powell TW. On the prediction of phonologic generalization learning patterns. *Journal of Speech and Hearing Disorders*. 1984; 49:309–317. [PubMed: 6748626]
- Estes KG, Edwards J, Saffran JR. Phonotactic constraints on infant word learning. *Infancy*. 2011; 16:180–197. [PubMed: 21297877]

- Ferguson CA, Farwell CB. Words and sounds in early language acquisition: English initial consonants in the first fifty words. *Language*. 1975; 51:419–439.
- Garlock VM, Walley AC, Metsala JL. Age-of-acquisition, word frequency and neighborhood density effects on spoken word recognition by children and adults. *Journal of Memory and Language*. 2001; 45:468–492.
- Gathercole SE, Adams A-M. Phonological working memory in very young children. *Developmental Psychology*. 1993; 29:770–778.
- Gierut, JA. Fundamentals of experimental design and treatment. In: Dinnsen, DA.; Gierut, JA., editors. *Optimality theory, phonological acquisition and disorders*. London: Equinox; 2008a. p. 93-118.
- Gierut, JA. Phonological disorders and the Developmental Phonology Archive. In: Dinnsen, DA.; Gierut, JA., editors. *Optimality theory, phonological acquisition and disorders*. London: Equinox; 2008b. p. 37-92.
- Gierut JA, Morrisette ML. Effect size in clinical phonology. *Clinical Linguistics & Phonetics*. 2011; 25:975–980. [PubMed: 21787149]
- Gierut JA, Morrisette ML. Age-of-word acquisition effects in treatment of children with phonological delays. *Applied Psycholinguistics*. 2012a; 33:121–144. [PubMed: 22408279]
- Gierut JA, Morrisette ML. Density, frequency and the expressive phonology of children with delay. *Journal of Child Language*. 2012b; 39:804–834. [PubMed: 22182669]
- Gierut JA, Morrisette ML. How to meet the neighbors: Modality effects on phonological generalization. *Clinical Linguistics & Phonetics*. 2014; 28:477–492. [PubMed: 25000373]
- Gierut JA, Morrisette ML, Champion AH. Lexical constraints in phonological acquisition. *Journal of Child Language*. 1999; 26:261–294. [PubMed: 11706466]
- Gierut JA, O'Connor KM. Precursors to onset clusters in acquisition. *Journal of Child Language*. 2002; 29:495–517. [PubMed: 12109361]
- Gilhooly KJ, Logie RH. Age-of-acquisition, imagery, concreteness, familiarity, and ambiguity measures for 1,944 words. *Behavior Research Methods, Instruments and Computers*. 1980; 12:395–427.
- Glass G. Integrating findings: The meta-analysis of research. *Review of Research in Education*. 1977; 5:351–379.
- Gnanadesikan, AE. Child phonology in optimality theory: Ranking markedness and faithfulness constraints. In: Stringfellow, A.; Cahana-Amitay, D.; Hughes, E.; Zukowski, A., editors. *Proceedings of the 20th annual Boston University conference on language development*. Vol. 1. Somerville, MA: Cascadilla Press; 1996. p. 237-248.
- Goldman, R.; Fristoe, M. *Goldman-Fristoe test of articulation*. 2nd. Circle Pines, MN: American Guidance Service; 2000.
- Hilgard, ER.; Bower, GH. *Theories of learning*. 4th. Englewood Cliffs, NJ: Prentice-Hall; 1975.
- Hoover JR, Storkel H. Grammatical treatment and specific language impairment: Neighbourhood density & third person singulars. *Clinical Linguistics & Phonetics*. 2013; 27:661–680. [PubMed: 23819675]
- Jusczyk PW, Luce PA, Charles-Luce J. Infants' sensitivity to phonotactic patterns in the native language. *Journal of Memory and Language*. 1994; 33:630–645.
- Kirk, SA.; McCarthy, JJ.; Kirk, WD. *Illinois test of psycholinguistic abilities—revised*. Chicago: University of Illinois Press; 1968.
- Krull V, Choi S, Kirk KI, Prusick L, French B. Lexical effects on spoken-word recognition in children with normal hearing. *Ear and Hearing*. 2010; 31:102–114. [PubMed: 19701087]
- Leonard LB, Miller CA, Grela B, Holland AL, Gerber E, Petucci M. Production operations contribute to the grammatical morpheme limitations of children with specific language impairment. *Journal of Memory and Language*. 2000; 43:362–378.
- Luce, PA. *Neighborhoods of words in the mental lexicon*. Bloomington, IN: Speech Research Laboratory, Indiana University; 1986.
- Macken MA. The child's lexical representation: The 'puzzle-puddle-pickle' evidence. *Journal of Linguistics*. 1980; 16:1–17.

- MacWhinney, B. The competition model. In: MacWhinney, B., editor. *Mechanisms of language acquisition*. Hillsdale, NJ: Erlbaum; 1987. p. 249-308.
- Maye, J.; Gerken, L. Learning phonemes without minimal pairs. In: Howell, SC.; Fish, SA.; Keith-Lucas, T., editors. *Proceedings of the 24th annual Boston University conference on language development*. Somerville, MA: Cascadilla Press; 2000. p. 522-533.
- Merriman WM, Marazita JM. The effect of hearing similar-sounding words on young 2-year-olds' disambiguation of novel noun reference. *Developmental Psychology*. 1995; 31:973-984.
- Metsala JL. An examination of word frequency and neighborhood density in the development of spoken word recognition. *Memory & Cognition*. 1997; 25:47-56. [PubMed: 9046869]
- Morrissette ML, Gierut JA. Lexical organization and phonological change in treatment. *Journal of Speech, Language, and Hearing Research*. 2002; 45:143-159.
- Munson B. Phonological pattern frequency and speech production in adults and children. *Journal of Speech, Language, and Hearing Research*. 2001; 44:778-792.
- Pierrehumbert JB. Stochastic phonology. *Glott International*. 2001; 5:195-207.
- Read K, Macauley M, Furay E. The Seuss boost: Rhyme helps children retain words from shared storybook reading. *First Language*. 2014; 34:354-371.
- Rice, K.; Avery, P. Variability in a deterministic model of language acquisition: A theory of segmental elaboration. In: Archibald, J., editor. *Phonological acquisition and phonological theory*. Hillsdale, NJ: Lawrence Erlbaum; 1995. p. 23-42.
- Robbins J, Klee T. Clinical assessment of oropharyngeal motor development in young children. *Journal of Speech and Hearing Disorders*. 1987; 52:271-277. [PubMed: 3455449]
- Roid, GH.; Miller, LJ. *Leiter international performance scale-revised*. Chicago: Stoelting; 1997.
- Rvachew S, Nowak M. The effect of target-selection strategy on phonological learning. *Journal of Speech, Language, and Hearing Research*. 2001; 44:610-623.
- Savage C, Lieven E, Theakston A, Tomasello M. Testing the abstractness of young children's linguistic representations: Lexical and structural priming of syntactic constructions in young children. *Developmental Science*. 2003; 6:557-567. [PubMed: 18259588]
- Savage C, Lieven E, Theakston A, Tomasello M. Structural priming as implicit learning in language acquisition: The persistence of lexical and structural priming in 4-year-olds. *Language Learning and Development*. 2006; 2:27-49.
- Shriberg LD, Austin D, Lewis B, McSweeney J, Wilson D. The percentage of consonants correct (PCC) metric: Extensions and reliability data. *Journal of Speech, Language, and Hearing Research*. 1997; 40:708-722.
- Shriberg LD, Kwiatkowski J. Phonological disorders II: A conceptual framework for management. *Journal of Speech and Hearing Disorders*. 1982; 47:242-256. [PubMed: 7186560]
- Shriberg LD, Kwiatkowski J. Developmental phonological disorders I: A clinical profile. *Journal of Speech and Hearing Research*. 1994; 37:1100-1126. [PubMed: 7823556]
- Shriberg LD, Kwiatkowski J, Gruber FA. Developmental phonological disorders II: Short-term speech-sound normalization. *Journal of Speech and Hearing Research*. 1994; 37:1127-1150. [PubMed: 7823557]
- Shriberg LD, Lohmeier HL, Campbell TF, Dollaghan CA, Green JR, Moore CA. A nonword repetition task for speakers with misarticulations: The Syllable Repetition Task (SRT). *Journal of Speech, Language, and Hearing Research*. 2009; 52:1189-1212.
- Sosa AV, Stoel-Gammon C. Lexical and phonological effects in early word production. *Journal of Speech, Language, and Hearing Research*. 2012; 55:596-608.
- Stoel-Gammon C. Relationships between lexical and phonological development in young children. *Journal of Child Language*. 2011; 38:1-34. [PubMed: 20950495]
- Stokes SF, Bleses D, Basbøll H, Lambersten C. Statistical learning in emerging lexicons: The case of Danish. *Journal of Speech, Language, and Hearing Research*. 2012; 55:1265-1273.
- Storkel HL. Restructuring of similarity neighborhoods in the developing mental lexicon. *Journal of Child Language*. 2002; 29:251-274. [PubMed: 12109371]
- Storkel HL. Do children acquire dense neighborhoods? An investigation of similarity neighborhoods in lexical acquisition. *Applied Psycholinguistics*. 2004a; 25:201-221.

- Storkel HL. Methods for minimizing the confounding effects of word length in the analysis of phonotactic probability and neighborhood density. *Journal of Speech, Language, and Hearing Research*. 2004b; 47:1454–1468.
- Storkel HL. A corpus of consonant-vowel-consonant real words and nonwords: Comparison of phonotactic probability, neighborhood density, and consonant age of acquisition. *Behavior Research Methods*. 2013; 45:1159–1167. [PubMed: 23307574]
- Vasilyeva M, Waterfall H. Beyond syntactic priming: Evidence for activation of alternative syntactic structures. *Journal of Child Language*. 2012; 39:258–283. [PubMed: 21729366]
- Walley AC. The role of vocabulary development in children’s spoken word recognition and segmentation ability. *Developmental Review*. 1993; 13:286–350.
- Williams, KT. Expressive vocabulary test. Circle Pines, MN: American Guidance Service; 1997.
- Zamuner TS. Phonotactic probabilities at the onset of language development: Speech production and word position. *Journal of Speech, Language, and Hearing Research*. 2009; 52:49–60.
- Zamuner TS, Gerken L, Hammond M. Phonotactic probabilities in young children’s speech production. *Journal of Child Language*. 2004; 31:515–536. [PubMed: 15612388]
- Zwitserslood, P. Form priming. In: Grosjean, F.; Frauenfelder, UH., editors. *A guide to spoken word recognition paradigms*. Hove: Psychology Press; 1997. p. 589-596.




Figure 1.

Mean percent generalization gain over baseline performance on the probe for neighbor versus nonneighbor conditions in Study 1, with standard error shown. Generalization is plotted at completion of treatment in imitative and spontaneous response modes, and longitudinally after withdrawal of treatment. Baseline performance was 1.9% mean accuracy for the neighbor condition and 8.8% mean accuracy for the nonneighbor condition.




Figure 2.

Mean percent gain in PCC-R scores over baseline performance for the neighbor and nonneighbor conditions in Study 1, with standard error shown. Gain in PCC-R scores are plotted at completion of the treatment protocol and longitudinally after withdrawal of treatment. Baseline PCC-R scores are reported in Table 1.




Figure 3. Effect size for neighbor and nonneighbor conditions in Study 1, with standard error shown.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Figure 4.

Mean percent generalization gain over baseline performance on the probe for the neighbor versus nonneighbor conditions in Study 2, with standard error shown. Generalization is plotted at completion of treatment in imitative and spontaneous response modes, and longitudinally after withdrawal of treatment. Baseline performance was 4.2% mean accuracy for the neighbor condition and 4.8% mean accuracy for the nonneighbor condition.




Figure 5. Mean percent gain in PCC-R scores over baseline performance for the neighbor and nonneighbor conditions in Study 2, with standard error shown. Gain in PCC-R scores are plotted at completion of the treatment protocol and longitudinally after withdrawal of treatment. Baseline PCC-R scores are reported in Table 4.




Figure 6. Effect size for neighbor and nonneighbor conditions in Study 2, with standard error shown.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 1

Diagnostic profiles of children enrolled in Study 1, with mean performance by experimental group.

	Neighbor			Nonneighbor				
	Child 1	Child 2	Child 3	Group Mean	Child 4	Child 5	Child 6	Group Mean
Age (years;months)	4;1	4;7	3;5	4;0	4;6	3;8	3;5	3;10
GFTA-2 ^a	55	54	64	58	76	79	57	71
N sounds excluded ^b	13	10	11	11	8	7	13	9
Oral-motor structure/function ^c	24/110	24/112	24/109	24/110	24/106	23/112	23/108	23/109
Nonverbal IQ ^a	123	95	105	108	124	106	122	117
Expressive vocabulary ^a	110	101	108	106	112	100	122	111
Receptive vocabulary ^a	110	93	102	102	103	107	107	106
Nonword repetition ^d	68	41	40	50	67	64	54	62
Working memory ^e	44	30	37	37	36	33	36	35
PCC-R ^f	38	50	26	38	67	60	29	52

Note: Perceptual discrimination of pre-exposure and treated stimuli was not tested. This would have introduced a confound because a child would have been exposed to the relevant stimuli prior to the experimental manipulation.

^a Standard scores on the *Goldman-Fristoe Test of Articulation-2* (Goldman & Fristoe, 2000), *Leiter International Performance Scale-Revised* (Roid & Miller, 1997), *Expressive Vocabulary Test* (Williams, 1997), and *Peabody Picture Vocabulary Test, Third Edition* (Dunn & Dunn, 1997).

^b Sounds excluded from the phonemic inventories of all children of Study 1 were /θ d f tʃ dʒ r/. Additionally, /ŋ k g f v s z l/ were excluded within and across children's inventories.

^c Oral-motor structure/function scores (Robbins & Klee, 1987).

^d Total percent accuracy on nonword repetition task (Dollaghan & Campbell, 1998; Gathercole & Adams, 1993).

^e Scaled scores on the Auditory Sequential Memory subtest of the *Illinois Test of Psycholinguistic Abilities-Revised* (Kirk, McCarthy, & Kirk, 1968) are interpreted relative to a $M=36$ ($SD=6$).

^f Percentage of Consonants Correct-Revised (Shriberg et al., 1997).

Table 2

Characteristics of the stimuli used in production training and in exposure to neighbors versus nonneighbors.

Characteristics	Production training	Neighbor	Nonneighbor
Age-of-word-acquisition ^a	3.10	2.99	2.83
Log frequency	2.77	2.53	2.81
Sum of segment frequency ^b	.81	.45	.51
Sum of biphone frequency ^b	.40	.28	.52

Note:

^a Age-of-word-acquisition ratings on a 7-point scale (Bird, Franklin, & Howard, 2001; Gilhooly & Logie, 1980).

^b Sum of segment frequency and sum of biphone frequency values represent z-score transformations to control for word length (Storkel, 2004b).

Results of the McNemar test of significance of change associated with pre-exposure to neighbors versus nonneighbors in Study 1.

Table 3

	Performance during treatment: Treated sound in treated words		Generalization from treatment: Treated sound in untreated probe words		<i>p</i>
	Baseline	Completion	Baseline	Completion	
Neighbor	0/18	18/18	0/63	8/63	.01
Nonneighbor	0/18	18/18	0/63	0/63	1.00

Note: Production of the treated sound in treated words during treatment is on the left; generalization of the treated sound to untreated probe words is on the right. Number of accurate productions of the treated sound/total words sampled is reported at baseline and completion of the treatment protocol.

Table 4

Diagnostic profiles of children enrolled in Study 2, with mean performance by experimental group.

	Neighbor			Nonneighbor		
	Child 1	Child 2	Child 3	Child 4	Child 5	Child 6
Age (years;months)	4;9	4;6	4;2	4;6	4;0	5;5
GFTA-2 ^a	53	59	56	56	70	40
N sounds excluded ^b	12	9	12	11	10	14
Oral-motor structure/function ^c	23/112	24/109	24/111	24/111	22/110	24/112
Nonverbal IQ ^a	114	114	123	117	121	105
Expressive vocabulary ^a	99	114	109	107	108	97
Receptive vocabulary ^a	86	117	99	101	105	100
Nonword repetition ^d	52	69	55	59	63	36
Working memory ^e	31	40	30	34	36	30
PCC-R ^f	43	55	47	48	60	52
						28
						47

Note: Perceptual discrimination of post-exposure and treated stimuli was not tested. This would have introduced a confound because a child would have been exposed to the relevant stimuli prior to the experimental manipulation.

^a Standard scores on the *Goldman-Fristoe Test of Articulation-2* (Goldman & Fristoe, 2000), *Leiter International Performance Scale-Revised* (Roid & Miller, 1997), *Expressive Vocabulary Test* (Williams, 1997), and *Peabody Picture Vocabulary Test, Third Edition* (Dunn & Dunn, 1997).

^b Sounds excluded from the phonemic inventories of all children of Study 2 were /θ d f tʃ dʒ r/. Additionally, /ŋ k g f v s z l j/ were excluded within and across children's inventories.

^c Oral-motor structure/function scores (Robbins & Klee, 1987).

^d Total percent accuracy on nonword repetition task (Dollaghan & Campbell, 1998; Gathercole & Adams, 1993).

^e Scaled scores on the Auditory Sequential Memory subtest of the *Illinois Test of Psycholinguistic Abilities-Revised* (Kirk, McCarthy, & Kirk, 1968) are interpreted relative to a $M=36$ ($SD=6$).

^f Percentage of Consonants Correct-Revised (Shriberg et al., 1997).

Results of the McNemar test of significance of change associated with post-exposure to neighbors versus nonneighbors in Study 2.

Table 5

	Performance during treatment: Treated sound in treated words		Generalization from treatment: Treated sound in untreated probe words		<i>p</i>
	Baseline	Spontaneous	Baseline	Spontaneous	
Neighbor	0/18	17/18	4/63	7/63	.38
Nonneighbor	0/18	12/18	2/63	8/63	.07

Note: Production of the treated sound in treated words during treatment is on the left; generalization of the treated sound to untreated probe words is on the right. Number of accurate productions of the treated sound/total words sampled is reported at baseline and completion of the treatment protocol.