

MPRA

Munich Personal RePEc Archive

Generalisation of Samet's (2010) agreement theorem

Bassel Tarbush

23 February 2011

Online at <https://mpa.ub.uni-muenchen.de/29067/>

MPRA Paper No. 29067, posted 27 February 2011 06:58 UTC

Generalisation of Samet's (2010) agreement theorem

Bassel Tarbush⁰

February 23, 2011

Abstract We develop a framework that allows us to reproduce the generalised agreement theorem of Samet (2010), and extend it to models with a non-partitional information structure, while highlighting the features that distinguish the result from the classic theorems found in the literature. Furthermore, we produce results that are similar to Samet's with some modified assumptions.

Keywords Agreeing to disagree, knowledge, common knowledge, belief, information, epistemic logic.

JEL classification D80, D83, D89.

1 Introduction

The agreement theorem of Aumann (1976) states that if agents have a common prior on some event, then if their posteriors are common knowledge, these posteriors must be equal, *even if the agents' updates are based on different information*. This was proved for posterior probabilities in the context of a *partitional* information structure.

This result was extended by many authors to generalised decision functions, instead of posterior probabilities (see Cave (1983), Bacharach (1985), Moses and Nachum (1990), Bonanno and Nehring (1998), Aumann and Hart (2006)). However, all these generalisations have relied on the imposition of some version of the Sure-Thing Principle as a condition on the decision functions. Informally, all versions of this principle attempt to capture the following intuition: "If I would perform some action when I know that p is the case, and I would perform the same action when I know that p is not the case, then I should also perform that same action when I do not know whether p is the case".

Samet (2010) also derives an agreement theorem in a partitional information structure with generalised decision functions. However, his approach differs significantly from the classic examples in the literature in that he does not use a standard version of the Sure-Thing Principle. Rather, Samet assumes an "interpersonal" Sure-Thing Principle (*ISTP*) which can informally be stated as: "If I have some information, but I know

⁰Department of Economics, University of Oxford, [bassel.tarbush\[at\]economics.ox.ac.uk](mailto:bassel.tarbush[at]economics.ox.ac.uk)
I would like to thank John Quah for invaluable help and Francis Dennig for very useful discussions.

that whatever information I have about something, you will be better informed about it than me, then if I know your action, I should perform that same action". So, unlike the standard versions of the principle, which are conditions over the decision function of a single agent, the *ISTP* is a condition imposed on the decision functions across agents.

We develop a framework which allows us to reproduce Samet's result in a partitional information structure. However, we are also able to keep track of some more subtle features of the result. For example, we show that Samet's result allows for decision functions to be based on interactive knowledge, whereas standard results require decision functions to be independent of such information. Furthermore, we extend Samet's result to a non-partitional information structure. Partitional information imply that agents can only know what is the case; in other words, agents cannot base their decisions on false information. But surely, it is perfectly plausible for rational agents to do so. So our extension effectively states that agents cannot agree to disagree even when their decision functions can be based on interactive knowledge (or belief) *and* possibly false information. Finally, Samet's results depend on the existence of an agent who is less informed than all other agents, called the dummy. We provide agreement theorems that replace this assumption with alternative ones.

2 Epistemic Logic

This section introduces standard concepts from epistemic logic.

Definition 1 (Basic syntax). Define a finite set of atomic *propositions*, \mathcal{P} , which consists of all propositions that cannot be further reduced. Let denote N the set of all agents. We then inductively define how to formulate all other *formulas* in our language, \mathcal{L} , via the following Bachus-Naur Form:

$$\psi ::= \mathcal{P} | \neg\psi | (\psi \wedge \phi) | (\psi \vee \phi) | (\psi \rightarrow \phi) | (\psi \leftrightarrow \phi) | \Box_{i \in N} \psi | C_G \subseteq N \psi$$

Note that \Box_i and C_G are modal operators, while $\neg, \wedge, \vee, \rightarrow, \leftrightarrow$ are the standard Boolean operators.

Definition 2 (Modal depth). The modal depth $md(\psi)$ of a formula ψ is the maximal length of a nested sequence of modal operators. This can be defined by the following recursion on our syntax rules:

$$\begin{aligned} md(p) &= 0 \text{ for any } p \in \mathcal{P} \\ md(\neg\psi) &= md(\psi) \\ md(\psi \wedge \phi) &= md(\psi \vee \phi) = md(\psi \rightarrow \phi) = md(\psi \leftrightarrow \phi) = \max(md(\psi), md(\phi)) \\ md(\Box_i \psi) &= 1 + md(\psi) \\ md(C_G \psi) &= 1 + md(\psi) \end{aligned}$$

So far, we have pure uninterpreted syntax. However, we can now introduce our semantics, to determine the truth or falsity of formulas.

Definition 3 (Kripke semantics). A *frame* is a pair $\langle \Omega, R_{i \in N} \rangle$, where Ω is a finite, non-empty set of *states* (or “possible worlds”), and $R_i \subseteq \Omega \times \Omega$ is a binary relation for each agent i , also called the *accessibility relation* for agent i . A *model* on a frame $\langle \Omega, R_{i \in N} \rangle$, is a triple $\mathcal{M} = \langle \Omega, R_{i \in N}, \mathcal{V} \rangle$, where $\mathcal{V} : \mathcal{P} \times \Omega \rightarrow \{0, 1\}$ is a *valuation map*.

Definition 4 (Truth). A formula ψ is *true* at state ω in model $\mathcal{M} = \langle \Omega, R_{i \in N}, \mathcal{V} \rangle$, denoted $\mathcal{M}, \omega \models \psi$, in virtue of the following inductive clauses:

| | |
|--|--|
| $\mathcal{M}, \omega \models p$ | iff $\mathcal{V}(p, \omega) = 1$ |
| $\mathcal{M}, \omega \models \neg\psi$ | iff not $\mathcal{M}, \omega \models \psi$ |
| $\mathcal{M}, \omega \models (\psi \wedge \phi)$ | iff $\mathcal{M}, \omega \models \psi$ and $\mathcal{M}, \omega \models \phi$ |
| $\mathcal{M}, \omega \models \Box_i \psi$ | iff $\forall \omega' \in \Omega$, if $\omega R_i \omega'$ then $\mathcal{M}, \omega' \models \psi$ |
| $\mathcal{M}, \omega \models C_G \psi$ | iff $\forall \omega' \in \Omega$ accessible from ω in a finite sequence of R_i ($i \in G \subseteq N$) steps, $\mathcal{M}, \omega' \models \psi$ |

The truth of formulas involving the other Boolean operators are similarly defined. Furthermore, note that if $\mathcal{M}, \omega \models C_G \psi$, then one can generate any formula of finite modal depth of the form $\Box_i \Box_j \dots \Box_r \psi$ with $i, j, \dots, r \in G$, and this formula will be true at ω in model \mathcal{M} .¹

Definition 5 (Component). For any $\omega \in \Omega$, we will denote the set of all states that are accessible from ω in a finite sequence of R_i ($i \in G$) steps, by $\Omega_G(\omega)$. We will call this set the *component* of ω .

Definition 6 (Validity). Formula ψ is valid in a model \mathcal{M} , denoted $\mathcal{M} \models \psi$ iff $\forall \omega \in \Omega$ in \mathcal{M} , $\omega \models \psi$. Formula ψ is valid in a frame $\langle \Omega, R_{i \in N} \rangle$, denoted $\langle \Omega, R_{i \in N} \rangle \models \psi$, iff $\forall \mathcal{M}$ over $\langle \Omega, R_{i \in N} \rangle$, $\mathcal{M} \models \psi$. Formula ψ is \mathcal{T} -valid (or valid), denoted $\models \psi$, iff $\forall \langle \Omega, R_{i \in N} \rangle \in \mathcal{T}$ (\mathcal{T} , a collection of frames), $\langle \Omega, R_{i \in N} \rangle \models \psi$.

We can identify collections of frames by the restrictions that we impose on the accessibility relations.

Definition 7 (Conditions on frames). We say that a frame $\langle \Omega, R_{i \in N} \rangle$ is,

| | |
|------------|--|
| Reflexive | if $\forall i \in N, \forall \omega \in \Omega, \omega R_i \omega$ |
| Symmetric | if $\forall i \in N, \forall \omega, \omega' \in \Omega$, if $\omega R_i \omega'$ then $\omega' R_i \omega$ |
| Transitive | if $\forall i \in N, \forall \omega, \omega', \omega'' \in \Omega$, if $\omega R_i \omega'$ and $\omega' R_i \omega''$ then $\omega R_i \omega''$ |
| Euclidean | if $\forall i \in N, \forall \omega, \omega', \omega'' \in \Omega$, if $\omega R_i \omega'$ and $\omega R_i \omega''$ then $\omega' R_i \omega''$ |
| Serial | if $\forall i \in N, \forall \omega \in \Omega, \exists \omega' \in \Omega, \omega R_i \omega'$ |

¹Note that the definition of the operator C_G is drawn from van Benthem (2010), where it is also mentioned that a more precise definition can be given: One can define a new accessibility relation R_G^* for the whole group G as the reflexive transitive closure of the union of all separate relations R_i ($i \in G$), and then simply let $\mathcal{M}, \omega \models C_G \psi$ if and only if $\forall \omega' \in \Omega$, if $\omega R_G^* \omega'$ then $\mathcal{M}, \omega' \models \psi$.

The system $S5$ consists of all frames that are reflexive, symmetric and transitive; and the system $KD45$ consists of all frames that are serial, transitive and Euclidean. The following formulas are validities in the respective frames, and in fact, the systems can be axiomatised in the sense that if the validities are assumed then they imply the desired restrictions on the accessibility relations:

| $S5$ axioms | $KD45$ axioms | Axiom names |
|---|---|------------------------|
| $\Box_i(\psi \rightarrow \phi) \rightarrow (\Box_i\psi \rightarrow \Box_i\phi)$ | $\Box_i(\psi \rightarrow \phi) \rightarrow (\Box_i\psi \rightarrow \Box_i\phi)$ | Distribution |
| $\Box_i\psi \rightarrow \psi$ | $\Box_i\psi \rightarrow \neg\Box_i\neg\psi$ | Veracity; Consistency |
| $\Box_i\psi \rightarrow \Box_i\Box_i\psi$ | $\Box_i\psi \rightarrow \Box_i\Box_i\psi$ | Positive introspection |
| $\neg\Box_i\psi \rightarrow \Box_i\neg\Box_i\psi$ | $\neg\Box_i\psi \rightarrow \Box_i\neg\Box_i\psi$ | Negative introspection |

It is standard to take the axioms of $S5$ as describing properties of (a rather strong notion of) knowledge. Thus, in $S5$, $\Box_i\psi$ is interpreted as "agent i knows that ψ ". In $KD45$ however, since veracity is dropped in favour of consistency, we are in a system in which to "know" that something is the case does not imply that it is true. The axioms of $KD45$ are thus rather seen as describing properties of a belief operator, so $\Box_i\psi$ is interpreted as "agent i believes that ψ ". These two systems mirror the partitioned and non-partitioned structures mentioned in the introduction.²

Similarly, the operator $C_G\psi$ is interpreted as "it is common knowledge to all the agents in G that ψ " in $S5$, and as "it is common belief to all the agents in G that ψ " in $KD45$.

3 Models with information and decisions

Let P be the set of all propositions which can describe "facts" about a state. If P is finite, then its closure under the standard Boolean operators, denoted P^* , is tautologically finite.³ Let Ψ_0^r be the set of all formulas of modal depth 0 up to r for an arbitrary $r \in \mathbb{N}_0$. Since P^* is finite, so is Ψ_0^r , so $|\Psi_0^r| = m$, for some $m \in \mathbb{N}$; and note that $\Psi_0^0 = P^*$.⁴

Definition 8 (New operators). For each agent $i \in N$ create a set of modal operators, $O_i = \{\Box_i, \hat{\Box}_i, \check{\Box}_i\}$, where for every formula ψ , $\hat{\Box}_i\psi := \Box_i\neg\psi$ and $\check{\Box}_i\psi := \neg(\Box_i\psi \vee \hat{\Box}_i\psi)$. The interpretation, for example in $S5$, is that $\hat{\Box}_i\psi$ stands for "agent i knows that it is not the case that ψ ", and $\check{\Box}_i\psi$ stands for "agent i does not know whether it is the case that ψ ". There are similar counterpart interpretations in $KD45$.

Definition 9 (Kens). Order the set Ψ_0^r into a vector of length m : $(\psi_1, \psi_2, \dots, \psi_m)$, and for each agent $i \in N$, create the sets

$$U_i = \{(\nu_i^1\psi_1 \wedge \nu_i^2\psi_2 \wedge \dots \wedge \nu_i^m\psi_m) \mid \forall n \in \{1, \dots, m\}, \nu_i^n \in O_i\}$$

²The philosophical grounds for these systems originated in Hintikka (1962), and for an extensive formal treatment, see Chellas (1980).

³In the sense that there is only a finite number of inequivalent formulas.

⁴If $P = \{p, q\}$, then one can generate 20 inequivalent formulas: 2 from p alone, 2 from q alone and 16 out of p and q together, so $|P^*| = 20$.

$$V_i = \{\nu_i \in U_i \mid \models \neg(\nu_i \leftrightarrow (p \wedge \neg p))\}$$

So, $\nu_i \in V_i$ is a *ken* for agent i , describing i 's information concerning every formula in Ψ_0^r . So, calling $\nu_i^n \psi_n$ the n^{th} entry of i 's ken, $\nu_i^n \psi_n$ states whether i knows that the formula ψ_n is the case, or knows that it is not the case, or does not know whether it is the case. Note that V_i is a restriction of U_i to the set of kens that are not logically equivalent to a contradiction; so only the logically consistent kens are considered.

The following lemma shows that at each state, there exists a ken for each agent which holds at that state, and moreover, that any two different kens must be contradictory at any given state.

Lemma 1. (i) $\forall \omega \in \Omega, \exists \nu_i \in V_i, \omega \models \nu_i$, (ii) $\forall \omega \in \Omega, \forall \nu_i, \mu_i \in V_i$, if $\nu_i \neq \mu_i$ then $\omega \models \neg(\nu_i \wedge \mu_i)$.

By the above lemma, there is a unique ken in V_i that holds at a given state. So for any $\nu_i \in V_i$, if $\omega \models \nu_i$, we can index the ken by the state, denoting it, $\nu(\omega)_i$.

Definition 10 (Informativeness). Create an order $\succsim \subseteq V_i \times V_j$ for all $i, j \in N$. We say that the ken ν_i is *more informative* than the ken μ_j , denoted $\nu_i \succsim \mu_j$, if and only if whenever i knows that ψ then j either also knows that ψ or does not know whether ψ , and whenever i does not know whether ψ , then so does j .⁵

Note that \succsim is not a complete order on kens. For example, consider any two kens ν_i and μ_i for agent i , in which the n^{th} entry is $\nu_i^n \psi_n = \square_i \psi_n$ and $\mu_i^n \psi_n = \hat{\square}_i \psi_n$. These two kens would not be comparable with \succsim .

Finally, note that $\nu_i \sim \mu_j$ denotes $\nu_i \succsim \mu_j$ and $\mu_j \succsim \nu_i$; which is interpreted as ν_i and μ_j carrying the *same* information, but seen from the perspectives of agents i and j respectively.

Definition 11 (Decision function). For each $i \in N$, $D_i : V_i \rightarrow \mathcal{A}$, is the *decision function* of agent i , where \mathcal{A} is a set of *actions*.

Definition 12 (Action function). For all $\nu_i \in V_i$, $\models \nu_i \rightarrow d_i^{D_i(\nu_i)}$

The *action function* d_i selects the action that is actually chosen at each state.⁶ " $D_i(\nu_i) = x$ " is read as "if i 's ken is ν_i , then i 's decision is to do x ", whereas " d_i^x " is read as " i performs action x ". So although the decision function, D_i , determines what the agent *would do* over *all* possible kens, $d_i^{D_i(\nu_i)}$ is the proposition describing the agent performing the action that her decision function requires her to take given the ken she has at each particular state.⁷

⁵Formally, (i) if $\nu_i^n \psi_n = \square_i \psi_n$ then ($\mu_j^n \psi_n = \square_j \psi_n$ or $\mu_j^n \psi_n = \dot{\square}_j \psi_n$), (ii) if $\nu_i^n \psi_n = \hat{\square}_i \psi_n$ then ($\mu_j^n \psi_n = \hat{\square}_j \psi_n$ or $\mu_j^n \psi_n = \dot{\square}_j \psi_n$), and (iii) if $\nu_i^n \psi_n = \dot{\square}_i \psi_n$ then ($\mu_j^n \psi_n = \dot{\square}_j \psi_n$).

⁶Lemma 1 guarantees that the action function is well-defined.

⁷Technically, we let all propositions of the form " $D_i(\nu_i) = x$ " live in a set \mathcal{D} , and all propositions of the form " d_i^x " live in a set \mathcal{Q} . Then the set of all propositions is $\mathcal{P} = \mathcal{D} \cup \mathcal{Q}$, so the valuation function is $\mathcal{V} : \mathcal{P} \times \Omega \rightarrow \{0, 1\}$.

3.1 Main assumption

We will assume that the Interpersonal Sure-Thing Principle is a formula, $ISTP$, that is valid in every model that we will consider.

Assumption 1 (Interpersonal Sure-Thing Principle).

$$\models IntSTP := \bigwedge_{i,j \in N} \bigwedge_{\nu_i \in V_i} \bigwedge_{\nu_j \in V_j} [\Box_i(\nu_i \rightarrow \nu_j \succsim \nu_i) \rightarrow (\Box_i(d_i^x) \rightarrow d_i^x)]$$

The above states that for any agents i and j , if i knows (believes) that her having ken ν_i implies that j 's ken is more informative than hers, then if i knows (believes) that j performs action x , then i performs action x .

4 Samet's (2010) result in $S5$

In $S5$, the accessibility relation R_i is an equivalence relation for each $i \in N$. Let $I_i(\omega) = \{\omega' \in \Omega \mid \omega R_i \omega'\}$ be the *information cell* of i at ω . One can verify that the set $\mathcal{I}_i = \{I_i(\omega) \mid \omega \in \Omega\}$ is a partition of the state space Ω .

The following lemma states that at any state in which the information cell of agent i is a subset of agent j 's cell at that state, then j 's ken is more informative than i 's ken at that state.

Lemma 2. *For any $\omega \in \Omega$ such that $I_i(\omega) \subseteq I_j(\omega)$, if $\omega \models \nu_i \wedge \nu_j$ then $\omega \models \nu_i \succsim \nu_j$.*

We will require two further lemmas.

Lemma 3. $\forall i \in G, \bigcup_{\omega' \in \Omega_G(\omega)} I_i(\omega') = \Omega_G(\omega)$.

Lemma 4. *If for some $\omega' \in I_i(\omega)$, $\omega' \models \nu_i$, then for all $\omega'' \in I_i(\omega)$, $\omega'' \models \nu_i$.*

Samet (2010) assumes that there always exists an "epistemic dummy": An agent whose information cell is equal to the entire component $\Omega_G(\omega)$.

Assumption 2 (Epistemic dummy). $\exists h \in G, I_h(\omega) = \Omega_G(\omega)$.

Theorem 1. *Suppose that assumptions 1 and 2 hold, and that the system is $S5$. Let $G = \{i, j, h\}$ with h the epistemic dummy. Then, $\models C_G(d_i^x \wedge d_j^y \wedge d_h^z) \rightarrow (x = y = z)$.*

4.1 Discussion

The intuition driving the result is that by assuming that there exists an epistemic dummy, one is assuming that there is an agent h whose performed action is based on a ken that is less informative than every other agents'. However, h knows the performed actions of the other agents, and knows that those actions are based on information that is more informative than her ken. She therefore models her choice on the performed actions of

each of the other agents. But if those more informed agents were taking different actions then she would have to simultaneously copy two different actions, which is impossible, thus the actions of the more informed agents must be the same.

In Tarbush (2011) it is shown that previous agreement theorems require the assumption that decision functions only be based on kens where Ψ_0^r is such that $r = 0$. That is, decisions cannot be based on *interactive* information.⁸ So, in previous results, agents *can* agree to disagree if say i bases her decision on what she knows about what j knows. However, one of the main distinguishing features of Samet's result is that this restriction does not need to be imposed.

Furthermore, when the "Disjoint Sure-Thing Principle" is imposed on decision functions in previous results (which emulates Bacharach's (1985) original condition), the language must be assumed to be "rich" enough to guarantee that information (or kens) are, in a sense, "disjoint".⁹ The implication is that whether or not the agreement results hold depends on the way in which the states are described! However, again, Samet's result requires no such condition.

Alternatively, we can derive an agreement theorem that is similar to Samet's but that does not require the assumption of an epistemic dummy.

Definition 13. Condition A: $\models \neg C_G \neg (\nu_i \sim \mu_j)$.
Condition B: $\models \neg C_G \neg (\nu_i \succsim \mu_j) \vee \neg C_G \neg (\mu_j \succsim \nu_i)$.

Condition A states that in every component there is some state at which i and j have equally informative kens; while condition B states that in every component, either there is some state in which i is more informed than j or there is a state in which j is more informed than i .¹⁰ Clearly condition A implies condition B. However, condition A neither implies nor is implied by the existence of an epistemic dummy.

Theorem 2. *Suppose that assumption 1 and condition B hold, and that the system is S5. Let $G = \{i, j\}$. Then, $\models C_G(d_i^x \wedge d_j^y) \rightarrow (x = y)$.*

5 Samet's (2010) result in KD45

We can now analyse the consequences of using a model for *belief* rather than knowledge. So we impose a KD45 frame rather than an S5 frame.

Essentially, the only difference between knowledge and belief that we will consider is that belief is not *infallible*. In S5, agents cannot know something that is false, because reflexivity implies that if one knows that p at some state, then p must be true

⁸As explained in the paper, this is in response to the criticism (Moses and Nachum (1990)) of the like-mindedness assumption of Bacharach (1985).

⁹The language in a component $\Omega_G(\omega)$ is said to be *rich* if and only if for all $i \in G$ and any pair $(\nu_i, \mu_i) \in \{(\nu(\omega')_i, \mu(\omega'')_i) | \omega', \omega'' \in \Omega_G(\omega)\}$ there is $n \in \{1, \dots, m\}$ such that $\nu_i^n = \square_i$ and $\mu_i = \hat{\square}_i$.

¹⁰Note that Condition A is in fact condition (1.b), and condition B is implied by (2.b) in Tarbush (2011).

at that state (Veracity). On the other hand, $KD45$ allows agents to believe what is false, and thus to base decision on *false* information, by dropping reflexivity. In fact, $S5 = KD45 + \text{reflexivity}$.

We can provide a description of the links between states in a $KD45$ frame: Some sets of states within Ω are “completely connected”, in the sense that the accessibility relation over states within such sets in an equivalence relation, so these sets have the same properties as information cells in $S5$; and, for each one of these completely connected sets there exists a (possibly empty) set of “associated” states that have arrows pointing from them to every state in the completely connected set, but with no arrow (by the same agent) pointing towards them. The set of all completely connected sets and their set of associated states exhaust the state space.

Formally, let $S_i(\omega) = \{\omega' \in \Omega \mid \omega E_i \omega'\}$, where E_i is an equivalence relation. We call this set of completely connected states the *information sink* of state ω for player i . Note, that this way of defining the sink guarantees that if $S_i(\omega) \neq \emptyset$ then $\omega \in S_i(\omega)$. Furthermore, we define ω 's *set of associated states* as $A_i(\omega) = \{\omega'' \in \Omega \mid \forall \omega''' \in S_i(\omega), \omega'' F_i \omega'''\}$, where F_i is now a simple arrow. So, note that now, for any agent i , we have that $R_i = E_i \cup F_i$. Finally, we can define $J_i(\omega) = S_i(\omega) \cup A_i(\omega)$, and note that $\mathcal{J}_i = \{J_i(\omega) \mid \omega \in \Omega\}$ exhausts the entire state space.

Proposition 1. *The above is a complete characterisation of the $KD45$ state space.*

We now use lemmas that are analogous to the ones used in $S5$.

Lemma 5. *For any $\omega \in \Omega$ such that $S_i(\omega) \subseteq S_j(\omega)$, if $\omega \models \nu_i \wedge \nu_j$ then $\omega \models \nu_i \succsim \nu_j$.*

Lemma 6. $\forall i \in G, \bigcup_{\omega' \in \Omega_G(\omega)} S_i(\omega') \subseteq \Omega_G(\omega) \subseteq \bigcup_{\omega' \in \Omega_G(\omega)} J_i(\omega')$.

Lemma 7. *If for some $\omega' \in J_i(\omega)$, $\omega' \models \nu_i$, then for all $\omega'' \in J_i(\omega)$, $\omega'' \models \nu_i$.*

We now require an assumption that is analogous to the epistemic dummy assumption.

Assumption 3 (Doxastic dummy). $\exists h \in G, \exists \omega' \in \Omega_G(\omega), \bigcup_{i \in G} \bigcup_{\omega'' \in \Omega_G(\omega)} S_i(\omega'') \subseteq S_h(\omega')$ and $J_h(\omega') = \Omega_G(\omega) \cup \{\omega\}$.

This assumption requires that some agent's (the dummy's) unique information sink be a superset of the union of the information sinks of every other agent in the component.

Theorem 3. *Suppose that assumptions 1 and 3 hold, and that the system is $KD45$. Let $G = \{i, j, h\}$ with h the doxastic dummy. Then, $\models C_G(d_i^x \wedge d_j^y \wedge d_h^z) \rightarrow (x = y = z)$.*

5.1 Discussion

The only substantial difference between theorem in $S5$ and the one is $KD45$ is the assumption made about the dummy. Note that the epistemic dummy assumption in $S5$ could have been stated as follows: $\exists h \in G, \bigcup_{i \in G} \bigcup_{\omega' \in \Omega_G(\omega)} I_i(\omega) \subseteq I_h(\omega)$ and




Figure 1: $\mathcal{M}' = \mathcal{M}$ plus dummy h

$I_h(\omega) = \Omega_G(\omega)$.¹¹ This provides a sense of how the doxastic dummy assumption is more general: Since the kens of any agent anywhere in the component essentially only depend on the kens in the sinks of that agent, it is enough for the dummy's sink to be a superset of the union of the sink of all other agents for the dummy to become the least informed agent.

Note that a different assumption could have been: $\exists h \in G, S_h(\omega) = \Omega_G(\omega) \cup \{\omega\}$. One can verify that this *implies* the doxastic dummy assumption. However, we see it as being unreasonably strong: It implies that if the "actual" state ω is not in the sink of any of the agents other than the dummy's, then it must at least be in the dummy's sink. In such a case, the dummy would be somewhat of a "wise fool" in the sense that all other agents would be deeming ω impossible, whereas the dummy does not rule out *any* possibility, including ω itself. This implication does not necessarily hold when the doxastic dummy assumption is taken as it is originally stated.

One rather worrying feature of Theorem 3, however, can be illustrated by the following example. Consider model \mathcal{M} in Figure 1 with $\omega \models p$ and $\omega' \models \neg p$. At every state of this model, i believes that $\neg p$ and at every state, j believes that p . In this model, the condition of "heterogeneity" fails, so all the agreement theorems mentioned in the introduction would concede that i and j can agree to disagree (see Tarbush (2011)).¹² Now, consider adding an epistemic dummy h to this model, to obtain model \mathcal{M}' . Heterogeneity would again fail, so the agents can again agree to disagree according to all the agreement theorems other than Samet's. However, according to Theorem 3, the agents cannot agree to disagree. But what drives the result in this case?

Agent i must surely perform his action as though he were certain that $\neg p$ is the case, since $\neg p$ is the only proposition that i believes, regardless of the state. Similarly, agent j must surely perform her action as though she were certain that p is the case. However, by the presence of h , the agents i and j must perform the same action. So the existence of the dummy must collapse the action that one would perform when p and when $\neg p$ to the *same* action.

One can interpret this in one of two ways: (i) The existence of the dummy can be seen as a constraint on the decision functions, requiring them to be independent of one's information regarding p . But this then makes agreement trivial. Or, (ii) the decision functions do depend on p , but the existence of the dummy implies that the more informed agents

¹¹Of course, by Lemma 3, this is equivalent to $\exists h \in G, I_h(\omega) = \Omega_G(\omega)$.

¹²Here, if $G = \{i, j, h\}$, heterogeneity can be stated as: $\models C_G(\nu_i \wedge \nu_j \wedge \nu_h) \rightarrow (\nu_i \sim \nu_j \sim \nu_h)$.

must nevertheless perform the same action. However, this must be the action that the agents would perform when they do not “know” whether p is true, *even though*, in this example, the more informed agents are effectively *certain* of their information regarding p .

As before, we can provide a further theorem without a doxastic dummy.

Definition 14. Condition C: For all $\omega \in \Omega$, there exists $\omega' \in \Omega_G(\omega)$ such that $S_i(\omega) = S_j(\omega)$.

Condition D: For all $\omega \in \Omega$, there exists $i \in G$ such that for some $\omega' \in \Omega_G(\omega)$, $S_i(\omega') \subseteq S_j(\omega')$.

One can see that conditions C and D are effectively the semantic counterparts of conditions A and B respectively.¹³ Note that C implies D, and that by Lemma 5, C implies A while D implies B.

Theorem 4. *Suppose that assumption 1 and condition D hold, and that the system is KD45. Let $G = \{i, j\}$. Then, $\models C_G(d_i^x \wedge d_j^y) \rightarrow (x = y)$.*

Note that this theorem would still imply agreement in model \mathcal{M}' represented in Figure 1. However, if we had assumed the stronger condition C then such a case would be ruled out.

¹³Note that Condition C is condition (1.a) and condition D is implied by condition (2.a) in Tarbush (2011).

Appendix

Proof of Lemma 1 (i) Consider an arbitrary $i \in N$ and $\omega \in \Omega$, and suppose that $\omega \models \psi$, for some formula $\psi \in \Psi_0^r$. It must be the case that either (i.a) $\forall \omega' \in \Omega$, if $\omega R_i \omega'$ then $\omega' \models \psi$, or (i.b) $\forall \omega' \in \Omega$, if $\omega R_i \omega'$ then $\omega' \models \neg \psi$, or (i.c) $\exists \omega', \omega'' \in \Omega$, such that $\omega R_i \omega'$ and $\omega R_i \omega''$, and $\omega' \models \psi$ and $\omega'' \models \neg \psi$ (i.e. neither (i.a) nor (i.b)).

If (i.a) is the case, then $\omega \models \Box_i \psi$. If (i.b) is the case, then $\omega \models \hat{\Box}_i \psi$, and finally, if (i.c) is the case, then $\omega \models \dot{\Box}_i \psi$. Therefore, in all cases, the operator over ψ belongs to the set O_i , and since this holds for any $\psi \in \Psi_0^r$, it holds for each entry of a ken. Furthermore, \models can only generate consistent lists of formulas, so kens cannot be inconsistent. This implies that a ken must exist that belongs to V_i .

(ii) Consider an arbitrary $i \in N$ and $\omega \in \Omega$. Let $\nu_i, \mu_i \in V_i$, and consider the n^{th} entry of each ken such that $\nu_i^n \psi_n \neq \mu_i^n \psi_n$. Case (ii.a): Suppose $\omega \models \nu_i^n \psi_n = \Box_i \psi_n$. So, $\forall \omega' \in \Omega$, if $\omega R_i \omega'$, then $\omega' \models \psi_n$. By definition, this rules out the possibility that also, $\omega \models \hat{\Box}_i \psi_n$, or $\omega \models \dot{\Box}_i \psi_n$. For cases (ii.b), $\omega \models \nu_i^n \psi_n = \hat{\Box}_i \psi_n$, and (ii.c), $\omega \models \nu_i^n \psi_n = \dot{\Box}_i \psi_n$, proceed analogously to (ii.a).

Proof of Lemma 2 Consider some arbitrary state $\omega \in \Omega$. Suppose $I_i(\omega) \subseteq I_j(\omega)$ and $\omega \models \nu_i \wedge \nu_j$. Consider the n^{th} entry of these kens.

(a) Suppose $\omega \models \nu_i^n \psi_n = \Box_i \psi_n$, and suppose that $\omega \models \nu_j^n \psi_n = \hat{\Box}_j \psi_n$. Then, $\forall \omega' \in I_j(\omega)$, $\omega' \models \neg \psi_n$. But if $I_i(\omega) \subseteq I_j(\omega)$, then $\forall \omega' \in I_i(\omega)$, $\omega' \models \neg \psi_n$, which contradicts the statement that $\omega \models \Box_i \psi_n$. Therefore, $\omega \models (\nu_j^n \psi_n = \Box_j \psi_n \vee \nu_j^n \psi_n = \dot{\Box}_j \psi_n)$.

Cases (b), $\omega \models \nu_i^n \psi_n = \hat{\Box}_i \psi_n$ and (c) $\omega \models \nu_i^n \psi_n = \dot{\Box}_i \psi_n$ can be dealt with analogously to case (a).

Proof of Lemma 3 Suppose $\omega'' \in \bigcup_{\omega' \in \Omega_G(\omega)} I_i(\omega')$. So, $\omega'' \in I_i(\omega')$ for some $\omega' \in \Omega_G(\omega)$. But, $\omega' R_i \omega''$, and there exists a sequence of R_i ($i \in G$) steps such that ω' is reachable from ω . Therefore, there exists a sequence, one step longer, such that ω'' is reachable from ω . So, $\omega'' \in \Omega_G(\omega)$. (And, note that $I_i(\omega'') \subseteq \Omega_G(\omega)$).

Suppose $\omega'' \in \Omega_G(\omega)$. Reflexivity guarantees that $\omega'' \in I_i(\omega'')$. So, for some $\omega^* \in \Omega_G(\omega)$, $\omega'' \in I_i(\omega^*)$, so $\omega'' \in \bigcup_{\omega' \in \Omega_G(\omega)} I_i(\omega')$.

Proof of Lemma 4 Suppose $\omega' \models \nu_i$ for some $\omega' \in I_i(\omega)$. Consider the n^{th} entry of the ken, namely, $\nu_i^n \psi_n$.

(a) Suppose $\omega' \models \nu_i^n \psi_n = \Box_i \psi_n$. Then, for all $\omega'' \in \Omega$, $\omega' R_i \omega''$ implies $\omega'' \models \psi_n$. So, for all $\omega'' \in I_i(\omega')$, $\omega'' \models \psi_n$. But since R_i is an equivalence relation, and $\omega' \in I_i(\omega)$, it follows that $I_i(\omega') = I_i(\omega)$. So, for all $\omega'' \in I_i(\omega)$, $\omega'' \models \psi_n$, from which it follows that for all $\omega'' \in I_i(\omega)$, $\omega'' \models \Box_i \psi_n$.

Case (b), $\omega' \models \nu_i^n \psi_n = \hat{\Box}_i \psi_n$ and (c), $\omega' \models \nu_i^n \psi_n = \dot{\Box}_i \psi_n$ are analogous to case (a).

Proof of Theorem 1 Suppose that assumptions 1 and 2 hold, and that the system is $S5$. Let $\omega \in \Omega$, and consider the set $\Omega_G(\omega)$. It must be the case that at ω , $\omega \models \nu(\omega)_h$. So by Lemma 4 and Assumption 2, for all $\omega' \in \Omega_G(\omega)$, $\omega' \models \nu(\omega)_h$. By Lemma 3, we

know that $\bigcup_{\omega' \in \Omega_G(\omega)} I_i(\omega') = \Omega_G(\omega)$. So for each state in each of i 's information cells, and therefore for each $\omega'' \in \Omega_G(\omega)$, it must be the case that $\omega'' \models \nu(\omega'')_i \succsim \nu(\omega)_h$ by Lemma 2. It follows that for all $\omega'' \in \Omega_G(\omega)$, $\omega'' \models \nu(\omega)_h \rightarrow \nu(\omega'')_i \succsim \nu(\omega)_h$ and $\omega'' \models \Box_h(\nu(\omega)_h \rightarrow \nu(\omega'')_i) \succsim \nu(\omega)_h$.

Finally, by the assumption that $\omega \models C_G(d_i^x)$, it follows that $\omega \models \Box_h(d_i^x)$. By Assumption 1, it follows that $\omega \models d_h^x$.

Reasoning similarly, between the dummy h and agent j , we find that $\omega \models d_h^y$. Therefore $\omega \models (x = y)$.

Proof of Theorem 2 Suppose that $\omega \models C_G(d_i^x \wedge d_j^y) \wedge (x \neq y)$. If condition B holds, then without loss of generality, there is some state $\omega' \in \Omega_G(\omega)$ such that $\omega' \models \nu(\omega')_i \succsim \mu(\omega')_j$. Now by the assumption of common knowledge of actions, it must be the case that for all $\omega'' \in \Omega_G(\omega)$, $\omega'' \models \Box_i d_j^y \wedge d_i^x$. By *ISTP*, it follows that $\omega'' \models \neg \Box_i(\nu(\omega'')_i \rightarrow \mu(\omega'')_j) \succsim \nu(\omega'')_i$. So, for every $\omega'' \in \Omega_G(\omega)$, there exists an $\omega''' \in \Omega_G(\omega)$ with $\omega'' R_i \omega'''$ such that $\omega''' \models \nu(\omega''')_i \wedge \neg(\mu(\omega''')_j \succsim \nu(\omega''')_i)$. But since $\omega' R_i \omega''$ and $\omega' \models \nu(\omega')_i \succsim \mu(\omega')_j$, we have a contradiction.

Proof of Proposition 1 Let “ i -arrow” refer to an arrow of i 's accessibility relation. Firstly, we can show that $R_i = E_i \cup F_i$. An arbitrary $\omega \in \Omega$ either has an i -arrow pointing to it or it does not. If it does not, by seriality, it points to another state. If it does, then there exists a state ω' that points to ω which itself points to some state ω'' by seriality. Transitivity implies that ω' points to ω'' and Euclideaness implies that ω'' points to ω . From here it is easy to prove that ω, ω' and ω'' are in an equivalence class.

Secondly, we show that if $J_i(\omega') \neq J_i(\omega'')$ then $J_i(\omega') \cap J_i(\omega'') = \emptyset$. Suppose $\omega \in J_i(\omega') \cap J_i(\omega'')$. If $\omega \in S_i(\omega') \cap S_i(\omega'')$ then $S_i(\omega')$ and $S_i(\omega'')$ are indistinguishable, and one can verify that $J_i(\omega') = J_i(\omega'')$. If $\omega \in S_i(\omega') \cap A_i(\omega'')$ then ω both does have and does not have an i -arrow pointing to it. Finally, if $\omega \in A_i(\omega') \cap A_i(\omega'')$ then by Euclideaness, ω' and ω'' are indistinguishable, and $J_i(\omega') = J_i(\omega'')$.

Thirdly, we can show that $\bigcup_{\omega \in \Omega} J_i(\omega) = \Omega$. Suppose $\omega' \in \bigcup_{\omega \in \Omega} J_i(\omega)$, then by the definitions of S_i and A_i , $\omega' \in \Omega$. On the other hand, suppose $\omega \in \Omega$. Then if there is an i -arrow pointing to ω , $\omega \in S_i(\omega) \subseteq J_i(\omega)$. If there is no i -arrow pointing to it, then by seriality, there is an ω' that ω points to, so $\omega \in A_i(\omega') \subseteq J_i(\omega')$. So, $\omega \in \bigcup_{\omega \in \Omega} J_i(\omega)$.

Proof of Lemma 5 Entirely analogous to the proof of Lemma 2.

Proof of Lemma 6 Suppose $\omega'' \in \bigcup_{\omega' \in \Omega_G(\omega)} S_i(\omega')$. So, $\omega'' \in S_i(\omega')$ for some $\omega' \in \Omega_G(\omega)$. But, $\omega' E_i \omega''$, and there exists a sequence of R_i ($i \in G$) steps such that ω' is reachable from ω . Therefore, there exists a sequence, one step longer, such that ω'' is reachable from ω . So, $\omega'' \in \Omega_G(\omega)$.

Suppose $\omega'' \in \Omega_G(\omega)$. Either ω'' has an i -arrow pointing towards it, in which case $\omega'' \in S_i(\omega'')$. So, $\omega'' \in S_i(\omega'') \cup A_i(\omega'') = J_i(\omega'')$, or, ω'' has no i -arrow pointing towards it, in which case, by seriality, there exists some ω''' such that $\omega'' \in A_i(\omega''')$. Note that ω''' must be in $\Omega_G(\omega)$ since it is reachable from ω'' . So, $\omega'' \in S_i(\omega''') \cup A_i(\omega''') = J_i(\omega''')$.

In either case, for some $\omega^* \in \Omega_G(\omega)$, $\omega'' \in J_i(\omega^*)$, so $\omega'' \in \bigcup_{\omega' \in \Omega_G(\omega)} J_i(\omega')$.

Proof of Lemma 7 Suppose $\omega' \models \nu_i$ for some $\omega' \in J_i(\omega)$. Firstly, suppose $\omega' \in S_i(\omega)$, and consider the n^{th} entry of the ken, namely, $\nu_i^n \psi_n$.

(a) Suppose $\omega' \models \nu_i^n \psi_n = \Box_i \psi_n$. Then, for all $\omega'' \in \Omega$, $\omega' E_i \omega''$ implies $\omega'' \models \psi_n$. So, for all $\omega'' \in S_i(\omega')$, $\omega'' \models \psi_n$. But since E_i is an equivalence relation, and $\omega' \in S_i(\omega)$, it follows that $S_i(\omega') = S_i(\omega)$. So, for all $\omega'' \in S_i(\omega)$, $\omega'' \models \psi_n$, from which it follows that for all $\omega'' \in S_i(\omega)$, $\omega'' \models \Box_i \psi_n$. Also, each $\omega''' \in A_i(\omega)$ has an arrow pointing to each state in $S_i(\omega)$, so for all $\omega^* \in S_i(\omega)$, if $\omega''' F_i \omega^*$, $\omega^* \models \psi_n$. So, for all $\omega''' \in A_i(\omega)$, $\omega''' \models \Box_i \psi_n$. It follows that for all $\omega'' \in J_i(\omega)$, $\omega'' \models \Box_i \psi_n$.

Case (b), $\omega' \models \nu_i^n \psi_n = \hat{\Box}_i \psi_n$ and (c), $\omega' \models \nu_i^n \psi_n = \check{\Box}_i \psi_n$ are analogous to case (a).

Now, suppose $\omega' \in A_i(\omega)$, and consider the n^{th} entry of the ken, namely, $\nu_i^n \psi_n$.

(d) Suppose $\omega' \models \nu_i^n \psi_n = \Box_i \psi_n$. Then, for all $\omega'' \in \Omega$, $\omega' F_i \omega''$ implies $\omega'' \models \psi_n$. So, for all $\omega'' \in S_i(\omega')$, $\omega'' \models \psi_n$. This implies that $\omega'' \models \Box_i \psi_n$ for all $\omega'' \in S_i(\omega)$, and $\omega''' \models \Box_i \psi_n$ for all other states $\omega''' \in A_i(\omega)$. It follows that for all $\omega'' \in J_i(\omega)$, $\omega'' \models \Box_i \psi_n$.

Case (e), $\omega' \models \nu_i^n \psi_n = \hat{\Box}_i \psi_n$ and (f), $\omega' \models \nu_i^n \psi_n = \check{\Box}_i \psi_n$ are analogous to case (d).

Proof of Theorem 3 Suppose that assumptions 1 and 3 hold, and that the system is *KD45*. Let $\omega \in \Omega$, and consider the set $\Omega_G(\omega)$. It must be the case that at ω , $\omega \models \nu(\omega)_h$. So by Lemma 7 and Assumption 3, for all $\omega' \in \Omega_G(\omega) \cup \{\omega\}$, $\omega' \models \nu(\omega)_h$. By Lemma 6, we know that $\bigcup_{\omega' \in \Omega_G(\omega)} S_i(\omega') \subseteq \Omega_G(\omega)$. So for each state ω'' in each of i 's information sinks, it must be the case that $\omega'' \models \nu(\omega'')_i \succsim \nu(\omega)_h$ by Lemma 5. However, this must also be true at every state ω''' that is in the component but not in any of i 's sinks (by Lemma 7). So, for all $\omega'' \in \Omega_G(\omega) \cup \{\omega\}$, $\omega'' \models \nu(\omega'')_i \succsim \nu(\omega)_h$. It follows that for all $\omega'' \in \Omega_G(\omega) \cup \{\omega\}$, $\omega'' \models \nu(\omega)_h \rightarrow \nu(\omega'')_i \succsim \nu(\omega)_h$ and $\omega'' \models \Box_h(\nu(\omega)_h \rightarrow \nu(\omega'')_i) \succsim \nu(\omega)_h$.

Finally, by the assumption that $\omega \models C_G(d_i^x)$, it follows that $\omega \models \Box_h(d_i^x)$. By Assumption 1, it follows that $\omega \models d_h^x$.

Reasoning similarly, between the dummy h and agent j , we find that $\omega \models d_h^y$. Therefore $\omega \models (x = y)$.

Proof of Theorem 4 Entirely analogous to the proof of Theorem 2.

References

- Aumann, R. J.: 1976, 'Agreeing to disagree'. *The annals of statistics* **4**(6), 1236–1239.
- Aumann, R. J. and S. Hart: 2006, 'Agreeing on decisions, The Hebrew University of Jerusalem'. *Center for Rationality, unpublished DP*.
- Bacharach, M.: 1985, 'Some extensions of a claim of Aumann in an axiomatic model of knowledge'. *Journal of Economic Theory* **37**(1), 167–190.

- Bonanno, G. and K. Nehring: 1998, 'Assessing the Truth Axiom under incomplete information'. *Mathematical social sciences* **36**(1), 3–29.
- Cave, J. A.: 1983, 'Learning to agree'. *Economics Letters* **12**(2), 147–152.
- Chellas, B. F.: 1980, *Modal logic*. Cambridge University Press Cambridge.
- Hintikka, J.: 1962, *Knowledge and belief: an introduction to the logic of the two notions*. Cornell University Press.
- Moses, Y. and G. Nachum: 1990, 'Agreeing to disagree after all'. In: *Proceedings of the 3rd conference on Theoretical aspects of reasoning about knowledge*. pp. 151–168.
- Samet, D.: 2010, 'Agreeing to disagree: The non-probabilistic case'. *Games and Economic Behavior* **69**(1), 169–174.
- Tarbush, B.: 2011, 'Agreeing to disagree with generalised decision functions'. *University of Oxford, Department of Economics, Working Paper*.
- van Benthem, J.: 2010, 'Modal Logic for Open Minds'. *Center for the Study of Language and Information*.