



Munich Personal RePEc Archive

Multiple bounded discrete choice contingent valuation: parametric and nonparametric welfare estimation and a comparison to the payment card

Christian A. Vossler

Department of Economics, University of Tennessee

2003

Online at <https://mpra.ub.uni-muenchen.de/38867/>

MPRA Paper No. 38867, posted 18. May 2012 12:47 UTC

MULTIPLE BOUNDED DISCRETE CHOICE CONTINGENT VALUATION: PARAMETRIC AND NONPARAMETRIC WELFARE ESTIMATION AND A COMPARISON TO THE PAYMENT CARD

Abstract

In multiple bounded discrete choice (MBDC) surveys, respondents indicate how certain they would be to vote in favor of a policy at different prices by choosing, for example, among “definitely yes”, “probably yes”, “not sure”, “probably no”, and “definitely no” response options for each price. In estimating non-market values from MBDC data, past researchers have made markedly different assumptions with respect to the assumed correlation of within-respondent decisions (one for each price) and the correspondence of stated payment certainty to actual behavioral intentions. The first objective of this paper is to provide guidance for future research efforts by discriminating between existing models and proposing new estimators that relax some important statistical assumptions of existing models. Contrary to a previous study, results in this paper suggest that within-respondent decisions should be treated as being perfectly correlated. The second objective is to examine whether it is worthwhile to collect the additional information on payment certainty, as it may place additional cognitive burden on respondents as well as data analysts. Using data from previous studies, MBDC is compared with the payment card, a related elicitation approach that does not gauge payment certainty. This comparison provides strong and systematic evidence that “definitely yes” and “probably yes” MBDC respondents would vote “yes” while other respondents would vote “no” in the absence of the certainty categories.

1. Introduction

Over three decades after its introduction, contingent valuation (CV) remains a popular method for estimating the willingness to pay (WTP) for nonmarket goods, especially those with a nonuse value component. Accepted practice is to ask a single dichotomous choice (DC) valuation question, to which respondents simply answer yes or no as to whether they are willing to pay the offered price for the nonmarket good. The method is favored on cognitive grounds as answering the valuation question is similar to voting in an election or making a market purchase. When properly framed, DC questions also have desirable incentive properties (see Carson, Groves, and Machina 2000). However, DC is notorious for extracting minimal information on the respondent's WTP. Further, researchers caution against assuming respondents are absolutely certain of their yes or no decision. It is realistic to presume, for example, that some respondents are indifferent to a yes or no vote (Opaluch and Segerson 1989) and that subjects have an inherent (and perhaps predictable) randomness in their preferences (Li and Mattsson 1995).

In order to address concerns about the accepted practice, a laundry list of alternative elicitation mechanisms have been proposed in attempt to gather more information on the respondent's WTP. The double-bounded DC elicitation format and its variants elicit yes/no answers to a sequence of two or more payment amounts (e.g., Cameron and Quiggin 1994). Champ et al. (1997), Johannesson et al. (1999), and others, ask "yes" DC respondents to indicate how certain they are that they would be willing to pay the stated price. Other researchers (e.g., Ready, Whitehead, and Blomquist 1995; Vossler and Kerkvliet 2003) incorporate payment certainty directly into the valuation question, rather than in a follow-up question. Because of strong arguments against open-ended questions (see Mitchell and Carson 1987), recent mechanisms predominantly include discrete choice valuation questions.

The multiple bounded discrete choice (MBDC) format, introduced in the recent literature by Welsh and Poe (1998), is the only format that asks the respondent about multiple payment amounts while collecting information on payment certainty. Specifically, for each price, the respondent indicates how certain she would be to vote in favor of a policy by choosing, for example, among “definitely yes”, “probably yes”, “not sure”, “probably no”, and “definitely no” response options. As such, within the realm of discrete choice valuation formats one can regard the MBDC approach as rather extreme in terms of the amount of information it attempts to elicit on the respondent’s WTP.

Since the MBDC approach is still in its infancy, important issues such as incentive compatibility and bid design effects are not yet fully explored.¹ However, along with the potential wealth of information collected from them, there are good reasons for considering MBDC questionnaires. First, use of MBDC questionnaires lessens the burden of optimal bid design endemic to single and double-bounded DC since they present each respondent with a full range of possible payment amounts (Alberini 1993; Kanninen 1995). Second, the double-bounded DC elicitation format and its variants present respondents with (unannounced) follow-up bids based on previous responses. A wealth of research suggests that anchoring, resentment, or other unintended response effects force the respondent’s underlying WTP distribution to shift between initial and follow-up DC valuation questions (Cameron and Quiggin 1994; Herriges and Shogren 1996; Alberini, Kanninen, and Carson 1997; Bateman et al. 2001; Whitehead 2002). Bateman et al. (in press) demonstrate that changing the choice set visible to respondents as they progress along a valuation exercise leads to anomalous behavior. In contrast, respondents behave consistently with economic theory when the entire choice set is visible throughout the experiment. Drawing from these results, it is

¹ Vossler et al. (2003c) provide an overview of findings to date.

unlikely that MBDC questionnaires are plagued with unintended response effects as they present respondents with all payment amounts at the same time.

While there is yet a solid premise for endorsing the MBDC format, given its potential advantages and the increasing number of field applications that are using this approach, further investigation is warranted. The main objectives of this paper are to thoroughly examine the gamut of statistical methods for analyzing MBDC survey data and to compare the MBDC approach with the payment card (PC), which can be regarded as a special case of the MBDC questionnaire that does not collect information on payment certainty. To meet these objectives, this study makes use of data from three existing studies that use both elicitation formats.

Given the breadth of elicited information, it is not surprising that several approaches for estimating WTP from MBDC responses already appear in the literature (Welsh and Poe 1998; Cameron et al. 2002; Alberini, Boyle, and Welsh 2003; Evans, Flores, and Boyle 2003). However, these analytical methods are somewhat at odds as important statistical and behavioral assumptions underlying them differ considerably. While the behavioral intention of the respondent is an open empirical question, the objective here is to provide guidance as to the appropriateness of underlying statistical assumptions and to add attractive estimators to the analyst's choice set.

In an effort to explore the impact of adding a certainty dimension, MBDC is compared with the PC, which also presents the respondent with a large set of potential payment amounts, but does not collect information on payment certainty.² If MBDC responses can be predictably parceled into yes and no decisions, then the added complexity of the MBDC approach arguably yields little benefit. Indeed, to the extent that respondents have cognitive difficulty with the MBDC format, the much simpler PC may be preferred *a priori*.

² While some PCs simply ask the respondent to circle their maximum WTP from a set of possible choices, other PC applications indeed elicit a yes or no decision for each payment amount.

The next section of the paper overviews existing methods for analyzing MBDC data, and raises important issues with respect to the interpretation and modeling of responses from this method. Section 3 introduces and provides motivation for alternative MBDC models. Section 4 briefly describes the data. Statistical assumptions and model performance is examined in Section 5. Section 6 presents comparisons of MBDC and PC WTP distributions. Concluding remarks appear in Section 7.

2. Review of Existing Methods for Analyzing MBDC Data

Four models for estimating non-market values from MBDC data appear in the recent literature: the Welsh-Poe interval model (Welsh and Poe 1998); the Dual Uncertainty Decision Estimator (DUDE) of Evans, Flores, and Boyle (2003); the binary choice random effects model (Alberini, Boyle and Welsh 2003); and the ordered choice model (Cameron et al. 2002; Alberini, Boyle, and Welsh 2003). These models are distinguishable in terms of (1) the assumed correlation between decisions from the same individual, and (2) the correspondence between the respondent's level of payment certainty and her assumed behavioral intentions.

Welsh and Poe (1998) recode categorical responses into simple yes/no decisions and model the resulting interval that bounds the respondent's WTP, employing the interval data model commonly used for analyzing PC (Cameron and Huppert 1989) and double-bounded DC (Hanemann, Loomis, and Kanninen 1991) data. Welsh and Poe (1998) estimate "definitely yes", "probably yes", and "not sure" versions of the model. The name of the model refers to the lowest certainty level recoded as yes. Implicit in this approach is that one WTP distribution underlies the respondent's multiple decisions: i.e., there is perfect response correlation.

Evans, Flores and Boyle (2003) develop the Dual Uncertainty Decision Estimator (DUDE), which extends the Welsh-Poe interval model by linking categorical responses to subjective payment probabilities. Similar to the Welsh-Poe model, a single

WTP distribution underlies within-subject responses. However, as an alternative to the simple yes/no recoding of responses, probability weights are assigned to payment certainty levels. The Welsh-Poe interval model represents a special case of the DUDE estimator where recoded yes responses receive a probability weight of 1.0 while no responses receive a weight of 0.0.

Alberini, Boyle, and Welsh (2003) estimate a binary choice probit model with random effects. Identical to the Welsh-Poe “probably yes” model, Alberini, Boyle, and Welsh (2003) treat “definitely yes” and “probably yes” as yes, and other responses as no. However, rather than using the yes/no responses to define the respondent’s WTP interval, each yes/no decision becomes a separate observation within a panel data framework. The multiple observations from the same individual are considered random draws from separate, but correlated WTP distributions: i.e., responses are freely correlated. A similar assumption underlies the bivariate probit model used for double-bounded CV data (Cameron and Quiggin 1994).

Cameron et al. (2002) and Alberini, Boyle, and Welsh (2003) estimate ordered logit and probit models, respectively. The ordered models treat the certainty categories as ordered response propensities and, from the data, estimate threshold parameters that define where respondents switch between certainty levels.³ As in the random effects probit, an observation is created from each of the respondent’s decisions. However, an underlying model assumption of Alberini, Boyle, and Welsh (2003) is that each within-respondent decision is a random draw from an *independent* WTP distribution – as though different respondents provided them. Cameron et al. (2002) realize the likely correlation between within-respondent decisions and, as a compromise, they weight each observation such that the effective sample size is equal to the number of respondents.

³ Alberini, Boyle, and Welsh (2003) let the thresholds be a linear function of respondent attributes, following the approach of Wang (1997). This formulation does not affect WTP in their study.

Table 1 summarizes the underlying assumptions of existing MBDC models. The log-likelihood functions for these models are included as Appendix A. I now outline the four research topics explored in this paper using data from three published studies.

Table 1. Overview of Existing MBDC Models

Model (citation)	Correlation of within-respondent decisions	Correspondence between payment certainty categories and behavioral intentions
Welsh-Poe (Welsh and Poe 1998)	Perfect correlation	Categorical responses recorded as simple yes and no decisions
DUDE (Evans, Flores, and Boyle 2003)	Perfect correlation	Probability weights assigned to categories
Binary choice random effects (Alberini, Boyle, and Welsh 2003)	Responses are freely correlated	Categorical responses recorded as simple yes and no decisions
Ordered choice (Cameron et al. 2002; Alberini, Boyle, and Welsh 2003)	Responses are independent	Certainty categories treated as ordered choices. Assumed that the propensity to vote yes switches from negative to positive within the “not sure” category

A. Assumptions regarding within-respondent decisions

MBDC respondents choose among the payment certainty categories for several possible payment amounts. Existing models assume that these within-respondent decisions are perfectly correlated, completely independent, or freely correlated (i.e., something between independence and perfect correlation). A natural research question is which assumption is appropriate. Evidence from double-bounded DC questionnaires suggests that WTP distributions of initial and follow-up DC responses are highly

correlated, but can be statistically different due to widely cited response-effects (see Cameron and Quiggin 1994). It is unclear whether this finding extends to MBDC data, as respondents see all payment amounts at the same time, rather than in an iterative manner.

Unlike double-bounded DC, there is no obvious way to test for shifts in the respondent's underlying WTP distribution across valuation questions. That is to say, the price associated with each decision does not vary across respondents and so one cannot estimate separate WTP functions for each of the respondent's decisions and then test for equality of these WTP functions. As an alternative approach, Alberini, Boyle, and Welsh (2003) test whether the correlation coefficient (ρ) in a binary choice random effects (probit) model is statistically different from zero. Similar to the bivariate probit model used for double-bounded DC data, the binary choice random effects model allows within-respondent decisions to be freely correlated and the correlation coefficient measures the degree of correlation. Failure to reject the hypothesis that the correlation coefficient is statistically different from zero lends support for treating within-subject decisions as stemming from independent WTP distributions. Although there is no reliable test of $\rho = 1$ (see Alberini 1995), a correlation coefficient near 1 provides evidence that within-subject responses can be treated as perfectly correlated.

Alberini, Boyle, and Welsh (2003) estimate the correlation coefficient in a random effects probit model to be 0.06, which – although statistically different from zero – suggests that the within-subject response correlation is very small such that assuming independence is unlikely to significantly distort the standard errors for estimated model parameters and corresponding WTP estimates. Alberini, Boyle, and Welsh (2003) appear startled by this result, as they state (footnote 13, p. 50):

“We were surprised to obtain such a low estimate of the correlation coefficient.... Perhaps this result reflects the routine's difficulty in identifying ρ . In this study ... our expectation [was] to find that ρ is close to one. This expectation was not borne out in the estimation results.”

This result carries with it serious implications. First, it would suggest that the Welsh-Poe and DUDE models are inappropriate. Second, and more importantly, it would imply that the MBDC approach causes considerable response-effects such as those that plague responses to follow-up DC questions. If separate WTP functions underlie each of the respondent's decisions, then there is little hope that valid WTP estimates can even be derived from MBDC questionnaires. That is to say, decisions from the individual are random and correspondingly meaningless. Clearly, the issue of within-subject response correlation warrants further investigation as it has serious implications for the validity of MBDC surveys.

B. On the use of random effects models

A binary choice random effects model conceptually serves as a compromise between the interval data models that assume, in essence, a perfect correlation between within-subject responses, and models that assume response independence. However, the standard random effects model assumes that the random effect is normally distributed. Further, the degree of correlation between any two WTP decisions from the same respondent is assumed to be equal (see Greene 2002, p. 690-693). In general, distributional assumptions play a key role in the estimation of WTP from discrete choice cross-section data models. It is therefore likely that estimation of WTP from discrete choice panel models is sensitive to the assumed distribution of the random effect. The restriction of equal correlation across within-subject decisions seems problematic. If the respondent's underlying WTP distribution does differ across decisions, it is unlikely that such revisions would occur at an equal rate. For instance, the respondent may "learn" about their underlying WTP as part of the process of responding to the valuation questions. The effects of a learning process would presumably diminish over decisions, violating the equal correlation assumption.

C. Estimation of WTP from Ordered Choice Models

The use of ordered choice models (e.g., ordered probit, ordered logit, etc.) has appeal since the analyst can avoid recoding responses into yes or no decisions, or assigning probability weights to certainty levels.⁴ In the absence of information on how certainty category responses reflect actual behavior, a reasonable expectation is for the switch between a “yes-ish” and a “no-ish” response to occur somewhere within the “not sure” interval. Unfortunately, unlike in a binary choice model, you cannot tell where the “propensity to say yes” passes from negative to positive. As such, the model implies a fitted interval of WTP and not a point estimate (Cameron et al. 2002). Specifically, the lower WTP bound is where respondents switch between “not sure” and “probably yes” and the upper bound is where respondents switch between “not sure” and “probably no”. To obtain mean and median WTP point estimates, Alberini, Boyle, and Welsh (2003) restrict the two estimated thresholds that bound the “not sure” category to be symmetric around zero and thus assume that the propensity to say yes passes from negative to positive at the midpoint of the “not sure” category. An important research question is whether this symmetry assumption is appropriate. Without making this assumption a range of possible values has to be considered, and the range of values tends to be rather wide for useful policy purposes.

D. Insensitivity of WTP estimates to alternative DUDE Model probability assignments

In their Benchmark DUDE Model, Evans, Flores, and Boyle (2003) average estimates from three psychology studies and derive subjective probability weights of

⁴ Note that it is not necessary to assume within-subject response independence to estimate an ordered choice model, as a random effects panel structure can also be assumed here.

0.75 and 0.15, respectively to “probably yes” and “probably no” responses. Probabilities of 1, 0.5, and 0 are assigned to the “definitely yes”, “not sure”, and “definitely no” categories, respectively. While holding constant the probability weights for “definitely yes”, “not sure”, and “definitely no” categories, they consider two other “symmetric” assignments. The Probably DUDE Model assigns weights of 0.6 and 0.4 to “probably yes” and “probably no” categories, respectively. In the Definitely DUDE Model, the same categories receive weights of 0.99 and 0.01. Applying these three DUDE Models to data from the first MBDC field study (Welsh et al. 1995) and assuming a normal distribution for WTP, they conclude that “within the class of symmetric (and approximately symmetric) assignments, the parameter estimates from the DUDE model are relatively insensitive to the specific probability assignment.” Since the “definitely yes”, “not sure”, and “definitely no” assignments seem defensible, an interesting research question is whether DUDE models applied to other data show a similar insensitivity to alternative probability assignments for “probably yes” and “probably no” categories.

3. New Methods for Analyzing MBDC Data

Indifference Interval Model

The main drawback of the Welsh-Poe and DUDE models is the need to recode “not sure” responses as either “yes” or “no”, or to assign a probability weight to this category. Since there is only one study that investigates the criterion validity of MBDC responses (Vossler et al. 2003b), many researchers may be skeptical about blindly second-guessing the respondent’s stated intentions. While the ordered choice model avoids the second guessing of responses, to obtain a WTP point estimate the researcher has to make some assumption about where in the “not sure” interval the propensity to say yes passes from negative to positive. However, under the premise that one WTP function drives all within-respondent decisions, we *know* the range of prices for which

the respondent may be “not sure”. In the studies to date, the majority of respondents do pick one or more of the three intermediate categories, allowing us to define the individual’s “not sure” price region with reasonable precision.

Let t^L denote the highest price the subject responded at least “probably yes” to and let t^U denote the lowest price the subject responded either “probably no” or “definitely no” to. Then, the transition between “yes” and “no” falls within the interval $[t^L, t^U]$. The likelihood function is analogous to that of the Welsh-Poe model and is included in Appendix A.

Nonparametric Estimation of Interval Models

Nonparametric estimators are useful as they are robust against misspecification of the response probability distribution. Nonparametric estimation seems especially well suited for MBDC data. Only in rare instances do respondents state a higher level of payment certainty for a higher payment amount. As a result, the cumulative response propensities are monotone decreasing with respect to price. This is in contrast to DC data, where it is common that the raw proportion of yes responses is not monotone (see Haab and McConnell 1997), and a rule for imposing monotonicity must be used to construct a valid nonparametric cumulative distribution function (cdf). Given the added parametric structure of the random effects and ordered choice models, nonparametric estimation is only feasible for MBDC interval data models, which now includes the Welsh-Poe, DUDE, and Indifference interval models.

Nonparametric estimation involves three steps: (1) estimating the value of the WTP cumulative density function (cdf) for each bid amount in the survey; (2) defining or estimating the upper and lower bounds of the cdf; and (3) a rule for interpolating between estimated discrete points of the cdf. The value of the WTP cdf at each bid can be estimated nonparametrically using techniques for survival analysis with censored data described by Turnbull (1974, 1976). Such nonparametric techniques have been

previously applied to single and double-bounded CV data (e.g., Kriström 1990; Carson, Wilks, and Imber 1994).

Let $k = 1, \dots, K$ denote the order bids are presented in the MBDC survey, F_k denote the value of the cdf for the k^{th} bid, and t_k denote the value of the k^{th} bid. For the Welsh-Poe and DUDE models likelihood (recall that the Welsh-Poe model is a special case of the DUDE model), the following log-likelihood is maximized with respect to F_1, \dots, F_K :

$$\ln L = \sum_{i=1}^n \left\{ [1 - P_i(WTP_i > t_1)] \ln[1 - F_1] + \sum_{i=1}^n [P_i(WTP_i > t_{k-1}) - P_i(WTP_i > t_k)] \ln[F_{k-1} - F_k] \right. \\ \left. = [P_i(WTP_i > t_k)] \ln[F_k] \right\} \quad [1]$$

where the P_i 's are the probability weights in the DUDE model and take on values of 1 or 0 in the Welsh-Poe model. For the Indifference Interval Model the log-likelihood is:

$$\ln L = \sum_{i=1}^n \ln[LB_i - UB_i] \quad [2]$$

where $LB_i = L_{i0} + \sum_{k=1}^K L_{ik} * F_{ik}$ and $UB_i = \sum_{k=1}^K U_{ik} * F_{ik}$. L_{ik} equals 1 if bid k is the highest price respondent i chose either the “definitely yes” or “probably yes” category for, and equals 0 otherwise; L_{i0} equals 1 if the respondent gave a “not sure”, “probably no” or “definitely no” response to the lowest bid, and equals 0 otherwise; and U_{ik} equals 1 if bid k is the lowest price respondent i chose either the “probably no” or “definitely no” category for, and equals 0 otherwise. Hence, to facilitate estimation, a set of dummy variables is created to identify the upper and lower bounds of the indifference interval for each respondent.

The likelihood functions [1] and [2] are straightforward analogs to the associated parametric models, as presented in Appendix A. In order to construct a valid cdf, both [1] and [2] are estimated subject to the restriction that $1 > F_1 > F_2 > \dots > F_K >$

0. As mentioned above, estimates of F_k will be generally non-increasing in price. Discussion of the case where $F_{k-1} = F_k$ for some k is relegated to a footnote.⁵

After estimating the F_k , one can construct a valid, nonparametric cdf and calculate mean and median WTP employing the same methods as for nonparametric DC models. The majority of previous DC applications use the Turnbull lower-bound estimator (see Haab and McConnell 1997), where the probability mass in each price interval is massed at the lower end-point of that interval. Rather than adapt this explicitly conservative approach, as questioned in Poe and Vossler (2002), I follow the approach of Kriström (1990). For the cdf to be valid, the upper bound (i.e., the highest price at which $F = 0$) and the lower bound (i.e., the lowest price for which $F = 1$) on WTP need to be chosen or estimated from the data. In the absence of knowledge regarding the upper bound, linear extrapolation can be used to estimate it:

$$\text{Upper WTP Bound } (t_{K+1}) = t_K + F_K \left(\frac{t_K - t_{K-1}}{F_{K-1} - F_K} \right) \quad [3]$$

To preclude negative WTP, one can assume $F = 1$ at $t_0 = \$0$. Alternatively, linear extrapolation or other methods are available for estimating the lower WTP bound.⁶

As an estimate of the cdf between bids (including the upper and lower WTP bound), Kriström uses linear interpolation, which coincides with the assumption that the probability mass between bids is uniformly distributed and results in a piecewise linear cdf. With the cdf constructed in this manner, the calculation for Mean WTP is:

⁵ If $F_{k-1}^* = F_k^*$, a quick solution is to decrease F_k by a negligible amount by decreasing P_i ($\text{WTP} > t_k$) by 0.00001 for one respondent. If $F_k = 0$, then the corresponding bid should be dropped from the model, and perhaps used as the upper WTP bound. Since no assumption is made regarding the value of the cdf between any two bids, it can be shown that when all individuals respond to all bids, the solution for (1)

has $F_k^* = 1/n \sum_{i=1}^n P_i (\text{WTP} > t_k)$, and hence if estimation problems are encountered the data can be easily checked to determine the source of the problem.

⁶ For instance, to allow for negative WTP one can assume symmetry around \$0 (see Kriström 1997).

$$\text{Mean WTP} = \sum_{k=0}^K 0.5(t_k + t_{k+1})(F_k - F_{K+1}) \quad [4]$$

where $F_{K+1} = 0$ and $F_0 = 1$. Median WTP is the price such that $F = 0.5$. Since the mean and median are linear functions of parameter estimates (i.e., the F_k), standard errors and confidence intervals can be calculated by conventional methods using the maximum likelihood parameters and covariance matrix. Without assuming normality, nonparametric bootstrap and simulation techniques are available for generating confidence intervals (see Poe, Severance-Lossin, and Welsh 1994).

Even if the end goal is to estimate a parametric model, the nonparametric cdf may be useful in discriminating between parametric distribution functions. For instance, the nonparametric distribution can be tested against parametric distributions using standard tests, such as the Smirnov Test (Conover 1980). Further, large differences between the nonparametric mean and median suggests that an asymmetric parametric distribution may be appropriate.

Robust Binary and Ordered Choice Models

As discussed in Section 2, estimating a random effects model involves imposing a rather rigid parametric structure on the data. Alternatively, a robust estimation approach is available that avoids such structure and is consistent with a random effects specification under the assumption that within-respondent decisions are freely correlated. Even if a random effects model is appropriate, a simple cross-section model that treats all within-respondent decisions as independent observations can be used to obtain consistent coefficient estimates; however, the estimated covariance matrix is incorrect (Maddala 1987). White's (1982) "sandwich" estimator provides a consistent estimate of the appropriate asymptotic covariance matrix. Let V represent the

(uncorrected) variance matrix from the cross-section model, usually estimated by the inverse of the Hessian matrix. The sandwich estimator is:

$$V_{robust} = V \frac{n}{n-1} \left[\sum_{i=1}^n \left(\sum_{k=1}^K \frac{\partial \ln L_{ik}}{\partial \beta} \right) \left(\sum_{k=1}^K \frac{\partial \ln L_{ik}}{\partial \beta} \right)' \right] V \quad [5]$$

where $\ln L_{ik}$ is the value of the (maximized) log-likelihood function for individual i and bid k . This covariance estimator can be generally applied to discrete choice models. As such, it serves as a “robust” alternative to both binary and ordered choice random effects models.

4. Data

In attempt to make general conclusions about the performance of various modeling approaches, and to corroborate prior findings, this paper analyzes data from three published studies. To the best of my knowledge, these are the only studies in the literature that use both MBDC and PC questionnaires. The first dataset (hereafter referred to as CLASSROOM) comes from a 1994 classroom experiment where students at Cornell University were asked their annual WTP for reduced fluctuations in Glenn Canyon Dam releases (Welsh and Poe 1998). The first MBDC field application utilized an extensive version of this survey (Welsh et al. 1995). Evans, Flores and Boyle (2003) apply the DUDE model to data from the field application.

The second dataset (POWER, hereafter) comes from a green electricity program survey conducted in Erie County, New York in 1996. In the survey, Niagara Mohawk customers were asked how much they were willing to pay per month to join Niagara Mohawk Power Company’s Green Choice™ program, a green electricity program that would fund a tree planting program and a landfill gas project that could replace fossil fuel generated for 1,200 households. This study was a large research effort that

compared MBDC, PC, open-ended, DC, conjoint analysis, and real purchase decisions. Cameron et al. (2002) and Poe et al. (2002) provide details on the study. While there were three MBDC survey versions, the focus in this paper is on the base survey version, which included bids that correspond closely with the PC survey.

The third dataset (ANGLER, hereafter) is from a survey of anglers who held a fishing license in Maine during 1994. Anglers were asked to value their fishing experiences for the 1994 open-water season (April 1, 1994 through September 30, 1994). Specifically, consumer surplus was elicited by asking respondents what they were willing to pay over and above their total trip expenditures. Split samples were presented with surveys that presented the array of bids in either ascending (lowest bid first) or descending (highest bid first) order. Responses from the version with bids in ascending order are used for comparability with the CLASSROOM and POWER datasets. Results of the study appear in Alberini, Boyle, and Welsh (2003).

5. MBDC Estimation Results and Model Selection

For parametric models, I adopt the framework of Cameron and James (1987) for estimating the parameters of a WTP function. Suppose WTP or some transformation of WTP, $f(WTP)$, is a linear function of covariates X and a random error term ε such that

$$f(WTP)_i = X_i\beta + \varepsilon_i \quad [6]$$

where i denotes the individual, β is a vector of parameters to be estimated, and ε has zero mean, a cumulative distribution function (cdf) denoted by $F(\varepsilon)$, and variance σ^2 . While WTP is not directly observed through responses to discrete choice questions, the presence of varying bids, coupled with a choice of $F(\varepsilon)$ allow direct estimation of β via maximum likelihood. Choices of $f(WTP)$ and $F(\varepsilon)$ jointly define a distribution for WTP.

In choosing an appropriate parametric distribution for WTP, I compared estimated normal, log-normal, logistic, log-logistic, and two-parameter Weibull response functions with corresponding nonparametric estimates under various behavioral assumptions. For illustrative purposes, consider Figures 1, 2, and 3, which display estimated parametric (normal, log-normal, and Weibull) and nonparametric functions corresponding to the Welsh-Poe “probably yes” model for the three datasets. In all cases, the normal distribution (the logistic is very similar) noticeably overestimates the cdf at intermediate bids, and underestimates the cdf for high bids. The log-normal distribution (the log-logistic is very similar) coincides with the nonparametric distribution for low to intermediate bids, but overestimates the cdf for high bids. The Weibull generally approximates the nonparametric cdf well. The Smirnov Test, which is based on the maximum distance between two cdfs, is used to test for differences between parametric and nonparametric functions. Using the normal distribution, the equality of distributions for each of the datasets is rejected at the 5% significance level. This is somewhat troubling given that in previous analysis of the datasets, the authors assume either a normal or logistic distribution for ε . The log-normal is rejected for the POWER dataset only, while the Weibull is never rejected. Overall, Weibull mean WTP estimates more closely mimic nonparametric estimates than do estimates from the normal and log-normal distributions. Distribution and welfare comparisons are qualitatively similar under alternative behavioral assumptions. Therefore, as a reasonable approximation, all presented parametric models assume a Weibull distribution. For the Weibull distribution, $f(WTP) = \ln(WTP)$ and ε is distributed extreme value, such that $F(\varepsilon) = \exp[-\exp(-\varepsilon)]$.




Figure 1. Estimated WTP Distributions for the “Probably Yes” Welsh-Poe Model, CLASSROOM Dataset




Figure 2. Estimated WTP Distributions for the "Probably Yes" Welsh-Poe Model, POWER Dataset




Figure 3. Estimated WTP Distributions for the “Probably Yes” Welsh-Poe Model, ANGLER Dataset

Table 2 presents “definitely yes”, “probably yes”, and “not sure” versions of the binary choice (Weibull) random effects model for the three datasets. Tables 3a, 4a, and 5a present binary choice models for the same three coding schemes, as well as ordered choice models. Tables 3b, 4b, and 5b present parametric and nonparametric interval models, including: “definitely yes”, “probably yes”, and “not sure” Welsh-Poe models; Benchmark, Probably, and Definitely DUDE models; and the Indifference Interval Model.

Random Effects models are estimated using LIMDEP econometric software. For this model, coefficients correspond with a reparameterization of the WTP function. See Appendix A for details. All other models are estimated with user-defined maximum likelihood routines in LIMDEP. Ninety-five percent confidence intervals for mean and median WTP point estimates are constructed using 10,000 random draws from the maximum likelihood coefficient vector and covariance matrix, following the approach of Krinsky and Robb (1986).

A. Examination of Within-Subject Response Correlation

As shown in Table 2, the correlation coefficient ranges from 0.959 to 0.985 in the various binary choice random effects models. This result is consistent with the expectation of negligible response-effects stemming from the presence of multiple bids, but is in stark contrast to Alberini, Boyle, and Welsh (2003), who estimate $\rho = 0.06$ for a probit version of the “probably yes” model using the ANGLER dataset. The difference in estimated correlation coefficients is not due to our omission of covariates used by Alberini, Boyle, and Welsh (2003), or the choice of parametric distribution. For completeness, I estimated a probit version of the “probably yes” random effects model – with and without covariates – and obtained correlation coefficients of 0.947 and 0.985, respectively. Estimated models are included as Appendix B. SAS and STATA provide similar estimates of the correlation coefficients in random effects models.

Table 2. Binary Choice Random Effects Models

CLASSROOM Dataset (n=2,403)			
	Definitely Yes	Probably Yes	Not Sure
γ_0	6.908 (1.156)	15.129 (2.437)	29.337 (5.204)
$-\gamma_t$	4.059 (0.633)	4.836 (0.786)	6.874 (1.230)
ρ	0.979 (0.006)	0.978 (0.007)	0.985 (0.005)
Mean WTP	21.03	54.26	136.48
Median WTP	5.01	21.17	67.65
POWER Dataset (n=2,942)			
	Definitely Yes	Probably Yes	Not Sure
γ_0	-1.888 (0.206)	1.884 (0.133)	1.459 (0.131)
$-\gamma_t$	2.635 (0.186)	3.230 (0.202)	3.529 (0.206)
ρ	0.959 (0.059)	0.970 (0.004)	0.971 (0.003)
Mean WTP	2.34	7.62	5.17
Median WTP	0.43	1.60	1.36
ANGLER Dataset (n=8,540)			
	Definitely Yes	Probably Yes	Not Sure
γ_0	14.751 (1.119)	21.365 (1.324)	26.102 (2.069)
$-\gamma_t$	3.909 (0.290)	4.480 (0.278)	4.846 (0.385)
ρ	0.976 (0.004)	0.975 (0.003)	0.979 (0.003)
Mean WTP	111.31	223.71	413.32
Median WTP	39.66	108.52	202.49

Notes: Standard errors are in parentheses. All estimated parameters are statistically different from zero at the 5% level. The two-parameter Weibull distribution is assumed for parametric models.

**Table 3a. MBDC Binary and Ordered Choice Models, CLASSROOM Dataset
(n=2,403)**

Binary Choice Models			
	Definitely Yes	Probably Yes	Not Sure
β	2.585	3.693	4.616
(std. error)	(0.057)	(0.042)	(0.047)
[robust std. error]	[0.124]	[0.097]	[0.101]
σ	1.581	1.233	1.173
(std. error)	(0.059)	(0.047)	(0.053)
[robust std. error]	[0.107]	[0.079]	[0.082]
Mean WTP	18.69	45.09	109.70
(95% C.I.)	(16.75, 20.93)	(41.19, 49.44)	(97.14, 124.36)
[robust 95% C.I.]	[14.63, 24.08]	[37.38, 54.86]	[88.88, 136.64]
Mean WTP	7.43	25.57	65.74
(95% C.I.)	(6.49, 8.50)	(23.26, 28.04)	(60.37, 71.50)
[robust 95% C.I.]	[5.64, 9.81]	[20.73, 31.57]	[53.58, 80.84]
Ordered Choice Model			
	Coefficient	Std. Error	Robust Std. Error
β	5.502	0.073	0.131
μ_1	0.800	0.051	0.073
μ_2	1.853	0.077	0.164
μ_3	2.937	0.096	0.120
σ	1.420	0.044	0.086
Mean WTP Interval	48.40 – 138.62		
Median WTP Interval	22.87 – 65.48		
Mean WTP ¹	81.91		
(95% C.I.)	(73.49, 91.25)		
[robust 95% C.I.]	[66.02, 102.51]		
Median WTP ¹	38.69		
(95% C.I.)	(35.52, 42.18)		
[robust 95% C.I.]	[30.84, 48.48]		

Notes: Standard errors are in parentheses. All estimated parameters are statistically different from zero at the 5% level. The two-parameter Weibull distribution is assumed for parametric models. ¹ Welfare measures calculated based on symmetry assumption – see text.

Table 3b. MBDC Interval Data Models, CLASSROOM Dataset (n=185)

Welsh-Poe Interval Models			
	Definitely Yes	Probably Yes	Not Sure
Parametric Estimates:			
β	2.600 (0.124)	3.697 (0.098)	4.669 (0.103)
σ	1.591 (0.097)	1.257 (0.077)	1.238 (0.088)
Mean WTP [95% C.I.]	19.13 [15.02, 24.46]	45.88 [37.90, 55.75]	119.87 [96.10, 150.51]
Median WTP [95% C.I.]	7.51 [5.71, 9.87]	25.45 [20.51, 31.50]	67.70 [54.58, 83.68]
Nonparametric Estimates:			
Mean WTP [95% C.I.]	19.33 [14.52, 24.26]	53.86 [40.87, 66.82]	128.61 [107.40, 149.72]
Median WTP [95% C.I.]	8.21 [6.49, 10.34]	28.10 [20.97, 34.98]	62.50 [47.42, 79.64]
DUDE Models			
	Benchmark DUDE	Probably DUDE	Definitely DUDE
Parametric Estimates:			
β	4.155 (0.129)	4.264 (0.150)	4.216 (0.107)
σ	1.597 (0.108)	1.830 (0.129)	1.337 (0.088)
Mean WTP [95% C.I.]	90.97 [68.71, 121.31]	122.30 [86.15, 176.17]	80.90 [64.85, 101.35]
Median WTP [95% C.I.]	35.51 [26.93, 46.59]	36.35 [26.47, 49.68]	41.51 [32.94, 52.12]
Nonparametric Estimates:			
Mean WTP [95% C.I.]	88.12 [70.35, 105.70]	95.21 [77.06, 113.55]	91.16 [72.72, 109.41]
Median WTP [95% C.I.]	33.12 [24.56, 43.91]	33.12 [24.56, 43.91]	40.90 [32.19, 47.57]
Indifference Interval Model			
	Parametric	Nonparametric	
β	4.146 (0.099)		
σ	1.157 (0.083)		
Mean WTP [95% C.I.]	68.06 [55.49, 83.97]	65.31 [54.33, 75.61]	
Median WTP [95% C.I.]	41.35 [33.54, 50.80]	43.49 [36.13, 53.17]	

Notes: Standard errors are in parentheses. All estimated parameters are statistically different from zero at the 5% level. The two-parameter Weibull distribution is assumed for parametric models.

Table 4a. MBDC Binary and Ordered Choice Models, POWER Dataset ($n=2,942$)

Binary Choice Models			
	Definitely Yes	Probably Yes	Not Sure
β	0.048	0.942	1.613
(std. error)	(0.070)	(0.048)	(0.051)
[robust std. error]	[0.151]	[0.113]	[0.126]
σ	2.053	1.617	1.796
(std. error)	(0.095)	(0.065)	(0.066)
[robust std. error]	[0.130]	[0.096]	[0.119]
Mean WTP	2.20	3.72	8.39
(95% C.I.)	(1.91, 2.57)	(3.31, 4.19)	(7.28, 9.72)
[robust 95% C.I.]	[1.72, 2.87]	[3.04, 4.57]	[6.34, 11.29]
Mean WTP	0.49	1.42	2.60
(95% C.I.)	(0.41, 0.59)	(1.26, 1.59)	(2.33, 2.90)
[robust 95% C.I.]	[0.35, 0.71]	[1.09, 1.84]	[1.97, 3.44]
Ordered Choice Model			
	Coefficient	Std. Error	Robust Std. Error
β	2.385	0.056	0.112
μ_1	0.751	0.043	0.104
μ_2	1.470	0.060	0.200
μ_3	2.334	0.076	0.140
σ	1.897	0.057	0.110
Mean WTP Interval	4.55 – 9.34		
Median WTP Interval	1.25 – 2.56		
Mean WTP ¹	6.52		
(95% C.I.)	(5.79, 7.35)		
[robust 95% C.I.]	[5.23, 8.24]		
Median WTP ¹	1.78		
(95% C.I.)	(1.59, 2.00)		
[robust 95% C.I.]	[1.29, 2.49]		

Notes: Standard errors are in parentheses. Except for β in the “definitely yes” model, all estimated parameters are statistically different from zero at the 5% level. The two-parameter Weibull distribution is assumed for parametric models. ¹ Welfare measures calculated based on symmetry of the fitted WTP interval – see text.

Table 4b. MBDC Interval Data Models, POWER Dataset (n=260)

Welsh-Poe Interval Models			
	Definitely Yes	Probably Yes	Not Sure
Parametric Estimates:			
β	-0.167 (0.166)	0.900 (0.120)	1.604 (0.135)
σ	2.400 (0.162)	1.764 (0.106)	1.954 (0.113)
Mean WTP [95% C.I.]	2.53 [1.76, 3.71]	4.00 [3.11, 5.20]	9.54 [7.11, 12.93]
Median WTP [95% C.I.]	0.35 [0.24, 0.52]	1.29 [0.98, 1.68]	2.43 [1.81, 3.25]
Nonparametric Estimates:			
Mean WTP [95% C.I.]	1.99 [1.51, 2.47]	3.57 [3.24, 4.20]	9.67 [6.67, 12.65]
Median WTP [95% C.I.]	0.50 [0.27, 0.88]	1.66 [1.35, 2.15]	2.93 [2.36, 4.13]
DUDE Models			
	Benchmark DUDE	Probably DUDE	Definitely DUDE
Parametric Estimates:			
β	1.180 (0.153)	1.257 (0.173)	1.266 (0.133)
σ	2.240 (0.135)	2.524 (0.156)	1.941 (0.113)
Mean WTP [95% C.I.]	8.21 [5.81, 11.79]	12.01 [7.97, 18.50]	6.72 [5.05, 9.06]
Median WTP [95% C.I.]	1.43 [1.02, 2.00]	1.39 [0.95, 2.04]	1.74 [1.30, 2.33]
Nonparametric Estimates:			
Mean WTP [95% C.I.]	7.31 [4.71, 9.91]	9.70 [6.35, 13.01]	6.18 [4.11, 8.24]
Median WTP [95% C.I.]	1.48 [1.15, 2.06]	1.43 [1.05, 2.06]	1.74 [1.34, 2.30]
Indifference Interval Model			
	Parametric	Nonparametric	
β	1.282 (0.121)		
σ	1.743 (0.111)		
Mean WTP [95% C.I.]	5.76 [4.41, 7.62]	5.25 [4.33, 6.18]	
Median WTP [95% C.I.]	1.90 [1.46, 2.47]	2.36 [1.72, 2.80]	

Notes: Standard errors are in parentheses. Except for β in the “definitely yes” model, all estimated parameters are statistically different from zero at the 5% level. The two-parameter Weibull distribution is assumed for parametric models.

**Table 5a. MBDC Binary and Ordered Choice Models, ANGLER Dataset
(n=8,540)**

Binary Choice Models			
	Definitely Yes	Probably Yes	Not Sure
β	4.524	5.257	5.880
(std. error)	(0.031)	(0.026)	(0.025)
[robust std. error]	[0.053]	[0.045]	[0.052]
σ	1.734	1.396	1.310
(std. error)	(0.034)	(0.026)	(0.025)
[robust std. error]	[0.056]	[0.045]	[0.050]
Mean WTP	146.49	237.77	420.15
(95% C.I.)	(136.71, 157.15)	(224.62, 251.85)	(396.95, 444.63)
[robust 95% C.I.]	[128.17, 168.47]	[212.58, 267.00]	[369.49, 480.88]
Mean WTP	48.87	115.08	221.40
(95% C.I.)	(45.46, 52.49)	(108.77, 121.63)	(210.31, 232.85)
[robust 95% C.I.]	[43.71, 54.78]	[105.62, 125.61]	[201.24, 244.10]
Ordered Choice Model			
	Coefficient	Std. Error	Robust Std. Error
β	6.542	0.030	0.064
μ_1	0.674	0.023	0.040
μ_2	1.355	0.032	0.089
μ_3	2.078	0.038	0.063
σ	1.520	0.023	0.046
Mean WTP Interval	241.28 – 476.82		
Median WTP Interval	102.49 – 202.55		
Mean WTP ¹	339.19		
(95% C.I.)	(320.87, 358.34)		
[robust 95% C.I.]	[297.80, 388.06]		
Median WTP ¹	144.05		
(95% C.I.)	(136.82, 151.78)		
[robust 95% C.I.]	[126.88, 164.09]		

Notes: Standard errors are in parentheses. All estimated parameters are statistically different from zero at the 5% level. The two-parameter Weibull distribution is assumed for parametric models. ¹ Welfare measures calculated based on symmetry of the fitted WTP interval – see text.

Table 5b. MBDC Interval Data Models, ANGLER Dataset (n=622)

Welsh-Poe Interval Model			
	Definitely Yes	Probably Yes	Not Sure
Parametric Estimates:			
β	4.500 (0.078)	5.224 (0.062)	5.844 (0.059)
σ	1.807 (0.057)	1.470 (0.046)	1.404 (0.045)
Mean WTP [95% C.I.]	151.80 [130.51, 176.85]	241.74 [214.49, 272.61]	429.89 [382.32, 483.83]
Median WTP [95% C.I.]	46.40 [39.26, 54.76]	108.35 [94.64, 123.90]	206.28 [181.25, 234.48]
Nonparametric Estimates:			
Mean WTP [95% C.I.]	171.17 [138.58, 203.90]	258.88 [222.87, 295.27]	549.05 [468.43, 631.66]
Median WTP [95% C.I.]	47.09 [40.30, 56.31]	100.00 [90.45, 121.36]	206.90 [179.65, 233.67]
DUDE Model			
	Benchmark	Symmetric I	Symmetric II
Parametric Estimates:			
β	5.519 (0.069)	5.620 (0.075)	5.551 (0.062)
σ	1.635 (0.053)	1.775 (0.059)	1.474 (0.047)
Mean WTP [95% C.I.]	366.24 [318.78, 421.15]	453.15 [387.62, 530.93]	336.37 [297.75, 380.33]
Median WTP [95% C.I.]	136.94 [117.77, 158.99]	143.94 [122.20, 169.29]	150.07 [131.02, 171.67]
Nonparametric Estimates:			
Mean WTP [95% C.I.]	392.45 [337.19, 448.73]	441.93 [384.33, 500.91]	383.62 [328.03, 440.37]
Median WTP [95% C.I.]	132.30 [105.97, 157.00]	137.31 [108.79, 164.37]	147.02 [122.89, 170.60]
Indifference Interval Model			
	Parametric	Nonparametric	
β	5.519 (0.060)		
σ	1.395 (0.046)		
Mean WTP [95% C.I.]	308.78 [274.20, 348.09]	269.96 [240.76, 299.22]	
Median WTP [95% C.I.]	149.57 [131.23, 170.22]	154.14 [136.90, 171.91]	

Notes: Standard errors are in parentheses. All estimated parameters are statistically different from zero at the 5% level. The two-parameter Weibull distribution is assumed for parametric models.

Given the very large and statistically significant correlation coefficients, models estimated under the premise of within-subject response independence are inappropriate. However, coefficient estimates from ordered or binary choice models where the responses are pooled but the correlation structure is ignored are consistent (Maddala 1987), and so the concern lies in the estimated standard errors. Tables 3a, 4a, and 5a report binary and ordered choice model coefficients and WTP estimates. Two sets of coefficient standard errors and corresponding 95% confidence intervals for WTP are reported. The first set assumes within-subject independence while the “robust” estimates are calculated using the “sandwich” estimator, which takes account of within-subject response correlation. In all cases, robust standard errors and 95% confidence intervals are noticeably larger, implying that inferences based on uncorrected standard errors are problematic. For the binary choice models using the CLASSROOM dataset, for example, the robust standard errors are approximately twice as large as the uncorrected errors.

B. Monte Carlo and Empirical Evidence on Model Performance

It is clear that the independence assumption is inappropriate, although the correlation coefficient estimates do not serve in and of themselves as a basis for choosing between random effects and interval data models. The correlation coefficients are very close to 1, but there is no reliable test of $\rho = 1$ (Alberini 1995). Even if there were such a test, concerns over distributional and other assumptions imposed in the random effects model call into question its usefulness.

As an alternative mode of investigation, simple Monte Carlo experiments were conducted to gain insight into the relative performance of random effects versus interval data models for analyzing MBDC data under the premise of perfect correlation of within-respondent decisions. Another feasible alternative is to estimate a binary choice model and use the “sandwich” covariance matrix, and so I consider this estimator as

well. The experiment is set up as follows. Numbers representing underlying WTP amounts for a sample of 100 individuals are randomly drawn from the normal distribution. For each individual a set of yes/no indicator variables are constructed by comparing the individual's WTP to a set of payment thresholds. Specifically, the set of bids is $\{0,5,10,15,20,25,30,35,40,45,50,55,60\}$, there are no response errors, and $\rho = 1$: a yes response is recorded if $WTP > \text{bid}$ and otherwise a no response is recorded.⁷ There are 13 bids and so this procedure produces 1,300 total observations. These observations are used to estimate mean WTP from binary probit and random effects probit models. The upper and lower WTP bounds for each individual are created by identifying the two consecutive survey bids that bound the individuals' WTP.⁸ An interval data model estimates mean WTP using the set of WTP bounds for the 100 individuals.

Table 6 presents results from two Monte Carlo experiments. In the first experiment, individual WTP values drawn from a normal distribution with a mean of \$30 and standard deviation of \$30. In the second experiment, the mean is \$50 with a standard deviation of \$30. Note that the expected mean WTP in experiment 1 is in the center of the bid distribution, while the mean and standard deviation of WTP in experiment 2 is such that many respondents have WTP amounts larger than the highest bid. This is analogous to a situation of a survey with a poor bid design. There are 500 replications in each experiment.

Several interesting results stem from the experiments. First, the correlation coefficients average 0.972 and 0.973, respectively, in the two experiments and the range of values across replications is small (0.960 to 0.984). Second, the mean-squared error (MSE) is approximately ten times larger for the random effects probit versus the other

⁷ Other experiments were run where the respondent's WTP is highly correlated across bids ($\rho=0.95$). There was no real difference in results, except for slightly lower random effect probit estimates of ρ .

⁸ As standard in interval models, the lower bound is $-\infty$ if WTP is less than the smallest bid and the upper bound is ∞ if WTP exceeds the highest bid.

two models. In a Monte Carlo comparison between the interval model and a bivariate probit for estimating WTP from double-bounded CV data, Alberini (1995) finds that the latter model is inferior in terms of MSE. Third, in Experiment 2, where the bid design is relatively poor, WTP from the random effects probit is 6% below the true value on average and is off by as much as \$13.90 (28%) in individual trials. In contrast, both the interval model and binary probit perform well in terms of accurately estimating mean WTP, with no evidence of systematic bias. The MSE of the interval model is approximately 20% smaller than the binary probit, but this is to be expected. Overall, it appears that the interval model is preferred under the premise that within-respondent decisions are highly correlated.

Table 6. Monte Carlo Results: Performance of the Probit, Random Effect Probit, and Interval Models

Experiment 1: Mean (β) = \$30, $\rho = 1$ (100 individuals, 500 Replications)			
	Probit	Random Effects Probit	Interval Model
Mean WTP	30.07	30.43	30.07
MSE	0.78	11.01	0.65
$\max \hat{\beta} - \beta $	3.19	9.90	2.55
ρ	0.000	0.972	1.000
Std. dev.		0.003	
Range		0.962 – 0.984	
Experiment 2: Mean (β)= \$50, $\rho = 1$ (100 individuals, 500 Replications)			
	Probit	Random Effects Probit	
Mean WTP	50.20	47.23	50.22
MSE	2.66	22.35	2.05
$\max \hat{\beta} - \beta $	5.58	13.90	4.66
ρ	0.000	0.973	1.000
Std. dev.		0.004	
Range		0.960 – 0.983	

Results from the Monte Carlo experiments do carry over to actual MBDC WTP estimates. Both the average (0.975) and range of coefficient values (0.959 to 0.985) estimated with actual MBDC data are strikingly close to the simulated values. While not a proof, this suggests that actual within-respondent decisions for the three datasets *could* be perfectly correlated. Quite generally, coefficient and WTP estimates are very similar between the binary choice and comparable Welsh-Poe interval models. Random effects model estimates are in the ballpark of those from comparable models but differ noticeably in instances. For the POWER dataset, estimated mean and median WTP is actually larger in the “probably yes” than in the “not sure” model, where more responses are treated as “yes”. Actual estimation results coupled with the relatively large MSE observed in Monte Carlo experiments serve to illustrate the routine’s apparent difficulty in accurately identifying WTP.

C. Comparison of WTP from Ordered Choice and Indifference Interval Models

Both the ordered choice model and the Indifference Interval model can estimate WTP without the need to recode or second-guess categorical responses. As discussed in Section 2, to estimate WTP from the ordered choice model the analyst must make the untestable assumption that the fitted WTP interval corresponding to the “not sure” category is symmetric. Such an assumption is not necessary for the Indifference Interval model. The key difference between modeling approaches is that the Indifference Interval estimator directly models the individual’s “not sure” price interval while the ordered choice model attempts to estimate the “not sure” interval from the data. Maintaining the hypothesis that a single underlying WTP distribution drives all within-respondent decisions, the Indifference Interval Model is presumably the preferable approach. Therefore, testing for equality of WTP between ordered choice and Indifference Interval models sheds light on whether the symmetry assumption is appropriate. For reasons given above, the focus of comparisons is between the “robust”

version of the ordered choice model and the indifference model. Since I use simulation methods to derive the distributions for our WTP point estimates, the method of convolutions (Poe, Severance-Lossin, and Welsh 1994) is appropriate for testing between two WTP point estimates. Both mean (CLASSROOM: $p_c = 0.230$; POWER: $p_p = 0.503$; ANGLER: $p_a = 0.250$) and median ($p_c = 0.677$; $p_p = 0.783$; $p_a = 0.336$) WTP estimates are not statistically different at the 5% level for all datasets. As such, this lends qualified support of the “symmetry” restriction needed to identify WTP point estimates in the robust ordered model. It is likely that in some applications, the individual’s “indifference interval” is not well defined. This can happen when many respondents are not willing to pay the first bid amount or otherwise do not choose the intermediate categories over the range of prices. In such situations, the robust ordered choice model is preferable.

D. Replication of Evans, Flores, and Boyle

Evans, Flores, and Boyle (2003) find that welfare estimates are insensitive to the different probability assignments of the Benchmark, Probably, and Definitely DUDE models. Using the method of convolutions, I conduct pair-wise tests of equality between mean or median WTP estimates from the different parametric specifications. Using the method of convolutions, I find that Probably DUDE and Definitely DUDE mean WTP estimates are statistically different at the 5% significance level for all datasets ($p_c = 0.050$; $p_p = 0.026$; $p_a = 0.004$). However, median WTP is never statistically different between these same models. In all possible cases, mean and median WTP estimates from the Benchmark DUDE model are not statistically different than estimates from the Probably and Definitely DUDE models. Overall, median WTP estimates are insensitive to alternative “probably yes” and “probably no” DUDE model probability assignments. However, when the goal is to estimate mean WTP, these same probability assignments indeed matter.

6. Comparing MBDC with the PC

The MBDC elicitation mechanism can be regarded as a PC with payment certainty as an added dimension. In this section, I examine how respondents behave in the absence of the payment certainty categories.⁹ This exploration is deliberately comprehensive as I consider a multitude of different MBDC models – each with differing behavioral assumptions, compare both WTP point estimates and estimated WTP functions, and employ both parametric and nonparametric approaches.

To limit the scope of comparisons, the focus is on the various interval data models as well as the robust ordered choice model. Note that binary choice models mimic the Welsh-Poe models. The (parametric) random effects models are ignored for obvious reasons. PC data are analyzed using the parametric and nonparametric versions of the interval data model.¹⁰ Comparisons are made between estimated nonparametric and various parametric distributions for the PC data, analogous to those described for MBDC data in Section 4. Again, I find that the Weibull distribution serves as a reasonable approximation for the underlying WTP distribution.

Table 7 presents estimated Weibull and nonparametric PC interval models. I first test for equality between PC and various MBDC parametric WTP functions, using the likelihood ratio test:

$$LR = -2[\ln L_{1=2} - (\ln L_1 + \ln L_2)] \sim \chi^2(r) \quad [7]$$

where $\ln L_1$ and $\ln L_2$ are values of the log-likelihood at solution for the independent PC and MBDC models, $\ln L_{1=2}$ is the value of the log-likelihood for a model that pools both datasets and estimates a common parameter vector (β, σ) , and r is the number of restrictions. Since the sample sizes differ between corresponding PC and MBDC

⁹ Despite several requests, our efforts to acquire the PC data from the ANGLER study were unsuccessful.

¹⁰ Estimated PC robust binary choice models are very similar to interval models.

samples in the POWER dataset, I use sampling weights such that each sample, rather than each observation, receives equal weight. The relative error variances are allowed to differ by data type, as any difference in error variance between data type would distort estimated parameters of a pooled model. When pooling the ordered choice and PC data, no symmetry restriction is needed as the PC data allow us to identify all four ordered thresholds (i.e., the normalization $\mu_0 = 0$ is not imposed; see Cameron 2002).

Table 7. Payment Card Models

Parametric Models		
	CLASSROOM Data	POWER Data
β	3.635 (0.096)	0.938 (0.098)
σ	1.241 (0.074)	1.548 (0.082)
Mean WTP [95% C.I.]	42.71 [35.52, 51.54]	3.51 [2.92, 4.24]
Median WTP [95% C.I.]	24.05 [19.46, 29.66]	1.45 [1.16, 1.81]
Nonparametric Models		
	CLASSROOM Data	POWER Data
Mean WTP [95% C.I.]	43.84 [35.65, 52.01]	3.57 [2.87, 4.27]
Median WTP [95% C.I.]	21.58 [17.53, 28.35]	1.36 [1.22, 1.94]
n	188	292

Notes: Standard errors are in parentheses. All estimated parameters are statistically different from zero at the 5% level. The two-parameter Weibull distribution is assumed for parametric models.

Employing a 5% significance level, I fail to reject the hypothesis of equal WTP functions *only* when comparing PC and “probably yes” Welsh-Poe interval models [CLASSROOM: $\chi_c^2(2)=0.205$, $p_c = 0.902$; POWER: $\chi_p^2(2) = 0.060$, $p_c = 0.970$]. While comparisons of welfare estimates are not as clean, I fail to reject both equal mean and median WTP between PC and “probably yes” models only [means: $p_c = 0.599$, $p_p = 0.423$; medians: $p_c = 0.713$, $p_p = 0.519$]. For the POWER dataset, there are several instances where MBDC medians are not different from the PC median. However, given

the results of LR tests and mean WTP tests, this result is merely coincidental as underlying response functions fundamentally differ.

To explore the robustness of the findings, the Smirnov Test is used to compare estimated nonparametric distribution functions.¹¹ Paralleling results for the LR tests, I fail to reject the hypothesis of equal WTP distributions between PC and “probably yes” models only and this result holds across datasets. The maximum distances between these WTP distributions are 6.6% and 8.4%, respectively, for the CLASSROOM and POWER datasets. Tests of WTP point estimates likewise coincide with the results from parametric tests, as I only fail to reject equality between PC and “probably yes” MBDC models for both point estimates.

Overall the evidence is clear – in the absence of the certainty response categories, otherwise “definitely yes” and “probably yes” respondents will vote yes, while “not sure”, “probably no”, and “definitely no” respondents will vote no. While presumption may be that some “not sure” respondents would vote yes and others vote no, neither the Indifference Interval, ordered choice model, nor any of the DUDE models match PC response functions or both mean and median WTP estimates.

So, what are the implications of this finding? It does suggest that, when not given the option, “not sure” respondents tend to vote “no”. This coincides closely with findings from CV survey comparisons with actual voting behavior (Carson, Hanemann, and Mitchell 1986; Champ and Brown 1997; Vossler et al. 2003a) and evidence from pre-election polls (Magelby 1989), where voters who are undecided before an election overwhelmingly vote “no” on Election Day. If we may extrapolate from these findings, it may be the case that “not sure” responses are an indication that the respondent would

¹¹ Since maximum likelihood estimates the discrete points of the nonparametric cdfs, we could use a likelihood ratio test of the hypothesis that the discrete points are equal across models. However, the large number of restrictions (one for each point) gives rise to concern over the power of such a test.

not actually pay the corresponding amount if faced with an analogous real purchase situation.

In the lone MBDC field validity test (Vossler et al. 2003b), from which the POWER data comes from, MBDC responses correspond with actual purchase decisions only when the vast majority of “definitely yes” and “probably yes” responses are treated as yes, and other responses are treated as no. Consistent with our findings, PC decisions match actual purchase behavior. If this finding is consistent in future validity studies, this suggests that virtually nothing is gained by inclusion of the payment certainty categories. Similarly, PCs are preferred as they reduce the cognitive burden of respondents as well as data analysts. Clearly, the issue of whether “not sure” responses reflect actual or fictional uncertainty introduced by the inclusion of such a response option in a contingent market survey warrants further research.

7. Concluding Remarks

In multiple bounded discrete choice (MBDC) surveys, respondents face multiple payment amounts for a nonmarket good and indicate their level of payment certainty for each amount. As it stands, this elicitation mechanism attempts to gather much more information on the respondent’s WTP than standard DC questions. Making use of three existing datasets, I explore the issue of data analysis and compare MBDC with the payment card (PC), which presents the respondent with multiple payment amounts but does not collect information on payment certainty.

Given the nature of the data, two important issues to consider when analyzing MBDC responses are how to treat the multiple responses from the same individual and how payment certainty levels correspond with real behavioral intentions. Contrary to past research, there is compelling evidence in favor of treating the within-respondent decisions as perfectly correlated, thus driven by a single underlying WTP distribution. Namely, the estimated correlation of within-respondent decisions is in the range of

0.959 to 0.985 using random effects models. Further, empirical WTP estimates and evidence from Monte Carlo experiments provide support for favoring interval models, which assume perfect correlation, over random effects models in terms of their ability to estimate WTP accurately. Thus, overall this study provides strong guidance for model selection. The set of existing analytic approaches is expanded by introducing nonparametric interval model estimators, an Indifference Interval model which serves as a alternative to the ordered choice model used in past research, and a procedure for correcting the covariance matrix from models that assume within-subject response independence, so that valid inferences may be drawn from such models.

Comparisons between PC and various MBDC models, which differ largely in terms of how payment certainty responses are treated, reveal a strong compatibility between PC and MBDC models that treat “definitely yes” and “probably yes” responses as yes and all other responses as no. In fact, this finding is consistent across datasets and holds for both parametric and nonparametric response functions as well as mean and median WTP estimates. For all other MBDC models, both parametric and nonparametric response functions are statistically different from corresponding PC functions. In the lone field validity test of MBDC surveys (Vossler et al. 2003b), this same recoding of certainty responses into yes/no decisions is needed for correspondence between survey responses and revealed behavior. Evidence from CV survey responses and actual voting behavior, as well as pre-election polls suggests likewise that the majority of “undecided” pre-election respondents vote no in actual elections. Evidence from field research, coupled with comparisons between PC and MBDC responses, suggests that all relevant information from the respondent is attainable through PC surveys. That is to say, the inclusion of the payment certainty categories may be unnecessary and simply serve to add complication for the respondent as well as the data analyst.

Appendix A: Log-Likelihood Functions for MBDC Models

A. Interval Models: within-subject responses are driven by a single underlying WTP distribution

Welsh-Poe Interval Model

Welsh and Poe (1998) analyze MBDC data using an interval model that is analogous to the models commonly used for PC (see Cameron and Huppert 1989) and multiple DC (see Carson and Mitchell 1987) data. Responses in a given certainty category are systematically recoded into simple yes/no answers. In the “definitely yes” model, “definitely yes” responses are recoded as yes and all other responses as no. In the “probably yes” model, both “definitely yes” and “probably yes” responses are treated as yes, and other responses as no. Based on the yes responses an interval is constructed that bounds the respondent’s WTP. The respondent’s WTP is bounded from above by the lowest bid they are not willing to pay, denoted t^U , and from below by the highest bid they are willing to pay, denoted t^L . It follows that $\Pr(\text{yes}) = \Pr[t^L \leq \text{WTP} \leq t^U] = F[f(t^L)/\sigma - X\beta/\sigma] - F[f(t^U)/\sigma - X\beta/\sigma]$. If the respondent answers yes to all prices: $\Pr(\text{yes}) = F[f(t^L)/\sigma - X\beta/\sigma]$, where t^L is the highest offered bid. When the respondent answers “no” to all prices: $1 - F[f(t^U)/\sigma - X\beta/\sigma]$, where t^U is equal to the lowest bid. The parameters of the WTP function are estimated by maximizing the following log-likelihood function:

$$\begin{aligned} \ln L = \sum_{i=1}^n \left\{ I_{1i} \ln F\left(\frac{f(t_i^L) - X_i\beta}{\sigma}\right) + I_{2i} \ln \left[F\left(\frac{f(t_i^L) - X_i\beta}{\sigma}\right) - F\left(\frac{f(t_i^U) - X_i\beta}{\sigma}\right) \right] \right. \\ \left. + I_{3i} \ln \left[1 - F\left(\frac{f(t_i^U) - X_i\beta}{\sigma}\right) \right] \right\} \end{aligned} \quad [\text{A1}]$$

where $I_{1i} = 1$ if the respondent answers yes to all bids, =0 otherwise; $I_{2i} = 1$ if the respondent's WTP lies between two offered bids, =0 otherwise; and $I_{3i} = 1$ if the respondent answers no to all bids, =0 otherwise.

Dual Uncertainty Decision Estimator (DUDE) Model

Motivated by research in psychology that suggests verbal probability statements convey subjective probabilities, Evans, Flores, and Boyle (2003) assign probability weights to the response categories. For instance, rather than recoding all “not sure” responses as either yes or no, a “not sure” response to t can be interpreted to mean that there is a 50% chance the individual would actually pay t . Assuming that the responses to the K bids are all driven by a single WTP distribution, a discrete cdf is constructed for the individual using a set of researcher-assigned probabilities for each response category. The name for the model, the Dual Uncertainty Decision Estimator (DUDE) reflects the notion that there are two sources of uncertainty: uncertainty surrounding the assignment of subjective probabilities and uncertainty regarding the log-likelihood contribution for each respondent given the probabilities. Using the optimal decision rule based on a quadratic loss function, the log-likelihood function is:

$$\begin{aligned} \ln L = \sum_{i=1}^N \{ & [1 - P_i(WTP_i > t_1)] \ln \left[1 - F \left(\frac{f(t_1) - X_i \beta}{\sigma} \right) \right] \\ & + \sum_{k=2}^K [P_i(WTP_i > t_{k-1}) - P_i(WTP_i > t_k)] \ln \left[F \left(\frac{f(t_{k-1}) - X_i \beta}{\sigma} \right) - F \left(\frac{f(t_k) - X_i \beta}{\sigma} \right) \right] \\ & + [P_i(WTP_i > t_K)] \ln \left[F \left(\frac{f(t_K) - X_i \beta}{\sigma} \right) \right] \} \end{aligned} \quad [A2]$$

where $P_i(WTP > t_k)$ is the subjective probability that the respondent's WTP is greater than bid t_k .

B. Binary and Ordered Choice Models

Binary Choice Model

After recoding categorical responses into yes/no decisions, as in the Welsh-Poe framework, standard techniques for analyzing DC data can be applied. That is, the K responses from the same individual are treated as separate observations, such that our sample size is nK , rather than n . The log-likelihood function is:

$$\ln L = \sum_{i=1}^n \left\{ Y_i \ln F\left(\frac{f(t_i) - X_i\beta}{\sigma}\right) + (1 - Y_i) \ln \left[1 - F\left(\frac{f(t_i) - X_i\beta}{\sigma}\right) \right] \right\} \quad [\text{A3}]$$

Estimated parameters are consistent, even if within-subject responses are correlated. The sandwich estimator in Section 3 corrects the covariance matrix in the presence of within-subject response correlation.

Ordered Choice Model

In order to avoid recoding uncertain responses into yes/no decisions or assigning probability weights, Alberini, Boyle, and Welsh (2003) estimate an ordered probit model and Cameron et al. (2002) estimate an ordered logit model. Each response option is retained as a separate category and, from the data, thresholds are estimated that imply bounds on WTP for responses within each category. The log-likelihood of the ordered probability model is:

$$\begin{aligned}
& \sum_{i=1}^N \sum_{k=1}^K \{ I_{ik}^{DY} \ln \left[F \left(\frac{f(t_{ik}) + \mu_3 - X_{ik}\beta}{\sigma} \right) \right] \right. \\
& + I_{ik}^{PY} \ln \left[\left(1 - F \left(\frac{f(t_{ik}) + \mu_3 - X_{ik}\beta}{\sigma} \right) \right) - \left(1 - F \left(\frac{f(t_{ik}) + \mu_2 - X_{ik}\beta}{\sigma} \right) \right) \right] \\
& + I_{ik}^{NS} \ln \left[\left(1 - F \left(\frac{f(t_{ik}) + \mu_2 - X_{ik}\beta}{\sigma} \right) \right) - \left(1 - F \left(\frac{f(t_{ik}) + \mu_1 - X_{ik}\beta}{\sigma} \right) \right) \right] \\
& + I_{ik}^{PN} \ln \left[\left(1 - F \left(\frac{f(t_{ik}) + \mu_1 - X_{ik}\beta}{\sigma} \right) \right) - \left(1 - F \left(\frac{f(t_{ik}) + \mu_0 - X_{ik}\beta}{\sigma} \right) \right) \right] \\
& \left. + I_{ik}^{DN} \ln \left[1 - F \left(\frac{f(t_{ik}) + \mu_0 - X_{ik}\beta}{\sigma} \right) \right] \right\}
\end{aligned} \tag{A4}$$

where $I_{ik}^j = 1$ if respondent i chose category j ($j = \text{DY} = \text{“definitely yes”}$, etc.) for bid k , and equals 0 otherwise; and the μ 's are threshold parameters to be estimated. The threshold parameters give us information on where respondents switch between two consecutive categories. Without assuming that the switch from yes to no occurs between two categories, a minimal restriction for estimating WTP is to make the two center thresholds (or all thresholds) symmetric around zero by forcing μ_2 equal to $-\mu_3$ and then using the standard formula for calculating WTP applies. The sandwich estimator in Section 3 corrects the covariance matrix in the presence of within-subject response correlation.

C. Binary and Ordered Choice Random Effects Models

Alberini, Boyle, and Welsh (2003) use a panel data approach to estimate the parameters of a WTP function. For each of the n respondents, each of the K decisions are recoded into “yes” and “no” responses. Pooling the nK responses, a random effects model can be estimated that allows the within-respondent decisions to be freely correlated. That is, it is assumed that each within-subject response comes from separate WTP distribution but there is an unobserved heterogeneity with respect to the individual. The WTP function is assumed to have the form:

$$f(WTP_{ik}) = X_{ik}\beta + v_{ik}, \quad \text{where } v_{ik} = u_i + \varepsilon_{ik}; \quad i = 1, \dots, N; \quad k = 1, \dots, K \quad [\text{A5}]$$

where u_i is the unobserved heterogeneity specific to individual i , is mean zero, and is uncorrelated with X and ε . Another key assumption is that the within-subject error correlation $[\text{Corr}(\varepsilon_{ij}, \varepsilon_{is})]$, denoted by ρ , is the same between any two bids. To simplify estimation, Alberini, Boyle, and Welsh (2003) specify [A5] in terms of a conventional binary choice model with random effects:

$$Y_{ik}^* = X_{ik}\gamma + f(t_{ik})\gamma_t + u_i + \varepsilon_{ik}, \quad i = 1, \dots, N; \quad k = 1, \dots, K \quad [\text{A6}]$$

where $Y_{ik}^* = 1$ for a “yes” response and equals 0 for a “no” response, and $f(t)$ is now included along with the regressors of the WTP function. Although any distribution can be assumed for u_i , it is most common to assume u_i is distributed normal and then apply the estimation procedure of Butler and Moffit (1982). Estimation is fairly complicated and involves using Gauss-Hermite quadrature to maximize the following log-likelihood function:

$$\ln L = \ln \left\{ \frac{1}{\sqrt{\pi}} \int_{-\infty}^{\infty} e^{-r_i^2} \left[\prod_{k=1}^K \{ y_{ik} F(\tilde{X}_{ik}\gamma + \theta r_i) + (1 - y_{ik})(1 - F(\tilde{X}_{ik}\gamma + \theta r_i)) \} \right] dr_i \right\} \quad [\text{A7}]$$

where $\tilde{X}_{ik} = [X_{ik}, f(t_{ik})]$, $\tilde{\gamma} = [\gamma, \gamma_t]$, $\theta = \sigma_u \sqrt{2}$, $r_i = u_i/\theta$, and, as before, $F(\cdot)$ is the distribution of ε and can be freely chosen. Alberini, Boyle, and Welsh (2003) assume $F(\varepsilon)$ is normal. The model can be readily extended to the ordered choice framework by substituting the following for the $\{.\}$ term in [A7]:

$$\left\{ I_{ik}^{DY} F(\tilde{X}_{ik}\tilde{\gamma} - \mu_3 + \theta r_i) + I_{ik}^{PY} \left[(1 - F(\tilde{X}_{ik}\tilde{\gamma} - \mu_3 + \theta r_i)) - (1 - F(\tilde{X}_{ik}\tilde{\gamma} - \mu_2 + \theta r_i)) \right] \right. \\ \left. + I_{ik}^{NS} \left[(1 - F(\tilde{X}_{ik}\tilde{\gamma} - \mu_2 + \theta r_i)) - (1 - F(\tilde{X}_{ik}\tilde{\gamma} - \mu_1 + \theta r_i)) \right] \right. \\ \left. + I_{ik}^{PN} \left[(1 - F(\tilde{X}_{ik}\tilde{\gamma} - \mu_1 + \theta r_i)) - (1 - F(\tilde{X}_{ik}\tilde{\gamma} - \mu_0 + \theta r_i)) \right] + I_{ik}^{DN} (1 - F(\tilde{X}_{ik}\tilde{\gamma} - \mu_0 + \theta r_i)) \right\} \quad [\text{A8}]$$

A parameter of key interest is the correlation coefficient, ρ , which is equal to $\sigma_u^2/(1+\sigma_u^2)$.

Appendix B. Parameter Estimates for “Probably Yes” Random Effects Probit Models, ANGLER Dataset

	Probit, with covariates	Probit
Constant	-0.238 (0.123)	9.713 (0.227)
Bid	-0.016 (0.000)	-0.031 (0.001)
ICEFISH	-0.155 (0.059)	
MARINEF	-1.024 (0.073)	
Age	0.065 (0.003)	
Male	1.212 (0.080)	
Income in thou. \$	-0.002 (0.001)	
Price per trip	0.002 (0.000)	
ρ	0.947 (0.003)	0.985 (0.001)

Notes: See Table II of Alberini, Boyle, and Welsh (2003) for description of covariates. Standard errors are in parentheses.

References

- Alberini, Anna. 1993. Optimal Designs and Discrete Choice Contingent Valuation Surveys: Single Bound, Double-Bounded, and Bivariate Models. *Journal of Environmental Economics and Management* **28**(3): 287-306.
- Alberini, Anna. 1995. Efficiency vs Bias of Willingness-to-Pay Estimates: Bivariate and Interval Data Models. *Journal of Environmental Economics and Management* **29**(2): 169-180.
- Alberini, Anna, Kevin Boyle, and Michael Welsh. 2003. Analysis of Contingent Valuation Data with Multiple Bids and Response Options Allowing Respondents to Express Uncertainty. *Journal of Environmental Economics and Management* **45**(1): 40-62.
- Alberini, Anna, Barbara Kanninen, and Richard T. Carson. 1997. Modeling Response Incentive Effects in Dichotomous Choice Contingent Valuation Data. *Land Economics* **73**: 309-324.
- Bateman, Ian J. Matthew Cole, Philip Cooper, Stavros Georgiou, David Hadley, and Gregory L. Poe. On Visible Choice Sets and Scope Sensitivity. *Journal of Environmental Economics and Management*, in press.
- Bateman, Ian J., Ian H. Langford, Andrew P. Jones, Geoffrey N. Kerr. 2001. Bound and Path Effects in Double and Triple Bounded Dichotomous Choice Contingent Valuation. *Resource and Energy Economics* **23**(3): 191-213.
- Butler, J. S. and Robert Moffit. 1982. A Computationally Efficient Quadrature Procedure for the One-Factor Multinomial Probit Model. *Econometrica* **50**(3): 761-764.
- Cameron, Trudy A. and D. D. Huppert 1989. OLS versus ML Estimation of Non-market Resource Values with Payment Card Interval Data. *Journal of Environmental Economics and Management* **17**: 230-246.
- Cameron, Trudy A. and Michael D. James. 1987. Efficient Estimation Methods for 'Closed-ended' Contingent Valuation Surveys. *The Review of Economics and Statistics* **69**: 269-276.
- Cameron, Trudy A., Gregory L. Poe, Robert G. Ethier, and William D. Schulze. 2002. Alternative Nonmarket Value-Elicitation Methods: Are Revealed and Stated Preferences the Same? *Journal of Environmental Economics and Management* **44**(3): 391-425.
- Cameron, Trudy A. and John Quiggin. 1994. Estimation Using Contingent Valuation Data from a "Dichotomous Choice with Follow-up" Questionnaire. *Journal of Environmental Economics and Management* **27**: 218-234.

- Carson, Richard T., Theodore Groves, and Mark J. Machina. 2000. Incentive and Information Properties of Preference Questions. Working paper, University of California, San Diego.
- Carson, Richard T., W. Michael Hanemann, and Robert C. Mitchell. 1986. The Use of Simulated Political Markets to Value Public Goods. Working paper, University of California, San Diego.
- Carson, Richard T., L. Wilks, and D. Imber. 1994. Valuing the Preservation of Australia's Kakadu Conservation Zone. *Oxford Economic Papers* **46**: 727-749.
- Champ, Patricia A. and Thomas C. Brown. 1997. A Comparison of Contingent and Actual Voting Behavior. *Proceedings from W-133 Benefits and Cost Transfer in Natural Resource Planning, 10th Interim Report*.
- Champ, Patricia A., Richard C. Bishop, Thomas C. Brown, and Daniel W. McCollum. 1997. Using Donation Mechanisms to Value Non-Use Benefits from Public Goods. *Journal of Environmental Economics and Management* **33**(2): 151-163.
- Conover, W. J. 1980. *Practical Nonparametric Statistics*, 2nd Ed. New York: John Wiley and Sons.
- Evans, Mary F., Nicholas E. Flores, and Kevin J. Boyle. 2003. Multiple Bounded Uncertainty Choice Data as Probabilistic Intentions. *Land Economics*, forthcoming.
- Greene, William H. 2002. *Econometric Analysis*, 5th edition. Upper Saddle River, N.J.: Prentice Hall.
- Haab, Timothy C. and Kenneth E. McConnell. 1997. Referendum Models and Negative Willingness to Pay: Alternative Solutions. *Journal of Environmental Economics and Management* **32**(2): 251-270.
- Hanemann, W. Michael, John Loomis, and Barbara Kanninen. 1991. Statistical Efficiency of Double-Bounded Dichotomous Choice Contingent Valuation. *American Journal of Agricultural Economics* **72**: 1255-1263.
- Herriges, Joseph A., and Jason F. Shogren. 1996. Starting Point Bias in Dichotomous Choice Valuation with Follow-Up Questioning. *Journal of Environmental Economics and Management* **30**(1): 112-131.
- Johanesson, M., Glenn C. Blomquist, Karen Blumenschein, Per-Olav Johansson, Bengt Liljas, and Richard M. O'Conor. 1999. Calibrating Hypothetical Willingness to Pay Responses. *Journal of Risk and Uncertainty* **8**:21-32.

- Kanninen, Barbara J. 1995. Optimal Experimental Design for Double Bounded Dichotomous Choice Contingent Valuation. *Land Economics* **69**: 138-146.
- Krinsky, Itzhak, and A. Leslie Robb. 1986. On Approximating the Statistical Properties of Elasticities. *Review of Economics and Statistics* **68**: 715-719.
- Kriström, Bengt. 1990. A Non-Parametric Approach to the Estimation of Welfare Measures in Discrete Response Valuation Studies. *Land Economics* **66**(2): 135-139.
- Kriström, Bengt. 1997. Spike Models in Contingent Valuation. *American Journal of Agricultural Economics* **79**(3): 1013-1023.
- Li, Chuan-Zhong, and Leif Mattson. 1995. Discrete Choice under Preference Uncertainty: An Improved Structural Model for Contingent Valuation. *Journal of Environmental Economics and Management* **28**: 256-269.
- Maddala, G. S. 1987. Limited Dependent Variable Models Using Panel Data. *Journal of Human Resources* **22**(3): 305-338.
- Magelby, Daniel. 1989. Opinion Formation and Opinion Change in Ballot Proposition Campaigns, in *Manipulating Public Opinion*, ed. M. Margolis and G Mauser. Pacific Grove, CA: Brooks/Cole Publishers.
- Mitchell, Robert C. and Richard T. Carson. 1989. *Using Surveys to Value Public Goods: The Contingent Valuation Method*. Washington, D.C.: Resources for the Future.
- Opaluch, John J. and Kathleen Segerson. 1989. Rational Roots of 'Irrational' Behavior: New Theories of Economic Decision-Making. *Northeastern Journal of Agricultural and Resource Economics* **18**(2):81-95.
- Poe, Gregory L., Jeremy R. Clark, Daniel Rondeau, and William D. Schulze. 2002. Provision Point Mechanisms and Field Validity Tests of Contingent Valuation. *Environmental and Resource Economics* **23**(1): 105-131.
- Poe, Gregory L., Eric K. Severance-Lossin, and Michael P. Welsh. 1994. Measuring the Difference (X-Y) of Simulated Distributions: A Convolutions Approach. *American Journal of Agricultural Economics* **76**(4): 904-915.
- Poe, Gregory L., and Christian A. Vossler. 2002. Monte Carlo Benchmarks for Discrete Response Valuation Methods: Comment. *Land Economics* **78**(4): 605-616.
- Ready, Richard C., John C. Whitehead, and Glenn C. Blomquist. 1995. Contingent Valuation when Respondents are Ambivalent. *Journal of Environmental Economics and Management* **29**(2): 181-196.

- Turnbull, Bruce W. 1974. Nonparametric Estimation of a Survivorship Function with Doubly Censored Data. *Journal of the American Statistical Association* **69**(345): 169-173.
- Turnbull, Bruce W. 1976. The Empirical Distribution Function with Arbitrarily Grouped, Censored and Truncated Data. *Journal of the Royal Statistical Society - Series B (Methodological)* **38**(3):290-295.
- Vossler, Christian A. and Joe Kerkvliet. 2003. A Criterion Validity Test of the Contingent Valuation Method: Comparing Hypothetical and Actual Voting Behavior for a Public Referendum. *Journal of Environmental Economics and Management* **45**(3): 631-649.
- Vossler, Christian A., Joe Kerkvliet, Stephen Polasky, and Olesya Gainutdinova. 2003a. Externally Validating Contingent Valuation: An Open-Space Survey and Referendum in Corvallis, Oregon. *Journal of Economic Behavior and Organization* **51**(2): 261-277.
- Vossler, Christian A., Gregory L. Poe, Robert G. Ethier, and Michael P. Welsh. 2003b. Payment Certainty in Discrete Choice Contingent Valuation Responses: Results from a Field Validity Test. *Southern Economic Journal* **69**(4): 886-902.
- Vossler, Christian A., Gregory L. Poe, Michael P. Welsh, and Robert G. Ethier. 2003c. Assessing Bid Position Bias in Multiple Bounded Discrete Choice Valuation Questions. Unpublished Manuscript.
- Wang, Hua. 1997. Treatment of "Don't-Know" Responses in Contingent Valuation Surveys: A Random Valuation Model. *Journal of Environmental Economics and Management* **32**(2): 219-232.
- Welsh, Michael P., Richard C. Bishop, M. L. Phillips, and R. M. Baumgartner. 1995. GCES Non-Use Value Study: Final Report. Report to Glen Canyon Environmental Studies, U.D. Bureau of Reclamation, Hagler Bailly Consulting Inc., Madison, WI.
- Welsh, Michael P. and Gregory L. Poe. 1998. Elicitation Effects in Contingent Valuation: Comparisons to a Multiple Bounded Discrete Choice Approach. *Journal of Environmental Economics and Management* **36**: 170-185.
- White, H. 1982. Maximum Likelihood Estimation of Misspecified Models. *Econometrica* **53**(1): 1-16.
- Whitehead, John C. 2002. Incentive Incompatibility and Starting-Point Bias in Iterative Valuation Questions. *Land Economics* **78**: 285-297.