

# MPRA

Munich Personal RePEc Archive

## **Social Learning and the Shadow of the Past**

Yuval Heller and Erik Mohlin

Bar Ilan University, Lund University

28 April 2017

Online at <https://mpa.ub.uni-muenchen.de/81434/>

MPRA Paper No. 81434, posted 18 September 2017 17:45 UTC

# Social Learning and the Shadow of the Past\*

Yuval Heller<sup>†</sup> and Erik Mohlin<sup>‡</sup>

September 13, 2017

## Abstract

In various environments new agents may base their decisions on observations of actions taken by a few other agents in the past. In this paper we analyze a broad class of such social learning processes, and study under what circumstances the initial behavior of the population has a lasting effect. Our results show that this question strongly depends on the expected number of actions observed by new agents. Specifically, we show that if the expected number of observed actions is: (1) less than one, then the population converges to the same behavior independently of the initial state; (2) between one and two, then in some (but not all) environments there are learning rules for which the initial state has a lasting impact on future behavior; and (3) more than two, then in all environments there is a learning rule for which the initial state has a lasting impact.

**Keywords:** Social learning, steady state, unique limiting behavior, path dependence. **JEL Classification:** C73, D83.

## 1 Introduction

Agents must often make decisions without knowing the cost and benefits of the possible choices. In such situations an inexperienced (or “newborn”) agent may learn from the experience of others, by basing his decision, on observations of a few actions taken by other agents in the past (see, e.g., the social learning models of [Ellison & Fudenberg, 1993, 1995](#); [Acemoglu \*et al.\*, 2011](#)). In other environments, agents interact with random opponents, and an agent may base his choice of action on a few observations of how his current opponent behaved in the past (as first described in [Rosenthal, 1979](#), and further developed and applied to various models of community enforcement in the Prisoner’s Dilemma game in ([Nowak & Sigmund, 1998](#); [Takahashi, 2010](#); [Heller & Mohlin, 2017](#))).

In this paper we analyze a broad class of social learning processes, and study under what circumstances the initial behavior of the population has a lasting influence on the population’s behavior in the long run. Our results show that the answer to this question strongly depends on the expected number of actions observed by new agents. Specifically, we show that: (1) if the mean sample size (expected number of actions observed by

---

\*The paper was previously titled “When Is Social Learning Path-Dependent?”. This paper replaces an obsolete working paper titled “Unique Stationary Behavior” that presented related results in a narrower setup. We thank Ron Peretz, Doron Ravid, Satoru Takahashi, Xiangqian Yang, and Peyton Young for valuable discussions and suggestions. Yuval Heller is grateful to the European Research Council for its financial support (ERC starting grant #677057). Erik Mohlin is grateful to Handelsbankens Forskningsstiftelser (grant #P2016-0079:1) and the Swedish Research Council (grant #2015-01751) for their financial support.

<sup>†</sup>Affiliation: Department of Economics, Bar Ilan University, Israel. E-mail: [yuval.heller@biu.ac.il](mailto:yuval.heller@biu.ac.il).

<sup>‡</sup>Affiliation: Department of Economics, Lund University, Sweden. E-mail: [erik.mohlin@nek.lu.se](mailto:erik.mohlin@nek.lu.se).

a new agent) is less than one then the population converges to the same behavior independently of the initial state; (2) if the mean sample size is between one and two then any environment allows for a rule with multiple steady states according to which agents learn from the experience of other agents, but only some environments allow for learning rules with multiple locally stable states; and (3) if the mean sample size is more than two then all environments admit a learning rule with multiple locally stable states, and the initial state determines which steady states will prevail.

**Overview of the Model** We consider an infinite population of agents (a continuum of mass one). Time is discrete and in every period each agent is faced with a choice among a fixed set of alternatives. The *population state* is a distribution of actions describing the aggregate behavior of agents in the population. In each period a fixed share of the agents die and are replaced with new agents. Each new agent observes a finite sequence of actions (called a *sample*) of random size. The sample may consist either of past actions of random agents in the population (as in the social learning models mentioned above) or past actions of the current, randomly drawn, opponent (as in the community enforcement models mentioned above).

An environment is a tuple that specifies all the above components. A learning rule specifies the distribution of actions played by new agents as a function of the observed sample. The agent keeps playing the same action throughout his life. The environment and the learning rule jointly induce a mapping between population states that determines a new population state for each initial state. A population state is a *steady state* if it is a fixed point of this mapping.

**Characterization of Multiple Steady States** Theorem 2 fully characterizes in which environments there exist learning rules that admit multiple steady states. Specifically, it shows that an environment allows for a learning rule that admits multiple steady states if and only if the mean sample size is strictly more than one (or if agents always observe a single action). In the opposite case, each learning rule admits a unique steady state, and, moreover, the population converges to the unique steady state from any initial state.

The intuition for the “only if” side is as follows. Consider two different initial population states of an environment with a mean sample size below one. The aggregate behavior of new agents may differ only to the extent in which they observe different distributions of samples. This implies that the distance between the distributions of actions played by new agents is bounded by the distance between the distributions of samples that they observe. One can show that this latter distance is bounded by the distance between the distributions of actions played by the incumbents multiplied by the mean sample size. Hence, if the mean sample size is less than one, the distance between the distributions of actions of new agents is strictly less than the distance between the distributions of actions of the incumbents, which implies that the mapping between population states is a contraction mapping.

The “if” side relies on constructing a specific learning rule, according to which agents play action  $a'$  if they observe action  $a'$  in their sample, and play action  $a''$  otherwise. One can show that such a learning rule always admits two different steady states provided that the expected number of observed actions is greater than one. We demonstrate that this learning rule (as well as all other learning rules used in the other results in the paper) may be consistent with Bayesian inference and the agents using best replies.

**Characterization of Multiple Locally Stable States** A steady state is locally stable if the population converges back to this state after any sufficiently small perturbation. Arguably, the initial state can be said

to have a lasting effect only when there are multiple locally stable states. In particular, in the construction of the above result (Theorem 2), only one of the steady states is locally stable. Moreover, one can show that a population converges to this state from almost all initial states. Our remaining results characterize when there are learning rules that admit multiple locally stable states.

Theorem 3 shows that in any environment in which the mean sample size is larger than two it is possible to define a learning rule that admits multiple locally stable states. According to this learning rule, each new agent (1) plays action  $a'$  if he observes action  $a'$  at least twice in his sample, (2) plays action  $a''$  if he never observes action  $a'$ , and (3) mixes between the two actions if he observes action  $a'$  exactly once. We show that this learning rule (with a well-calibrated mixing probability) admits two locally stable states, one in which  $a'$  is never played, and the other in which it is played with a positive probability.

Our next two results show that when the mean sample size is between one and two, then some (but not all) environments allow for learning rules that admit multiple locally stable states. Specifically, we show that if each new agent observes at most two actions, and there are two feasible actions, then any learning rule admits a unique locally stable state, and, moreover, the population converges to this state from almost all initial states. The intuition is that when the sample size is at most two, then the mapping induced by the learning process can be represented as a polynomial of degree two. Hence, the mapping can have at most two steady states, and it is relatively straightforward to show that at most one of these states can be locally stable.

Finally, we show that an environment with two feasible actions  $a', a''$  in which some agents observe a single action, while others observe three actions, and each new agent chooses the frequently observed action in his sample, admits two locally stable states: one in which all agents choose action  $a'$ , and another in which everyone chooses action  $a''$  (in addition, to an unstable state in which half of the population plays each action).

**Extensions** Our results so far have not assumed anything about the agents' learning rules. Obviously, additional information on the learning rules, may allow us to achieve stronger results. Next, we present a simple notion that measures how responsive a learning rule is to different samples. For example, a learning rule might be relatively unresponsive due to new agents having strong priors about which action is best. We use this notion of responsiveness to define the effective sample size of a learning process (which is always weakly smaller than the simple mean sample size). Next, we apply the notion of effective sample size to derive a tighter upper bound for learning processes that admit unique steady states.

Finally, we extend our model and main results to (1) heterogeneous populations in which agents are endowed with different types, and the various types differ in their sample sizes and learning rules, (2) non-stationary environments, in which the distribution of sample sizes and the agents' learning rules depend on calendar time, and (3) stochastic shocks that influence the learning rules of all agents (at the aggregate level).

**Related Literature** Various papers have studied different aspects of the question of when the initial behavior of the population has lasting effects on social learning processes. Most of this literature focuses on specific learning rules, according to which new (or revising) agents myopically best reply to the empirical frequency of the observed actions. Arthur (1989) (see related models and extensions in Arthur, 1994; Kaniovski & Young, 1995; Smith & Sorensen, 2014) studies games in which agents sequentially choose which competing technology to adopt, and he shows that social learning is path-dependent if the technologies have increasing returns.

Kandori *et al.* (1993) and Young (1993a) study models of finite large populations that are involved in a social learning process, and agents occasionally make mistakes (e.g., an agent adopts a technology that is not

his myopic best reply to his sampled information). They show that the path dependency of the social learning process vanishes when infinite time horizons are considered. In many cases, when the probability of mistakes is sufficiently small the population spends almost all the time in a unique “stochastically stable state,” which is independent of the initial state. A key difference between our model and theirs is that we model an infinite population, rather than a large finite population. In Section 6, we discuss the relations between the present paper and the literature on stochastic stability, and, in particular, the implications of our results for finite large populations.

Banerjee & Fudenberg (2004) study a model with a continuum of agents in which a fixed share of new agents in each period choose one of two technologies. There are two possible states of nature, and each technology has a higher quality in one of these states. Each agent, after he observes  $l$  past actions and a noisy signal about the quality of each technology, chooses the technology with the higher expected quality, conditional on the information that he has observed. Banerjee & Fudenberg show that when  $l \geq 2$  the behavior of the population converges to everyone choosing the efficient technology, while if  $l = 1$  the population converges to an inefficient state in which only some of the agents choose the (ex-post) better technology.

Sandholm (2001) shows that when each new agent observes  $k$  actions and the game admits a  $\frac{1}{k}$ -dominant action  $a^*$  (i.e., action  $a^*$  is the unique best reply against any mixed strategy assigning a mass of at least  $\frac{1}{k}$  to  $a^*$ ), then social learning converges to this action regardless of the initial state. Recently, Oyama *et al.* (2015) strengthened this result by extending it to iterated  $p$ -dominant actions, and by showing that global convergence is fast.

Our model differs from all the above-mentioned research in that we study general environments and arbitrary learning rules. Specifically, we ask what properties of the agents’ sampling procedures imply that *any* learning rule admits a unique steady state and global convergence to this state, whereas the existing literature focuses on the dynamic behavior induced by a specific learning rule (in most of the literature, the agents myopically best reply to specific payoffs, such as those induced by competing technologies with increasing returns).

**Structure** We present motivating examples in Section 2. The basic model is described in Section 3. Section 4 presents our main results. In Section 5 we define and apply the notion of a responsiveness of a learning rule. We conclude in Section 6. Appendix A extends the basic model to heterogeneous populations, non-stationary processes, and common stochastic shocks. Technical proofs are presented in Appendix B–.

## 2 Motivating Examples

In this section we present three motivating examples, which will be revisited further below to demonstrate the applicability of our model and the implications of our results. In all the examples the population is modeled as a continuum of mass one, and time is discrete. The main example deals with social learning with competing technologies, while the other two examples study situations in which agents are randomly matched to play a two-player game.

**Example 1** (Main Motivating Example: Competing Technologies with Increasing Returns<sup>1</sup>). Consider a population in which in each period a share  $\beta \in (0, 1)$  of the incumbent agents die, and are replaced with new

<sup>1</sup>The example is similar to the model of Banerjee & Fudenberg (2004), except that the technologies have increasing returns, rather than having unknown inherent different qualities.

agents. Each new agent chooses one of two competing technologies  $a'$  and  $a''$ , which he adopts for the rest of his life. The technologies have increasing returns. A share of 99% of the new agents observe the technology followed by a single random incumbent, and then they choose to adopt this observed technology.

We consider two cases for what the remaining 1% of the new agents observe before they choose a technology (as summarized in Table 1):

1. They observe nothing, and in this case half of the new agents adopt technology  $a'$ , and the other half adopt technology  $a''$ .
2. They observe the technologies adopted by three random incumbents, and in this case each new agent adopts the technology chosen by the majority of his sample.

Let  $\alpha_1 \in [0, 1]$  describe the share of agents who use technology  $a'$  initially (in the first period). One can show that in Case (1), in which the mean sample size of a new agent is slightly less than one, the population converges to a unique steady state in which half of the agents follow each technology. By contrast, in Case (2), in which the mean sample size is slightly more than one, the initial behavior of the population has a lasting effect. Specifically, the population converges to everyone following technology  $a'$  if initially a majority of the agents followed technology  $a'$  (i.e., if  $\alpha_1 > 50\%$ ), and the population converges to everyone following technology  $a''$  if  $\alpha_1 < 50\%$ .

Table 1: Summary of the Two Cases in Example 1

Case	Probability of Observing			Mean Sample Size	Convergence and steady states
	0 actions	1 action	3 actions		
1	1%	99%	-	0.99	Global convergence to 50%–50%
2	-	99%	1%	1.02	Convergence to $a'$ if $\alpha_1 > 0.5$ ; convergence to $a''$ if $\alpha_1 < 0.5$ .

We conclude this example by noting that the described behavior is consistent with each new agent playing a unique best reply, if we make the following further assumptions. Nature privately chooses the initial share of agents who follow each technology in the first period. In each later period, new agents have a symmetric common prior on this initial share (e.g., the common prior might be that  $\alpha_1$  is uniformly distributed on<sup>2</sup>  $[0, 1]$ ). The payoff of each new agent is increasing in the current share of agents who follow the same technology (i.e., the technologies have increasing returns), and hence agents have an incentive to play the action that they believe that the majority is playing. In addition, half of the agents have a weak preference for technology  $a'$ , and the remaining half have a weak preference for technology  $a''$ . These idiosyncratic preferences are much smaller than the payoff differences due to the increasing returns. For example, assume that if 51% of the population follow technology  $a'$ , then an agent prefers technology  $a'$  regardless of his own idiosyncratic preference.

**Example 2** (Community Enforcement in the Prisoner’s Dilemma). Consider a population such that in each round each agent is randomly matched with three opponents, and plays a Prisoner’s Dilemma with each of them. In round one, each agent defects with probability  $\alpha$  in each match. In any later round and match, with a probability of 95% each agent observes two actions played in the previous period by the current opponent

<sup>2</sup>As argued by Banerjee & Fudenberg (2004, p. 5), the aggregate uncertainty about the initial population state may reflect the choices of a group of “early adopters” whose preferences are uncertain even at the aggregate level.

(i.e., actions played by the current opponent against two of his three opponents in the previous period). With the remaining probability of 5% each agent observes  $k$  actions played by the current opponent in the previous period. We consider two cases: (1)  $k = 1$ , and (2)  $k = 3$ . All agents follow the same behavior (in both cases): (I) an agent defects if he observes his partner defecting more times than cooperating, (II) an agent cooperates if he observes his partner cooperating more times than defecting, and (III) an agent defects with probability 51% if he observes the partner defecting and cooperating an equal number of times.

One can show that in both cases the environment admits two steady states: one in which all agents cooperate, and one in which all agents defect. In the first case ( $k = 1$ , in which the mean sample size is slightly below 2), only the state in which all agents defect is locally stable. Specifically, the population converges to everyone defecting for any  $\alpha > 0$ . By contrast, in the second case ( $k = 3$ , in which the mean sample size is slightly above 2), both steady states are locally stable. In particular, one can show that the population converges to everyone defecting if  $\alpha > 31\%$ , and it converges to everyone cooperating if  $\alpha < 31\%$ .

**Example 3** (Rock-Paper-Scissors). Consider a population in which each agent is randomly matched in each round to play the rock-paper-scissors game. Each player has three pure actions ( $r, p, s$ ), and each action is the unique best reply to the previous action (modulo 3). In the initial round  $t = 1$  the aggregate distributions of actions is  $\hat{\gamma} \in \Delta(r, p, s)$ . In each later round, each agent observes the opponent’s action in the previous round with probability  $q$ , and best replies to the observed action. With the remaining probability of  $1 - q$  the agent observes no actions, and plays the mixed action  $\gamma' \in \Delta(r, p, s)$ .

Observe that the parameter  $q$  is equal to the mean sample size of a random agent. If  $q = 1$  it is immediate that the population’s behavior cycles “around” permutations of the initial behavior (as is common in evolutionary models of rock-paper-scissors; see, e.g., the analysis in [Cason et al., 2014](#)). Formally, let  $t \in \{0, 1, 2, \dots\}$ , then in round  $3 \cdot t + 1$  (resp.,  $3 \cdot t + 2$ ,  $3 \cdot t + 3$ ) agents play  $r$  with a probability of  $\hat{\gamma}(r)$  (resp.,  $\hat{\gamma}(s)$ ,  $\hat{\gamma}(p)$ ),  $p$  with a probability of  $\hat{\gamma}(p)$  (resp.,  $\hat{\gamma}(r)$ ,  $\hat{\gamma}(s)$ ), and  $s$  with a probability of  $\hat{\gamma}(s)$  (resp.,  $\hat{\gamma}(p)$ ,  $\hat{\gamma}(r)$ ). By contrast, when  $q < 1$ , one can show that the population converges to the following unique behavior (regardless of the initial behavior  $\hat{\gamma}$ ):

$$\Pr(r) = \frac{\gamma'(r) + p\gamma'(s) + q^2\gamma'(p)}{1 + q + q^2}, \Pr(p) = \frac{\gamma'(p) + q\gamma'(r) + q^2\gamma'(s)}{1 + q + q^2}, \Pr(s) = \frac{\gamma'(s) + q\gamma'(p) + q^2\gamma'(r)}{1 + q + q^2}.$$

Note that when  $q$  is close to one, the unique behavior is close to the uniform mixed profile that assigns a probability of  $\frac{1}{3}$  to each action.

### 3 Model

Throughout the paper we restrict attention to distributions with a finite support. Given a (possibly infinite) set  $X$ , let  $\Delta(X)$  denote the set of distributions over this set that have a finite support. With slight abuse of notation we use  $x \in X$  to denote the degenerate distribution  $\mu \in \Delta(X)$  that assigns probability one to  $x$  (i.e., we write  $\mu \equiv x$  if  $\mu(x) = 1$ ). We use  $\mathbb{N}$  to denote the set of natural numbers including zero.

**Population state.** Consider an infinite population of agents. More precisely, the population consists of a continuum of agents with mass one. Time is discrete and in every period (or “round”) each agent is faced with a choice among a fixed set of alternatives  $A$ . Let  $A$  be a finite set of at least two actions (i.e.,  $|A| \geq 2$ ).

The *population state* (or *state* for short) is identified with the aggregate distribution of actions played in the population, denoted  $\gamma \in \Delta(A)$ . Let  $\Gamma$  denote the set of all population states.

**New/Revising agents.** In each period, a share of  $0 < \beta \leq 1$  of the agents exit the population and are replaced with new agents, while the remaining  $1 - \beta$  share of the agents play the same action as they played in the past (see, e.g., Banerjee & Fudenberg, 2004). Each new agent chooses an action based on a sample of a few actions of incumbents. The agent then keeps playing this chosen action throughout his active life, possibly because the initial choice requires a substantial action-specific investment, and it is too costly for an agent to reinvest in a different action later on. The model can also be interpreted as describing a fixed population in which each agent reevaluates his action only every  $\frac{1}{\beta}$  periods (under this interpretation, when the sample is non-empty, the first observed action after a revision might be interpreted as the revising agent's own past action).<sup>3</sup>

**Sample.** Each new agent observes a finite sequence of actions (or *sample*). The size of the observed sample is a random variable with a distribution  $\nu \in \Delta(\mathbb{N})$ . Let  $M$  denote the set of all feasible samples, i.e.,  $M = \cup_{l \in \text{supp}(\nu)} A^l$ , where  $A^0 = \{\emptyset\}$  is a singleton consisting of the empty sample  $\emptyset$ .

Let  $\bar{l} = \max(\text{supp}(\nu)) < \infty$  be the maximal sample size. Note that  $M$  is finite in virtue of the finite-support assumption. For each sample size  $l \in \mathbb{N}$ , let  $\psi_l : \Gamma \rightarrow \Delta(A^l)$  denote the distribution of samples observed by each agent in the population (or *sampling rule* for short), conditional on the sample having size  $l$ . A typical sample of size  $l$  is represented by the vector  $\vec{a} = (a_1, \dots, a_l)$ .

We assume that each agent independently samples different agents, and observes a random action played by each of these agents. This kind of sampling is common in models of social learning (see, e.g., Ellison & Fudenberg, 1995; Banerjee & Fudenberg, 2004). Formally, we define for each sample size  $l \in \mathbb{N}$ , each state  $\gamma \in \Gamma$ , and each sample  $(a_1, \dots, a_l)$ ,

$$\psi_{l,\gamma}(a_1, \dots, a_l) = \prod_{1 \leq i \leq l} \gamma(a_i). \quad (1)$$

**Environment.** An *environment* is a tuple  $E = (A, \beta, \nu)$  that includes the three components described above: a finite set of actions  $A$ , a fraction of new agents at each stage  $\beta$ , and a distribution of sample sizes  $\nu$ .

Given environment  $E = (A, \beta, \nu)$ , let  $\mu_l$  denote the *mean sample size*, i.e., the expected number of actions observed by a random agent in the population. Formally:

$$\mu_l = \sum_{l \in \text{supp}(\nu)} \nu(l) \cdot l.$$

**Learning rule and stationary learning process.** Each new agent chooses his action in the new population state by following a stationary (i.e., time-independent) *learning rule*  $\sigma : M \rightarrow \Delta(A)$ . That is, a new agent who observes sample  $m \in M$  plays action  $a$  with probability  $\sigma_m(a)$ . The remaining  $1 - \beta$  agents play the same action as in the previous round.

A *learning process* is a pair  $P = (E, \sigma)$  consisting of an environment and a learning rule.

---

<sup>3</sup>The learning process is unaffected by having the first observed action be the agent's own past action, due to the fact that the population is a continuum, and the behavior of the revising agents coincides with the aggregate behavior in the population.

**Population dynamics.** An initial state and a learning process uniquely determine a new state. To see this note that since the number of messages  $M$ , and actions  $A$  are finite, whereas the population is a continuum, the *probability* that an agent observes a message  $m$  and switches to an action  $a$  is equal to the *fraction* of agents who observe a message  $m$  and switch to an action  $a$ . For this reason we say that the learning process is *deterministic*, despite the fact that the choice of an individual agent may be stochastic.

Time is discrete in our model. Let  $f_P : \Gamma \rightarrow \Gamma$  denote the mapping between states induced by a single step of the learning process  $P$ . That is,  $f_P(\hat{\gamma})$  is the new state induced by a single step of the process  $P$ , given an initial state  $\hat{\gamma}$ . Similarly, for each  $t > 1$ , let  $f_P^t(\hat{\gamma})$  denote the state induced after  $t$  steps of the learning process  $P$ , given an initial state  $\hat{\gamma}$  (e.g.,  $f_P^2(\hat{\gamma}) = f_P(f_P(\hat{\gamma}))$ ,  $f_P^3(\hat{\gamma}) = f_P(f_P(f_P(\hat{\gamma})))$ , etc.).

**$L_1$ -distance.** Throughout the paper we measure distances with the  $L_1$ -distance (norm). Specifically, let the  $L_1$ -distance between two distributions of samples  $\psi_{l,\gamma}, \psi_{l,\gamma'} \in \Delta(A^l)$  of size  $l$ , be defined as follows:

$$\|\psi_{l,\gamma} - \psi_{l,\gamma'}\|_1 = \sum_{m \in A^l} |\psi_{l,\gamma}(m) - \psi_{l,\gamma'}(m)|.$$

Similarly the  $L_1$ -distance between two distributions of actions  $\gamma, \gamma' \in \Delta(A)$  is defined as follows:

$$\|\gamma - \gamma'\|_1 = \sum_{a \in A} |\gamma(a) - \gamma'(a)|.$$

**Steady States and Stability** We say that  $\gamma^*$  is a *steady state* with respect to the stationary learning process  $P$ , if it is a fixed point of the induced mapping  $f_P$ , i.e., if  $f_P(\gamma^*) = \gamma^*$ .

Steady state  $\gamma^*$  is (asymptotically) locally stable if a population beginning near  $\gamma^*$  remains close to  $\gamma^*$ , and eventually converges to  $\gamma^*$ . Formally, for each  $\epsilon > 0$  there exists  $\delta > 0$  such that  $\|\hat{\gamma} - \gamma^*\| < \delta$  implies that:

$$(1) \ \|f_P^t(\hat{\gamma}) - f_P^t(\gamma^*)\| < \epsilon \ \forall t \geq 1 \quad , (2) \ \lim_{t \rightarrow \infty} f_P^t(\hat{\gamma}) = \gamma^*.$$

Steady state  $\gamma^*$  is an (*almost-*)*global attractor*, if the population converges to  $\gamma^*$  from any (interior) initial state, i.e., if

$$\lim_{t \rightarrow \infty} f_P^t(\hat{\gamma}) = \gamma^* \text{ for all } \hat{\gamma} \in \Gamma \ (\hat{\gamma} \in \text{Int}(\Gamma)),$$

where  $\text{Int}(\Gamma)$  denotes the set of totally mixed distributions of actions (distributions that assign positive probability to all actions).

We conclude by demonstrating how the model captures motivating Examples 1–3.

**Example 1** (Competing Technologies revisited). The environment in which agents choose to adopt one of two competing increasing returns technologies is modeled by a learning process  $P = (\{a', a''\}, \beta, \nu, \sigma)$ , in which  $\{a', a''\}$  is the set of competing technologies,  $\beta \in (0, 1)$  is the share of new agents that join the population in each round, and the learning rule of the agent is defined as

$$\sigma(\vec{a}) = \begin{cases} 0.5 \cdot a' + 0.5 \cdot a'' & \vec{a} = \emptyset \\ a' & \vec{a} \in \{a', (a', a', a'), (a'', a', a'), (a', a'', a'), (a', a', a'')\} \\ a'' & \text{otherwise.} \end{cases}$$

The initial population state is given by  $\hat{\gamma}(a') = \alpha$ . Finally, the distribution of sample size is:

$$\text{Case I: } \nu(l) = \begin{cases} 1\% & l = 0 \\ 99\% & l = 1 \end{cases} \quad \text{Case II: } \nu(l) = \begin{cases} 1\% & l = 3 \\ 99\% & l = 1. \end{cases}$$

Observe that the mean sample size ( $\mu_l$ ) is equal to 0.99 in Case I, and is equal to 1.02 in Case II.

**Example. 2** (Prisoner's Dilemma revisited). The environment in which agents play the Prisoner's Dilemma is modeled by a learning process

$$P = (\{c, d\}, \beta = 1, \nu, \sigma),$$

where  $\nu(2) = 95\%$ , and in Case (1)  $\nu(1) = 5\%$ , while in Case (2)  $\nu(3) = 5\%$ . In Case (1) the learning rule is given by:

$$\sigma(c, c) = \sigma(c) = c, \quad \sigma(d, d) = \sigma(d) = d, \quad \sigma(c, d) = \sigma(d, c) = 51\% \cdot d + 49\% \cdot c.$$

In Case (2) the learning rule is given by:

$$\sigma(c, c) = \sigma(c, c, c) = \sigma(c, c, d) = \sigma(c, d, c) = \sigma(d, c, c) = c,$$

$$\sigma(d, d) = \sigma(d, d, d) = \sigma(d, d, c) = \sigma(d, c, d) = \sigma(c, d, d) = d, \quad \sigma(c, d) = \sigma(d, c) = 51\% \cdot d + 49\% \cdot c.$$

Observe that  $\mu_l = 1.95$  in Case (1), and  $\mu_l = 2.05$  in Case (2).

**Example 3** (Rock-Paper-Scissors revisited) The environment in which agents play the rock-paper-scissors game is modeled by a learning process  $P = (\{r, p, s\}, \beta = 1, \nu, \sigma)$ , and the initial population state is given by  $\hat{\gamma}$ . The distribution of the sample size and the learning rule are given by

$$\nu(l) = \begin{cases} 0 & 1 - p \\ 1 & p, \end{cases} \quad \sigma(a) = \begin{cases} r & a = s \\ p & a = r \\ s & a = p. \end{cases} \quad \sigma(\emptyset) = \gamma'.$$

Observe that the mean sample size is equal to  $p$  (i.e.,  $\mu_l = p$ ).

## 4 Main Results

### 4.1 Upper Bound on the Distance between New States

Our first result shows that the distance between two new states is at most  $1 - \beta + \beta \cdot \mu_l$  times the distance between the two initial states. The intuition is as follows. Consider two different initial population states  $\gamma$  and  $\gamma'$ . The incumbents who do not die (a share of  $1 - \beta$ ) continue to behave as before, and hence the distance between the distributions of actions of those agents remains  $\|\gamma - \gamma'\|$ . The aggregate behavior of new agents may differ only to the extent that they face different distributions of samples. This implies that the distance between the distributions of actions played by new agents is bounded by the distance between

the distributions of samples that they observe. By using the triangle inequality one can show that this latter distance is bounded by the distance between the distributions of actions played by the incumbents ( $\|\gamma - \gamma'\|$ ) multiplied by the mean sample size ( $\mu_l$ ). Finally, due to another use of a triangle inequality, this implies that the distance between the new population states is at most  $(1 - \beta + \beta \cdot \mu_l) \cdot \|\gamma - \gamma'\|_1$ . Formally,

**Theorem 1.** *Let  $P = (A, \beta, \nu, \sigma)$  be a learning process, and let  $\gamma \neq \gamma' \in \Gamma$  be two population states. Then:*

$$\|f_P(\gamma) - f_P(\gamma')\|_1 \leq (1 - \beta + \beta \cdot \mu_l) \cdot \|\gamma - \gamma'\|_1,$$

with a strict inequality if there exists an  $l > 1$  such that  $\nu(l) > 0$ .

(Sketch of proof. Formal proof is presented for the more general result of Theorem 8 in Appendix B.1.)

The distance between the final population states is bounded as follows:

$$\|(f_P(\gamma)) - (f_P(\gamma'))\|_1 \leq \beta \cdot \sum_{l \in \mathbb{N}} \nu(l) \cdot \|\psi_{l,\gamma} - \psi_{l,\gamma'}\|_1 + (1 - \beta) \cdot \|\gamma - \gamma'\|_1. \quad (2)$$

The intuition of this inequality is as follows. The first part of the RHS of Eq. (2) reflects the actions played by the  $\beta$  new agents. The social learning stage may induce different behaviors for new agents who observe samples of size  $l$  only if they observe different samples. Thus, taking the weighted average of the distances between samples yields the bound on how much the aggregate behaviors of the new agents may differ (i.e.,  $\sum_{l \in \mathbb{N}} \nu(l) \cdot \|\psi_{l,\gamma} - \psi_{l,\gamma'}\|_1$ ). Finally, the mixed average of this expression and the behavior of the incumbents, gives the total bound on the difference between the final population states.

Next, observe that the distance between distributions of samples is bounded by the sample size times the distance between the distributions of actions:

$$\|\psi_{l,\gamma} - \psi_{l,\gamma'}\|_1 \leq l \cdot \|\gamma - \gamma'\|_1,$$

with a strict inequality if  $l > 1$ . This is so because the event that two samples of size  $l$  differ is (a non-disjoint) union of the  $l$  events: the first action in the samples differ, the second action in the samples differs, ..., the last  $l^{\text{th}}$  action in the samples differ.

Substituting the second inequality in (2) yields:

$$\|(f_P(\gamma)) - (f_P(\gamma'))\|_1 \leq \beta \cdot \sum_{l \in \mathbb{N}} \nu(l) \cdot l \cdot \|\gamma - \gamma'\|_1 + (1 - \beta) \cdot \|\gamma - \gamma'\|_1 =$$

$$\left( \beta \cdot \left( \sum_{l \in \mathbb{N}} \nu(l) \cdot l \right) + (1 - \beta) \right) \cdot \|\gamma - \gamma'\|_1 = (\beta \cdot \mu_l + 1 - \beta) \cdot \|\gamma - \gamma'\|_1 = (1 - \beta + \beta \cdot \mu_l) \cdot \|\gamma - \gamma'\|_1,$$

with a strict inequality if there exists an  $l > 1$  such that  $\nu(l) > 0$ . □

Observe that  $1 - \beta + \beta \cdot \mu_l < 1$  iff  $\mu_l < 1$ . Recall that mapping  $f$  is a *weak contraction* (or shrinking) if  $\|(f(\gamma)) - (f(\gamma'))\| < \|\gamma - \gamma'\|$  for each  $\gamma \neq \gamma'$ . Theorem 1 implies that  $f_P$  is a weak contraction mapping if either (1)  $\mu_l < 1$ , or (2)  $\mu_l = 1$  and<sup>4</sup>  $\nu(1) < 1$ . The fact that the mapping  $f_P$  is a weak contraction mapping

---

<sup>4</sup>Note that  $\mu_l = 1$  and  $\nu(1) < 1$  jointly imply that there exists  $l > 1$  such that  $\nu(l) > 0$ .

implies that  $f_P$  admits a global attractor.<sup>5</sup> Formally:

**Corollary 1.** *Let  $P = (A, \beta, \nu, \sigma)$  be a stationary learning process satisfying (1)  $\mu_l < 1$ , or (2)  $\mu_l = 1$  and  $\nu(1) < 1$ . Then  $f_P$  is a weak contraction mapping, which implies that (1)  $f_P$  admits a unique steady state  $\gamma^*$ , and (2) this unique steady state  $\gamma^*$  is a global attractor (i.e.,  $\lim_{t \rightarrow \infty} f_P^t(\hat{\gamma}) = \gamma^*$  for each  $\hat{\gamma} \in \Gamma$ ).*

## 4.2 Full Characterization of Environments that Admit Multiple Steady States

Our main result fully characterizes which environments allow for learning rules for which the past casts a long shadow. Specifically, it shows that an environment allows a learning rule that admits multiple steady states iff  $\mu_l > 1$  (alternatively if all agents sample exactly one action). In the opposite case ( $\mu_l \leq 1$ ) each learning rule admits a unique steady state, and, moreover, the population converges to the unique steady state from any initial state. Formally:

**Theorem 2.** *Let  $E = (A, \beta, \nu)$  be an environment. The following two conditions are equivalent:*

1.  $\mu_l > 1$ , or  $\nu(1) = 1$ .
2. *There exists a learning rule  $\sigma^*$ , such that the learning process  $(E, \sigma^*)$  admits two different steady states.*

*Proof.* Corollary 1 immediately implies that  $\neg 1 \Rightarrow \neg 2$ . We are left with the task of showing that  $1 \Rightarrow 2$ .

Case A: Assume that  $\nu(1) = 1$  (i.e., each new agent in the population observes a single action). Consider the learning rule in which each agent plays the action that he observed, i.e.,  $\sigma^*(a) = a$ . Let  $\gamma$  be an arbitrary population state. Observe that  $\gamma$  is a steady state of the learning process  $(E, \sigma^*)$  because:

$$(f_P(\gamma))(a) = \gamma(a).$$

Case B: Assume that  $\mu_l > 1$ . Let  $a$  and  $a'$  be different actions ( $a \neq a' \in A$ ). Let  $\sigma^*$  be a learning rule according to which each agent plays action  $a^*$  if he has observed action  $a^*$  at least once, and plays action  $a'$  otherwise, that is,

$$\sigma^*(a^l) = \begin{cases} a^* & \exists i, \text{ s.t., } a_i^l = a^* \\ a' & \text{otherwise.} \end{cases}$$

It is immediate that the population state in which all agents play action  $a'$  (i.e.,  $\gamma(a') = 1$ ) is a steady state of the learning process  $(E, \sigma^*)$ . We now show that there exists  $x > 0$ , such that the population state  $\gamma^x$  in which all agents play action  $a^*$  with probability  $x$ , and play action  $a'$  with the remaining probability of  $1 - x$  (i.e.,  $\gamma^x(a^*) = x$  and  $\gamma^x(a') = 1 - x$ ) is another steady state of the learning process  $(E, \sigma^*)$ . Observe that the state  $\gamma^x$  is consistent with the learning process  $(E, \sigma^*)$  if and only if

$$(f_P(\gamma^x))(a^*) = \sum_{l \in \text{supp}(\nu)} \nu(l) \cdot \frac{1}{|A|^l} \cdot \sum_{\vec{a} \in A^l} \mathbf{1}_{(\exists i \text{ s.t., } a_i = a^*)} = \sum_{l \in \text{supp}(\nu)} \nu(l) \cdot \left(1 - (1 - x)^l\right) \equiv g(x). \quad (3)$$

Observe that: (1)  $g(x)$  (defined in (3) above) is continuous and differentiable, (2) the derivative of  $g(x)$  is given by  $g'(x) = \sum_{l \in \text{supp}(\nu)} \nu(l) \cdot l \cdot (1 - x)^{l-1}$ , (3)  $g'(0) = \sum_{l \in \text{supp}(\nu)} \nu(l) \cdot l = \mu_l > 1$ , (4)  $g(0) = 0$ , and

<sup>5</sup>See Pata (2014, Theorem 1.7) for a formal proof that any weak contraction mapping on a compact metric space admits a global attractor (see also the sketch of the proof in Munkres, 2000, Section 28, Exercise 7). We thank Xiangqian Yang for kindly referring us to these proofs.

(5)  $g(1) \leq 1$ . These observations imply by the intermediate value theorem that there is  $x^* > 0$  such that  $g(x^*) = x^*$ , and hence  $\gamma^{x^*}$  is an additional steady state of the learning process  $(E, \sigma^*)$ .  $\square$

*Remark 1.* We note that the learning rules constructed in the proof above can be consistent with Bayesian inference and best-replying in plausible setups. The learning rule in Case A (playing the observed action) induces a Nash equilibrium in a setup with competing technologies with increasing returns and uncertainty about the initial population state, such as the setup presented in Example 1.

The learning rule in Case B induces a Nash equilibrium in the following setup of two competing technologies with uncertainty about their quality. There are two states of the world. In State 1 technology  $a^*$  has a higher quality, and in state 2 technology  $a'$  has a higher quality. The technology with the higher quality yields a payoff of one to an agent who follows it, and the technology with the lower quality yields a payoff of zero. State 1 has a prior probability of 60%. In state 1, 10% of the agents follow technology  $a^*$  in the first period, and the remaining agents follow technology  $a$ . In state 2, all agents follow technology  $a'$  in period one (i.e., the setup has a payoff-determined initial popularity à la Banerjee & Fudenberg, 2004). Observe that the unique Nash equilibrium in this setup is for an agent to play  $a^*$  when observing  $a^*$  at least once (as in this case the agent knows for sure that action  $a^*$  has a higher quality), and to play  $a'$  otherwise (as in this case the posterior probability that action  $a'$  has a higher quality is at least 60%).

Similarly, one can design plausible setups, in which the learning rules presented in all other constructions in the paper are consistent with Bayesian inference and best-replying (omitted for brevity).

### 4.3 Any Environment with $\mu_l > 2$ Admits Multiple Locally Stable States

Theorem 2 shows that any environment with a sample size larger than one admits multiple steady states, but it does not address the question of whether these steady states are locally stable. In particular, the learning rule presented in the proof of Theorem 2 (Case B) admits two steady states  $\gamma_0$  and  $\gamma_{x^*}$ . It is relatively simple to see that the state  $\gamma_{x^*}$  is an almost global attractor: the population converges to  $\gamma_{x^*}$  from any initial state  $\hat{\gamma}$  that assigns a positive probability to action  $a^*$  (see the related continuous-time analysis in Oyama *et al.* (2015, Sections 3.2 and 3.3)). Hence,  $\gamma_0$  is not locally stable. In the next two sections we characterize necessary and sufficient conditions for environments to admit multiple locally stable states.

The following result shows that in any environment with a mean sample size larger than 2 it is possible to define a learning rule that admits multiple locally stable states. According to this learning rule, each new agent (1) plays action  $a'$  if he observes action  $a'$  at least twice in his sample, (2) plays action  $a''$  if he never observes action  $a'$ , and (3) plays action  $a'$  with probability  $q$  and action  $a''$  with probability  $1 - q$ , if he observes action  $a'$  exactly once.

**Theorem 3.** *Let  $E = (A, \beta, \nu)$  be an environment satisfying  $\mu_l > 2$ . There exists a learning rule  $\sigma^*$ , such that the learning process  $(E, \sigma^*)$  admits two different locally stable states.*

The sketch of the proof is as follows (the formal proof is presented in Appendix B.2). If the incumbents play action  $a'$  with a frequency of  $x \ll 1$ , then the share of new agents who play action  $a'$  is  $q \cdot \mu_l \cdot x + (1 - 2 \cdot q) \cdot O(x^2)$  (the first term reflects the fact that the probability that a new agent plays action  $a'$  is approximately  $q$  times the expected number of times in which action  $a'$  is observed, namely,  $\mu_l \cdot x$ ; the second term “corrects” the fact that when a new agent observes action  $a'$  twice he plays action  $a'$  with probability one rather than with probability  $2 \cdot q$ ). Choosing  $q < \frac{1}{\mu_l}$  implies that a population in which very few agents play  $a'$  converges to

no one playing  $a'$ . Choosing  $q$  sufficiently close to  $\frac{1}{\mu_l} < \frac{1}{2}$  implies that a population in which a few more agents play action  $a'$  converges to a larger share of agents playing action  $a'$  (due to the second-order term,  $(1 - 2 \cdot q) \cdot O(x^2)$ , being positive).

#### 4.4 Some Environments with $1 < \mu_l < 2$ Admit Multiple Locally Stable States


In this section we show that some (but not all) environments in which the mean sample size is between one and two allow for a learning rule that admits multiple locally stable states.

Theorem 4 presents a family of environments with a mean sample size of up to two, in which every learning rule admits at most one locally stable state. Specifically, we show that in any environment in which (1) there are two feasible actions ( $|A| = 2$ ), and (2) each new agent observes at most 2 actions, any learning rule admits at most one locally stable state.

**Theorem 4.** *Let  $E = (A = \{a', a''\}, \beta, \nu)$  be an environment. Assume that  $\nu(l) = 0$  for each  $l > 2$ . Then for any learning rule  $\sigma$ , the learning process  $(E, \sigma)$  admits at most one locally stable state.*

The sketch of the proof of Theorem 4 is as follows (the formal proof is presented in Appendix B.3). In environments with two actions, the state can be identified with a number  $x \in [0, 1]$  representing the frequency of agents playing the first action. Recall that any steady state is a solution to the equation  $f_\sigma(x) = x$ , where  $f_\sigma(x)$  is the dynamic mapping induced by learning rule  $\sigma$ . The fact that the maximal sample size is two implies that  $f_\sigma(x)$  is a polynomial of degree two. This implies that there are at most two steady states solving  $f_\sigma(x) = x$ . Simple geometric arguments regarding the intersection points of a parabola and the  $45^\circ$  line imply that at most one of these steady states can be locally stable (as illustrated in Figure 1).

Figure 1: Illustrations for the Intersections of a Parabola and the  $45^\circ$  Line



Theorem 4 presents a family of environments (which extends Case (2) in Example 1) in which the mean sample size is between one and two, such that a simple “follow the majority” rule admits multiple locally stable states. Specifically, in these environments (1) there are two feasible actions, (2) some agents observe a single action and the remaining agents observe three actions, and (3) each agent plays the frequently observed action in his sample. It is relatively straightforward to see that this process admits two locally stable steady states: one in which all agents play the first action, and one in which all agents play the other action. In addition, the

state in which half of the agents play each action is an unstable steady state. The formal proof of Theorem 4 is given in Appendix B.4.

**Theorem 5.** *Let  $E = (A = \{a', a''\}, \beta, \nu)$  be an environment. Assume that  $\nu(1) < 1$  and  $\nu(1) + \nu(2) = 1$ . Then there exists a learning rule  $\sigma^*$ , such that  $(E, \sigma^*)$  admits multiple locally stable states.*

## 4.5 Summary of Main Results

Combining the various results of this section shows that the environment's mean sample size has important implications for determining whether the initial behavior of the population has an influence on the long-run behavior of the population.

**Corollary 2.** *Let  $E$  be an environment with an expected sample size  $\mu_l$ .*

1. *If  $\mu_l < 1$  (or  $\mu_l = 1$  and  $\nu(1) \neq 1$ ), then any learning rule admits a unique steady state that is globally stable.*
2. *If  $1 < \mu_l \leq 2$ , then there exists a learning rule that admits multiple steady states. By contrast, the multiplicity of locally stable states depends on other details of the environment. That is, for each  $1 < \mu_l \leq 2$  there exist environments  $E', E''$  with mean sample size  $\mu_l$ , such that Environment  $E'$  generates a learning rule that admits multiple locally stable states, while environment  $E''$  does not.*
3. *If  $\mu_l > 2$ , then there exists a learning rule that admits multiple locally stable states.*

## 5 Responsiveness and Effective Sample Size

In this section, we present simple notions of responsiveness and expected effective sample size, and use them to derive a (weakly) tighter upper bound for processes that admit global attractors (relative to the upper bound presented in Theorem 1).

### 5.1 Definitions

Fix a learning process  $P = (A, \beta, \nu, \sigma)$ . For each sample size  $l \in \text{supp}(\nu)$ , and each action  $a \in A$ , let  $\underline{\sigma}_l(a)$  ( $\bar{\sigma}_l(a)$ ) be the minimal (maximal) probability that learning rule  $\sigma$  assigns to action  $a$  after observing a sample of size  $l$ , i.e.,

$$\underline{\sigma}_l(a) = \min_{m \in A^l} \sigma_m(a) \quad (\bar{\sigma}_l(a) = \max_{m \in A^l} \sigma_m(a)).$$

Let  $r_l$  denote the maximal *responsiveness* of new agents to changes in observed samples of size  $l$ , which is defined as follows:

$$r_l = \min \left( 1, \frac{1}{2} \cdot \sum_{a \in A} (\bar{\sigma}_l(a) - \underline{\sigma}_l(a)) \right), \quad (4)$$

and let  $r_0 = 0$ . The responsiveness effectively limits the maximal influence of different samples of length  $l$  on the behavior of agents to be at most  $r_l \leq 1$ . One reason a learning rule may have limited responsiveness is that new agents might have strong priors about the best action, which are influenced only to a limited extent

by their observed samples. Observe that when there are two actions (i.e.,  $A = \{a, b\}$ ), then  $l$  is simply the difference between the maximal and minimal probability assigned to each action, i.e.,

$$r_l = \bar{\sigma}_l(a) - \underline{\sigma}_l(a) = \bar{\sigma}_l(b) - \underline{\sigma}_l(b) \quad (A = \{a, b\}). \quad (5)$$

When there are more than two actions,  $\frac{1}{2} \cdot \sum_{a \in A} (\bar{\sigma}_l(a) - \underline{\sigma}_l(a))$  may be larger than one. We bound  $r_l$  from above by one in Eq.(4) because, any change of sample cannot affect an agent's mixed behavior by more than one (as measured by the  $L_1$ -distance over the set of mixed actions).

We call the product of the sample size and the responsiveness,  $r_l \cdot l$  the *effective sample size*. Let  $\mu_l^e \in \mathbb{R}^+$  denote the *effective sample size*, i.e.,

$$\mu_l^e = \sum_{l \in \text{supp}(\nu)} \nu(l) \cdot r_l \cdot l.$$

It is immediate that the effective sample size is always weakly smaller than the mean sample size in the population; i.e.,  $\mu_l^e \leq \mu_l$ .

## 5.2 A Tighter Bound on the Distance between New States

Our main result in this section shows that the distance between two new states is at most  $(1 - \beta + \beta \cdot \mu_l^e)$  times the distance between the two initial states. This bound is (weakly) tighter than the one presented in Theorem 1, as we replace expected sample size  $\mu_l$  with the (weakly) smaller effective sample size  $\mu_l^e$ . Formally,

**Theorem 6.** *Let  $P = (A, \beta, \nu, \sigma)$  be a stationary learning process, and let  $\gamma \neq \gamma' \in \Gamma$  be two population states. Then:*

$$\|f_P(\gamma) - f_P(\gamma')\|_1 \leq (1 - \beta + \beta \cdot \mu_l^e) \cdot \|\gamma - \gamma'\|_1,$$

where the inequality is strict if there exist a type  $\theta$  and an  $l > 1$  such that  $\nu_\theta(l) > 0$ .

*Proof.* The key step of the proof is to show the following inequality:

$$\|(f_P(\gamma)) - (f_P(\gamma'))\|_1 \leq \beta \cdot \sum_{l \in \mathbb{N}} \nu(l) \cdot r_l \cdot \|\psi_{l,\gamma} - \psi_{l,\gamma'}\|_1 + (1 - \beta) \cdot \|\gamma - \gamma'\|_1. \quad (6)$$

Inequality (6) is the same as (2) in the proof of Theorem 1, except for the factor of  $r_l \leq 1$  on the RHS. All other arguments of the proof of Theorem 1 remain the same. We prove (6) in Lemma 6 in Appendix B.5.

Observe that  $(1 - \beta + \beta \cdot \mu_l^e) < 1$  iff  $\mu_l^e < 1$ , and in this case  $f_P$  is a contraction mapping, which implies that  $f_P$  admits a global attractor. This allows us to strengthen Corollary 1 as follows.  $\square$

**Corollary 3.** *Let  $P = (A, \beta, \nu, \sigma)$  be a learning process satisfying (1)  $\mu_l^e < 1$ , or (2)  $\mu_l^e = 1$  and there is a type  $\theta \in \Theta$  such that  $\nu_\theta(1) < 1$ . Then  $f_P$  is a contraction mapping, which implies that (1)  $f_P$  admits a unique steady state  $\gamma^*$ , and (2) this unique steady state  $\gamma^*$  is a global attractor (i.e.,  $\lim_{t \rightarrow \infty} f_P^t(\hat{\gamma}) = \gamma^*$  for each  $\hat{\gamma} \in \Gamma$ ).*

We demonstrate the implications of Corollary 3 in the following example.

**Example 4. 1** Consider a population in which in each period a share of  $\beta \in (0, 1)$  of the incumbent agents die, and are replaced with new agents. A population state describes the share of agents who use each of

two competing technologies,  $a_1$  and  $a_2$ . Each new agent observes the technology followed by a single random incumbent. Assume that the learning rule used by the agents implies that each new agent plays (on average) action  $a_1$  with a probability of  $\bar{\alpha} \in [0, 1]$  after observing action  $a'$ , and with a probability of  $\underline{\alpha} < \bar{\alpha}$  after observing action  $a''$ . Observe that the effective number of observations,  $\mu_l^e$ , is equal to:

$$\mu_l^e = \mu_{\theta}^e = r_{\theta, l=1} \cdot 1 = \frac{1}{2} \cdot \sum_{a \in A} (\bar{\sigma}_{\theta, l}(a) - \underline{\sigma}_{\theta, l}(a)) = \frac{1}{2} \cdot ((\bar{\alpha} - \underline{\alpha}) + ((1 - \underline{\alpha}) - (1 - \bar{\alpha}))) = \bar{\alpha} - \underline{\alpha},$$

which is strictly less than one if  $\bar{\alpha} < 1$  or  $\underline{\alpha} > 0$ . Corollary 3 implies that the learning process converges to a global attractor (which is the unique steady state) whenever  $\bar{\alpha} < 1$  or  $\underline{\alpha} > 0$ .<sup>6</sup>

Our final result demonstrates that our bound of the effective sample size being less than one is tight. Specifically, it shows that given any environment in which the expected sample size  $\mu_l > 1$ , and any number  $1 < y \leq \mu_l$ , there is a learning rule with an effective sample size of  $\mu_l^e = y$  with multiple steady states. Formally:

**Theorem 7.** *Let  $E = (A, \beta, \nu)$  be an environment satisfying  $\mu_l > 1$ . Let  $1 < y \leq \mu_l$ . Then there exists a learning rule  $\sigma$ , such that the learning process  $(E, \sigma)$  admits two different steady states, and satisfies  $\mu_l^e = y$ .*

*Proof.* Let  $a$  and  $a'$  be different actions ( $a \neq a' \in A$ ). Let  $\sigma^*$  be a learning rule according to which each agent plays action  $a^*$  with a probability of  $\frac{y}{\mu_l}$  if he has observed action  $a^*$  at least once, and plays action  $a'$  otherwise, that is,

$$\sigma^*(a^l) = \begin{cases} \frac{y}{\mu_l} \cdot a^* + \left(1 - \frac{y}{\mu_l}\right) \cdot a' & \text{if } \exists i, \text{ s.t., } a_i^l = a^* \\ a' & \text{otherwise.} \end{cases}$$

Observe that the responsiveness of  $(E, \sigma)$  is equal to  $x$  because:

$$\begin{aligned} \mu_l^e &= \sum_{l \in \text{supp}(\nu)} \nu(l) \cdot r_l \cdot l = \sum_{l \in \text{supp}(\nu)} \nu(l) \cdot \frac{1}{2} \cdot \sum_{a \in A} (\bar{\sigma}_l(a) - \underline{\sigma}_l(a)) \cdot l = \\ & \sum_{l \in \text{supp}(\nu)} \nu(l) \cdot \frac{1}{2} \cdot \left( \left( \frac{y}{\mu_l} - 0 \right) + \left( 1 - \left( 1 - \frac{y}{\mu_l} \right) + 0 + \dots + 0 \right) \right) \cdot l = \\ & \sum_{l \in \text{supp}(\nu)} \nu(l) \cdot \frac{y}{\mu_l} \cdot l = \frac{y}{\mu_l} \cdot \sum_{l \in \text{supp}(\nu)} \nu(l) \cdot l = \frac{y}{\mu_l} \cdot \mu_l = y. \end{aligned}$$

It is immediate that the uniform population state in which all agents play action  $a'$  (i.e.,  $\gamma(a') = 1$ ) is a steady state of the learning process  $(E, \sigma^*)$ . An analogous argument to the one presented in Case B of the proof of Theorem 2 shows that there exists  $x > 0$  such that the uniform population state  $\gamma^x$  in which all agents play action  $a^*$  with probability  $x$ , and play action  $a'$  with the remaining probability of  $1 - x$ , is another steady state of the learning process  $(E, \sigma^*)$ .  $\square$

<sup>6</sup>One can show that in this global attractor a share  $\frac{\alpha}{1+\alpha-\alpha}$  of the agents play action  $a_1$ . If  $\bar{\alpha} = 1$  and  $\underline{\alpha} > 0$ , then any population state is steady.

## 6 Concluding Remarks

**Extensions** The basic model assumes that all agents share the same distribution of sample sizes, and the same learning rule. In many applications the population might be heterogeneous, i.e., the population includes various groups that differ in their sampling procedures and learning rules (see, e.g., [Ellison & Fudenberg, 1993](#); [Munshi, 2004](#); [Young, 1993b](#)). In [Appendix A.1](#) we formally extend our model and results to heterogeneous populations.

The basic model assumes that the learning rule is stationary. In [Appendix A.2](#) we extend our model and results to time-dependent learning rules, and we characterize when a non-stationary environment admits a unique sequence of states, such that it converges to this sequence of states from any initial population state.

Finally, we further extend the model in [Appendix A.3](#) to stochastic shocks that influence the learning rules of all agents (on the aggregate level), and we characterize when the initial population state may have a lasting effect in such environments.

**Repeated Interactions without Calendar Time.** In many real-life situations agents are randomly matched within a community, and these interactions have been going on since time immemorial. Modeling such situations as repeated games with a definite starting point and strategies that can be conditioned on calendar time may be a problematic, as it seems implausible that agents would be aware of the the exact time that has transpired since the starting point, and be aware of the very distant history of play of other agents. An alternative approach is to model behavior in such situations as steady states of environments without a calendar time (see, e.g., ([Rosenthal, 1979](#); [Okuno-Fujiwara & Postlewaite, 1995](#); [Heller & Mohlin, 2017](#)), and the working paper version of [Phelan & Skrzypacz, 2006](#)).

An interesting question about such environments is whether the distribution of strategies used by the players to choose their actions as a function of their observations is sufficient to uniquely determine the steady states, or whether the same distribution of rules may admit multiple steady states. Our main result shows that the former is true whenever the expected number of observed actions is less than one, while if the expected number of observed actions is more than one, then there is always a distribution of rules with multiple steady states.

**Large Finite Populations.** Our model studies infinite populations, and it is important to know what the implications of our results are for large finite populations. The key difference between an infinite and a finite population, is that in the former, the law of large numbers implies that the new state of the population is a deterministic function of the initial state and the learning rule. By contrast, in finite populations the new population state is a random variable. If the finite population is sufficiently large then we expect the resulting stochastic process to be close to the deterministic process over finite time horizons. However, when time goes to infinity, rare random events will occasionally take the population away from one (locally stable) steady state towards another steady state (see [Sandholm, 2011](#) for a textbook overview of the deterministic approximation of stochastic evolutionary processes).

When dealing with large finite populations, one may therefore interpret our main result ([Theorem 2](#)) as follows. In environments in which  $\mu_l < 1$ , all learning processes admit a unique globally stable state  $\gamma^*$ . The population is highly likely to quickly converge to state  $\gamma^*$ , and will almost always remain very close to this state. In the rare event that the realized observations of many agents substantially differ from their expected

values, the population may temporarily move away from  $\gamma^*$ , but with a very high probability the population will quickly converge back to  $\gamma^*$ .

In environments in which  $\mu_l > 1$ , there are learning rules that admit multiple steady states. The fact that the population is finite and that the new population state is a random variable will typically quickly take the population away from steady states that are not locally stable. If the environment admits multiple locally stable states, then the initial state is highly likely to determine which of these locally stable states the population converges to in the medium run. Moreover the population will likely stay there for a significant amount of time.<sup>7</sup>

**Observations of Action Profiles.** In [Heller & Mohlin \(2017\)](#) we investigate environments in which an agent may observe action profiles played in past interactions by the current opponent against her past opponents. All of our results can be extended to this setup, with relatively minor adjustments to the proofs. Specifically one should count an observation of an action profile (in a two-player game) as two actions when calculating the expected number of observed actions  $\mu_l$ . Our main result still holds in this setup: an environment allows a profile of learning rules that admits multiple steady states, essentially, if and only if  $\mu_l \leq 1$ .

## A Extensions

### A.1 Heterogeneous Population

The basic model assumes that all agents share the same distribution of sample sizes, and the same learning rule. In many applications the population might be heterogeneous, i.e., the population includes various groups that differ in their sampling procedures and learning rules. A few examples of such models with heterogeneous populations can be found in: (1) [Ellison & Fudenberg \(1993\)](#), who study competing technologies where each technology is better for some of the players and these different tastes induce different learning rules (see also [Munshi, 2004](#)); (2) [Young \(1993b\)](#), who studies social learning in a bargaining model in which agents differ in the size of their samples; and (3) [Heller & Mohlin \(2017\)](#), who in a companion paper analyze community enforcement in which the population includes several types of agents, and each type uses a different strategy.

#### A.1.1 Model with Heterogeneous Population

In what follows we introduce heterogeneous populations that include different types, and we redefine the notions of population state, environment, and learning process to deal with this heterogeneity.

**Population state.** Let  $\Theta$  denote a finite set of types with a typical element  $\theta$ . Let  $\lambda_\theta$  denote the mass of agents of type  $\theta$  (or  $\theta$ -agents). For simplicity, we assume that  $\lambda$  has full support. We redefine a *population state* (or *state* for short) to be a vector  $\gamma = (\gamma_\theta)_{\theta \in \Theta}$ , where each  $\gamma_\theta \in \Delta(A)$  denotes the aggregate distribution of actions played by  $\theta$ -agents. Let  $\bar{\gamma} \in \Delta(A)$  denote the average distribution of actions in the population (i.e.,

---

<sup>7</sup>The literature on stochastic evolutionary game theory (starting with the pioneering works of [Foster & Young, 1990](#); [Kandori et al., 1993](#); [Young, 1993a](#); see [Young, 2015](#), for a recent survey) studies situations the long-run behavior in environments with multiple locally stable states, and in which there is a small level of noise in the agents' behavior. We think that it would be interesting to extend the methodology of this literature in order to apply it to the setup analyzed in this paper. It might be that such future research can characterize various cases in which, if the population size is sufficiently large, in the long run the population will spend almost all of the time in one of these locally stable states.

$\bar{\gamma}(a) = \sum_{\theta} \lambda_{\theta} \gamma_{\theta}(a)$  for each action  $a \in A$ ). A population state is *uniform* if all types play the same aggregate distribution of actions, i.e., if  $\gamma_{\theta}(a) = \bar{\gamma}(a)$  for each type  $\theta \in \Theta$  and action  $a \in A$ . We redefine  $\Gamma$  to denote the set of all populations with heterogeneous types.

**New/Revising agents.** In each period, a share of  $0 < \beta \leq 1$  of the agents of each type die and are replaced with new agents (or, alternatively, are randomly selected to reevaluate their choice), while the remaining  $1 - \beta$  share of the agents of each type play the same action as they played in the past.

**Sample.** Each new agent observes a finite sequence of actions (or *sample*). The size of the sample observed by type  $\theta$  is a random variable with a distribution  $\nu_{\theta} \in \Delta(\mathbb{N})$ . Let  $M$ , the set of all feasible samples, be redefined as:  $M = \cup_{\theta \in \Theta} \cup_{l \in \text{supp}(\nu_{\theta})} A^l$ . Let  $\bar{l} = \max_{l \in (\cup_{\theta \in \Theta} \text{supp}(\nu_{\theta}))} l < \infty$  be the maximal sample size. For each sample size  $l \in \mathbb{N}$ , let  $\psi_l : \Gamma \rightarrow \Delta(A^l)$  denote the distribution of samples observed by each agent in the population (or *sampling rule* for short), conditional on the sample having size  $l$ . A typical sample of size  $l$  is represented by the vector  $\vec{a} = (a_1, \dots, a_l)$ .

We analyze two kinds of sampling methods in heterogeneous populations:

1. *Observing different random agents:* Each agent independently samples different agents, and observes a random action played by each of these agents. This kind of sampling is a common modeling choice in situations in which an agent's payoff depends not on the behavior of a specific sub-group of opponents, but on the agent's own action, the state of nature, and, possibly, the aggregate behavior of the population (see, e.g., [Ellison & Fudenberg, 1995](#); [Banerjee & Fudenberg, 2004](#)). Formally, we define for each sample size  $l \in \mathbb{N}$ , each state  $\gamma \in \Gamma$ , and each sample  $(a_1, \dots, a_l)$ ,

$$\psi_{l,\gamma}(a_1, \dots, a_l) = \prod_{1 \leq i \leq l} \bar{\gamma}(a_i). \quad (7)$$

2. *Observing a single random type:* Each agent randomly draws a type  $\bar{\theta}$ , and then the agent samples different agents of type  $\bar{\theta}$ , and observes a random action played by each of these  $\bar{\theta}$ -agents. This kind of observation is relevant to models in which the agent is randomly matched with an opponent, and may sample some actions played in the previous period by agents with the same type as the opponent. Formally, we define for each size  $l \in \mathbb{N}$ , each state  $\gamma \in \Gamma$ , and each sample  $(a_1, \dots, a_l)$ ,

$$\psi_{l,\gamma}(a_1, \dots, a_l) = \sum_{\theta \in \Theta} \lambda_{\theta} \cdot \prod_{1 \leq i \leq l} \gamma_{\theta}(a_i). \quad (8)$$

In the case of  $\beta = 1$ , this sampling method has another interpretation that is common in models of strategic interactions among randomly matched agents (e.g., [Rosenthal, 1979](#); [Nowak & Sigmund, 1998](#); [Heller & Mohlin, 2017](#)). According to this interpretation, each agent is involved in  $n \geq \bar{l}$  interactions in each period. In each of these interactions the agent is randomly matched with a different opponent, and the agent observes a sample of random actions played by the opponent in the previous round. The random type of the opponent is distributed according to  $\lambda_{\theta}$ , and each of the actions played by the opponent of type  $\theta$  in the previous round is distributed according to  $\gamma_{\theta}$ .

Observe that both cases, i.e., (7) and (8), coincide in two special setups: (1) when the population state is uniform (as in the basic model), or (2) when agents observe at most one action (i.e.,  $\bar{l} = 1$ ).

*Remark 2.* Our results work also in a setup in which some types use the first sampling method, while other types use the second sampling method.

**Environment.** We redefine an *environment* as a tuple

$$E = (A, \Theta, \beta, \psi_l, (\lambda_\theta, \nu_\theta)_{\theta \in \Theta})$$

that includes the six components described above: a finite set of actions  $A$ , a finite set of types  $\Theta$ , a fraction of new agents at each stage  $\beta$ , a sampling rule  $\psi_l$  (satisfying either (7) or (8)), a distribution over the set of types  $\lambda$ , and a profile of distributions of sample sizes  $(\nu_\theta)_{\theta \in \Theta}$ .

Given environment  $E = (A, \Theta, \beta, \psi_l, (\lambda_\theta, \nu_\theta)_{\theta \in \Theta})$ , let  $\mu_l$ , the *mean sample size*, be redefined as the expected number of actions observed by a random agent in the population. Formally:

$$\mu_l = \sum_{\theta \in \Theta} \lambda_\theta \sum_{l \in \text{supp}(\nu_\theta)} \nu_\theta(l) \cdot l.$$

**Learning rule and stationary learning process.** Each new  $\theta$ -agent chooses his action in the new population state by following a stationary (i.e., time-independent) *learning rule*  $\sigma_\theta : M \rightarrow \Delta(A)$ . That is, a new  $\theta$ -agent who observes sample  $m \in M$  plays action  $a$  with probability  $\sigma_{\theta, m}(a)$ . The remaining  $1 - \beta$  incumbent agents play the same action as in the previous round. A profile of learning rules  $(\sigma_\theta)_{\theta \in \Theta}$  is *uniform* if all types use the same learning rule, i.e., if  $\sigma_\theta = \sigma_{\theta'}$  for each type  $\theta, \theta' \in \Theta$ .

A stationary learning process (or *learning process* for short) is a pair

$$P = (E, (\sigma_\theta)_{\theta \in \Theta}) = (A, \Theta, \beta, \psi_l, (\lambda_\theta, \nu_\theta, \sigma_\theta)_{\theta \in \Theta}),$$

consisting of an environment and a learning rule.

As in the basic model, let  $f_P : \Gamma \rightarrow \Gamma$  denote the mapping between states induced by a single step of the learning process  $P$ .

**$L_1$ -distance.** Each population state  $\gamma \in \Gamma$  corresponds to a distribution  $q_\gamma \in \Delta(\Theta \times A)$  as follows:  $q_\gamma(\theta, a) = \lambda_\theta \cdot \gamma_\theta(a)$ . We define the distance between two population states  $\gamma, \gamma' \in \Gamma$  as the  $L_1$ -distance between the corresponding distributions  $q_\gamma, q_{\gamma'} \in \Delta(\Theta \times A)$ :

$$\|\gamma - \gamma'\|_1 = \|q_\gamma - q_{\gamma'}\|_1 = \sum_{\theta \in \Theta} \sum_{a \in A} |\lambda_\theta \cdot \gamma_\theta(a) - \lambda_\theta \cdot \gamma'_\theta(a)| = \sum_{\theta \in \Theta} \lambda_\theta \cdot \|\gamma_\theta - \gamma'_\theta\|_1.$$

### A.1.2 Generalizing Results

In what follows we formally show how to generalize the first result (Theorem 1) to heterogeneous populations.

**Theorem 8.** (*Generalization of Theorem 1*) Let  $P = (A, \Theta, \beta, \psi_l, (\lambda_\theta, \nu_\theta, \sigma_\theta)_{\theta \in \Theta})$  be a stationary learning process, and let  $\gamma \neq \gamma' \in \Gamma$  be two population states. Then:

$$\|f_P(\gamma) - f_P(\gamma')\|_1 \leq (1 - \beta + \beta \cdot \mu_l) \cdot \|\gamma - \gamma'\|_1,$$

with a strict inequality if there exist a type  $\theta$  and an  $l > 1$  such that  $\nu_\theta(l) > 0$ .

The intuition is similar to Theorem 1. The proof is presented in Appendix B.

Similarly to the generalization of Theorem 1 above, one can generalize in a straightforward way all the other results of the paper to the setup of a heterogeneous population (proofs omitted for brevity).

## A.2 Non-Stationary Learning Process

In this section we further extend the model to deal with non-stationary deterministic learning processes, in which the process explicitly depends on calendar time, and we show how to generalize our results to this setup.

**Adaptations to the model.** For each period  $t \geq 1$ , let  $\beta_t \in [0, 1]$  denote the random share of agents who revise their actions in period  $t$ . For each type  $\theta \in \Theta$  and period  $t \geq 1$ , let  $\nu_\theta^t \in \Delta(\mathbb{N})$  denote the distribution of sample sizes of type  $\theta$  in period  $t$ . To simplify the notation we assume that the support of the sample sizes of each type is independent of the period, i.e.,  $\text{supp}(\nu_\theta^{t_1}) = \text{supp}(\nu_\theta^{t_2}) := \text{supp}(\nu_\theta)$  for each type  $\theta \in \Theta$  and periods  $t_1, t_2 \geq 1$ . As in the basic model, let  $M$  denote the set of all feasible sample sizes. A *non-stationary environment* is a tuple

$$E = \left( A, \Theta, (\beta_t)_{t \in \mathbb{N}}, \psi_l, (\lambda_\theta)_{\theta \in \Theta}, (\nu_\theta^t)_{\theta \in \Theta, t \geq 1} \right).$$

Given a non-stationary environment, let  $\mu_l^t$  denote the expected number of actions observed in period  $t$ , i.e.,  $\mu_l^t = \sum_{\theta \in \Theta} \lambda_\theta \sum_{l \in \text{supp}(\nu_\theta)} \nu_\theta^t(l) \cdot l$ .

Given a non-stationary environment  $E$ , let  $\bar{\mu}_l$  be the upper limit of the geometric mean of  $1 - \beta^t \cdot (1 - \mu_l^t)$  as  $t$  goes to infinity, i.e.,

$$\bar{\mu}_l = \limsup_{t \rightarrow \infty} \sqrt[t]{\prod_{t \leq t_0} (1 - \beta^t \cdot (1 - \mu_l^t))}.$$

For each type  $\theta \in \Theta$  and period  $t \geq 1$ , let  $\sigma_\theta^t : M \rightarrow \Delta(A)$  denote the non-stationary learning rule of new  $\theta$ -agents in period  $t$ . A *non-stationary learning process* is a pair consisting of a non-stationary environment and a non-stationary learning rule, i.e.,

$$P = \left( E, (\sigma_\theta^t)_{\theta \in \Theta, t \geq 1} \right) = \left( A, \Theta, (\beta_t)_{t \geq 1}, \psi_l, (\lambda_\theta)_{\theta \in \Theta}, (\nu_\theta^t, \sigma_\theta^t)_{\theta \in \Theta, t \geq 1} \right).$$

As in the basic model, a non-stationary learning process  $P$  and an initial state uniquely determine a new state in each period  $t$ . Let  $f_P^t(\hat{\gamma}) \in \Gamma$  denote the state induced after  $t$  stages of the non-stationary learning process  $P$ .

A sequence of states  $(\gamma_t^*)_{t \in \mathbb{N}}$  is a *global attractor* of the non-stationary learning process  $P$ , if

$$\lim_{t \rightarrow \infty} \|f_P^t(\hat{\gamma}) - \gamma_t^*\|_1 = 0$$

for each initial state  $\hat{\gamma} \in \Gamma$ .

**Adapted results.** Minor adaptations to the proof of Theorem 8 and a simple inductive argument immediately imply that the distance between two states at time  $t_0$  is at most  $\prod_{t \leq t_0} (1 - \beta^t + \beta^t \cdot \mu_l^t)$  the initial distance. Formally:

**Corollary 4.** Let  $P = \left( A, \Theta, (\beta_t)_{t \geq 1}, \psi_l, (\lambda_\theta)_{\theta \in \Theta}, (\nu_\theta^t, \sigma_\theta^t)_{\theta \in \Theta, t \geq 1} \right)$  be a non-stationary learning process, let  $\hat{\gamma}, \hat{\gamma}' \in \Gamma$  be two population states, and let  $\hat{t} \geq 1$ . Then:

$$\left\| f_P^{\hat{t}}(\hat{\gamma}) - f_P^{\hat{t}}(\hat{\gamma}') \right\|_1 \leq \|\hat{\gamma} - \hat{\gamma}'\|_1 \cdot \prod_{t \leq \hat{t}} (1 - \beta^t + \beta^t \cdot \mu_l^t).$$

This, in turn, immediately implies that in any non-stationary environment in which  $\bar{\mu}_l < 1$ , any profile of non-stationary learning rules admits a global attractor. Formally:

**Corollary 5.** Let  $E = \left( A, \Theta, (\beta_t)_{t \geq 1}, \psi_l, (\lambda_\theta)_{\theta \in \Theta}, (\nu_\theta^t)_{\theta \in \Theta, t \geq 1} \right)$  be a non-stationary environment satisfying  $\bar{\mu}_l < 1$ . Then for any profile of non-stationary learning rules  $(\sigma_\theta^t)_{\theta \in \Theta, t \geq 1}$ , the non-stationary learning process  $P = \left( E, (\sigma_\theta^t)_{\theta \in \Theta, t \geq 1} \right)$  admits a global attractor.

The example presented in Case A of the proof of Theorem 2 demonstrates that the above bound of  $\bar{\mu}_l < 1$  is binding in the sense that there is an environment with  $\bar{\mu}_l = 1$  that admits a profile of learning rules with multiple steady states.

The adaptation of the remaining results to the time-dependent setup is similar (proof omitted for brevity).

### A.3 Process with Common Shocks

In this section we further extend our model to deal also with common stochastic shocks to the learning rules.

**Additional adaptations to the model.** In what follows we further adapt the model of Section A.2 by allowing common stochastic shocks to the learning rules of the agents.

Let  $(\Omega, \mathcal{F}, p)$  be an arbitrary probability space. Each element  $\omega \in \Omega$  represents the state of nature, which determines the realizations of all common shocks to the learning rules in all periods. For each type  $\theta \in \Theta$  and period  $t \in \mathbb{N}$ , let  $\sigma_\theta^t : \Omega \times M \rightarrow \Delta(A)$  denote the state-dependent learning rule of new  $\theta$ -agents in period  $t$ .

Our interpretation of the state-dependent learning rule  $\sigma_\theta^t$  is as follows. The state of nature determines a distribution of noisy signals from which each new agent draws a signal. Based on this noisy signal as well as on the sample of past actions (and on information about calendar time), each new agent chooses an action. The choices of actions (which depend on the noisy signals) and the distribution of noisy signals jointly generate a distribution of actions that depend only on the state of nature (and on calendar time), which is captured by the state-dependent learning rule  $\sigma_\theta^t$ .

A *learning process* with common shocks is a pair consisting of a non-stationary environment and a state-dependent learning rule, i.e.,  $P = \left( E, (\sigma_\theta^t)_{\theta \in \Theta, t \geq 1} \right) = \left( A, \Theta, (\beta_t)_{t \geq 1}, \psi_l, (\lambda_\theta)_{\theta \in \Theta}, (\nu_\theta^t, \sigma_\theta^t)_{\theta \in \Theta, t \geq 1} \right)$ .

Learning processes with common shocks are important in modeling situations in which there are stochastic factors that influence the learning rules of all new agents in period  $t$ . For example, [Ellison & Fudenberg \(1995\)](#) model a situation in which new agents in period  $t$  choose between two agricultural technologies, and each such new agent observes a noisy signal about the expected payoff of each technology conditional on the weather in period  $t$  (which is common to all agents), where the (unknown) state of nature determines the weather in all periods.

The state of nature, the learning process, and the initial population state uniquely determine the population state in each period. Let  $f_P^t(\omega)(\hat{\gamma}) \in \Gamma$  denote the population state induced after  $t$  stages of the non-stationary learning process  $P$ , given an initial population state  $\hat{\gamma}$ , and state of nature  $\omega \in \Omega$ .

We say that a sequence of state-dependent population states  $(\gamma_t^*)_{t \geq 1}$ , where  $\gamma_t^* : \Omega \rightarrow \Gamma$ , is a *state-dependent global attractor* of the learning process with common shocks  $P$  if, for each  $\omega \in \Omega$ ,  $\lim_{t \rightarrow \infty} \|f_P^t(\omega)(\hat{\gamma}) - \gamma_t^*(\omega)\|_1 = 0$  for each initial state  $\hat{\gamma} \in \Gamma$ .

Example 5 below demonstrates how to apply the extended model to a social learning process with competing technologies with common shocks:

**Example 5** (Competing Technologies with Common Shocks). Consider a stochastic environment in which there are two possible regimes  $\{1, 2\}$ . There are two technologies:  $a_1$  and  $a_2$ . Technology  $a_1$  is advantageous in regime 1, while technology  $a_2$  is advantageous in regime 2. There is a uniform common prior about the regime in round 1. In each subsequent round, the regime is the same as in the previous round with probability 99%, and it is a new regime with probability 1%. In each round, a share of 25% of the incumbents die, and are replaced with new agents. Each new agent observes the action of a single random incumbent and a noisy signal about the current regime, and based on these observations, the agent chooses one of the two technologies. Assume that the learning rule used by the agents implies that each new agent plays action  $a_1$ :

1. with a probability of 95% after observing action  $a_1$  in regime 1;
2. with a probability of 80% after observing action  $a_1$  in regime 2;
3. with a probability of 20% after observing action  $a_2$  in regime 1;
4. with a probability of 5% after observing action  $a_2$  in regime 2.

One can show that the environment admits a unique steady state that is a state-dependent global attractor. The induced aggregate behavior of the population converges towards playing action  $a_1$  with an average probability of 80% in regime 1, and it converges towards playing action  $a_1$  with an average probability of 20% in regime 2.

This learning process with common shocks is modeled as

$$P = \left( \{a_1, a_2\}, \{\theta\}, (\beta_t \equiv 25\%)_{t \in \mathbb{N}}, \psi_l, \lambda_\theta, (\nu_\theta^t \equiv 1, \sigma_\theta^t)_{t \geq 1} \right).$$

The set of states of nature  $\Omega = \{(\omega_n)_{n \in \mathbb{N}}\}$  is the set of infinite binary sequences, where each  $\omega_n \in \{1, 2\}$  describes the regime in round  $n$ . The definition of  $(\mathcal{F}, p)$  is derived from the Markovian process determining the regime in each round in a standard way. Given state  $\omega = (\omega_n)_{n \in \mathbb{N}}$ , let the learning rule be defined as follows:

$$\sigma_\theta(a_1, \omega) = \begin{cases} 95\% & a = a_1 \text{ and } \omega_t = 1 \\ 80\% & a = a_1 \text{ and } \omega_t = 2 \\ 20\% & a = a_2 \text{ and } \omega_t = 1 \\ 5\% & a = a_2 \text{ and } \omega_t = 2. \end{cases}$$

**Adapted Results.** Minor adaptations to the proof of Theorem 8 imply that the distance between two states at time  $\hat{t}$  is at most  $\prod_{t \leq \hat{t}} (1 - \beta^t + \beta^t \cdot \mu_t^t)$  the initial distance. Formally:

**Corollary 6.** Let  $P = \left( A, \Theta, (\beta_t)_{t \geq 1}, \psi_l, (\lambda_\theta)_{\theta \in \Theta}, (\nu_\theta^t, \sigma_\theta^t)_{\theta \in \Theta, t \geq 1} \right)$  be a learning process with common shocks, let  $\hat{\gamma}, \hat{\gamma}' \in \Gamma$  be two population states, and let  $\hat{t} \in \mathbb{N}$ . Then, for each  $\omega \in \Omega$ ,

$$\left\| f_P^{\hat{t}}(\omega)(\hat{\gamma}) - f_P^{\hat{t}}(\omega)(\hat{\gamma}') \right\|_1 \leq \|\hat{\gamma} - \hat{\gamma}'\|_1 \cdot \prod_{t \leq \hat{t}} (1 - \beta^t + \beta^t \cdot \mu_l^t).$$

An immediate corollary is that any environment with common shocks in which  $\bar{\mu}_l < 1$ , given any profile of learning rules, admits a state-dependent global attractor. That is, in the long run, the population's behavior depends only on the state of nature, but it is independent of the initial population state in time zero. Formally:

**Corollary 7.** Let  $E = \left( A, \Theta, (\beta_t)_{t \geq 1}, \psi_l, (\lambda_\theta)_{\theta \in \Theta}, (\nu_\theta^t)_{\theta \in \Theta, t \geq 1} \right)$  be an environment satisfying  $\bar{\mu}_l < 1$ . Then for any profile of stochastic learning rules  $(\sigma_\theta^t)_{\theta \in \Theta, t \geq 1}$ , the learning process with common shocks  $P = \left( E, (\sigma_\theta^t)_{\theta \in \Theta, t \geq 1} \right)$  admits a state-dependent global attractor.

The adaptation of the remaining results to the time-dependent setup is similar to the adaptation above (proof omitted for brevity).

## B Formal Proofs

### B.1 Proof of Theorem 8 (Upper Bound Result; Generalization of Theorem 1)

The distance between the final population states is bounded as follows (where the second inequality is strict if  $\nu_\theta(l) > 0$  for some  $\theta \in \Theta$  and  $l \geq 2$ ):

$$\begin{aligned} \|(f_P(\gamma))_\theta - (f_P(\gamma'))_\theta\|_1 &\leq \beta \cdot \sum_{\theta \in \Theta} \lambda_\theta \cdot \sum_{l \in \mathbb{N}} \nu_\theta(l) \cdot \|\psi_{l,\gamma} - \psi_{l,\gamma'}\|_1 + (1 - \beta) \cdot \|\gamma - \gamma'\|_1 \leq \\ &\beta \cdot \sum_{\theta \in \Theta} \lambda_\theta \cdot \sum_{l \in \mathbb{N}} \nu_\theta(l) \cdot l \cdot \|\gamma - \gamma'\|_1 + (1 - \beta) \cdot \|\gamma - \gamma'\|_1 = \\ &\left( \beta \cdot \left( \sum_{\theta \in \Theta} \lambda_\theta \cdot \sum_{l \in \mathbb{N}} \nu_\theta(l) \cdot l \right) + (1 - \beta) \right) \cdot \|\gamma - \gamma'\|_1 = (\beta \cdot \mu_L + 1 - \beta) \cdot \|\gamma - \gamma'\|_1 = (1 - \beta \cdot (1 - \mu_l)) \cdot \|\gamma - \gamma'\|_1. \end{aligned}$$

The first inequality is proven in Lemma 1. The second inequality (is strict if  $\nu_\theta(l) > 0$  for some  $\theta \in \Theta$  and  $l \geq 2$ ) is implied by the inequality

$$\|\psi_{l,\gamma} - \psi_{l,\gamma'}\|_1 \leq l \cdot \|\gamma - \gamma'\|_1 \quad (\text{with a strict inequality if } l \geq 2),$$

which is proven in Lemma 4.

### Proofs of the various Lemmas used in the Proof of Theorem 8

**Lemma 1.** For each learning environment  $E$  and states  $\gamma \neq \gamma' \in \Gamma$ ,

$$\|(f_P(\gamma)) - (f_P(\gamma'))\|_1 \leq \beta \cdot \sum_{\theta \in \Theta} \lambda_\theta \cdot \sum_{l \in \mathbb{N}} \nu_\theta(l) \cdot \|\psi_{l,\gamma} - \psi_{l,\gamma'}\|_1 + (1 - \beta) \cdot \|\gamma - \gamma'\|_1.$$

*Proof.*

$$\begin{aligned}
\|(f_P(\gamma)) - (f_P(\gamma'))\|_1 &= \sum_{\theta \in \Theta} \lambda_\theta \cdot \|(f_P(\gamma))_\theta - (f_P(\gamma'))_\theta\|_1 \leq \\
&\sum_{\theta \in \Theta} \lambda_\theta \cdot \left( \beta \cdot \sum_{l \in \mathbb{N}} \nu_\theta(l) \cdot \|\psi_{l,\gamma} - \psi_{l,\gamma'}\|_1 + (1 - \beta) \cdot \|\gamma_\theta - \gamma'_\theta\|_1 \right) = \\
\beta \cdot \sum_{\theta \in \Theta} \lambda_\theta \cdot \sum_{l \in \mathbb{N}} \nu_\theta(l) \cdot \|\psi_{l,\gamma} - \psi_{l,\gamma'}\|_1 + (1 - \beta) \cdot \sum_{\theta \in \Theta} \lambda_\theta \cdot \|\gamma_\theta - \gamma'_\theta\|_1 &= \\
\beta \cdot \sum_{\theta \in \Theta} \lambda_\theta \cdot \sum_{l \in \mathbb{N}} \nu_\theta(l) \cdot \|\psi_{l,\gamma} - \psi_{l,\gamma'}\|_1 + (1 - \beta) \cdot \|\gamma - \gamma'\|_1, &
\end{aligned}$$

where the inequality is due to Lemma 2.  $\square$

**Lemma 2.** For each social learning environment  $E$ , type  $\theta \in \Theta$ , and each two states  $\gamma \neq \gamma' \in \Gamma$ :

$$\|(f_P(\gamma))_\theta - (f_P(\gamma'))_\theta\|_1 \leq \beta \cdot \sum_{l \in \mathbb{N}} \nu_\theta(l) \cdot \|\psi_{l,\gamma} - \psi_{l,\gamma'}\|_1 + (1 - \beta) \cdot \|\gamma_\theta - \gamma'_\theta\|_1.$$

*Proof.*

$$\|(f_P(\gamma))_\theta - (f_P(\gamma'))_\theta\|_1 = \sum_{a \in A} |(f_P(\gamma))_\theta(a) - (f_P(\gamma'))_\theta(a)| =$$

$$\begin{aligned}
&\sum_{a \in A} \left| \left( \sum_{l \in \text{supp}(\nu_\theta)} \beta \cdot \nu_\theta(l) \sum_{m \in A^l} \psi_{l,\gamma}(m) \cdot \sigma_{\theta,m} + (1 - \beta) \cdot \gamma_\theta \right) (a) \right. \\
&\left. - \left( \beta \cdot \sum_{l \in \text{supp}(\nu_\theta)} \nu_\theta(l) \cdot \sum_{m \in A^l} \psi_{l,\gamma'}(m) \cdot \sigma_{\theta,m} + (1 - \beta) \cdot \gamma'_\theta \right) (a) \right| =
\end{aligned}$$

$$\sum_{a \in A} \left| \beta \cdot \sum_{l \in \text{supp}(\nu_\theta)} \nu_\theta(l) \cdot \sum_{m \in A^l} (\psi_{l,\gamma}(m) - \psi_{l,\gamma'}(m)) \cdot \sigma_{\theta,m}(a) + (1 - \beta) \cdot (\gamma_\theta(a) - \gamma'_\theta(a)) \right| \leq \quad (9)$$

$$\sum_{a \in A} \left( \beta \cdot \sum_{l \in \text{supp}(\nu_\theta)} \nu_\theta(l) \cdot \left| \sum_{m \in A^l} (\psi_{l,\gamma}(m) - \psi_{l,\gamma'}(m)) \cdot \sigma_{\theta,m}(a) \right| + (1 - \beta) \cdot |\gamma_\theta(a) - \gamma'_\theta(a)| \right) =$$

$$\beta \cdot \sum_{l \in \text{supp}(\nu_\theta)} \nu_\theta(l) \cdot \sum_{a \in A} \left| \sum_{m \in A^l} (\psi_{l,\gamma}(m) - \psi_{l,\gamma'}(m)) \cdot \sigma_{\theta,m}(a) \right| + (1 - \beta) \cdot \sum_{a \in A} |\gamma_\theta(a) - \gamma'_\theta(a)| \leq \quad (10)$$

$$\beta \cdot \sum_{l \in \text{supp}(\nu_\theta)} \nu_\theta(l) \cdot \|\psi_{l,\gamma} - \psi_{l,\gamma'}\|_1 + (1 - \beta) \cdot \|\gamma_\theta - \gamma'_\theta\|_1,$$

where the (9) is a triangle inequality, and (10) is due to Lemma 3.  $\square$

**Lemma 3.** For each social learning environment  $E$ , each size  $l \in \mathbb{N}$ , each type  $\theta \in \Theta$ , and any two states  $\gamma \neq \gamma' \in \Gamma$ :

$$\sum_{a \in A} \left| \sum_{m \in A^l} (\psi_{l,\gamma}(m) - \psi_{l,\gamma'}(m)) \cdot \sigma_{\theta,m}(a) \right| \leq \|\psi_{l,\gamma} - \psi_{l,\gamma'}\|_1.$$

*Proof.*

$$\begin{aligned}
\sum_{a \in A} \left| \sum_{m \in A^l} (\psi_{l,\gamma}(m) - \psi_{l,\gamma'}(m)) \cdot \sigma_{\theta,m}(a) \right| &\leq \sum_{a \in A} \sum_{m \in A^l} |\psi_{l,\gamma}(m) - \psi_{l,\gamma'}(m)| \cdot \sigma_{\theta,m}(a) \\
&= \sum_{m \in A^l} |\psi_{l,\gamma}(m) - \psi_{l,\gamma'}(m)| \cdot \sum_{a \in A} \sigma_{\theta,m}(a) \\
&= \sum_{m \in A^l} |\psi_{l,\gamma}(m) - \psi_{l,\gamma'}(m)| \cdot 1,
\end{aligned}$$

where the inequality is a triangle inequality.  $\square$

**Lemma 4.** *For each social learning environment  $E$ , type  $\theta \in \Theta$ , sample size  $l \in \mathbb{N}$ , and states  $\gamma \neq \gamma' \in \Gamma$*

$$\|\psi_{l,\gamma} - \psi_{l,\gamma'}\|_1 \leq l \cdot \|\gamma - \gamma'\|_1,$$

with a strict inequality if  $l > 1$ .

*Proof.* Case I - Observing different random agents:

$$\|\psi_{l,\gamma} - \psi_{l,\gamma'}\|_1 = \sum_{\vec{a} \in A^l} |\psi_{l,\gamma}(\vec{a}) - \psi_{l,\gamma'}(\vec{a})| = \quad (11)$$

$$\sum_{\vec{a} \in A^l} \left| \prod_{1 \leq i \leq l} \bar{\gamma}(a_i) - \prod_{1 \leq i \leq l} \bar{\gamma}'(a_i) \right| = \quad (12)$$

$$\sum_{\vec{a} \in A^l} \left| \sum_{1 \leq i \leq l} (\bar{\gamma}(a_i) - \bar{\gamma}'(a_i)) \cdot \prod_{i < j \leq l} \bar{\gamma}(a_j) \cdot \prod_{1 \leq k < i} \bar{\gamma}'(a_k) \right| \leq \quad (< \text{if } l > 1) \quad (13)$$

$$\sum_{\vec{a} \in A^l} \left( \sum_{1 \leq i \leq l} |\bar{\gamma}(a_i) - \bar{\gamma}'(a_i)| \cdot \prod_{i < j \leq l} \bar{\gamma}(a_j) \cdot \prod_{1 \leq k < i} \bar{\gamma}'(a_k) \right) =$$

$$\sum_{1 \leq i \leq l} \left( \sum_{\vec{a} \in A^l} |\bar{\gamma}(a_i) - \bar{\gamma}'(a_i)| \cdot \prod_{i < j \leq l} \bar{\gamma}(a_j) \cdot \prod_{1 \leq k < i} \bar{\gamma}'(a_k) \right) =$$

$$\sum_{1 \leq i \leq l} \left( \sum_{a_i \in A} |\bar{\gamma}(a_i) - \bar{\gamma}'(a_i)| \right) \cdot \left( \sum_{(a_{i+1}, \dots, a_l) \in A^{l-i}} \prod_{i < j \leq l} \bar{\gamma}(a_j) \right) \cdot \left( \sum_{(a_1, \dots, a_{i-1}) \in A^{i-1}} \prod_{1 \leq k < i} \bar{\gamma}'(a_k) \right) = \quad (14)$$

$$\sum_{1 \leq i \leq l} \left( \sum_{a_i \in A} |\bar{\gamma}(a_i) - \bar{\gamma}'(a_i)| \right) \cdot 1 \cdot 1 = \sum_{1 \leq i \leq l} (\|\bar{\gamma} - \bar{\gamma}'\|_1) = l \cdot \|\bar{\gamma} - \bar{\gamma}'\|_1 \leq l \cdot \|\gamma - \gamma'\|.$$

Eq. (11) is due to the independence of different observations. Eq. (12) is implied by adding to the sum elements that cancel out. Specifically, let  $b_i = \bar{\gamma}(a_i)$  and  $c_i = \bar{\gamma}'(a_i)$ ; then due to a “telescoping series”

argument (in which each new element appears once with a positive sign and once with a negative sign):<sup>8</sup>

$$\begin{aligned}
& \prod_{1 \leq i \leq l} \bar{\gamma}(a_i) - \prod_{1 \leq i \leq l} \bar{\gamma}'(a_i) = \prod_{1 \leq i \leq l} b_i - \prod_{1 \leq i \leq l} c_i = \\
& (b_1 \cdot \dots \cdot b_l - c_1 \cdot b_2 \cdot \dots \cdot b_l) + (c_1 \cdot b_2 \cdot \dots \cdot b_l + c_1 \cdot c_2 \cdot b_3 \cdot \dots \cdot b_l) - c_1 \cdot c_2 \cdot b_3 \cdot \dots \cdot b_l + \dots + c_1 \cdot \dots \cdot c_l = \\
& (b_1 - c_1) \cdot b_2 \cdot \dots \cdot b_l + (b_2 - c_2) \cdot b_3 \cdot \dots \cdot b_l \cdot c_1 + (b_3 - c_3) \cdot b_4 \cdot \dots \cdot b_l \cdot c_1 \cdot c_2 \dots + (b_l - c_l) \cdot c_2 \cdot \dots \cdot c_l = \\
& = \sum_{1 \leq i \leq l} \left( (b_i - c_i) \cdot \prod_{i < j \leq l} b_j \cdot \prod_{1 \leq j < i} c_j \right) = \sum_{1 \leq i \leq l} (\bar{\gamma}(a_i) - \bar{\gamma}'(a_i)) \cdot \prod_{i < j \leq l} \bar{\gamma}(a_j) \cdot \prod_{1 \leq k < i} \bar{\gamma}'(a_k).
\end{aligned}$$

Eq. (13) is a triangle inequality, and it is strict if  $l > 1$  because the sum inside the “ $\left| \right|$ ” in (13) includes both positive and negative elements. Eq. (14) holds because each sum adds the probabilities of disjoint and exhausting events. The final inequality is implied by Lemma 5.

Case II - Observing a single random type:

$$\|\psi_{l,\gamma} - \psi_{l,\gamma'}\|_1 = \sum_{\vec{a} \in A^l} |\psi_{l,\gamma}(\vec{a}) - \psi_{l,\gamma'}(\vec{a})| = \quad (15)$$

$$\sum_{\vec{a} \in A^l} \left| \sum_{\theta \in \Theta} \lambda_\theta \cdot \left( \prod_{1 \leq i \leq l} \gamma_\theta(a_i) - \prod_{1 \leq i \leq l} \gamma'_\theta(a_i) \right) \right| = \quad (16)$$

$$\sum_{\vec{a} \in A^l} \left| \sum_{\theta \in \Theta} \lambda_\theta \cdot \left( \sum_{1 \leq i \leq l} (\gamma_\theta(a_i) - \gamma'_\theta(a_i)) \cdot \prod_{i < j \leq l} \gamma_\theta(a_j) \cdot \prod_{1 \leq j < i} \gamma'_\theta(a_j) \right) \right| \leq \quad (< \text{if } l > 1) \quad (17)$$

$$\sum_{\vec{a} \in A^l} \sum_{\theta \in \Theta} \lambda_\theta \cdot \left( \sum_{1 \leq i \leq l} |\gamma_\theta(a_i) - \gamma'_\theta(a_i)| \cdot \prod_{i < j \leq l} \gamma_\theta(a_j) \cdot \prod_{1 \leq j < i} \gamma'_\theta(a_j) \right) =$$

$$\sum_{1 \leq i \leq l} \sum_{\theta \in \Theta} \lambda_\theta \cdot \left( \sum_{\vec{a} \in A^l} |\gamma_\theta(a_i) - \gamma'_\theta(a_i)| \cdot \prod_{i < j \leq l} \gamma_\theta(a_j) \cdot \prod_{1 \leq j < i} \gamma'_\theta(a_j) \right) =$$

$$\sum_{1 \leq i \leq l} \sum_{\theta \in \Theta} \lambda_\theta \cdot \left( \sum_{a_i \in A} |\gamma_\theta(a_i) - \bar{\gamma}'_\theta(a_i)| \right) \cdot \left( \sum_{(a_{i+1}, \dots, a_l) \in A^{l-i}} \prod_{i < j \leq l} \gamma_\theta(a_j) \right) \cdot \left( \sum_{(a_1, \dots, a_{i-1}) \in A^{i-1}} \prod_{1 \leq j < i} \gamma'_\theta(a_j) \right) = \quad (18)$$

$$\sum_{1 \leq i \leq l} \sum_{\theta \in \Theta} \lambda_\theta \cdot \left( \sum_{a_i \in A} |\gamma_\theta(a_i) - \bar{\gamma}'_\theta(a_i)| \right) \cdot 1 \cdot 1 = \sum_{1 \leq i \leq l} \sum_{\theta \in \Theta} \lambda_\theta \cdot \|\gamma_\theta - \gamma'_\theta\|_1 = \sum_{1 \leq i \leq l} \|\gamma - \gamma'\|_1 = l \cdot \|\gamma - \gamma'\|_1.$$

Eq. (15) is due to the different observations being independent conditional on the observed type  $\theta$ . Eq. (16) is implied by adding to the sum elements that cancel out (i.e., a “telescoping series”). Eq. (17) is a triangle inequality, and it is strict if  $l > 1$  because the sum inside the “ $\left| \right|$ ” in (17) includes both positive and negative elements. Eq. (18) holds because each sum adds the probabilities of disjoint and exhausting events.  $\square$

**Lemma 5.**  $\|\bar{\gamma} - \bar{\gamma}'\|_1 \leq \|\gamma - \gamma'\|_1$  for each two states  $\gamma \neq \gamma' \in \Gamma$ .

<sup>8</sup>We use the convention that a product of an empty set (e.g.,  $\prod_{1 \leq j < 1}$ ) is equal to one.

*Proof.*

$$\begin{aligned} \|\gamma - \gamma'\|_1 &= \sum_{\theta \in \Theta} \lambda_\theta \cdot \|\gamma_\theta - \gamma'_\theta\|_1 = \sum_{\theta \in \Theta} \lambda_\theta \cdot \sum_{a \in A} |\gamma_\theta(a) - \gamma'_\theta(a)| = \sum_{a \in A} \sum_{\theta \in \Theta} \lambda_\theta \cdot |\gamma_\theta(a) - \gamma'_\theta(a)| \geq \\ &\sum_{a \in A} \left| \sum_{\theta \in \Theta} \lambda_\theta (\gamma_\theta(a) - \gamma'_\theta(a)) \right| = \sum_{a \in A} \left| \sum_{\theta \in \Theta} \lambda_\theta \gamma_\theta(a) - \sum_{\theta \in \Theta} \lambda_\theta \gamma'_\theta(a) \right| = \sum_{a \in A} |\bar{\gamma}(a) - \bar{\gamma}'(a)| = \|\bar{\gamma} - \bar{\gamma}'\|_1, \end{aligned}$$

where the various equalities are immediately implied by the definitions on the  $L_1$ -norm and  $\bar{\gamma}$ , and the inequality is a triangle inequality.  $\square$

## B.2 Proof of Theorem 3 ( $\mu_l > 2$ )

For each  $0 < q < \frac{1}{\mu_l}$  define  $\sigma_q$  as the learning rule according to which each agent plays action  $a^*$  if he has observed action  $a^*$  at least twice, plays action  $a'$  if he has not observed action  $a^*$ , and he plays action  $a^*$  with probability  $q$  and action  $a'$  with the remaining probability  $1 - q$ ; that is, for each  $a^l \in A^l$ ,

$$\sigma^*(a^l) = \begin{cases} a^* & |\{i | a_i^l = a^*\}| \geq 2 \\ q \cdot a^* + (1 - q) \cdot a' & |\{i | a_i^l = a^*\}| = 1 \\ a' & |\{i | a_i^l = a^*\}| = 0. \end{cases}$$

Observe that new agents play only  $a^*$  or  $a'$ . Further note that the probability that a new agent plays  $a^*$  depends only on the frequency with which the incumbents play action  $a^*$  (and not on any other aspect of the population state). Thus, by a slight abuse of notation, we identify a state  $\gamma \in \Delta(A)$  with the frequency  $x$  of agents who choose action  $a^*$ , i.e.,  $x := \gamma(a)$  (as the actions played by the remaining  $1 - x$  of the agents do not play any role in the dynamics, and in the long run each agent plays either action  $a^*$  or  $a'$ ). The mapping induced by the environment  $P_q = (E, \sigma_q)$  is given by the function  $f_q : [0, 1] \rightarrow [0, 1]$  (neglecting terms that are  $O(x^3)$ ):

$$f_q(x) := f_{(P_q)}(x) = (1 - \beta) \cdot x + \beta \cdot \left( q \cdot \mu_l \cdot x + (1 - 2 \cdot q) \cdot \sum_{2 \leq l \in \text{supp}(\nu)} \nu(l) \cdot \binom{l}{2} \cdot x^2 + O(x^3) \right). \quad (19)$$

The argument for (19) is as follows. The term of  $(1 - \beta) \cdot x$  describes the behavior of incumbents who have not died. The terms multiplying  $\beta$  represent the behavior of new agents. The first of these terms ( $q \cdot \mu_l \cdot x$ ) derives from the fact that the first-order approximation for the probability that a new agent plays action  $a'$  is  $q$  times the expected number of times that action  $a'$  is observed ( $\mu_l \cdot x$ ), since action  $a'$  is almost always observed once in a sample. The second term multiplying  $\beta$  in (19) reflects the correction required to adjust the above first-order approximation due to the fact that an agent who observes action  $a'$  twice in the sample plays action  $a'$  with probability 1, rather than with probability  $2 \cdot q$ . Hence, the additional probability of playing  $a^*$  conditional on observing  $a^*$  twice is  $(1 - 2 \cdot q)$ . The probability of observing  $a^*$  twice in a random sample is  $\sum_{2 \leq l \in \text{supp}(\nu)} \nu(l) \cdot \binom{l}{2} \cdot x^2 + O(x^3)$ . Finally, note that the probability of observing  $a^*$  three or more times is negligible (i.e.,  $O(x^3)$ ), so that the remaining adjustment required for (19) to coincide with the dynamic mapping induced by  $P_q$  is  $O(x^3)$ .

It is immediate from the definition of learning rule  $\sigma^*$  that  $f_q(x)$  is strictly increasing in  $x$ . Recall that state  $x$  is a steady state iff  $f_q(x) = x$ . Observe that:  $f_q(0) = 0$  and  $f'_q(0) = q \cdot \mu_l < 1$  for each  $q < \frac{1}{\mu_l}$ . Fix  $q < \frac{1}{\mu_l}$ . The previous observations imply that there exists  $\bar{x} > 0$  such that  $f_q(x) < x$  and  $f'_q(x) < 1$  for each  $x \in (0, \bar{x})$ . This implies that  $f_q^t(x) < x$  and  $\lim_{t \rightarrow \infty} f_q^t(x) = 0$  for each  $x \in (0, \bar{x})$ , and hence state 0 is locally stable.

Assume that  $\nu(0) = \nu(1) = 0$ . Then, the definition of learning rule  $\sigma^*$  implies that  $f_q(1) = 1$ , and that for each  $\epsilon \ll 1$  it holds that

$$f_q(1 - \epsilon) = (1 - \beta) \cdot (1 - \epsilon) + \beta \cdot (1 - \nu(2) \cdot 2 \cdot \epsilon \cdot (1 - q) + O(\epsilon^2)).$$

This is so because when  $x = 1 - \epsilon$  and  $\epsilon \ll 1$ , a new agent plays action  $a'$  with probability  $1 - q$  when observing action  $a^*$  once in a sample of size two (the probability of this event is given by  $\nu(2) \cdot 2 \cdot \epsilon$ ). Moreover, observing  $a^*$  once in a longer sample (or not observing  $a^*$  at all) is a rare event with a probability of  $O(\epsilon^2)$ . This implies that for each  $q < 0.5$ , there is  $\bar{\epsilon} > 0$ , such that for each  $x > 1 - \bar{\epsilon}$ : (1)  $f_q^t(x) > x$  for each  $t$ , and (2)  $\lim_{t \rightarrow \infty} f_q^t(x) = 1$ . This shows that the state 1 is locally stable.

We are left with the case in which  $\nu(0) > 0$  or  $\nu(1) > 0$ . Observe that (1)  $\lim_{q \rightarrow \frac{1}{\mu_l}} f'_q(0) = 1$  and (2)  $f''_q(0) > 0$  for each  $q < \frac{1}{\mu_l}$ . This implies (by a Taylor approximation around  $x = 0$ ) that there exists  $(q^*, \hat{x})$  satisfying: (1)  $0 < \hat{x} \ll 1$ , (2)  $q^* < \frac{1}{\mu_l}$ , (3)  $f_{q^*}(\hat{x}) = \hat{x}$ , (4)  $f_{q^*}(x) < x$  for each  $x \in (0, \hat{x})$ , and (5)  $f'_{q^*}(\hat{x}) > 1$ . This implies that  $\hat{x}$  is an (unstable) steady state.

Next observe that  $f_{q^*}(\hat{x}) = \hat{x}$ ,  $f'_{q^*}(\hat{x}) > 1$ , and  $f_{q^*}(1) < 1$ . These observations, due to the intermediate value theorem and standard arguments, imply that there exists  $\hat{x} < x^* < 1$ , such that  $f_{q^*}(x^*) = x^*$  and  $f'_{q^*}(x^*) < 1$ . This, in turn, implies that there exists  $\bar{\epsilon} > 0$ , such that for each  $x \in (x^* - \bar{\epsilon}, x^* + \bar{\epsilon})$ : (1)  $f_{q^*}(x) < x$  if  $x < x^*$ , (2)  $f_{q^*}(x) > x$  if  $x > x^*$ , and (3)  $f'_{q^*}(x) < 1$ . These observations imply (due to the monotonicity of  $f_{q^*}$ ) that for each  $x^* \neq x \in (x^* - \bar{\epsilon}, x^* + \bar{\epsilon})$ : (1)  $f_q^t(x)$  is strictly between  $x$  and  $x^*$  for each  $t$ , and (2)  $\lim_{t \rightarrow \infty} f_q^t(x) = x^*$ . Hence, state  $x^*$  is locally stable.

### B.3 Proof of Theorem 4 (Two Feasible Actions, $\mu(l) = 0 \forall l > 2$ )

Let  $E = (A = \{a', a''\}, \beta, \nu)$  be an environment satisfying  $\nu(l) = 0$  for each  $l > 2$ . Let  $\sigma$  be an arbitrary learning rule. The fact that there are two feasible actions (i.e.,  $|A| = 2$ ) implies that we can identify a population state with a number  $x \in [0, 1]$  representing the frequency of agents who play action  $a'$ . Let  $f_\sigma(x)$  be the dynamic mapping induced by learning rule  $\sigma$ . The fact that the maximal length of the sample observed by new agents is two implies that  $f_\sigma(x)$  is a polynomial of degree at most two. Specifically, the explicit formula for  $f_\sigma(x)$  is given by:

$$\begin{aligned} f_\sigma(x) &= (1 - \beta) \cdot x + \beta \cdot [\nu(0) \sigma(\emptyset)(a') + \nu(1) \cdot (x \cdot \sigma(a')(a') + (1 - x) \cdot \sigma(a'')(a')) + \\ &\quad \nu(2) \cdot (x^2 \cdot \sigma(a', a')(a') + (1 - x)^2 \cdot \sigma(a'', a'')(a') + x \cdot (1 - x) \cdot (\sigma(a', a'')(a') + \sigma(a'', a')(a')))] = \\ &= b \cdot x^2 + c \cdot x + d, \end{aligned}$$

where  $b, c, d \in \mathbb{R}$ . Recall that  $x^*$  is a steady state iff  $f(x^*) = x^*$ . We conclude the proof by looking at three exhaustive cases. Cases 2 and 3 are illustrated in Figure 1 in Section 4.4.

1. Case 1:  $b = 0$  (recall, that  $b$  is the parameter multiplying  $x^2$  in the formula for  $f_\sigma(x)$ ). If  $f_\sigma(x) \equiv x$  (i.e.,

if  $f_\sigma(x) = x$  for each  $x$ ), then any state is steady, but none is locally stable. Otherwise, the equation  $f_\sigma(x) = x$  has at most one solution, and, hence,  $\sigma^*$  has at most one locally stable state.

2. Case 2:  $b > 0$ . The equation  $f_\sigma(x) = x$  has at most two solutions. Assume that it has two solutions in the interval  $[0, 1]$  (otherwise, it is immediate that  $\sigma$  admits at most one locally stable state). Simple geometric arguments (regarding the incidence points of a parabola satisfying  $f_\sigma(1) \leq 1$  and the  $45^\circ$  line) imply that one of these solutions must be one (i.e.,  $f_\sigma(1) = 1$ ), and that  $f'_\sigma(1) > 1$ . By standard continuity arguments there exists a sufficiently small  $\bar{\epsilon} > 0$  such that  $f'_\sigma(x) > 1$  for each  $x > 1 - \bar{\epsilon}$ . This implies that for each  $x > 1 - \bar{\epsilon}$ : (1)  $f_\sigma^t(x) < x$ , and (2) if  $\lim_{t \rightarrow \infty} f_\sigma^t(x)$  exists then it must satisfy  $\lim_{t \rightarrow \infty} f_\sigma^t(x) < 1 - \bar{\epsilon}$ . Hence, state 1 cannot be locally stable, and the learning rule  $\sigma$  admits at most one locally stable state.
3. Case 3:  $b < 0$ . Assume that the equation  $f_\sigma(x) = x$  has two solutions in the interval  $[0, 1]$  (otherwise, it is immediate that  $\sigma$  admits at most one locally stable state). Simple geometric arguments (regarding the points of intersection of a parabola bounded with positive values and the  $45^\circ$  line) imply that one of these solutions must be zero (i.e.,  $f_\sigma(0) = 0$ ), and that  $f'_\sigma(0) > 1$ . By standard continuity arguments there exists a sufficiently small  $\bar{\epsilon} > 0$  such that  $f'_\sigma(x) > 1$  for each  $x \in (0, \bar{\epsilon})$ . This implies that for each  $x \in (0, \bar{\epsilon})$ : (1)  $f_\sigma^t(x) > x$ , and (2) if  $\lim_{t \rightarrow \infty} f_\sigma^t(x)$  exists then it must satisfy  $\lim_{t \rightarrow \infty} f_\sigma^t(x) > \bar{\epsilon}$ . This implies that state  $x^*$  cannot be locally stable, and hence learning rule  $\sigma$  admits at most one locally stable state.

#### B.4 Proof of Theorem 5 ( $\nu(1) + \nu(3) = 1$ ), “Follow Majority” Rule)

Let  $E = (A = \{a', a''\}, \beta, \nu)$  be an environment, such that  $\nu(l) = 0$  for each  $l \notin \{1, 3\}$  and  $\nu(1) < 1$ . Let  $\sigma^*$  be the learning rule in which each new agent follows the frequently observed action in his sample, i.e.,

$$\sigma^*(a') = \sigma^*(a', a', a') = \sigma^*(a', a', a'') = \sigma^*(a', a'', a') = \sigma^*(a'', a', a') = a', \text{ and}$$

$$\sigma^*(a'') = \sigma^*(a'', a'', a'') = \sigma^*(a'', a'', a') = \sigma^*(a'', a', a'') = \sigma^*(a', a'', a'') = a''.$$

We identify a state with the number  $x \in [0, 1]$  representing the frequency of agents who play action  $a'$ . Let  $f_{\sigma^*}(x)$  be the dynamic mapping induced by learning rule  $\sigma^*$ . The explicit formula for  $f_{\sigma^*}(x)$  is given by

$$f_{\sigma^*}(x) = (1 - \beta) \cdot x + \beta \cdot (\nu(1) \cdot x + \nu(3) \cdot (x^3 + 3 \cdot x^2 \cdot (1 - x))) = \nu(1) \cdot x + \nu(3) \cdot (3 \cdot x^2 - 2 \cdot x^3),$$

and its derivative is given by

$$f'_{\sigma^*}(x) = (1 - \beta) + \beta \cdot \nu(1) + \beta \cdot \nu(3) \cdot (6 \cdot x - 6 \cdot x^2) = (1 - \beta) + \beta \cdot \nu(1) + \beta \cdot 6 \cdot \nu(3) \cdot x \cdot (1 - x).$$

Observe that: (1)  $f_{\sigma^*}(x)$  is strictly increasing, (2)  $f_{\sigma^*}(x^*) = x^*$  for three values of  $x$ : 0, 0.5, 1, (3)  $f'_{\sigma^*}(0) = f'_{\sigma^*}(1) = \nu(1) < 1$ , and (4)  $f'_{\sigma^*}(0.5) = \nu(1) + 1.5 \cdot \nu(3) > 1$ . These observations imply (by arguments analogous to those in the proof of B.3 above) that the process  $(E, \sigma^*)$  admits three steady states: two locally stable states, 0 and 1, and the locally unstable state 0.5.

## B.5 Lemma Required for the Proof of Theorem 6 (Bound with Responsiveness)

**Lemma 6.** *For each social learning environment  $E$ , each size  $l \in \mathbb{N}$ , and any two states  $\gamma \neq \gamma' \in \Gamma$ :*

$$\sum_{a \in A} \left| \sum_{m \in A^l} (\psi_{l,\gamma}(m) - \psi_{l,\gamma'}(m)) \cdot \sigma_m(a) \right| \leq r_l \cdot \|\psi_{l,\gamma} - \psi_{l,\gamma'}\|_1.$$

*Proof.* We begin with a preliminary definition. Let  $A_{\gamma > \gamma'}^l \subseteq A^l$  be the set of samples that have higher probabilities given state  $\gamma$  than given state  $\gamma'$ , i.e.,

$$A_{\gamma > \gamma'}^l = \{m \in A^l \mid \psi_{l,\gamma}(m) > \psi_{l,\gamma'}(m)\}.$$

We now prove the lemma:

$$\begin{aligned} & \sum_{a \in A} \left| \sum_{m \in A^l} (\psi_{l,\gamma}(m) - \psi_{l,\gamma'}(m)) \cdot \sigma_m(a) \right| = \\ & \sum_{a \in A} \left| \sum_{m \in A_{\gamma > \gamma'}^l} (\psi_{l,\gamma}(m) - \psi_{l,\gamma'}(m)) \cdot \sigma_m(a) - \sum_{m \in A_{\gamma' > \gamma}^l} (\psi_{l,\gamma'}(m) - \psi_{l,\gamma}(m)) \cdot \sigma_m(a) \right| \leq \\ & \sum_{a \in A} \left| \sum_{m \in A_{\gamma > \gamma'}^l} (\psi_{l,\gamma}(m) - \psi_{l,\gamma'}(m)) \cdot \bar{\sigma}_l(a) - \sum_{m \in A_{\gamma' > \gamma}^l} (\psi_{l,\gamma'}(m) - \psi_{l,\gamma}(m)) \cdot \underline{\sigma}_l(a) \right| = \\ & \sum_{a \in A} \left| \bar{\sigma}_l(a) \cdot \sum_{m \in A_{\gamma > \gamma'}^l} (\psi_{l,\gamma}(m) - \psi_{l,\gamma'}(m)) - \underline{\sigma}_l(a) \cdot \sum_{m \in A_{\gamma' > \gamma}^l} (\psi_{l,\gamma'}(m) - \psi_{l,\gamma}(m)) \right| = \quad (20) \\ & \sum_{a \in A} \left| (\bar{\sigma}_l(a) - \underline{\sigma}_l(a)) \cdot \sum_{m \in A_{\gamma > \gamma'}^l} (\psi_{l,\gamma}(m) - \psi_{l,\gamma'}(m)) \right| = \\ & \sum_{a \in A} (\bar{\sigma}_l(a) - \underline{\sigma}_l(a)) \cdot \sum_{m \in A_{\gamma > \gamma'}^l} (\psi_{l,\gamma}(m) - \psi_{l,\gamma'}(m)) = \\ & \sum_{a \in A} (\bar{\sigma}_l(a) - \underline{\sigma}_l(a)) \cdot 0.5 \cdot \left( \sum_{m \in A^l} |(\psi_{l,\gamma}(m) - \psi_{l,\gamma'}(m))| \right) = \\ & 0.5 \cdot \sum_{a \in A} (\bar{\sigma}_l(a) - \underline{\sigma}_l(a)) \cdot \|\psi_{l,\gamma} - \psi_{l,\gamma'}\|_1. \end{aligned}$$

Equality (20) is implied by the fact that  $\psi_{l,\gamma}$  and  $\psi_{l,\gamma'}$  are both distributions, and the sum of the differences in the probabilities that they assign to samples of size  $l$  must be equal to zero. Thus we have shown that

$$\sum_{a \in A} \left| \sum_{m \in A^l} (\psi_{l,\gamma}(m) - \psi_{l,\gamma'}(m)) \cdot \sigma_m(a) \right| \leq 0.5 \cdot \sum_{a \in A} (\bar{\sigma}_l(a) - \underline{\sigma}_l(a)) \cdot \|\psi_{l,\gamma} - \psi_{l,\gamma'}\|_1, \quad (21)$$

which together with Lemma 3 implies that the LHS of (21) is weakly smaller than  $r_l \cdot \|\psi_{l,\gamma} - \psi_{l,\gamma'}\|_1$ .  $\square$

## References

- Acemoglu, Daron, Dahleh, Munther A., Lobel, Ilan, & Ozdaglar, Asuman. 2011. Bayesian learning in social networks. *The Review of Economic Studies*, **78**(4), 1201–1236.
- Arthur, W. Brian. 1989. Competing technologies, increasing returns, and lock-in by historical events. *The economic journal*, **99**(394), 116–131.
- Arthur, W. Brian. 1994. *Increasing returns and path dependence in the economy*. University of michigan Press.
- Banerjee, Abhijit, & Fudenberg, Drew. 2004. Word-of-mouth learning. *Games and Economic Behavior*, **46**(1), 1–22.
- Cason, Timothy N., Friedman, Daniel, & Hopkins, Ed. 2014. Cycles and instability in a rock–paper–scissors population game: A continuous time experiment. *The Review of Economic Studies*, **81**(1), 112–136.
- Ellison, Glenn, & Fudenberg, Drew. 1993. Rules of thumb for social learning. *Journal of Political Economy*, **101**, 612–643.
- Ellison, Glenn, & Fudenberg, Drew. 1995. Word-of-mouth communication and social learning. *The Quarterly Journal of Economics*, **110**(1), 93–125.
- Foster, Dean, & Young, Peyton. 1990. Stochastic evolutionary game dynamics. *Theoretical Population Biology*, **38**(2), 219–232.
- Heller, Yuval, & Mohlin, Erik. 2017. *Observations on Cooperation*. Mimeo.
- Kandori, Michihiro, Mailath, George J., & Rob, Rafael. 1993. Learning, mutation, and long run equilibria in games. *Econometrica*, **61**(1), 29–56.
- Kaniovski, Yuri M., & Young, H. Peyton. 1995. Learning dynamics in games with stochastic perturbations. *Games and Economic Behavior*, **11**(2), 330–363.
- Munkres, James R. 2000. *Topology*. 2nd edn. Prentice Hall.
- Munshi, Kaivan. 2004. Social learning in a heterogeneous population: technology diffusion in the Indian Green Revolution. *Journal of development Economics*, **73**(1), 185–213.
- Nowak, Martin A., & Sigmund, Karl. 1998. Evolution of indirect reciprocity by image scoring. *Nature*, **393**(6685), 573–577.
- Okuno-Fujiwara, Masahiro, & Postlewaite, Andrew. 1995. Social norms and random matching games. *Games and Economic Behavior*, **9**(1), 79–109.
- Oyama, Daisuke, Sandholm, William H., & Tercieux, Olivier. 2015. Sampling best response dynamics and deterministic equilibrium selection. *Theoretical Economics*, **10**(1), 243–281.
- Pata, Vittorino. 2014. *Fixed point theorems and applications*. Mimeo.
- Phelan, Christopher, & Skrzypacz, Andrzej. 2006. *Private monitoring with infinite histories*. Tech. rept. Federal Reserve Bank of Minneapolis.

- Rosenthal, Robert W. 1979. Sequences of games with varying opponents. *Econometrica*, **47**(6), 1353–1366.
- Sandholm, William H. 2001. Almost global convergence to p-dominant equilibrium. *International Journal of Game Theory*, **30**(1), 107–116.
- Sandholm, William H. 2011. *Population Games and Evolutionary Dynamics*. Cambridge, MA: MIT Press.
- Smith, Lones, & Sorensen, Peter Norman. 2014. Rational social learning by random sampling.
- Takahashi, Satoru. 2010. Community enforcement when players observe partners' past play. *Journal of Economic Theory*, **145**(1), 42–62.
- Young, H. Peyton. 1993a. The evolution of conventions. *Econometrica*, **61**(1), 57–84.
- Young, H. Peyton. 1993b. An evolutionary model of bargaining. *Journal of Economic Theory*, **59**(1), 145–168.
- Young, H. Peyton. 2015. The evolution of social norms. *Annual Reviews of Economics*, **7**(1), 359–387.