

The absence of LPA1 receptor results in lipidome dysregulation and Neuropeptide-Y underexpression

Sara Tabbai¹, Román D. Moreno-Fernandez¹, Margarita Pérez-Martín², Cristina Rosell-Valle³, Estela Castilla-Ortega⁴, María-Fernández⁵, Jerold Chun⁶, Fernando Rodríguez de Fonseca⁴, Guillermo Estivill-Torrús⁷, Luis J. Santín¹, Carmen Pedraza¹.

(1) *Departamento de Psicobiología y Metodología en las CC, Instituto de Investigación Biomédica de Málaga (IBIMA), Universidad de Málaga, Málaga, Spain*

(2) *Departamento de Biología Celular, Genética y Fisiología. Facultad de Ciencias. Instituto de Investigación Biomédica de Málaga (IBIMA), Universidad de Málaga, Málaga, Spain*

(3) *Unidad de Producción de Reprogramación Celular, GMP-Iniciativa Andaluza en Terapia Avanzadas, Junta de Andalucía, Spain*

(4) *Unidad de Gestión Clínica de Salud Mental, Instituto de Investigación Biomédica de Málaga (IBIMA), Hospital Regional Universitario de Málaga, Málaga, Spain*

(5) *Departamento de Fisiología y Medicina Deportiva, Instituto de Investigación Biomédica de Málaga (IBIMA), Universidad de Málaga, Málaga, Spain*

(6) *Sanford Burnham Prebys Medical Discovery Institute, La Jolla, CA, USA*

(7) *Unidad de Gestión Clínica de Neurociencias, Instituto de Investigación Biomédica de Málaga (IBIMA), Hospital Regional Universitario de Málaga, Málaga, Spain*

Abstract

LPA1 receptor is one of the six characterized G protein-coupled receptors (LPA₁₋₆) through which lysophosphatidic acid acts as an intercellular signaling molecule. It has been shown that the LPA1 receptor is involved in emotional regulation and, when depleted, has a key role in vulnerability to stress. In this sense, maLPA1-null mice, a knockout model for LPA1 receptor has been recently proposed as a model of anxious depression. Here, we sought to elucidate the effect of the genetic depletion of this receptor of LPA₁ receptor in both lipidome and Neuropeptide-Y (NPY) signaling, two factors associated with adaptive stress regulation. For that purpose, we measured the lipidomic profile of wild-type mice and maLPA1-null mice in both hippocampus and serum. In addition, through immunohistochemical procedures we quantified NPY⁺ cells in hippocampus, basolateral amygdala (BLA) and central amygdala (CeA). Interestingly, the comparative lipidomics analysis revealed differences in certain subspecies which are related to LPA1 receptor functionality. Regarding NPY, we found a reduction in BLA, but

not in hippocampus. Overall, both lipid abnormalities and amygdalar dysfunction of NPY can be related to lower resources in stress coping and, in turn, higher vulnerability to the noxious effect of stress that might lead to anxiety and depressive-like states.

Funding: Andalusian Ministry of Economy, Innovation, Science and Employment (SEJ1863 to C.P) and of Health (Nicolas Monardes programme, to G.E-T); the Spanish Ministry of Economy and Competitiveness (PSI2013-44901-P to L.J.S. and C.P.). Author R.D. M-F holds a Grant of the Spanish Ministry of Education, Culture and Sports (FPU14/01610). Author S.T. holds a Grant of the Andalusian Ministry of Economy, Innovation, Science and Employment (FPDI 2014). Author E.C-O holds a grant from the Spanish Ministry of Economy and Competitiveness co-funded by European Research Development Fund (UE-ERDF) (code: PSI2015-73156-JIN). I Plan Propio de Investigación y Transferencia, Universidad de Málaga. Campus de Excelencia. Andalucía Tech.