



V Encuentro de Oceanografía Física 2018 (EOF)
Vigo (Spain), 20-22th June 2018

BOTTOM-UP CONTROL OF SARDINE AND ANCHOVY POPULATION CYCLES IN THE CANARY CURRENT: INSIGHTS FROM AN END-TO-END MODEL SIMULATION

J. C. Sánchez-Garrido¹, F. E. Werner², J. Fiechter³, A. Ramos⁴, E. Curchitser⁵, K. Rose⁶, J. G. Lafuente¹, J. Arístegui⁷, S. Hernández-León⁷ and A. R. Santana⁴

¹GOFIMA, Applied Physics II, University of Malaga, Spain
jcsanchez@ctima.uma.es, glafuente@ctima.uma.es

²NOAA Fisheries, 1315, Silver Spring, USA
cisco.werner@noaa.gov

³Institute of Marine Sciences, University of California, Santa Cruz, USA
fiechter@ucsc.edu

⁴Fac. Ciencias del Mar, Univ. Las Palmas de Gran Canaria, Las Palmas, Spain
antonio.ramos@ulpgc.es, angel.santana@ulpgc.es

⁵Department of Environmental Sciences, Rutgers University, New Brunswick, USA
enrique@marine.rutgers.edu

⁶University of Maryland Center for Environmental Science, Cambridge, USA
krose@umces.edu

⁷Instituto de Oceanografía y Cambio Global, ULPGC, Spain
shernandezleon@ulpgc.es, javier.aristegui@ulpgc.es

Abstract: Sardine and anchovy can exhibit dramatic decadal-scale shifts in abundance in response to climate variability. Sharpe declines of these populations entail particularly serious commercial and ecological consequences in eastern boundary current ecosystems, where they sustain major world fisheries and provide the forage for a broad variety of predators. Understanding the mechanisms and environmental forcing that drive the observed fish variability remains a challenging problem. The modelling study presented here provides an approach that bridges a comprehensive database with an end-to-end modelling framework enabling the investigation of the sources of variability of sardine and anchovy in the Canary Current System. Different biological traits and behaviour prescribed for sardine and anchovy gave rise to different distribution and displacements of the populations, but to a rather synchronous variability in terms of abundance and biomass, in qualitative agreement with historical landing records. Analysis of years with anomalously high increase and decline of the adult population points to food availability (instead of temperature or other environmental drivers) as the main environmental factor determining recruitment for both sardine (via spawning and survival of feeding age-0 individuals) and anchovy (via survival of feeding age-0). Consistent with this, the two species thrive under enhanced upwelling-favourable winds, but only up to some threshold of the wind velocity beyond which larval drift mortality exceeds the positive effect of the extra food supply. Based on the analysis of the simulation, we found that anchovy larvae

are particularly vulnerable to enhanced wind-driven advection, and as such do better with more moderate upwelling than sardines.

Key words: physical-biological interactions, small pelagic fisheries, eastern boundary currents, biophysical models, end-to-end models

References:

Jose C. Sanchez-Garrido, Werner, F. E., Fiechter, J., Ramos, A., Curchitser, E., Rose, K. A., García-Lafuente, J., Arístegui, J., Hernández-León, S., Rodríguez-santana, A. (2018). Decadal-scale variability of sardine and anchovy reproduced by an end-to-end coupled simulation of the Canary Current ecosystem. Submitted to Progress in Oceanography.

Rose, K. A. , Fiechter, J., Curchitser, E. N., Hedstrom, K., Bernal, M., Creekmore, S., Haynie, A., Ito, S., Lluch-Cota, S., Megrey, B. A., Edwards, C. A., Checkley, D., Koslow, T., McClatchie, S., Werner, F., MacCall, A., Agostini, V., (2015). Demonstration of a fully-coupled end-to-end model for small pelagic fish using sardine and anchovy in the California Current, Progress in Oceanography, 138, 348-380.

Fiechter, J., Kenneth A. R, Curchitser, E. N., Hedstrom, K. S, (2015). The role of environmental controls in determining sardine and anchovy population cycles in the California Current: Analysis of an end-to-end model, Progress in Oceanography, 138, 381-398.