

DYNAMIC ECONOMETRIC MODELS

Vol. 11 – Nicolaus Copernicus University – Toruń – 2011

Piotr Fiszeder

Nicolaus Copernicus University in Toruń

Minimum Variance Portfolio Selection for Large Number of Stocks – Application of Time-Varying Covariance Matrices

Abstract. An evaluation of the efficiency of different methods of the minimum variance portfolio selection was performed for seventy stocks from the Warsaw Stock Exchange. Eight specifications of multivariate GARCH models and six other methods were used. The application of all considered GARCH-class models was more efficient in stocks allocation than the implementation of the other analyzed methods. The simple specifications of multivariate GARCH models, whose parameters were estimated in two stages, like the DCC and CCC models were the best performing models.

Key words: multivariate GARCH models, time-varying covariance matrix, portfolio selection.

Introduction

The selection of estimators of the population mean, variance and covariance of financial returns and closely connected with it the selection of forecasting methods of the population mean, variance and covariance of returns plays a vital role in the construction of efficient portfolios. There is a consensus among both financial market practitioners and scientists, that financial returns are difficult to forecast and the portfolio construction process according to the criteria proposed by Markowitz (1952, 1959) is very sensitive to the selection of an estimator of the expected value of returns (see Michaud, 1989; Best, Grauer, 1991; Chopra, Ziemba, 1993). Small differences in estimates of the expected returns often lead to a meaningful portfolio reconstruction (see e.g. Jobson, Korkie, 1980). It is believed that the estimation of population variances and covariances of returns is easier than the estimation of population means (Merton 1980; Nelson, 1992), however the selection of estimators of population variances and covariances of returns has also a significant impact on an asset allocation (see e.g. Litterman, Winkelmann, 1998; Chan, Karceski, Lakonishok, 1999). Traditionally used estimators of the population mean, variance and co-

variance, namely the arithmetic mean and empirical variance and covariance calculated on the basis of available historical data do not give the best results in efficient portfolios construction (see e.g. Jorion, 1991; Sheedy, Trevor, Wood, 1999; Johannes, Polson, Stroud, 2002; Flavin, Wickens, 2006).

The classical approach to the selection of efficient portfolios is static, i.e. a chosen period or moment is considered, but not their sequence. Such an approach ignores therefore the variability of conditional variances and covariances of returns. A dynamic approach to the selection of efficient portfolios for homogeneous assets, i.e. stocks, based on the forecasts of variances and covariances of returns constructed from multivariate GARCH models is presented in this paper. There is an extensive literature on time-varying conditional variances of returns but also on time-varying conditional covariances and correlation coefficients between financial series. The multivariate GARCH process allows to describe both time-varying conditional variances and covariances of returns. If variances and covariances of returns are not constant, then the forecasts based on multivariate GARCH models should give additional benefits in the selection of efficient portfolios. The literature on the application of GARCH models in construction of portfolios is poor, and the results of such analyses, due to the complexity of the problem, are still fragmentary. In most of the studies the univariate GARCH model is applied or a very limited number of assets are used. Among the few investigations in which multivariate GARCH models were applied for large portfolios the following papers can be mentioned: Engle, Shephard (2001), Engle, Colacito (2006), Osiewalski, Pajor (2010).

The primary purpose of this study is to evaluate the effectiveness of different methods of the minimum variance portfolio selection, mainly with the application of various multivariate specifications of GARCH models. The adopted approaches to the portfolio construction differ only by forecasting methods of the covariance matrix of returns. The paper is an extension of the author's earlier work presented in Fiszeder (2004, 2007). The plan for the rest of the paper is as follows. Section 1 outlines the way in which efficient portfolios are selected. In section 2 competing methods of the covariance matrix estimation are presented. Section 3 contains the effectiveness analysis of the minimum variance portfolio selection for seventy stocks quoted on the Warsaw Stock Exchange (WSE) and section 4 presents the conclusions.

1. Dynamic Process of Portfolio Selection with Application of GARCH Models

It is difficult to evaluate the influence of the choice of the covariance matrix estimator against the selection of the population mean estimator in construction of efficient portfolios. One of the ways to eliminate the influence of the choice of the mean estimator is the minimum variance portfolio selection. Shares of

individual assets in the minimum variance portfolio depend solely on the covariance matrix, that is why such procedure is used in the paper.

For a given t , based on all the available data from the period $[1, t]$ parameters of a GARCH model are estimated¹. The forecast of the covariance matrix based on the estimated model is formulated at time $t + \tau$, where the τ is the forecast horizon. The constructed forecast is used for the selection of efficient portfolio. Let $\mathbf{W}'_{t+\tau|t,p} = (w_{1t+\tau|t,p}, w_{2t+\tau|t,p}, \dots, w_{Nt+\tau|t,p})$, where $w_{it+\tau|t,p}$ is the share of asset i in the portfolio at time $t + \tau$, $\mathbf{H}_{t+\tau|t,p}$ is the forecast of the conditional covariance matrix of returns at time $t + \tau$. The variance of portfolio returns is then equal to $\mathbf{W}'_{t+\tau|t,p} \mathbf{H}_{t+\tau|t,p} \mathbf{W}_{t+\tau|t,p}$. In order to select the minimum variance portfolio (the global minimum) the following quadratic programming problem has to be solved:

$$\mathbf{W}'_{t+\tau|t,p} \mathbf{H}_{t+\tau|t,p} \mathbf{W}_{t+\tau|t,p} \rightarrow \min, \quad (1)$$

subject to the constraint:

$$\mathbf{W}'_{t+\tau|t,p} \mathbf{1} = 1, \quad (2)$$

where $\mathbf{1}$ is the $N \times 1$ vector of ones.

When short selling is not allowed the boundary conditions have to be additionally imposed:

$$w_{it+\tau|t,p} \geq 0 \quad \text{for } i = 1, 2, \dots, N. \quad (3)$$

The whole procedure is repeated for successive periods (with the arrival of subsequent data).

If short selling is allowed, then shares of assets in the minimum variance portfolio are defined by the following formula:

$$\mathbf{W}_{t+\tau|t,p} = \frac{1}{C_{t+\tau|t,p}} \mathbf{H}_{t+\tau|t,p}^{-1} \mathbf{1}, \quad (4)$$

where $C_{t+\tau|t,p} = \mathbf{1}' \mathbf{H}_{t+\tau|t,p}^{-1} \mathbf{1}$.

The variance of the minimum variance portfolio is then equal to $V_{t+\tau|t,p} = 1/C_{t+\tau|t,p}$.

Other efficient portfolios can be selected by finding the minimum variance portfolio subject to a minimum expected return.

¹ If conditional expected values of returns are different from zero, then parameters of conditional mean equations should also be estimated.

2. Specifications of Multivariate GARCH Models

Eight parameterizations of multivariate GARCH models were applied in the analysis: Scalar BEKK, Integrated, CCC, Orthogonal, DCC, Integrated DCC, DECO-DCC and additionally Scalar BEKK with Student-t innovations. The results obtained for the GARCH models were compared with the outcomes for six other methods: equal shares in all stocks, the unconditional covariance matrix of returns, the rolling covariance matrix², the 25-day rolling covariance matrix, the exponentially weighted moving average estimate of the covariance matrix (hereafter the EWMA covariance matrix)³, the EWMA covariance matrix with a smoothing parameter set to 0.94 (hereafter the RiskMetrics⁴).

Only some basic information about the considered multivariate models is given below. More details can be found for example in Bauwens, Laurent, Rombouts (2006) or Silvennoinen, Teräsvirta (2009).

Let assume that $\boldsymbol{\varepsilon}_t$ can be either returns with the mean zero or residuals from filtered time series:

$$\boldsymbol{\varepsilon}_t \mid \boldsymbol{\psi}_{t-1} \sim N(\mathbf{0}, \mathbf{H}_t), \quad (5)$$

where $\boldsymbol{\psi}_{t-1}$ is the set of all information available at time $t-1$ and \mathbf{H}_t is the $N \times N$ symmetric conditional covariance matrix.

The estimation of parameters of the general form of a multivariate GARCH model, VECH model⁵ (Kraft, Engle, 1983), is very difficult even for a small number of assets. For this reason simpler parameterizations of multivariate GARCH models were used in the study. Considered were only those specifications which ensure the positive definiteness of the covariance matrix.

Baba, Engle, Kraft, Kroner (1990) introduced the following form the so-called BEKK(p,q) model⁶ (see Engle, Kroner, 1995):

$$\mathbf{H}_t = \mathbf{C}\mathbf{C}' + \sum_{i=1}^q \mathbf{D}_i \boldsymbol{\varepsilon}_{t-i} \boldsymbol{\varepsilon}_{t-i}' \mathbf{D}_i' + \sum_{j=1}^p \mathbf{E}_j \mathbf{H}_{t-j} \mathbf{E}_j', \quad (6)$$

where \mathbf{C} , \mathbf{D}_i and \mathbf{E}_j are $N \times N$ parameter matrices and \mathbf{C} is an upper triangular matrix.

² The value of a rolling window was chosen to minimize the variance of a portfolio in a pre-sample.

³ The value of a smoothing parameter was chosen to minimize the variance of a portfolio in a pre-sample.

⁴ The RiskMetrics methodology was developed in the investment bank J. P. Morgan for measuring market risk with VaR. The value of a smoothing parameter in the EWMA model was often set by financial market practitioners to 0.94 for daily data.

⁵ The name of the model comes from the application of the vech operator for the conditional covariance matrix.

⁶ The name of the model is formed by the first letters of the authors surnames.

The BEKK model is also too complex for large portfolios and the imposition of restrictions on parameters in estimation is necessary. The so-called Scalar BEKK model can be obtained by replacing the matrices \mathbf{D}_i and \mathbf{E}_j by the scalars $d_i^{1/2}$ and $e_j^{1/2}$. Additionally the variance targeting approach (see Engle, Mezrich, 1996) was used. For example for $q = p = 1$ instead of the product $\mathbf{C}\mathbf{C}'$ the following formula can be substituted:

$$\mathbf{C}\mathbf{C}' = (1 - d_1 - e_1)\mathbf{S}, \tag{7}$$

where \mathbf{S} is the sample covariance matrix given as $\mathbf{S} = \frac{1}{t} \sum_{i=1}^t \boldsymbol{\varepsilon}_i \boldsymbol{\varepsilon}_i'$. Two specifications of the conditional distribution of $\boldsymbol{\varepsilon}_i$ in (5) were considered for the Scalar BEKK model, namely multivariate normal and Student-t.

A further simplification of the model can be obtained by the assumptions $\mathbf{C}\mathbf{C}' = 0$ and $\sum_{i=1}^q d_i + \sum_{j=1}^p e_j = 1$. This formulation is often called the Integrated multivariate model⁷. For $q = p = 1$ the model has only one parameter.

In the Constant Conditional Correlations (CCC) model of Bollerslev (1990), which is outside the BEKK class, the time-varying conditional covariances are parameterized to be proportional to the product of the corresponding conditional standard deviations:

$$\mathbf{H}_t = \mathbf{D}_t \boldsymbol{\Gamma} \mathbf{D}_t, \tag{8}$$

where \mathbf{D}_t is a $N \times N$ diagonal matrix $\mathbf{D}_t = \text{diag}(h_{1t}^{1/2}, h_{2t}^{1/2}, \dots, h_{Nt}^{1/2})$ with h_{it} defined as any univariate GARCH model, and $\boldsymbol{\Gamma}$ is a $N \times N$ matrix of the time-invariant conditional correlations.

Alexander and Chibumba (1996) introduced the Orthogonal GARCH model defined as:

$$\mathbf{V}^{-1/2} \boldsymbol{\varepsilon}_t = \mathbf{u}_t = \boldsymbol{\Lambda}_m \mathbf{f}_t, \tag{9}$$

$$\mathbf{H}_t = \mathbf{V}^{1/2} \mathbf{V}_t \mathbf{V}^{1/2}, \tag{10}$$

where $\mathbf{V} = \text{diag}(v_1, v_2, \dots, v_N)$, with v_i the population variance of ε_{it} , $\boldsymbol{\Lambda}_m$ is a matrix of dimension $N \times m$ given by $\boldsymbol{\Lambda}_m = \mathbf{P}_m \text{diag}(l_1^{1/2}, l_2^{1/2}, \dots, l_m^{1/2})$, $l_1 \geq l_2 \geq \dots \geq l_m > 0$ being the m largest eigenvalues of the population correlation matrix of \mathbf{u}_t , and \mathbf{P}_m the $N \times m$ matrix of associated (mutually orthogonal) eigenvectors, $\mathbf{f}_t = (f_{1t} f_{2t} \dots f_{mt})'$ is a random process such that $E_{t-1}(\mathbf{f}_t) = 0$, the

⁷ The name of the model comes from an analogous parameterization of the univariate IGARCH model.

conditional covariance matrix of \mathbf{f}_t is equal to $\mathbf{Q}_t = \text{diag}(\sigma_{f_{1t}}^2, \sigma_{f_{2t}}^2, \dots, \sigma_{f_{mt}}^2)$, $\sigma_{f_{it}}^2$ is defined as a univariate GARCH model (for $i = 1, 2, \dots, m$) and the conditional covariance matrix of \mathbf{u}_t is equal to $\mathbf{V}_t = \mathbf{\Lambda}_m \mathbf{Q}_t \mathbf{\Lambda}'_m$.

The Dynamic Conditional Correlation (DCC) model of Engle (2002) can be defined as:

$$\mathbf{H}_t = \mathbf{D}_t \mathbf{R}_t \mathbf{D}_t, \quad (11)$$

$$\mathbf{R}_t = \mathbf{Q}_t^{*-1} \mathbf{Q}_t \mathbf{Q}_t^{*-1}, \quad (12)$$

$$\mathbf{Q}_t = (1 - \sum_{i=1}^Q \alpha_i - \sum_{j=1}^P \beta_j) \mathbf{S} + \sum_{i=1}^Q \alpha_i (\mathbf{z}_{t-i} \mathbf{z}'_{t-i}) + \sum_{j=1}^P \beta_j \mathbf{Q}_{t-j}, \quad (13)$$

where $\mathbf{D}_t = \text{diag}(h_{1t}^{1/2}, h_{2t}^{1/2}, \dots, h_{Nt}^{1/2})$, h_{kt} can be defined as a univariate GARCH model (for $k = 1, 2, \dots, N$), \mathbf{z}_t is a vector of standardized values of ε_{kt} , i.e. $z_{kt} = \varepsilon_{kt} / \sqrt{h_{kt}}$, \mathbf{R}_t is the time-varying conditional correlation matrix of \mathbf{z}_t , \mathbf{S} is the sample covariance matrix of \mathbf{z}_t , \mathbf{Q}_t^* is a diagonal matrix composed of the square root of the diagonal elements of \mathbf{Q}_t and the parameters have to satisfy

the condition $0 < \sum_{i=1}^Q \alpha_i + \sum_{j=1}^P \beta_j < 1$.

Besides the standard DCC model two other modifications were also considered. When the sum of the parameters is equal to one $\sum_{i=1}^Q \alpha_i + \sum_{j=1}^P \beta_j = 1$ the DCC model is called the Integrated DCC model. Engle and Kelly (2008) introduced the DECO-DCC (dynamic equicorrelation) model. In this parameterization of the DCC model the equality of all pairwise conditional correlations at each time is assumed.

3. Evaluation of Portfolio Performance for Seventy Stocks Quoted on the WSE

The presented approach for the minimum variance portfolio selection was evaluated for Polish stocks. The investigated period was November 17, 2000⁸ to June 30, 2009 (2158 daily returns). All stocks quoted in the specified period on the Warsaw Stock Exchange were considered in the analysis. The companies for which the percent of non-trading days was higher than 5% were omitted in order to avoid the problem of non-synchronous trading. In total seventy stocks were analyzed, however all the models used in the paper can be applied for

⁸ Since the introduction of the new trading system Warset.

a much larger number of companies. An evaluation of portfolio performance was based on data from January 2004 to June 2009 (1380 observations). The following five steps were performed: (1) the estimation of parameters of all considered models⁹ (at the beginning for data from November 17, 2000 to December 30, 2003)¹⁰, (2) the construction of one-day ahead forecasts of the conditional covariance matrix, (3) the minimum variance portfolio selection (4) the ex post calculation of the portfolio variance as a square of the realized portfolio return, (5) the extension of the sample with one observation. All the steps were repeated 1380 times. Every time the estimation of parameters was performed for increasing sample size. For each model and method the mean of the portfolio variances was calculated. The results are presented in Table 1 (a square root of the mean is given).

Table 1. Estimates of the standard deviations of returns for the minimum variance portfolios

Portfolio designation	Standard deviation ($\times 10^{-2}$)	Rank
Integrated DCC	0.9257	1
DCC	0.9316	2
CCC	0.9326	3
DECO-DCC	0.9843	4
Scalar BEKK Student-t	1.0030	5
Scalar BEKK	1.0065	6
Orthogonal 70 factors	1.0854	7
Integrated	1.1155	8
Unconditional covariance matrix	1.1155	8
Rolling covariance matrix	1.2634	10
Equal shares in all stocks	1.3538	11
RiskMetrics	1.6549	12
EWMA covariance matrix	2.9846	13
25-day rolling covariance matrix	29.7629	14

The best performing model in the selection of the minimum variance portfolio was the Integrated DCC model. On the other hand the worst performing method was the 25-day rolling covariance matrix.

An application of all the GARCH-class models was more effective in allocation of stocks than the implementation of the other analyzed methods. The first four positions in the ranking (see Table 1) were occupied by the models, whose parameters were estimated in two steps and in the first stage parameters of a univariate GARCH model were estimated.

The Integrated model had the worst rank among all the applied GARCH models, however in this case the model was reduced to the unconditional covar-

⁹ Codes written by the author in the Gauss programming language were applied.

¹⁰ Logarithmic returns were used in the study. Because daily data were used, the logarithmic return on a portfolio is very close to the weighted average of the logarithmic returns on the individual assets.

iance matrix of returns¹¹. The Orthogonal GARCH model, which assumes that returns depend on common independent factors extracted by principal component analysis ranked the one before last among the GARCH models. This result follows, first of all, from the loss of important information.

The methods used by financial market practitioners, i.e. the rolling covariance matrix and the exponentially weighted moving average estimate of the covariance matrix took a distant positions in the ranking, even behind the unconditional covariance matrix of returns. The very poor result of the rolling covariance matrix with a low value of the rolling window equal to 25 deserves to be highlighted. The value of the smoothing parameter, set often for daily data in the RiskMetrics methodology to 0.94 is not the optimal value for the Polish stock market (it is definitely too low).

For a large number of assets like in this study, probably none of the restrictions, assumed by the considered GARCH models, is met. However, one can attempt to indicate the restrictions, which have a stronger negative influence on the performance of portfolio, in the case when they are not met. For example, the assumption that models describing conditional variances and covariances have the same parameters for all series¹² (like in the Scalar BEKK and Integrated models) is probably too strong. It also seems that the presupposition of the constancy of conditional correlation coefficients is less restrictive.

The statistical significance of the observed differences between the performance of the different models was not verified, however some of them are relevant from the economic point of view. For example, the difference between the first in the ranking, namely the Integrated DCC model and the unconditional covariance matrix of returns means a decrease of about 3 percentage points of the standard deviation per year.

Additionally, values of the Schwarz information criterion (SIC) were calculated, when it was possible to evaluate a joint likelihood function (see Table 2). The ranking of the models according to the SIC is similar to the one for the portfolio performance evaluation (Table 1). The exceptions are the Scalar BEKK model with Student-t innovations and the Integrated model which take better position in the SIC ranking. The better performance of GARCH models with Student-t innovations in the rankings based on the information criteria is common in other studies (see Fiszeder, 2009). The rankings of models constructed on the basis of the information criteria provide useful clues for the selection of models for the minimum variance portfolio construction. It has to be remembered, however, that some characteristics are crucial in the evaluation of the general fit of the model in a sample (like for instance a type of conditional density), but their influence on the performance of portfolio is not so important.

¹¹ Estimates of the parameter on the lagged conditional covariance matrix were equal to one.

¹² It means, that the volatility dynamics of all series is very similar.

Table 2. Ranking of the models based on the Schwarz information criterion

Portfolio designation	SIC	Rank
Integrated DCC	-679060.02	1
DCC	-666933.43	2
Scalar BEKK Student-t	-661928.17	3
DECO-DCC	-658854.85	4
Integrated	-656773.56	5
Scalar BEKK	-645459.24	6
RiskMetrics	2052174.1	7

It is interesting to compare the results of this analysis with the similar study for twenty stocks (Fiszeder, 2007). The GARCH models, whose parameters were estimated in one stage (like the Scalar BEKK and Integrated models) took further positions in the ranking. It seems that the assumption that models describing conditional variances have the same parameters for all series becomes more and more restrictive for a larger number of assets. Furthermore, the differences between evaluations of portfolios, obtained with the use of the applied methods, were greater for the seventy stocks. The reason for such results is probably the higher differentiation of the seventy stocks, among which are both huge but also very small companies. In the study for the twenty stocks only the biggest companies were considered. This comparison for different numbers of stocks clearly shows that the received results depend on the properties of financial time series.

Conclusions

The dynamic approach to the selection of efficient portfolios for a large number of homogeneous assets, i.e. stocks, based on the forecasts of variances and covariances of returns constructed from multivariate GARCH models has been presented in this paper. An evaluation of the efficiency for different methods of the minimum variance portfolio selection was performed for the seventy stocks from the Warsaw Stock Exchange. The eight specifications of multivariate GARCH models and the six other methods were used.

Capturing time-varying variances and covariances of stock returns does not always increase the efficiency of the asset allocation process. The application of all the considered GARCH-class models was more efficacious in the allocation of stocks than the implementation of the other analyzed methods, including the methods employed often by financial market practitioners. The simple specifications of multivariate GARCH models, whose parameters were estimated in two stages, like the DCC and CCC models, were the best performing models.

References

- Alexander, C., Chibumba, A. (1996), Multivariate Orthogonal Factor GARCH, University of Sussex Discussion Papers in Mathematics.
- Baba, Y., Engle, R. F., Kraft, D. F., Kroner, K. F. (1990), Multivariate Simultaneous Generalized ARCH, Department of Economics, University of California at San Diego, Working Paper.
- Bauwens, L., Laurent, S., Rombouts, J., V., K. (2006), Multivariate GARCH Models: A Survey, *Journal of Applied Econometrics*, 21, 1, 79–110.
- Best, M. J., Grauer, R. R. (1991), On the Sensitivity of Mean-Variance-Efficient Portfolios to Changes in Asset Means: Some Analytical and Computational Results, *Review of Financial Studies*, 4, 315–342.
- Bollerslev, T. (1990), Modelling the Coherence in Short-Run Nominal Exchange Rates: A Multivariate Generalized ARCH Approach, *Review of Economics and Statistics*, 72, 498–505.
- Chan, L. K. C., Karceski, J., Lakonishok, J. (1999), On Portfolio Optimization: Forecasting Covariances and Choosing the Risk Model, *Review of Financial Studies*, 12, 937–974.
- Chopra, V. K., Ziemba, W. T. (1993), The Effect of Errors in Means, Variances and Covariances on Optimal Portfolio Choice, *Journal of Portfolio Management*, 19, 6–11.
- Engle, R. F. (2002), Dynamic Conditional Correlation – A Simple Class of Multivariate GARCH Models, *Journal of Business and Economic Statistics*, 20, 339–350.
- Engle, R. F., Colacito, R. (2006), Testing and Valuing Dynamic Correlations for Asset Allocation, *Journal of Business and Economic Statistics*, 24, 2, 238–253.
- Engle, R. F., Kelly, B. (2008), Dynamic Equicorrelation, Working Paper, Stern School of Business, New York.
- Engle, R. F., Kroner, K. F. (1995), Multivariate Simultaneous Generalized ARCH, *Econometric Theory*, 11, 122–150.
- Engle, R. F., Mezrich, J. (1996), GARCH for Groups, *Risk*, 9, No. 8, 36–40.
- Engle, R. F., Sheppard, K. (2001), Theoretical and Empirical Properties of Dynamic Conditional Correlation Multivariate GARCH, NBER Working Paper No. 8554.
- Fiszeder, P. (2004), Dynamiczna alokacja aktywów – Model Markowitza (Dynamic Assets Allocation – Markowitz Model), *Rynki finansowe – prognozy a decyzje (Financial Markets – Forecasts and Decisions)*, *Acta Universitatis Lodzianis, Folia Oeconomica*, 177, Uniwersytet Łódzki, Łódź.
- Fiszeder, P. (2007), Konstrukcja portfeli efektywnych z zastosowaniem wielorównaniowych modeli GARCH (Efficient Portfolios Construction with Application of Multivariate GARCH Models), *Folia Oeconomica Cracoviensia*, 48, 47–68.
- Fiszeder, P. (2009), *Modele klasy GARCH w empirycznych badaniach finansowych, (The Class of GARCH Models in Empirical Finance)*, Wydawnictwo UMK, Toruń.
- Flavin, T. J., Wickens, M. R. (2006), Optimal International Asset Allocation with Time-Varying Risk, *Scottish Journal of Political Economy*, 53, 5, 543–564.
- Jobson, J. D., Korkie, B. (1980), Estimation for Markowitz Efficient Portfolios, *Journal of American Statistical Association*, 75, 544–554.
- Johannes, M., Polson, N., Stroud, J. (2002), Sequential Optimal Portfolio Performance: Market and Volatility Timing, Columbia University, Working Paper.
- Jorion, P. (1991), Bayesian and CAPM Estimators of the Means: Implications for Portfolio Selection, *Journal of Banking and Finance*, 15, 717–727.
- Kraft, D. F., Engle, R. F. (1983), Autoregressive Conditional Heteroskedasticity in Multiple Time Series, Department of Economics, UCSD, Working Paper.
- Litterman, R., Winkelmann, K. (1998), Estimating Covariance Matrices, Risk Management Series, Goldman Sachs.
- Markowitz, H. M. (1952), Portfolio Selection, *Journal of Finance*, 7, 77–91.
- Markowitz, H. M. (1959), *Portfolio Selection: Efficient Diversification of Investments*, Yale University Press, New Haven, CT.
- Merton, R. C. (1980), On Estimating the Expected Return on the Market: An Exploratory Investigation, *Journal of Financial Economics*, 8, 323–361.

- Michaud, R. O. (1989), The Markowitz Optimization Enigma: Is 'Optimized' Optimal?, *Financial Analysts Journal*, 45, 31–42.
- Nelson, D. B. (1992), Filtering and Forecasting with Misspecified ARCH Models I. Getting the Right Variance with the Wrong Model, *Journal of Econometrics*, 52, 61–90.
- Osiewalski, J., Pajor, A. (2010), Bayesian Value-at-Risk and Expected Shortfall for a Portfolio (Multi- and Univariate Approaches), Paper presented at The V Symposium FENS 2010, Warsaw.
- Sheedy, E., Trevor, R., Wood, J. (1999), Asset-Allocation Decisions When Risk is Changing, *Journal of Financial Research*, 22, 3, 301–315.
- Silvennoinen, A., Teräsvirta, T. (2009), *Multivariate GARCH Models*, in Andersen T. G., Davis R. A., Kreiss J. P., Mikosch T. (ed.), *Handbook of Financial Time Series*, 201–229, Springer, New York.

Konstrukcja portfeli o minimalnej wariancji dla dużej liczby spółek – zastosowanie zmiennych w czasie macierzy kowariancji

Z a r y s t r e ś c i. W pracy dokonano oceny efektywności różnych metod tworzenia portfeli o minimalnej wariancji, w tym przede wszystkim z wykorzystaniem różnych specyfikacji wielorównaniowych modeli GARCH. Badanie zostało przeprowadzone dla 70 spółek notowanych na GPW w Warszawie. Zastosowano osiem parametryzacji modelu GARCH: skalarny BEKK, zintegrowany, CCC, ortogonalny dla 70 czynników, DCC, zintegrowany DCC, DECO-DCC, skalarny BEKK z warunkowym rozkładem t Studenta oraz sześć innych metod: równe udziały dla wszystkich aktywów, bezwarunkowa macierz kowariancji stóp zwrotu, ruchoma macierz kowariancji, ruchoma macierz kowariancji ze stałą wygładzania równą 25, metoda wyrównywania wykładniczego dla macierzy kowariancji oraz metoda wyrównywania wykładniczego dla macierzy kowariancji z parametrem wygasania równym 0,94.

S ł o w a k l u c z o w e: wielorównaniowe modele GARCH, zmieniająca się w czasie macierz kowariancji, konstrukcja portfela.

