

Characterization of Cfa1, a Monofunctional Acyl Carrier Protein Involved in the Biosynthesis of the Phytotoxin Coronatine

Heather Seidle,¹ Vidhya Rangaswamy,^{2†} Robin Couch,¹ Carol L. Bender,²
and Ronald J. Parry^{1*}

Department of Chemistry, Rice University, Houston, Texas 77005,¹ and Department of Entomology and Plant Pathology, Oklahoma State University, Stillwater, Oklahoma 74078²

Received 27 October 2003/Accepted 4 January 2004

Cfa1 was overproduced in *Escherichia coli* and *Pseudomonas syringae*, and the degree of 4'-phosphopantethinylation was determined. The malonyl-coenzyme A:acyl carrier protein transacylase (FabD) of *P. syringae* was overproduced and shown to catalyze malonylation of Cfa1, suggesting that FabD plays a role in coronatine biosynthesis. Highly purified Cfa1 did not exhibit self-malonylation activity.

Coronatine (COR) is a non-host-specific chlorosis-inducing phytotoxin produced by several *Pseudomonas syringae* pathogens (1). COR consists of two moieties joined by an amide linkage: the polyketide coronafacic acid (CFA) and coronamic acid (Fig. 1) (1). The CFA polyketide synthase (PKS) genes include two genes (*cfa6* and *cfa7*) that encode modular type I PKS proteins as well as three genes (*cfa1*, *cfa2*, and *cfa3*) that encode monofunctional proteins similar to the acyl carrier protein (ACP), dehydratase, and β -ketoacyl:acyl carrier protein synthase, respectively, of type II fatty acid synthases (FASs) and PKSs (25, 27). The genes coding for the CFA portion of COR are located on a single long transcript driven by a promoter upstream of the *cfl* gene that is regulated by a three-component regulatory system present in the COR gene cluster (17, 24).

A biosynthetic hypothesis for CFA formation was derived from analysis of the CFA PKS (12, 27) and the isolation of a potential biosynthetic intermediate, 2-[1-oxo-2-cyclopenten-2-ylmethyl]-butanoic acid (19, 23, 35). This hypothesis postulates that Cfa1 functions as the carrier in the assembly of a cyclic carboxylic acid that is subsequently transferred from Cfa1 to the ACP domain of Cfa6 via the loading didomain. The important role assigned to Cfa1 and the lack of information concerning PKS ACPs of *Pseudomonas* prompted us to investigate the properties of Cfa1.

Cloning of *cfa1*, *fabD*, and *acp* from *P. syringae*. To clone *cfa1* into the FLAG vectors pFLAG-CTC (Sigma) and pSFFLAG-CTC (4), *cfa1* was amplified from pSAY10 (38) using the following primers: 5' GCCAAGCTTGTGAACGAGATC (HindIII site in bold) and 5' GCCGAATTCCTTGGCCTGCAG (EcoRI site in bold). The PCR product was ligated into the HindIII-EcoRI-digested vectors, and the ligation mixtures were used to transform *Escherichia coli* DH5 α or DH10B, resulting in pFLAG-Cfa1 and pSFFLAG-Cfa1, respectively. For overproduction of Cfa1 in *P. syringae* pv. glycinea PG4180, pFLAG-Cfa1 was linearized with

BamHI and ligated into BamHI-digested pRK415 (13), resulting in pRKFLAG-Cfa1.

Primers were designed to amplify the *fabD* gene from *P. syringae* FF5 (9) based upon the sequence of *fabD* in *Pseudomonas aeruginosa* (<http://Pseudomonas.bit.uq.edu.au/>) and *P. syringae* pv. tomato DC3000 (<http://www.tigr.org>). The forward primer for *fabD* amplification was 5' CCCAAGCTTATGTCTGCATCCCTCGCATTC (HindIII site in bold), and the reverse primer containing a BglII site (bold) was 5' GTTAGATCTGGCCAGTGCCGCGCTG. *P. syringae* FF5 genomic DNA was used as a template for PCR. The PCR product was digested with HindIII and BglII and ligated into HindIII-BglII-digested pSFFLAG-CTC.

The primers designed to amplify the FAS ACP gene from *P. syringae* FF5 were based on the sequence of the FAS ACP gene in *P. syringae* pv. tomato DC3000. The forward primer contained an XhoI site (bold): GTCCTCGAGATGAGCACCATCGAAGAG. The reverse primer contained a BglII site (bold) to allow insertion into the pSFFLAG-CTC vector with removal of the stop codon: GTAGATCTATAAGCCTGATGGCTG. The PCR product was digested with BglII and XhoI and ligated into pSFFLAG-CTC digested with the same enzymes.

The deduced translation products of the cloned *P. syringae* FF5 *fabD* and *acp* genes exhibited high similarities to the corresponding genes from the COR producer *P. syringae* pv. tomato DC3000 and from *P. aeruginosa* as well as lower similarities to homologues found in *E. coli* and *Streptomyces coelicolor* (Table 1). *P. syringae* FF5 was selected as the source of the *fabD* and FAS *acp* genes, since unpublished studies have shown that introduction of the entire COR gene cluster into this strain leads to COR production (Y. Zhao, V. Rangaswamy, and C. L. Bender, unpublished data). *P. syringae* FF5 therefore possesses all the ancillary enzymes required for COR production. In addition, *P. syringae* FF5 provides a convenient host strain to overproduce proteins from the COR pathway (4).

Overproduction of Cfa1-FLAG in *E. coli*. A starter culture of *E. coli* DH5 α containing pFLAG-Cfa1 in terrific broth (TB) plus ampicillin (100 μ g/ml) was used to inoculate 100 ml of the same medium, and cells were grown at 37°C to an optical density at 600 nm (OD₆₀₀) of 0.5. Isopropyl- β -D-thiogalacto-

* Corresponding author. Mailing address: Department of Chemistry, MS60, Rice University, Houston, TX 77005. Phone: (713) 348-2446. Fax: (713) 348-5155. E-mail: parry@rice.edu.

† Present address: Reliance Life Sciences, Jannagar 361140, Gujarat, India.




FIG. 1. Structures of COR, CFA, and coronamic acid (CMA) and presumed formation of COR by action of a ligase.

pyranoside (IPTG) was then added to a final concentration of 0.5 mM, and the cells were incubated an additional 3 h before harvesting. The cells were lysed with 8 ml of CellLytic B-cell lysis agent (Sigma) containing lysozyme (10 mg/ml) and DNase I (0.02 mg/ml). The suspension was shaken gently for 15 min and then centrifuged at $10,000 \times g$ for 15 min. The supernatant was applied to a column of M2 immunoaffinity resin (Sigma) (1 ml). The eluant was reapplied to the column three times, and the column was then washed with 36 ml of Tris-buffered saline (TBS) buffer (50 mM Tris-HCl [pH 7.4], 150 mM MgCl_2). The C-terminal FLAG-tagged Cfa1 protein (Cfa1-FLAG) was eluted from the resin using FLAG peptide (100 $\mu\text{g}/\text{ml}$; TBS buffer) as described in the FLAG manual (Sigma).

Overproduction and purification of Cfa1, FabD, and ACP in *P. syringae*. The pSFFLAG-Cfa1 construct was electroporated into *P. syringae* FF5 cells, and the pRKFLAG-Cfa1 plasmid was mobilized into *P. syringae* pv. glycinea PG4180 using the helper plasmid pRK2013 (2). An overnight culture (20 ml) of *P. syringae* FF5 containing pSFFLAG-Cfa1 in King's medium B (KMB) (14) plus ampicillin (100 $\mu\text{g}/\text{ml}$) was used to inoculate 1 liter of KMB, which was then incubated at 28°C and 250 rpm. After 5 h, the incubation temperature was reduced to 18°C, and cells were grown for 16 h or to an OD_{600} of 0.5 to 1.0. The cells were then induced with IPTG (1 mM), incubated for 24 h at 18°C, and harvested by centrifugation ($8,000 \times g$), and the pellets were stored at -80°C . *P. syringae* PG4180 containing pRKFLAG-Cfa1 was grown in 1 liter of TB plus tetracycline (25 $\mu\text{g}/\text{ml}$) at 18 or 28°C to an A_{600} of 0.5, induced with 1 mM IPTG, and incubated an additional 16 h at the initial incubation temperature (18 or 28°C). Cells were harvested as described for FF5, and pellets were stored at -80°C . Starter cultures of *P. syringae* transformants containing *fabD* and *acp*

in pSFFLAG-CTC were grown overnight as described above, added to 1 liter of KMB with ampicillin, and grown at 28°C until the OD_{600} was 0.5 to 0.8. The cultures were then induced with IPTG (1 mM) and incubated for 16 h at 28°C. The cells were harvested by centrifugation and frozen at -80°C . Frozen pellets were thawed to room temperature and resuspended in a lysis buffer (3 ml per g of cells) containing Tris buffer (100 mM; pH 8), EDTA (150 mM), and lysozyme (0.32 mg/ml). The mixture was incubated for 30 min at 37°C. After incubation, a solution (0.3 ml per g of cells) containing NaCl (150 mM), MgCl_2 (100 mM), CaCl_2 (100 mM), and DNase I (0.2 mg/ml) was added, and the mixture was incubated for 15 min at room temperature with stirring and then centrifuged at $26,000 \times g$. Supernatants were applied three times to a column of M2 resin (1.5 ml), which was washed with TBS (three times, 20 ml). FLAG-tagged proteins were eluted with 4 ml of FLAG peptide (100 $\mu\text{g}/\text{ml}$) and concentrated to a volume of approximately 200 μl with an Amicon concentrator. Proteins were further purified on a Mono Q column equilibrated with 100 mM Na_2HPO_4 , 2 mM EDTA, and 2 mM dithiothreitol. A NaCl gradient (0 to 500 mM) in the same buffer was used to elute the proteins. The purity of the eluted proteins was assessed by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE), and the total protein concentrations were determined with the Advanced protein assay reagent (Cytoskeleton Inc.) using bovine serum albumin or gamma globulin as the standard. The overproduced proteins were analyzed by SDS-PAGE. The mass of Cfa1-FLAG was approximately 12 kDa, which is close to the predicted size of 11.5 kDa (Fig. 2, lanes 2 and 3). The Cfa1-FLAG purified from *P. syringae* FF5 was further purified on a Mono Q column, although no contaminating proteins were visible by SDS-PAGE (Fig. 2, lane 2). After purification, the mass of FabD-FLAG was found to be 34.2 kDa by SDS-PAGE (Fig. 2, lane 4) and 58.7 kDa by native PAGE (data not shown), suggesting that the native protein is a dimer, as reported for FabD from *Streptomyces glaucescens* (10). ACP-FLAG appeared at a mass of approximately 22 kDa by SDS-PAGE (Fig. 2, lane 5), a value which is significantly larger than the expected one (10.76 kDa). Similar behavior has been reported for the FAS ACP of *P. aeruginosa* (15) and is consistent with the fact that some ACPs run anomalously on SDS-PAGE due to their highly charged nature and low hydrophobic amino acid content (15, 21, 32). The predicted mass was obtained for ACP-FLAG when the protein was analyzed by mass spectrometry.

TABLE 1. Amino acid similarity of *P. syringae* FF5 FabD and FAS ACP to some related proteins

Organism	% Identity to <i>P. syringae</i> FabD	Protein identification no.	% Identity to <i>P. syringae</i> FAS ACP	Protein identification no.
<i>P. syringae</i> pv. tomato	97	NP_793606	97	P80923
<i>P. aeruginosa</i>	80	NP_251658.1	88	NP_251656.1
<i>E. coli</i>	58	NP_287226.1	87	NP_287228.1
<i>S. coelicolor</i> A3(2)	37	CAB62719.1	41	CAB62721.1




FIG. 2. SDS-PAGE (8 to 25% gradient) of recombinant Cfa1 expressed in *P. syringae* FF5 at 18°C (lane 2), Cfa1 expressed in *P. syringae* pv. glycinea PG4180 at 18°C (lane 3), FabD expressed in *P. syringae* FF5 at 28°C (lane 4), and FAS ACP expressed in *P. syringae* FF5 at 28°C (lane 5). All proteins contain a C-terminal FLAG tag. Proteins were purified by affinity chromatography followed by anion exchange chromatography (Mono Q), except for lanes 2 and 3, which show Cfa1 purified only by affinity chromatography. The molecular masses of the standards (lanes 1 and 6) were 200, 116, 97.4, 66, 45, 31, 21.5, 14.4, and 6.5 kDa.

MALDI-TOF analysis of Cfa1 and ACP. Cfa1-FLAG obtained from *E. coli* DH5 α was analyzed by positive-ion matrix-assisted laser desorption ionization–time-of-flight (MALDI-TOF) mass spectrometry to determine the degree to which the protein had been converted into its holo form by addition of a 4'-phosphopantetheine moiety. The protein exhibited a peak at mass of 11,516 Da for the apo protein and no peak corresponding to the holo protein (data not shown). The expected sizes for the apo and holo proteins are 11,516.3 and 11,855.3 Da, respectively. Cfa1-FLAG expressed in *P. syringae* FF5 exhibited a small peak at a mass of 11,511 Da, corresponding to the apo protein, and a larger peak at a mass of 11,854 Da, corresponding to the holo protein (data not shown). The ratio between the areas of the two peaks indicated that the ratio of apo to holo protein was ca. 5:95. Similarly, MALDI-TOF analysis of Cfa1-FLAG overproduced at 18°C in the COR producer *P. syringae* pv. glycinea PG4180 showed that about 90% of the protein was in the holo form (data not shown). However, when Cfa1-FLAG was overproduced in PG4180 at 28°C, the ratio of apo to holo Cfa1-FLAG was 2:1. The MALDI-TOF spectrum of the ACP-FLAG protein exhibited peaks for both the apo and holo forms, with only about 20% of the protein present in the holo form (data not shown). This might be the consequence of a 4'-phosphopantetheinyl transferase (PPTase) reaction rate that is slow relative to the rate of ACP-FLAG production. Since the primary purpose for ACP-FLAG production was to check the overproduced FabD for malonyltransferase activity, the factors affecting the ratio of apo to holo ACP-FLAG were not further investigated.

FabD malonyl-CoA transferase assay with Cfa1 and FAS ACP. The malonyl-coenzyme A (CoA) transferase assay was carried out in a total volume of 50 μ l, which contained 25 mM Tris (pH 7), 1 mM dithiothreitol, 10 μ M FabD, and Cfa1-




FIG. 3. Loading of [2-¹⁴C]malonate onto Cfa1 and FAS ACP from *P. syringae* FF5 by FabD. The concentrations of enzymes used were 25 μ M Cfa1 or FAS ACP and 10 μ M FabD. Reaction mixtures were incubated for 10 min at room temperature, and reactions were stopped by the addition of TCA. The precipitated protein was resuspended in Tris buffer and added to 10 ml of ScintiVerse scintillation cocktail for counting. The results obtained with Cfa1 and ACP further purified by Mono Q (MQ) chromatography (A) or not further purified (B) are shown. The percent loading values for Cfa1 and ACP are corrected to account for the amount of malonyl-CoA on FabD (where added) and for the percentage of holo protein as determined by MALDI-TOF. Assays were done in triplicate.

FLAG and ACP-FLAG concentrations of 10 to 100 μ M. To start the assay, 50 μ M [2-¹⁴C]malonyl-CoA (specific activity, 27.5 mCi/mmol) was added. The reaction mixture was incubated for 5 to 10 min at 25°C, and the reaction was stopped by the addition of 10% trichloroacetic acid (TCA) (400 μ l) and bovine serum albumin (10 mg/ml; 10 μ l). The pellet obtained after centrifugation was washed three times with 10% TCA. The final pellet was resuspended in Tris buffer (150 μ l) and added to 10 ml of aqueous scintillation cocktail for radioactivity determination. In additional experiments, the malonyl-CoA concentration was varied from 10 to 250 μ M and the concentration of Cfa1-FLAG was maintained at 20 μ M. Inaccurate data were obtained at malonyl-CoA concentrations of less than 10 μ M due to the low levels of radioactivity incorporated into protein. All kinetic parameters were measured in the linear range of the reaction for specific enzyme and substrate concentrations, and the experiments were carried out in triplicate. Incubations were also carried out in the absence of Cfa1 or ACP in order to monitor the ability of FabD to load malonyl-CoA. These reactions were treated in a similar manner to those described above.

The overproduced *Pseudomonas* FabD-FLAG was able to malonylate both the ACP-FLAG and Cfa1-FLAG proteins. When the assays were performed with ACP-FLAG and Cfa1-FLAG that were purified only by affinity chromatography, both proteins also exhibited an apparent ability to “self-load” malonate (Fig. 3B). However, when Cfa1 and ACP were further purified by Mono Q chromatography, they no longer exhibited self-malonylation activity (Fig. 3A). Curiously, the addition of FabD-FLAG to the impure Cfa1-FLAG or ACP-FLAG appeared to decrease the rate of self-loading. This behavior can be explained if the Cfa1-FLAG and ACP-FLAG were contaminated with a small amount of native FabD that possesses a specific activity that is significantly greater than that of the overproduced proteins, whose specific activity could be reduced by the presence of the FLAG tag. An analysis of the

reaction between FabD-FLAG and excess malonyl-CoA in the absence of Cfa1-FLAG or ACP-FLAG showed that FabD-FLAG was about 25% loaded with malonate after 10 min. Determination of kinetic parameters for the FabD-FLAG-catalyzed malonylation of Cfa1-FLAG revealed K_m values of <10 and $22.05 \pm 1.10 \mu\text{M}$ for malonyl-CoA and holo-Cfa1, respectively, and K_{cat} values of 0.03 ± 0.002 and $0.08 \pm 0.004 \text{ min}^{-1}$ for malonyl-CoA and holo-Cfa1, respectively. The only kinetic data available for comparison are those for malonylation of TcmM by FabD of *S. glaucescens*. The TcmM malonylation reaction was significantly faster than that observed for Cfa1 (8). Kinetic data for the FabD-catalyzed malonylation of other PKS ACPs will be needed before conclusions can be drawn from these differences.

The type II ACPs are small, acidic proteins that are post-translationally converted from the enzymatically inactive apo form of the protein to the active holo form by transfer of a 4'-phosphopantetheine moiety from CoA to a conserved serine residue (37). The enzymes that catalyze this transfer, known as PPTases, have been shown to constitute a superfamily (16). Three PPTases are known to be present in *E. coli* (31), while recent investigations have indicated that *P. aeruginosa* contains a single PPTase (PcpS) that modifies both fatty acid ACPs and the peptidyl carrier protein domain associated with nonribosomal peptide synthetases (7). In previous investigations where type II ACPs from *Streptomyces* were produced in *E. coli*, the apo ACP was the predominant form of the protein observed (5, 6, 32). Interestingly, some FAS ACPs from actinomycetes are produced in *E. coli* largely in the holo form (29, 30). When Cfa1 was overproduced in *E. coli* in the present study, the protein was recovered primarily in its apo form. However, when Cfa1 was overproduced in *P. syringae* FF5 at 18°C, the protein was recovered predominantly in the holo form. Similar results were obtained when Cfa1 was overproduced in *P. syringae* PG4180 at 18°C, while overproduction at 28°C in the latter strain yielded protein that was about a 2:1 mixture of apo to holo protein. The reasons for the decrease in the amount of holo Cfa1 at the higher temperature are unclear. However, several studies suggest that COR production is thermoregulated at both the transcriptional and posttranslational levels (3, 33). Collectively, these results suggest that a suitable PPTase for Cfa1 exists in *P. syringae*, whereas Cfa1 is a poor substrate for the *E. coli* PPTases. Thus, *E. coli* may not be a suitable host for the heterologous expression of enzymatically active *Pseudomonas* PKSs, unless a PPTase with broad specificity (7, 20, 31) is simultaneously coexpressed. Our results are consistent with reports that the peptidyl carrier protein from the pyoluteorin pathway of *Pseudomonas fluorescens* Pf-5 and the thiolation domains of nonribosomal peptide synthetase proteins from the pyochelin pathway of *P. aeruginosa* are not posttranslationally modified when these proteins are overproduced in *E. coli* (26, 36).

Type II PKSs generally lack a malonyl CoA:ACP transacylase to catalyze the transfer of a malonyl extender unit onto the PKS ACP (8), although the aromatic polyketide auricin is an exception to this generalization (22). For this reason, the potential role of the FAS malonyl CoA:ACP transacylase (FabD) in the biosynthesis of type II polyketides has been examined by several research groups. For example, the FAS FabDs of *S. glaucescens* and *S. coelicolor* can catalyze the transfer of a

malonyl group to the ACP components of the tetracenomyacin and actinorhodin PKSs, respectively (8, 28, 34, 39). In our experiments, the *P. syringae* FabD catalyzed the malonylation of Cfa1, which provides evidence that FabD can function in CFA biosynthesis. Some controversy concerning the role of FabD in type II polyketide biosynthesis has arisen because many PKS ACPs appear to self-malonylate (11, 18, 39). This dispute was recently resolved for TcmM, an ACP in *S. glaucescens*, by a rigorous set of experiments showing that the self-malonylation is an artifact of the expression and purification protocols (8). In the case of Cfa1, the affinity-purified protein isolated from *E. coli* also exhibited significant self-malonylation activity (Fig. 3B), even though the appearance of the protein on SDS-PAGE suggested a high degree of purity (Fig. 2). Additional purification of the affinity-purified Cfa1 by anion-exchange chromatography eliminated the self-malonylation activity (Fig. 3A), which is consistent with results obtained for TcmM. Our study thus provides additional evidence that the self-malonylation of ACPs is an artifact and further supports a role for FabD in CFA biosynthesis.

Nucleotide sequence accession number. The nucleotide sequences for *fabD* and *acp* of *P. syringae* FF5 have been deposited in the GenBank database under accession numbers AY391840 and AY391841.

We acknowledge support from the National Science Foundation (MCB 9807774 to R.P. and C.L.B. and IBN-0130693 to C.L.B.), the Oklahoma Center for the Advancement of Science and Technology (AR031-005 to C.L.B.), and The Robert A. Welch Foundation (C-0729 to R.P.).

Preliminary sequence data were obtained from The Institute for Genomic Research website at <http://www.tigr.org>. Sequencing of *P. syringae* pv. tomato DC3000 at The Institute for Genomic Research was accomplished with support from the National Science Foundation Plant Genome Program.

REFERENCES

- Bender, C. L., F. Alarcon-Chaidez, and D. C. Gross. 1999. *Pseudomonas syringae* phytotoxins: mode of action, regulation, and biosynthesis by peptide and polyketide synthetases. *Microbiol. Mol. Biol. Rev.* **63**:266–292.
- Bender, C. L., S. A. Young, and R. E. Mitchell. 1991. Conservation of plasmid DNA sequences in coronatine-producing pathovars of *Pseudomonas syringae*. *Appl. Environ. Microbiol.* **57**:993–999.
- Budde, I. P., B. H. Rohde, C. L. Bender, and M. S. Ullrich. 1998. Growth phase and temperature influence promoter activity, transcript abundance, and protein stability during biosynthesis of the *Pseudomonas syringae* phytotoxin coronatine. *J. Bacteriol.* **180**:1360–1367.
- Couch, R., H. Seidle, and R. J. Parry. 2002. Construction of expression vectors to produce affinity-tagged proteins in *Pseudomonas*. *BioTechniques* **32**:1230–1236.
- Cox, R. J., T. S. Hitchman, K. J. Byrom, I. S. C. Findlow, J. A. Tanner, J. Crosby, and T. J. Simpson. 1997. Post-translational modification of heterologously expressed *Streptomyces* type II polyketide synthase acyl carrier proteins. *FEBS Lett.* **405**:267–272.
- Crosby, J., D. H. Sherman, M. J. Bibb, W. P. Revill, D. A. Hopwood, and T. J. Simpson. 1995. Polyketide synthase acyl carrier proteins from *Streptomyces*: expression in *Escherichia coli*, purification and partial characterisation. *Biochim. Biophys. Acta* **1251**:32–42.
- Finking, R., J. Solsbacher, D. Konz, M. Schobert, A. Schafer, D. Jahn, and M. A. Marahiel. 2002. Characterization of a new type of phosphopantetheinyl transferase for fatty acid and siderophore synthesis in *Pseudomonas aeruginosa*. *J. Biol. Chem.* **277**:50293–50302.
- Florova, G., G. Kazanina, and K. A. Reynolds. 2002. Enzymes involved in fatty acid and polyketide biosynthesis in *Streptomyces glaucescens*: role of FabH and FabD and their acyl carrier protein specificity. *Biochemistry* **41**:10462–10471.
- Garde, S., and C. L. Bender. 1991. DNA probes for detection of copper resistance genes in *Xanthomonas campestris* pv. vesicatoria. *Appl. Environ. Microbiol.* **57**:2435–2439.
- Han, L., S. Lobo, and K. A. Reynolds. 1998. Characterization of beta-

- ketoacyl-acyl carrier protein synthase III from *Streptomyces glaucescens* and its role in initiation of fatty acid biosynthesis. *J. Bacteriol.* **180**:4481–4486.
11. Hitchman, T. S., J. Crosby, K. J. Byrom, R. J. Cox, and T. J. Simpson. 1998. Catalytic self-acylation of type II polyketide synthase acyl carrier proteins. *Chem. Biol.* **5**:35–47.
 12. Jiralerspong, S., V. Rangaswamy, C. L. Bender, and R. J. Parry. 2001. Analysis of the enzymatic domains in the modular portion of the coronafacic acid polyketide synthase. *Gene* **270**:191–200.
 13. Keen, N. T., S. Tamaki, D. Kobayashi, and D. Trollinger. 1988. Improved broad-host-range plasmids for DNA cloning in gram-negative bacteria. *Gene* **70**:191–197.
 14. King, E. O., M. K. Ward, and D. E. Raney. 1954. Two simple media for the demonstration of pyocyanin and fluorescein. *J. Lab. Clin. Med.* **44**:301–307.
 15. Kutchma, A. J., T. T. Hoang, and H. P. Schweizer. 1999. Characterization of a *Pseudomonas aeruginosa* fatty acid biosynthetic gene cluster: purification of acyl carrier protein (ACP) and malonyl-coenzyme A:ACP transacylase (FabD). *J. Bacteriol.* **181**:5498–5504.
 16. Lambolt, R. H., A. M. Gehring, R. S. Flugel, P. Zuber, M. LaCelle, M. A. Marahiel, R. Reid, C. Khosla, and C. T. Walsh. 1996. A new enzyme superfamily—the phosphopantetheinyl transferases. *Chem. Biol.* **3**:923–936.
 17. Liyanage, H., D. A. Palmer, M. Ullrich, and C. L. Bender. 1995. Characterization and transcriptional analysis of the gene cluster for coronafacic acid, the polyketide component of the phytotoxin coronatine. *Appl. Environ. Microbiol.* **61**:3843–3848.
 18. Matharu, A. L., R. J. Cox, J. Crosby, K. J. Byrom, and T. J. Simpson. 1998. MCAT is not required for in vitro polyketide synthesis in a minimal actinorhodin polyketide synthase from *Streptomyces coelicolor*. *Chem. Biol.* **5**:699–711.
 19. Mitchell, R. E., H. Young, and M. J. Liddell. 1995. Isolation and structural characterization of 2-[1-oxo-2-cyclopenten-2-ylmethyl]-butanoic acid, a polyketide product of coronatine-producing *Pseudomonas* sp. *Tetrahedron Lett.* **36**:3237–3240.
 20. Mootz, H. D., R. Finking, and M. A. Marahiel. 2001. 4'-Phosphopantetheine transfer in primary and secondary metabolism of *Bacillus subtilis*. *J. Biol. Chem.* **276**:37289–37298.
 21. Morbidoni, H. R., D. de Mendoza, and J. E. Cronan, Jr. 1996. *Bacillus subtilis* acyl carrier protein is encoded in a cluster of lipid biosynthesis genes. *J. Bacteriol.* **178**:4794–4800.
 22. Novakova, R., J. Bistakova, D. Homerova, B. Rezuchova, and J. Kormanec. 2002. Cloning and characterization of a polyketide synthase gene cluster involved in biosynthesis of a proposed angucycline-like polyketide auricin in *Streptomyces aureofaciens* CCM 3239. *Gene* **297**:197–208.
 23. Parry, R. J., S. Jiralerspong, S. V. Mhaskar, L. Alemany, and R. Willcott. 1996. Investigations of coronatine biosynthesis. Elucidation of the mode of incorporation of pyruvate into coronafacic acid. *J. Am. Chem. Soc.* **118**:703–704.
 24. Penalzo-Vazquez, A., and C. L. Bender. 1998. Characterization of CorR, a transcriptional activator which is required for biosynthesis of the phytotoxin coronatine. *J. Bacteriol.* **180**:6252–6259.
 25. Penfold, C. N., C. L. Bender, and J. G. Turner. 1996. Characterization of genes involved in biosynthesis of coronafacic acid, the polyketide component of the phytotoxin coronatine. *Gene* **183**:167–173.
 26. Quadri, L. E., T. A. Keating, H. M. Patel, and C. T. Walsh. 1999. Assembly of the *Pseudomonas aeruginosa* nonribosomal peptide siderophore pyochelin: in vitro reconstitution of aryl-4,2-bisthiazoline synthetase activity from PchD, PchE, and PchF. *Biochemistry* **38**:14941–14954.
 27. Rangaswamy, V., S. Jiralerspong, R. Parry, and C. L. Bender. 1998. Biosynthesis of the *Pseudomonas* polyketide coronafacic acid requires monofunctional and multifunctional polyketide synthase proteins. *Proc. Natl. Acad. Sci. USA* **95**:15469–15474.
 28. Revill, W. P., M. J. Bibb, and D. A. Hopwood. 1995. Purification of a malonyltransferase from *Streptomyces coelicolor* A3(2) and analysis of its genetic determinant. *J. Bacteriol.* **177**:3946–3952.
 29. Revill, W. P., M. J. Bibb, and D. A. Hopwood. 1996. Relationships between fatty acid and polyketide synthases from *Streptomyces coelicolor* A3(2): characterization of the fatty acid synthase acyl carrier protein. *J. Bacteriol.* **178**:5660–5667.
 30. Revill, W. P., and P. R. Leadlay. 1991. Cloning, characterization, and high-level expression in *Escherichia coli* of the *Saccharopolyspora erythraea* gene encoding an acyl carrier protein potentially involved in fatty acid biosynthesis. *J. Bacteriol.* **173**:4379–4385.
 31. Sanchez, C., L. Du, D. J. Edwards, M. D. Toney, and B. Shen. 2001. Cloning and characterization of a phosphopantetheinyl transferase from *Streptomyces verticillus* ATCC 15003, the producer of the hybrid peptide-polyketide antitumor drug bleomycin. *Chem. Biol.* **8**:725–738.
 32. Shen, B., R. G. Summers, H. Gramajo, M. J. Bibb, and C. R. Hutchinson. 1992. Purification and characterization of the acyl carrier protein of the *Streptomyces glaucescens* tetracenomycin C polyketide synthase. *J. Bacteriol.* **174**:3818–3821.
 33. Smirnova, A. V., L. Wang, B. Rohde, I. Budde, H. Weingart, and M. S. Ullrich. 2002. Control of temperature-responsive synthesis of the phytotoxin coronatine in *Pseudomonas syringae* by the unconventional two-component system CorRPS. *J. Mol. Microbiol. Biotechnol.* **4**:191–196.
 34. Summers, R. G., A. Ali, B. Shen, W. A. Wessel, and C. R. Hutchinson. 1995. Malonyl-coenzyme A:acyl carrier protein acyltransferase of *Streptomyces glaucescens*: a possible link between fatty acid and polyketide biosynthesis. *Biochemistry* **34**:9389–9402.
 35. Tao, T., and R. J. Parry. 2001. Determination by enantioselective synthesis of the absolute configuration of CPE, a potential intermediate in coronatine biosynthesis. *Org. Lett.* **3**:3045–3047.
 36. Thomas, M. G., M. D. Burkart, and C. T. Walsh. 2002. Conversion of L-proline to pyrrolyl-2-carboxyl-S-PCP during undecylprodigiosin and pyoluteorin biosynthesis. *Chem. Biol.* **9**:171–184.
 37. Walsh, C. T., A. M. Gehring, P. H. Weinreb, L. E. Quadri, and R. S. Flugel. 1997. Post-translational modification of polyketide and nonribosomal peptide synthases. *Curr. Opin. Chem. Biol.* **1**:309–315.
 38. Young, S. A., S. K. Park, C. Rodgers, R. E. Mitchell, and C. L. Bender. 1992. Physical and functional characterization of the gene cluster encoding the polyketide phytotoxin coronatine in *Pseudomonas syringae* pv. *glycinea*. *J. Bacteriol.* **174**:1837–1843.
 39. Zhou, P., G. Florova, and K. A. Reynolds. 1999. Polyketide synthase acyl carrier protein (ACP) as a substrate and a catalyst for malonyl ACP biosynthesis. *Chem. Biol.* **6**:577–584.