



Evaluation of the Antimicrobial Activity of Endophytic Bacterial Populations From Chinese Traditional Medicinal Plant Licorice and Characterization of the Bioactive Secondary Metabolites Produced by *Bacillus atrophaeus* Against *Verticillium dahliae*

OPEN ACCESS

Edited by:

Paula García-Fraile,
Academy of Sciences of the Czech
Republic (ASCR), Czechia

Reviewed by:

Learn-Han Lee,
Monash University Malaysia, Malaysia
Birinchi Kumar Sarma,
Banaras Hindu University, India

*Correspondence:

Osama A. A. Mohamad
dr.osamaabusaud@hotmail.com
Wen-Jun Li
liwenjun3@mail.sysu.edu.cn

† These authors have contributed
equally to this work.

Specialty section:

This article was submitted to
Microbial Symbioses,
a section of the journal
Frontiers in Microbiology

Received: 01 January 2018

Accepted: 20 April 2018

Published: 09 May 2018

Citation:

Mohamad OAA, Li L, Ma J-B,
Hatab S, Xu L, Guo J-W,
Rasulov BA, Liu Y-H, Hedlund BP
and Li W-J (2018) Evaluation of the
Antimicrobial Activity of Endophytic
Bacterial Populations From Chinese
Traditional Medicinal Plant Licorice
and Characterization of the Bioactive
Secondary Metabolites Produced by
Bacillus atrophaeus Against
Verticillium dahliae.
Front. Microbiol. 9:924.
doi: 10.3389/fmicb.2018.00924

Osama A. A. Mohamad^{1,2*†}, Li Li^{1,3†}, Jin-Biao Ma¹, Shaimaa Hatab⁴, Lin Xu⁵,
Jian-Wei Guo^{1,6}, Bakhtiyor A. Rasulov^{1,7}, Yong-Hong Liu¹, Brian P. Hedlund³ and
Wen-Jun Li^{1,8*}

¹ Key Laboratory of Biogeography and Bioresource in Arid Land, Xinjiang Institute of Ecology and Geography, Chinese Academy of Sciences, Urumqi, China, ² Environmental Science Department, Institute of Environmental Studies, Arish University, El-Arish, Egypt, ³ School of Life Sciences, University of Nevada, Las Vegas, Las Vegas, NV, United States, ⁴ Department of Food Science and Technology, College of Environmental Agricultural Sciences, Arish University, El-Arish, Egypt, ⁵ Key Laboratory of Hexi Corridor Resources Utilization, Hexi University, Zhangye, China, ⁶ Key Laboratory of Crops with High Quality and Efficient Cultivation and Security Control, Yunnan Higher Education Institutions, Honghe University, Mengzi, China, ⁷ Institute of Genetics and Plant Experimental Biology, Uzbekistan Academy of Sciences, Tashkent, Uzbekistan, ⁸ State Key Laboratory of Biocontrol and Guangdong Provincial Key Laboratory of Plant Resources, School of Life Sciences, Sun Yat-sen University, Guangzhou, China

Endophytic bacteria associated with medicinal plants possess unique strategies that enhance growth and survival of host plants, many of which are mediated by distinctive secondary metabolites. These bacteria and their secondary metabolites are important subjects for both basic and applied research aimed at sustainable agriculture. In the present study, 114 endophytic strains isolated from the wild ethnomedicinal plant *Glycyrrhiza uralensis* (licorice) were screened for their *in vitro* antimicrobial activities against common fungal pathogens of tomato (*Fusarium oxysporum* f. sp., *Fulvia fulva*, *Alternaria solani*), cotton (*Fusarium oxysporum* f. sp. *Vesinfectum*, *Verticillium dahliae*), pomegranate (*Ceratocystis fimbriata*), *Cymbidium* (*Colletotrichum gloeosporioides*), and Tsao-ko (*Pestalotiopsis microspora* and *Fusarium graminearum*) and the common bacteria *Staphylococcus aureus*, *Bacillus cereus*, *Salmonella enteritidis*, and *Escherichia coli*. Several *Bacillus* strains, particularly *Bacillus atrophaeus* and *Bacillus mojavensis*, had a broad spectrum of antifungal and antibacterial activity. A total of 16 strains, selected based on broad antimicrobial activity, were shown to contain at least one putative secondary metabolite-encoding gene (i.e., polyketide synthase or non-ribosomal peptide synthetase) and/or one lytic enzyme (i.e., protease, cellulase, lipase, chitinase), which may be important mediators of antagonistic activity against pathogens. Five strains, representing *Bacillus atrophaeus* and *Bacillus mojavensis*, were selected for plant growth chamber experiments based on strong *in vitro* antifungal activities.

All five strains significantly reduced disease severity in *Arabidopsis thaliana* plants challenged with *V. dahlia* infection. Gas-chromatography/mass-spectrometry analysis of cell-free extracts of *Bacillus atrophaeus* strain XEG150 showed that at least 13 compounds were produced only during co-cultivation with *V. dahlia*, including putative compounds known to have antimicrobial activity, such as 1,2-benzenedicarboxylic acid, bis (2-methylpropyl) ester; 9,12-octadecadienoic acid (Z,Z)-, methyl ester; 9-octadecenoic acid, methyl ester, (E)-; and decanedioic acid, bis(2-ethylhexyl) ester. To our knowledge, this study is the first to report that bacteria isolated from *G. uralensis* have biocontrol abilities. Our findings provide new insights into the antimicrobial activities of natural endophytes, particularly *B. atrophaeus*, and suggest this species may be a promising candidate as a biocontrol agent to confer resistance to *Verticillium* wilt disease and other phytopathogens in cotton and other crops.

Keywords: medicinal plants, endophytes, environmental microbiology, biological control, *Verticillium dahliae*, *Bacillus atrophaeus*, Licorice

INTRODUCTION

Fungal disease is the main threat to both crop yields and global food security (Fisher et al., 2012). Vascular wilts are devastating plant diseases that cause major losses to food crops and destroy natural ecosystems (Yadeta and Thomma, 2013). Two major genera of pathogenic fungi, *Fusarium* and *Verticillium*, enter their host plants through the roots or are transmitted by beetles and cause vascular wilts. Both are characterized by a wide host range (Juzwik et al., 2008; Harwood et al., 2011). The signs of *Verticillium* wilts are similar to those of *Fusarium*, starting with yellowing of the older leaves, followed by chlorosis and necrosis. As a result, vascular discoloration and stunting may be visible (Ting, 2014). In China, approximately 3 million hectares of cotton crops are affected by *Verticillium* wilt, accounting for an annual loss of yield of 10–30% (Bibi et al., 2013). In addition, no fungicides are registered for controlling *Verticillium* wilt disease in cotton (Göre et al., 2009). Xinjiang Province, located in the northwest of China, produces 11% of the global cotton fiber yield and is disproportionately harmed by *Verticillium* wilt disease (Peng et al., 2016).

Controlling vascular wilt pathogens can be challenging due to the fact that there are no efficient approaches to treat infected plants. Chemical fungicides are costly, inefficient, and have adverse impacts on the environment and human health (Qing et al., 2015). Increasing concern about the environmental and human health impacts of traditional fungicides has spawned intense interest in the development of safer alternatives. Biological control, an eco-friendly alternative, is mediated by microbial antagonists that possess unique traits that enable them to inhibit the growth of fungal pathogens (Walker et al., 1998; Chernin and Chet, 2002; Berg and Hallmann, 2006). Several mechanisms are responsible for these antagonistic activities, including inhibition of pathogen growth via antibiotics, toxins, surface-active compounds (antibiosis), and extracellular digestive enzymes such as proteases, cellulases and chitinases (Chernin and Chet, 2002; de Souza et al., 2003). However, several studies of biocontrol agents have reported dissimilarities between the

antagonistic effects *in vitro* and the corresponding *in situ* efficacy (Berg et al., 2000). Therefore, additional research is needed to better understand the basis by which plant-associated bacteria suppress fungal diseases *in vivo* (Santhanam et al., 2014), not only for basic science purposes, but also for the development of improved biocontrol approaches to support sustainable eco-friendly crop production (Weller, 1988; Sturz et al., 2000).

Endophytic bacteria have the capability to systematically colonize plant tissues and establish a symbiotic relationship with the host, which makes them highly efficient biocontrol agents (Bakker et al., 2013). Several reports have investigated bacterial endophytes as possible biocontrol agents against diverse pathogenic fungi (Lacava et al., 2007; Erdogan and Benlioglu, 2010; Egamberdieva et al., 2017a,b). Recently, several studies have described bacterial endophytes with biological control activity on a number of crops as potential sources of antimicrobial metabolites. The host plants in these studies have included *Solanum trilobatum* (Bhuvaneshwari et al., 2013), *Nicotiana attenuata* (Santhanam et al., 2014), *Solanum trilobatum melongena*, and *Solanum torvum* (Achari and Ramesh, 2014).

In view of the importance of endophytes to plant health, and increased focus on traditional herbal remedies as alternatives to synthetic pharmaceuticals, recent studies have begun to probe the importance of endophytic bacteria to medicinal plants, particularly those growing in unusual or stressed environments (Strobel and Daisy, 2003; Vieira et al., 2011; Liu Y.-H. et al., 2016; Egamberdieva et al., 2017a; Liu et al., 2017). Many of these studies have fingered the genus *Bacillus*, due to its widespread abundance in different plants, its broad-spectrum antimicrobial activities (Zheng et al., 2013; Jiang et al., 2015; Gao et al., 2017), and its ability to form endospores that are highly resistant to abiotic stresses such as UV light, desiccation, and extremes of pH, salinity, and temperature (Horikoshi, 2008).

Bacteria associated with medicinal plants have rarely been explored with regard to antagonistic activity against plant pathogens (Bakker et al., 2013; Bhuvaneshwari et al., 2013; Egamberdieva et al., 2017a). Licorice (*Glycyrrhiza uralensis*) is a popular traditional Chinese medicine. Licorice contains bioactive

compounds such as phenolics, flavonoids, triterpene saponins, and coumarins (Zhang and Ye, 2009). According to Chinese medicine theory, licorice has many important pharmacological activities, including antimicrobial and antiviral activity, histamine inhibition, anti-inflammatory activity, detoxification, and antioxidant and antitumor activities (Liao et al., 2012). Despite the economic interest and broad pharmacological effects of the popular medicinal usage of *G. uralensis*, very little research has been conducted on its endophytes or the potential for its endophytes to be used as biocontrol agents (Asl and Hosseinzadeh, 2008). Therefore, the objectives of the present study were: (1) to screen a diverse collection of endophytic bacteria (18 genera and 34 species) isolated from wild populations of *G. uralensis* for activity against a variety of pathogens *in vitro*; (2) to evaluate selected isolates for their biological control efficiency against the vascular wilt pathogen *V. dahliae* *in vivo*; and (3) to identify prevalent volatile organic compounds (VOCs) produced by endophytes only in the presence of *V. dahliae*, which are likely to be among the effectors of the antimicrobial properties.

MATERIALS AND METHODS

Endophytic Bacteria and Culture Condition

A collection of 114 endophytes were previously isolated from wild populations of *G. uralensis* (Li et al., 2018) from three areas in Xinjiang province, representing 18 genera and 34 species. All isolates were submitted to NCBI GenBank under Accession Number (KY127308 – KY127422) and used in this study. Bacterial isolates were stored in 20% glycerol at the Key Laboratory of Biogeography and Bioresource in Arid Land, Xinjiang Institute of Ecology and Geography, Chinese Academy of Sciences under -20°C . Bacteria were routinely cultured on ISP₂ growth medium, and incubated at $28 \pm 2^{\circ}\text{C}$ for 48–72 h.

Antibacterial Activity

A modification of the agar disk diffusion method for detecting antagonism was used against the four common bacteria listed in **Table 1** (Nie et al., 2012). The common bacteria and endophytes were each pre-cultured overnight, and 5 mL^{-1} of each culture was centrifuged at $604 \times g$ for 5 min. The pellets were resuspended in sterile phosphate buffered saline (PBS) in a laminar air flow cabinet and density adjusted to 10^8 colony forming units (CFU) mL^{-1} by using Densicheck plus (BioMérieux, United States). A total of $200 \mu\text{L}$ of the common bacteria cell concentrate was inoculated and evenly spread by sterile cotton swaps onto the surface of the medium, and then four 5-mm-diameter pieces of sterile filter paper were placed on each corner of the petri dish. A total of $10 \mu\text{L}$ of each endophyte strain was then added dropwise to the filter paper. All plates were wrapped with parafilm, incubated at $37 \pm 2^{\circ}\text{C}$ for 24 h and observed for the inhibition of the common bacteria (Cho et al., 2007). Antibacterial activity was assessed by measuring the diameter of the clear zone of growth inhibition. An equivalent

TABLE 1 | Common bacteria used in this study.

Scientific Name	Strain	Gram reaction	Source
<i>Staphylococcus aureus</i> (SA)	10786	+	China Center of Industrial Culture Collection (CICC)
<i>Bacillus cereus</i> (BC)	10451	+	China Center of Industrial Culture Collection (CICC)
<i>Salmonella enteritidis</i> (SE)	10982	–	China Center of Industrial Culture Collection (CICC)
<i>Escherichia coli</i> (EC)	GUM1.705	–	Microbial Culture Collection Center of Guangdong (GIMCC)

volume of sterile phosphate buffered saline (PBS) instead of the endophytic bacteria was used as a negative control.

Antagonistic Assays of Antifungal Activities *in Vitro*

The antifungal activity of each bacterial endophyte was screened for antagonism against the pathogenic fungi in **Table 2** by the plate diffusion method (Sun et al., 2017). Bacterial strains were grown in ISP₂ medium at 28°C overnight. The fungal pathogens were grown on potato dextrose agar (PDA) plates. A 5-mm agar plug containing 6-day-old mycelial growth was placed at the center of a 9-cm modified culture PDA plate, which favors for the growth of both endophytes and the pathogen. A concentrate of each endophyte was placed onto the agar surface at 8 equidistant points, 2.5 cm from the plate periphery (Loqman et al., 2009). All plates were wrapped with parafilm and incubated at $28 \pm 2^{\circ}\text{C}$ for 3–5 days and observed for the inhibition of the pathogen. Plates with pathogenic fungi alone served as a control. The percentage of growth inhibition was calculated by measuring the diameter of the inhibition zone by using the following formula:

$$\text{Inhibition rate (\%)} = \frac{F_{cd} - T_{cd}}{F_{cd} - F_0} \times 100$$

where F_{cd} is the fungal colony diameter on the control PDA base plate, T_{cd} is the fungal colony diameter on the experimental PDA

TABLE 2 | Fungal pathogens used in this study.

Strain	Scientific name	Host Plant and disease	Source
F1	<i>Fusarium oxysporum</i> f. sp.	Tomato, <i>Fusarium</i> wilt	Xinjiang
F2	<i>Fulvia fulva</i> (Cooke) Cif.	Tomato, Leaf mildew	Xinjiang
F3	<i>Alternaria solani</i> Sorauer	Tomato, Early Blight	Xinjiang
F4	<i>Fusarium oxysporum</i> f. sp. Vesinfectum	Cotton, <i>Fusarium</i> wilt	Xinjiang
F5	<i>Verticillium dahliae</i> Kleb.	Cotton, <i>Verticillium</i> wilt	Xinjiang
F6	<i>Ceratocystis fimbriata</i>	Pomegranate, Pomegranate wilt	Yunnan
F7	<i>Colletotrichum gloeosporioides</i>	Cymbidium sinense, Anthracnose	Yunnan
F8	<i>Pestalotiopsis microspora</i>	Tsao-ko, pseudostem black spot disease	Yunnan
F9	<i>Fusarium graminearum</i>	Tsao-ko, Leaf spot disease	Yunnan

base plate, and F_0 is the diameter of the test fungus agar disks (5 mm) (Aeron et al., 2011). Each experiment was performed with three replicates, and the analysis was repeated to ensure consistency of the results.

Screening for Natural Product Biosynthetic Gene Clusters by PCR Method

A total of 16 strains were used for screening for natural product biosynthetic gene clusters by PCR. Three sets of degenerate primers targeting biosynthetic genes were used for PCR amplification: KSF (5'-GTSCCGTSSCRTGSSHYTCSA-3') and KSR (5'-CGCTCCATGGAYCCSCARCA-3'), targeting polyketide synthase (*PKS*)-I KS and methyl malonyl transferase domains (Kun et al., 2010); KSαF (5'-TSGCSTGCTTGGAYGCSATC-3') and KSαR (5'-TGGAANCCGCCGAABCCGCT-3'), targeting *PKS*-II KSα genes (Metsä-Ketelä et al., 1999); and A3F (5'-GCS TACSYSATSTACACSTCSGG-3') and A7R (5'-SASGTCVCCS GTSCGGTAS-3'), targeting non-ribosomal peptide synthetase (*NRPS*) genes. The reactions were performed in a (BIO-RAD C1000 Thermal Cycler) in a total volume 25 μ l consisting of 50 ng of genomic DNA, 10 pmol of each primer, 2.5 mM of each deoxynucleotide triphosphates, 1X PCR buffer, and 1 U of Taq DNA polymerase. Gradient PCR was performed under the following conditions: initial denaturation step at 95°C for 5 min, followed by 32 cycles of denaturation at 96°C for 1 min, annealing at 56, 62.1, and 52.5°C for *PKS*-I, *PKS*-II, and *NRPS* genes, respectively, for 1 min, and extension at 72°C for 2 min, with a final extension step at 72°C for 10 min. The amplified PCR products were analyzed by electrophoresis on 1% agarose gels with TAE buffer. A negative control without DNA template was included with each PCR.

Digestive Enzymes

A total of 16 isolates were used for screening their ability to produce digestive enzymes. Cellulase activity was assayed with modified DSMZ¹ medium 65 without CaCO₃ and supplemented with carboxymethyl cellulose (5 g L⁻¹; Sigma-Aldrich) in place of glucose. After incubation for 3–4 days at 28°C, plates were stained with a Congo red solution and destained with a NaCl solution (Li et al., 2018). A clear or lightly colored halo around the colonies indicated a positive reaction. Protease activity was assayed with YEM agar medium containing 5% (v/v) skim milk. After incubation for 3–4 days at 28°C, a clear halo around the bacterial colonies due to hydrolysis of milk indicated a positive reaction. Lipase enzyme activity was assayed with modified Sierra lipolysis agar supplemented with beef extract (3 g L⁻¹) and ferrous citrate (0.2 g L⁻¹). After autoclaving, 50 mL of Victoria Blue B solution (0.1 g 150 mL⁻¹) and 10 mL of Tween-80 was added to the medium. After 5–6 days incubation at 28°C, white calcium precipitates around the bacterial colonies indicated a positive reaction (Li et al., 2018).

Colloidal chitin was prepared from commercial chitin by the method of Agrawal and Kotasthane, 2012. Chitin was hydrolyzed

in concentrated HCl by stirring at 4°C overnight, followed by extraction of colloidal chitin in 200 mL of ice-cold 99% ethanol, neutralization at room temperature overnight, and centrifugation at 1677 \times g for 10 min at 4°C. The pellet was washed with sterile distilled water by centrifugation at 2415 \times g for 5 min at 4°C till the smell of alcohol was completely removed and the pH was 7. The colloidal chitin had a soft pasty consistency with 90% moisture and was stored at 4°C until further use. Chitinase detection medium composed of (L⁻¹) 0.3 g of MgSO₄·7H₂O, 3.0 g of (NH₄)₂SO₄, 2.0 g of KH₂PO₄, 1.0 g of citric acid monohydrate, 15 g of agar, 200 μ L of Tween-80, 4.5 g of colloidal chitin, and 0.15 g of bromocresol purple and then autoclaved at 121°C for 15 min. To test for chitinase activity, inoculated plates were incubated at 25 \pm 2°C and were observed for formation of a colored zone. For all the tests mentioned above, sterile nutrient agar was used as a control for bacterial growth. All experiments were performed twice with three replicates for each isolate.

Evaluation of Biocontrol Efficacy in Pot Experiments Under Greenhouse Conditions

Five strains, XEGI33, XEGI38, XEGI44, XEGI50, and XEGI78, were selected for greenhouse experiments based on antagonistic activity against *V. dahliae* *in vitro* and presence of at least one biosynthetic gene and at least one digestive enzyme. *Arabidopsis thaliana* was used as a model plant and *V. dahliae* was used as a model pathogen (Maldonado-González et al., 2015). *A. thaliana* seeds with uniform shape and size were surface-sterilized with 99% ethanol for 0.5 min, and then washed with sterile distilled water 5–6 times. The seeds were placed on MS plates at 25 \pm 2°C for 3–5 days. *Arabidopsis thaliana* seedlings with true stage leaves were then transplanted singly into pots of 8 cm diameter containing 60 g of sterilized soil. After 2 days of transplantation, plants were divided into five groups and each group was given different treatments. *V. dahliae* was cultivated in advance in Czapek-Dox broth at 28°C for 4 days. After 72 h of transplantation, 10 mL of the fungal mycelia suspension (10⁸ CFU mL⁻¹) was spread on the soil surface above fine roots by using a sterile syringe, and then after another 48 h, 10 mL of an endophytic bacteria suspension (10⁸ CFU mL⁻¹) was added, as generally described by Chen et al., 2014. Two controls were used in this experiment: seedlings treated with sterile water (CK⁺), and seedlings inoculated with the pathogen alone (CK⁻). The pots were placed in a growth chamber with the following conditions: 25–30°C, 60% humidity, and 16 h of daylight alternating with 8 h of darkness (Jiang et al., 2015). Three replicates were done for each treatment. Plant phenotypes were observed after 10 days of pathogen inoculation, while the development of wilt disease signs was recorded after 40 days.

Disease Assessment

A disease index, based on yellowing and chlorosis of cotyledons and leaves after 40 days, was used to classify disease signs for each leaf into six grades (i.e., grade 0, 1, 2, 3, 4, and 5) (Zhang et al., 2012), and then the percentage of affected leaves was calculated and categorized (\leq 25, 25–35, 35–45, 45–55, and

¹http://www.dsmz.de/microorganisms/medium/pdf/DSMZ_Medium65.pdf

55–80%). The final disease index (DI) was calculated according to the following formula: $DI = [(\sum \text{disease grades} \times \text{number of infected}) / (\text{total checked plants} \times 5)] \times 100$ (Zhang et al., 2012). The DI represents a comprehensive and objective measure of plant health, with high DI values corresponding to serious infections.

Isolation and Purification of Antimicrobial Agents

Strain XEGI50 was inoculated into 500 mL⁻¹ of ISP₂ broth at 28°C for 10 days with agitation at 120 rpm and used as a control (1). *V. dahliae* was cultivated in 500 mL⁻¹ in Czapek broth at 28°C for 10 days with agitation at 120 rpm and used as a control (2). The antibiosis experiment was carried out by co-cultivation of strain XEGI50 with *V. dahliae* in 500 mL⁻¹ of broth medium at 28°C for 12 days with agitation at 120 rpm. All cells were collected by centrifugation at 5000 × *g* for 10 min. The cell-free supernatant was divided into equal volumes. After that, the supernatant was adjusted to pH 7 and pH 3 with 500 mL⁻¹ of 1 N HCl and an equal volume (1:1) of ethyl acetate was added and mixed by vigorous shaking for 30 min and allowed to settle. The organic solvent phase was collected and evaporated at 40°C under vacuum, using a rotary evaporator model (IKA, HB10 basic). The ethyl acetate extract was dissolved in 5 mL of Tris-Cl buffer (0.02 M, pH 7.0) and used for gas-chromatography/mass-spectrometry (GC-MS).

Identification and Antibacterial Evaluation of Bioactive Compounds

The GC-MS analysis of the cell-free extracts was performed using a gas chromatograph (Model 7890A, Agilent, Palo Alto, CA, United States) equipped with a split-splitless injector, an Agilent model 7693 autosampler, and an Agilent HP-5MS fused silica column (5% phenyl-methylpolysiloxane, 30 m length, 0.25 mm I.D., film thickness 0.25 mm). Injecting volume was 1 μL, and the GC conditions included programmed heating from 50 to 300°C at 10°C/min, followed by 10 min at 300°C. The injector was maintained at 280°C. Helium was the carrier gas, at 1.0 mL min⁻¹, and the split mode was 5:1. The GC was fitted with a quadrupole mass spectrometer with an Agilent model 5975 detector. The MS conditions were as follows: ionization energy, 70 eV; electronic impact ion source temperature, 230°C; quadrupole temperature, 150°C; scan rate, 3.2 scan/s; mass range, 50–1000 μ. The compounds were identified based on the match with their mass spectra and retention indices with the NIST/Wiley 275 library (Wiley, New York, NY, United States). Relative abundance of each feature was calculated from Total Ion Chromatogram (TIC) computationally.

Intelligent Live Digital Imaging of Strain XEGI50 and *V. dahliae*

The morphological response of XEGI50 to *V. dahliae* was observed under a laser microscope (Olympus SZX2-ILLT, Japan) at different magnifications. Bacteria were incubated in ISP₂ medium. *V. dahliae* was grown on PDA medium. A 6-day-old mycelial disk (5 mm) was placed at the center of a 7 cm modified

culture PDA plate. The bacteria were placed at four corners on the bacterial lawn at four equidistant points of 2.5 cm from the plate periphery. All plates were wrapped with parafilm, incubated at 28 ± 2°C for 6 days, and observed for the inhibition of the pathogen. Plates with pathogenic fungi alone served as control.

RESULTS

Antibacterial Activity


A total of 114 endophytes isolated from wild populations of *G. uralensis*, representing 18 genera and 34 species, were screened for their ability to inhibit four common bacteria, representing the Gram-positive phylum *Firmicutes* and the Gram-negative phylum *Proteobacteria* (Table 1). Of the 114 isolates examined, 56 (49.1%) displayed inhibitory activity, ranging from 8.6 to 11.8 mm against *S. aureus*, *B. cereus*, and *S. enteritidis*; in contrast, only 14 (12.3%) were antagonistic to *E. coli*, ranging from 7.5 to 10.3 mm (Supplementary Figures S1A,B). These 14 strains belonged to 6 different genera, namely *Bacillus*, *Microbacterium*, *Brevibacterium*, *Phyllobacterium*, *Pantoea*, and *Stenotrophomonas*. The genus *Bacillus* showed the highest antimicrobial activity against the selected Gram-positive and Gram-negative bacteria (Figure 1A).

Antifungal Activity

The inhibitory effect of all endophytic isolates was tested against the nine fungal phytopathogens listed in Table 2. The endophytes varied in their ability to inhibit the growth of the fungi, with the percentage of inhibition ranging from 12.3 to 75.3% (Supplementary Figures S1C,D). A total of 44, 75, and 34 strains (38.6, 65.8, and 29.8%) were antagonistic to the tomato pathogens *F. oxysporum* f. sp. (F1), *F. fulva* (F2), and *A. solani* (F3), respectively. For *F. oxysporum* f. sp., the largest inhibition zone (72.1%) was observed for strain XEGI174, belonging to *Bacillus halotolerans*. For *F. fulva*, the largest inhibition zone (72.0%), was observed for *Bacillus atrophaeus* strain XEGI151. And for *A. solani*, the largest inhibition zone (63.0%) was observed for strain XEGI115, belonging to *Brevibacterium frigiditolerans*.

For the pathogens causing cotton wilt diseases, the antibiosis assay demonstrated that 48 and 83 endophytic strains (42.1 and 72.8%) were antagonistic to *F. oxysporum* f. sp. Vesinfectum (F4) and *V. dahliae* (F5), respectively. For *F. oxysporum* f. sp. Vesinfectum, the largest inhibition zone (70.1%) was observed for strain XEGI9, belonging to *Nocardioideis alkalitolerans*; for *V. dahliae*, the largest inhibition zone (75.5%) was observed for strain XEGI50, belonging to *B. atrophaeus*.

For the pathogens *Ceratocystis fimbriata*, *Colletotrichum gloeosporioides*, *Fusarium graminearum*, and the recently described pathogen *Pestalotiopsis microspora* (Guo J. W. et al., 2016), the antibiosis assay demonstrated that 60, 45, 57, and 40 strains (52.6, 39.5, 50.0, and 35.1%) were antagonistic to *C. fimbriata* (F6), *C. gloeosporioides* (F7), *P. microspora* (F8), and *F. graminearum* (F9), respectively. For *C. fimbriata*, the largest inhibition zone (55.4%) was observed for strain XEGI146, which belonged to *Bacillus mojavensis*. For *C. gloeosporioides*, the largest inhibition zone (48.0%) was observed for *B. atrophaeus* strain



XEGI10. For *P. microspora*, the largest inhibition zone (47.9%) was observed for strain XEGI44, belonging to *Bacillus mojavenis*; and for *F. graminearum*, the largest inhibition zone (34.0%) was observed for strain XEGI39, which belonged to *B. atrophaeus*. Based on our investigation, members of the genus *Bacillus* had the highest antagonistic activity of any of the bacterial genera (Figure 1B), and *V. dahliae* was the most susceptible fungal pathogen.

Analysis of Putative Biosynthetic Genes

The presence of putative biosynthetic genes encoding *PKS*s and *NRPS* of peptide antibiotics were investigated in the 16 strains exhibiting the strongest antimicrobial activities by using 3 sets of degenerate PCR primers. Among the 16 strains, 12 were positive for *PKS*s genes (Table 3); the translated amino acid sequences of these *PKS-I* genes shared moderate to high amino acid identity (54–99%) with those from members of the phylum *Actinobacteria*. A total of seven were positive for *NRPS* genes (Table 3). The *NRPS* sequences shared 40–70% amino acid sequence identities with peptide synthetase genes of the genus *Bacillus*, except the sequence from *Microbacterium paraoxydans* strain XEGI12, which was most closely related to enzymes of genus *Microbacterium*. Moreover, of the 16 isolates, 7 strains were positive for amplification of all 3 biosynthetic genes; 2 strains were positive for 2 genes; 4 strains were positive for only 1 gene; and only 3 strains were negative for all 3 *PKS* and *NRPS* genes (Table 3 and Supplementary Figure S2).

Digestive Enzyme Activity

The same 16 strains were assayed for protease, cellulase, lipase, and chitinase, which are potentially involved in lysis of phytopathogens or modulation of virulence factors. Some of the

tested strains produced one or more lytic enzyme, as assessed by a change in pH in plates containing soluble chitin and bromocresol purple (chitinase) or by the diameter of the halo zone on media containing skim milk (protease), carboxymethyl cellulose (cellulase), or Sierra lipolysis agar (lipase) (Table 3 and Supplementary Figure S3). Proteases and lipases were the most common enzymes detected. The *Bacillus* strains were more active in these assays than the other two genera tested.

Biological Control of *V. dahliae* in Growth Chambers

An infection time course of *A. thaliana* and *V. dahliae* was developed by evaluating plant disease signs over a 5-week period. The first signs developed within 7 days of inoculation, including yellowing of leaves, and leaf chlorosis, which began with older leaves and progressed to younger leaves. Severe signs were seen on the leaves of the plants challenged with *V. dahliae* in the absence of endophytes (Figure 2). After 5 weeks, the control plantlets under pathogen-challenged conditions showed severe signs, while those inoculated with endophytes showed mild signs of disease. Although all tested endophytes conferred some degree of *Verticillium* wilt resistance, the distribution of disease grades varied dramatically (Figure 3A). The disease severity index (DSI) in plantlets treated with strains XEGI33, XEGI38, XEGI44, XEGI50, and XEGI78 were each significantly reduced (44.5, 50.0, 44.5, 33.3, and 48.6%), compared to those grown in the absence of the biocontrol agent, which was 78.0% (Figure 3B). XEGI50 conferred the best protection (33.3% DSI) by suppressing yellowing other and wilt signs. This result suggested that XEGI50 could slow disease development, and the expression of signs were delayed compared with other treatments.

TABLE 3 | Presence of biosynthetic genes (*PKSI*, *PKSII*, and *NRPS*) and digestive enzyme activity of the 16 most active strains.

Strain	Accession Number	Closest species in 16S rRNA gene sequences database	<i>PKSI</i>	<i>PKS II</i>	<i>NRPS</i>	Protease ^a	Cellulase ^b	Lipase ^c	Chitinase
XEGI10	KY127316	<i>Bacillus atrophaeus</i>	+	+	+	+++	-	+	+
XEGI12	KY127318	<i>Microbacterium paraoxydans</i>	+	+	+	-	-	-	-
XEGI14	KY127320	<i>Bacillus atrophaeus</i>	+	+	+	++	-	++	-
XEGI15	KY127321	<i>Brevibacterium frigoritolerans</i>	-	-	-	-	-	++	-
XEGI16	KY127322	<i>Bacillus mojavensis</i>	+	-	-	++	+	+++	-
XEGI33	KY127337	<i>Bacillus atrophaeus</i>	+	+	+	++	-	-	+
XEGI35	KY127339	<i>Bacillus atrophaeus</i>	+	+	-	++	-	-	-
XEGI38	KY127341	<i>Bacillus atrophaeus</i>	+	+	-	+++	-	-	-
XEGI44	KY127347	<i>Bacillus mojavensis</i>	-	+	-	+++	-	+	-
XEGI45	KY127348	<i>Bacillus atrophaeus</i>	+	-	-	+	+	+	-
XEGI46	KY127349	<i>Bacillus mojavensis</i>	+	+	+	+	-	+	-
XEGI50	KY127353	<i>Bacillus atrophaeus</i>	+	+	+	+++	-	++	+
XEGI56	KY127359	<i>Bacillus mojavensis</i>	+	-	-	++	+	+	+
XEGI74	KY127370	<i>Bacillus halotolerans</i>	-	-	-	+	++	++	-
XEGI78	KY127374	<i>Bacillus atrophaeus</i>	+	+	+	++	-	+	-
XEGI95	KY127390	<i>Bacillus tequilensis</i>	-	-	-	+	+	+	-

^aProtease production: “-” no production; “+” weak halo around the colony (0.50–1.50 cm); “++” clear halo around the colony (1.50–2.00 cm); “+++” strong halo around the colony (2.00–3.00 cm). ^bCellulase production: “-” no production; “+” weak halo around the colony (1.00–1.50 cm); “++” clear halo around the colony (1.50–2.51 cm). ^cLipase production: “-” no production; “+” weak halo around the colony (1.00–1.50 cm); “++” clear halo around the colony (1.50–2.00 cm); “+++” strong halo around the colony (2.00–2.50 cm).

Detection of Bioactive Compounds by GC-MS Analysis

To determine the prevalent organic compounds produced by the most bioactive strain, *B. atrophaeus* XEGI50, ethyl acetate extracts of cell supernatant buffered at pH 7 and pH 3 were concentrated from cultures of strain XEGI50, *V. dahlia*, and a co-culture. GC-MS analysis showed that different features (putative compounds) were produced at pH 7 and pH 3. Features were tentatively identified based on comparison of spectra available through the National Institute of Standards and Technology (NIST) database, and biological activities were interpreted primarily based on Dr. Duke's Phytochemical and Ethnobotanical Databases created by Dr. Jim Duke of the Agricultural Research Service/USDA.

The GC-MS analysis of crude extracts buffered at pH 7 from strain XEGI50 cultivated alone revealed at least 21 features (Supplementary Table S1) and 36 features at pH 7 (Supplementary Table S2). Nine distinctive features in the pH 7 extract, compared to pH 3, were present at RT 3.897, 4.171, 17.302, 19.849, 20.701, 21.511, 22.301, 22.374, and 24.331, suggesting the presence of benzene, 1,3-dimethyl-; o-xylene; dibutyl phthalate; heptadecane; eicosane; tetracosane; pentacosane; bis (2-ethylhexyl) phthalate; and decanedioic acid, bis (2-ethylhexyl) ester (Figure 4A). The pH 3 extract showed three major peaks that were different from the pH 7 extract at RT 3.908, 18.954, and 19.859, suggestive of p-xylene; heneicosane; and docosane (Figure 4B).

The GC-MS analysis of crude extracts from *V. dahlia* cultivated alone showed 32 compounds in the pH 7 extract





FIGURE 2 | Defense response to *V. dahliae* in *Arabidopsis thaliana* plants (A) *A. thaliana* seedling; (B) *A. thaliana* without inoculation of *V. dahliae*; (C) Response of *A. thaliana* to *V. dahliae*; (D) Response of *A. thaliana* to *V. dahliae* inoculated with different endophytic strains after 5 weeks.

(Supplementary Table S3) and 17 compounds in the pH 3 extract (Supplementary Table S4). Six major features were obtained from the pH 7 extract at RT 3.802, 3.918, 4.181, 17.302, 20.711,



and 22.384, suggestive of ethylbenzene; p-xylene; benzene 1,3-dimethyl-; dibutyl phthalate; heptadecane; and bis (2-ethylhexyl) phthalate (Figure 4C). The pH 3-buffered extract from *V. dahliae* showed one major peak are different from the crude extract at pH 7 at RT 22.374, suggestive of phthalic acid, di (2-propylpentyl) ester (Figure 4D).

The GC-MS resolved several features in the extracts of XEGI50 and *V. dahliae* mixture: 37 compounds from the pH 7 extract (Supplementary Table S5) and 32 compounds from pH 3 extract (Supplementary Table S6). A total of 12 major peaks were obtained from the pH 7 extract at RT 3.813, 3.929, 4.192, 11.568, 17.397, 18.712, 18.796, 19.848, 20.711, 21.521, 22.426, and 24.352, suggestive of ethylbenzene; p-xylene; dimethyl phthalate; 1,2-benzenedicarboxylic acid; bis (2-methyl propyl) ester; 9,12-octadecadienoic acid (Z,Z)- methyl ester; 9-octadecenoic acidmethyl ester, (E)-; eicosane; heptadecane; tetracosane; bis (2-ethylhexyl) phthalate; and decanedioic acid, bis (2-ethylhexyl) ester (Figure 4E). The pH 3 extract from the co-culture contained most of the same compounds, and one additional feature at RT 4.192, suggestive of o-xylene (Figure 4F).

Many compounds tentatively identified in the GC-MS analysis are known to have antimicrobial activity. Therefore, to determine the prevalent compounds that were induced by co-cultivation, putative compounds were compared between the three different datasets. At least 13 compounds were detected only in the co-culture, consisting of mainly fatty acid esters, phenols, alkanes, alkenes, and aromatic chemicals. Four of them showed high peaks at RT 17.397, 18.712, 18.796, and 24.35, suggestive of 1,2-benzenedicarboxylic acid butyl 2-ethylhexyl ester; 9,12-octadecadienoic acid (Z,Z)-, methyl ester; 9-octadecenoic acid, methyl ester, (E)-; and decanedioic acid, bis (2-ethylhexyl) ester (Figures 4E,F). In addition, several minor peaks were present, including butanoic acid, 2-hydroxy-3-methyl-; eicosane9-cyclohexyl-, heptadecanoic acid, 16-methyl-, methyl ester; hexadecanoic acid, methyl ester; hexanedioic acid,


dioctyl ester; naphthalene, 1-methyl-; naphthalene, 2-methyl-; n-decanoic acid; pentanoic acid, 4-methyl-; and picolinamide (Supplementary Tables S5, S6).

Defense Response of Strain XEGI50 to *V. dahlia* via Laser Microscopy

The results showed that *V. dahliae* could not grow after 7 days of incubation with antagonistic strains (Figures 5Aa,b). The morphological response of XEGI50 to *V. dahliae* was observed under a laser microscope at different magnifications (1.25 \times , 2.5 \times , 4 \times). The microscopic characteristics of strains XEGI50 based on laser microscopy showed that at 1.25 \times magnification the endophytic strain XEGI50 was able to control the growth of the fungal mycelium after 5 days of cultivation (Figure 5B) and at 4 \times , a white powder appeared only on the side facing the pathogenic fungi, we hypothesize that strain XEGI50 may secrete some antifungal compound (Figure 5C). In addition, at the same magnification of 2.5 \times , strain XEGI50 looked intense at the bacteria/fungi interface, but expanded in the direction away from *V. dahliae* (Figure 5D).

DISCUSSION

Biological control of plant pathogens using microorganisms can be a safe, cost-effective, and efficient method for suppressing plant diseases. Endophytes associated with medicinal plants are rich sources of secondary metabolites with antimicrobial activity, and they spend their whole life cycle within plant tissues without causing any infections or signs of disease (Bacon and White, 2000; Saikkonen et al., 2004). In addition, it has also been documented that endophytes associated with medicinal plants may produce the same metabolites *in vitro* and within host plant tissue (Kusari et al., 2013; Dos Santos et al., 2016). In the present study, we analyzed the antimicrobial activity of a



mycelial growth of diverse vascular wilts caused by pathogenic fungi (Cho et al., 2002; Ebrahimi et al., 2010; Jiang et al., 2015; Akinsanya et al., 2015; Gao et al., 2017).

In our study, 16 strains showing antagonistic activity toward at least 7 pathogenic fungi and at least 1 common bacterium, were screened for the presence of *PKS-I*, *PKS-II*, and *NRPS* gene clusters to determine whether their genomes contain these genes. These 16 strains consisted of 1 strain each of *Microbacterium paraoxydans* and *B. frigoritolerans*, and 14 strains of *Bacillus*, belonging to *Bacillus atrophaeus* (8 strains), *Bacillus mojavensis* (four strains), *Bacillus halotolerans* (1 strain), and *Bacillus tequilensis* (1 strain). Only 12, 10, and 7 strains were positive for PCR amplification of fragments of *PKS-I*, *PKS-II*, and *NRPS* genes, respectively (Table 3). Several isolates had antimicrobial activity, but biosynthetic genes were not successfully amplified; they were *B. frigoritolerans* XEGI15, *Bacillus halotolerans* XEGI74, and *Bacillus tequilensis* XEGI95. The absence of amplification products may be due to the lack of natural product biosynthetic genes or because they contain divergent or novel genes that are not recognized by the degenerate primers used in this study (Courtois et al., 2003; Finking and Marahiel, 2004). Moreover, not all *NRPS* genes are involved in the biosynthesis of bioactive secondary metabolites, and there might be additional types of bioactive agents or mechanisms that may be involved in the generation of antimicrobial activities (Schneemann et al., 2010; Liu L. et al., 2016).

The 16 most antagonistic strains each produced at least 1 digestive enzyme, such as protease, cellulase, lipase, and chitinase (Table 3). Thus, these endophytes may protect the plant from fungi and insects by degrading the fungal cell

wall or cell membrane, by degrading cell membrane proteins or extracellular virulence factors, or by stimulating systemic resistance in plants (Frankowski et al., 2001). In our previous investigation, endophytic bacteria isolated from the medicinal plants *Ferula songorica*, *Hypericum perforatum*, and *Ferula sinkiangensis* secreted similar digestive enzymes (Liu Y.-H. et al., 2016; Liu et al., 2017). Moreover, similar work done by Egamberdieva et al. (2017a) reported that endophytic bacteria associated with the medicinal plant *Ziziphora capital* were able to produce chitinolytic enzymes.

Verticillium dahliae, which infects cotton and several other plants, causes wilt diseases and crop losses of varying severity as well as natural ecosystems (Erdogan and Benlioglu, 2010). The signs of *Verticillium* wilt disease start with yellowing followed by chlorosis and necrosis of leaves (Ting, 2014). *Arabidopsis thaliana* is an excellent tool to identify traits involved in *V. dahliae* biocontrol by endophytic bacteria (Maldonado-González et al., 2015). In the current study, five endophytic bacterial strains, XEGI33, XEGI38, XEGI44, XEGI50, and XEGI78, representing *B. atrophaeus* and *B. mojavensis*, were selected for growth chamber experiments to suppress *V. dahliae* pathogenesis in *A. thaliana*. These five strains were chosen based on *in vitro* effects against a large number of pathogens. All five strains seemed to colonize *A. thaliana* and significantly reduced the DSI (Figure 3). In accordance with these results, several previous studies have shown that endophytic *Bacillus* species control fungal pathogens, including *B. mojavensis*, *B. subtilis* (Bacon and Hinton, 2002; Bacon et al., 2005; Cazorla et al., 2007), *B. tequilensis*, *B. velezensis*, *B. amyloliquefaciens* (Alvarez et al., 2012; Akinsanya et al., 2015; Gao et al., 2017), and *B. megaterium*, (Cho et al., 2002; Lin et al., 2013). In addition, Guo Y. et al. (2016) isolated and characterized *B. atrophaeus* strain OSY-7LA and showed that it exhibited a strong antagonistic activity against *Listeria innocua*, a food-borne pathogen that can survive at extreme pH, temperature, and high salt concentration.

Among the endophytes, *B. atrophaeus* XEGI50 was selected for exometabolomic studies by GC-MS, based on its ability to decrease the DSI and suppress the growth of *V. dahliae*. A total of 13 features that were expressed only in co-cultures of XEGI50 and *V. dahliae* were tentatively identified as compounds with known antimicrobial, antiprastic, antitumor, and anticancer properties. Among these compounds, four of them were major peaks in cell-free extracts from the co-culture, suggesting they play an important role in antimicrobial activities; these compounds were 1,2-benzenedicarboxylic acid, butyl 2-ethylhexyl ester (Kavitha et al., 2010); 9,12-octadecadienoic acid (*Z,Z*-), methyl ester (Sermakkani and Thangapandian, 2012); 9-octadecenoic acid, methyl ester, (*E*-); and decanedioic acid, bis (2-ethylhexyl) ester (Tambekar et al., 2014). In addition, several putative antimicrobial compounds were identified as minor peaks: eicosane, 9-cyclohexyl (Hsouana et al., 2011); heptadecanoic acid, 16-methyl-, methyl ester (Kandasamy et al., 2012); hexadecanoic acid, methyl ester (Chandrasekaran et al., 2011); hexanedioic acid, dioctyl ester (Rodríguez-Meizoso et al., 2010); naphthalene, 1-methyl; naphthalene, 2-methyl- (Rokade and Sayyed, 2009); and pentanoic acid, 4-methyl- (Sharma et al., 2016). Since these compounds were not produced by pure cultures of XEGI50,

they were likely induced by the presence of fungal pathogens such as *V. dahliae*, and very likely play a role in antagonism of pathogens.

The genus *Bacillus* is well known for the natural production of secondary metabolites with antibacterial and antifungal activities and has a strong potential to control plant diseases (Lodewyckx et al., 2002; Cavaglieri et al., 2005; Radhakrishnan et al., 2017). This study further illustrates its potential role as a biological agent for controlling phytopathogens. In recent years, the development of biological agents derived from *Bacillus* isolates, such as “Avogreen” (Korsten et al., 1997; Janisiewicz and Korsten, 2002) and “Shemer” (Droby, 2005), has been shown to be effective biocontrols against some plant diseases. In the present study, we provide insights about plant beneficial traits of culturable endophytic bacteria associated with the medicinal plant *G. uralensis*. Our results may provide a new biological control agent for controlling *V. dahliae* and improve our understanding of the biocontrol mechanism of natural endophytes belonging to the genus *Bacillus*. These results support the development of natural products that may minimize the need for the application of chemical fungicides, which would be an environmentally friendly approach and preserve biological resources in a sustainable agricultural system.

CONCLUSION

Our study revealed that natural endophytes of natural populations of the medicinal plant *G. uralensis* have a variety of antimicrobial activities *in vitro* and *in vivo*. The genus *Bacillus*, particularly *B. atrophaeus* and *B. mojavensis*, were the most effective biocontrol agents, with most strains exhibiting broad antibacterial and antifungal activities. Most of these bacteria contained genes for *PKS* and non-ribosomal proteins, both known to encode antimicrobial compounds, as well as extracellular digestive enzymes that may destroy or neutralize a variety of pathogens, including chitinases, cellulases, lipases, and proteases. Strain XEGI50, which belongs to *Bacillus atrophaeus*, was the most effective at reducing disease signs in *A. thaliana* in plant growth chambers. XEGI50 produces at least 13 compounds when co-cultivated with *V. dahlia*, many of which were putatively identified as compounds with known antimicrobial effects. To our knowledge, this is the first report establishing that *B. atrophaeus* produces bioactive compounds with antimicrobial activity. Future studies are needed to unequivocally identify these compounds, to establish their effects individually in model plant systems, and to better understand genetic and biochemical pathways for synthesis of these compounds.

AUTHOR CONTRIBUTIONS

OAAM, LL and W-JL participated in the design of the study, performed all the experiments and the interpretation of results, and wrote the manuscript. J-BM and SH participated in the

antimicrobial experiments *in vitro* condition and provided the pathogenic bacteria, and also conducted the greenhouse experiments and data analysis. LX and J-WG conducted to antifungal activity *in vitro* condition and also provided some of the pathogenic fungi. BR did the GC-MS analyses and data analysis. Y-HL helped for preparing enzymes activity test, microscopic analysis, and PCR works. BH revised the revision and improve the language and structure of the manuscript. W-JL and OAAM revised the manuscript and supervised the hall experiments. All authors edited and critically revised the manuscript.

FUNDING

This research was supported by the National Key Research and Development Program of China (2017YFD0200503), National Natural Science Foundation of China (Grant No. 31650110479), Xinjiang Uygur Autonomous Region Regional Coordinated Innovation Project (Shanghai Cooperation Organization Science and Technology Partnership Program) (Grant No. 2017E01031), and West Light Foundation of the Chinese Academy of Sciences (Grant Nos. XBBS201305 and XBBS201201). W-JL was also supported by Guangdong Province Higher Vocational Colleges and Schools Pearl River Scholar Funded Scheme (2014). LL was supported by China Scholarship Council to study in the United States (Grant No. 201509655013). OAAM was supported by Available Position Talented Young Scientists Program of Ministry of Science and Technology of China (Grant No. P-EG-16-01), and also funded by Chinese Academy of Sciences President's International Fellowship Initiative (Grant No. 2016PB024).

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fmicb.2018.00924/full#supplementary-material>

FIGURE S1 | *In vitro* evaluation of antagonistic activity of endophytic bacterial isolates associated with *G. uralensis*. (A) *Bacillus cereus* (BC); (B) *Salmonella enteritidis* (SE); (C) *Verticillium dahliae*; (D) *Fusarium oxysporum*.

FIGURE S2 | Agarose gel showing PCR results for *PKS-I*, *PKS-II*, and *NRPS* from selected strains. (M: Marker 1,500 bp).

FIGURE S3 | Ability of endophytes strains to produce lytic enzymes. (A) breakdown of chitin into N-acetyl glucosamine causes an increase in pH and a change from yellow to purple zone around bacterial colony; (B) control; (C) clear zone formation in cellulose medium around bacterial colony; (D) Clear zone formation in proteases medium around bacterial colony.

TABLE S1 | GC-MS identified components of the crude extract of XEGI50 at pH7 (Volatile compounds are listed in ascending order of Retention Time).

TABLE S2 | GC-MS identified components of the crude extract of XEGI50 at pH3 (Volatile compounds are listed in ascending order of Retention Time).

TABLE S3 | GC-MS identified components of the crude extract of *V. dahliae* at pH7 (Volatile compounds are listed in ascending order of Retention Time).

TABLE S4 | GC-MS identified components of the crude extract of *V. dahliae* at pH3 (Volatile compounds are listed in ascending order of Retention Time).

TABLE S5 | GC-MS identified components of the antibiosis crude extract of XEGI50 and *V. dahliae* mixture at pH7 (Volatile compounds are listed in ascending order of Retention Time).

TABLE S6 | GC-MS identified components of the antibiosis crude extract of XEGI50 and *V. dahliae* mixture at pH3 (Volatile Compounds are listed in ascending order of Retention Time).

REFERENCES

- Achari, G. A., and Ramesh, R. (2014). Diversity, biocontrol, and plant growth promoting abilities of xylem residing bacteria from solanaceous crops. *Int. J. Microbiol.* 2014, 1–14. doi: 10.1155/2014/296521
- Aeron, A., Dubey, R., Maheshwari, D., Pandey, P., Bajpai, V. K., and Kang, S. C. (2011). Multifarious activity of bioformulated *Pseudomonas fluorescens* PS1 and biocontrol of *Sclerotinia sclerotiorum* in Indian rapeseed (*Brassica campestris* L.). *Eur. J. Plant Pathol.* 131, 81–93. doi: 10.1007/s00248-009-9531-y
- Agrawal, T., and Kotasthane, A. S. (2012). Chitinolytic assay of indigenous *Trichoderma* isolates collected from different geographical locations of Chhattisgarh in Central India. *Springerplus* 1:73. doi: 10.1186/2193-1801-1-73
- Akinsanya, M. A., Goh, J. K., Lim, S. P., and Ting, A. S. Y. (2015). Diversity, antimicrobial and antioxidant activities of culturable bacterial endophyte communities in *Aloe vera*. *FEMS Microbiol. Lett.* 362:fnv184. doi: 10.1093/femsle/fnv184
- Alvarez, F., Castro, M., Principe, A., Borioli, G., Fischer, S., Mori, G., et al. (2012). The plant-associated *Bacillus amyloliquefaciens* strains MEP218 and ARP23 capable of producing the cyclic lipopeptides iturin or surfactin and fengycin are effective in biocontrol of sclerotinia stem rot disease. *J. Appl. Bacteriol.* 112, 159–174. doi: 10.1111/j.1365-2672.2011.05182.x
- Asl, M. N., and Hosseinzadeh, H. (2008). Review of pharmacological effects of *Glycyrrhiza* sp. and its bioactive compounds. *Phytother. Res.* 22, 709–724. doi: 10.1002/ptr.2362
- Bacon, C., Hinton, D., and Snook, M. (2005). Tentative identification of *Bacillus mojavensis* antifungal inhibitor. *Phytopathology* 95:S5.
- Bacon, C. W., and Hinton, D. M. (2002). Endophytic and biological control potential of *Bacillus mojavensis* and related species. *Biol. Control* 23, 274–284. doi: 10.1006/bcon.2001.1016
- Bacon, C. W., and White, J. (2000). *Microbial Endophytes*. Boca Raton, FL: CRC Press.
- Bakker, P. A., Berendsen, R. L., Doornbos, R. F., Wintermans, P. C., and Pieterse, C. M. (2013). The rhizosphere revisited: root microbiomics. *Front. Plant Sci.* 4:165. doi: 10.3389/fpls.2013.00165
- Berg, G., and Hallmann, J. (2006). "Control of plant pathogenic fungi with bacterial endophytes," in *Microbial Root Endophytes*, eds B. J. E. Schulz, C. J. C. Boyle, and T. N. Sieber (Berlin: Springer), 53–69.
- Berg, G., Kurze, S., Buchner, A., Wellington, E. M., and Smalla, K. (2000). Successful strategy for the selection of new strawberry-associated rhizobacteria antagonistic to *Verticillium wilt*. *Can. J. Microbiol.* 46, 1128–1137. doi: 10.1139/w00-101
- Bhuvaneshwari, S., Madhavan, S., and Panneerselvam, A. (2013). Enumeration of endophytic bacteria from *Solanum trilobatum* L. *World J. Pharm. Res.* 3, 2270–2279.
- Bibi, N., Zhang, G., Li, F., Fan, K., Yuan, S., and Wang, X. (2013). Utilization of Vd toxin for rapid screening of cotton germplasm against *Verticillium dahliae*. *Pak. J. Bot.* 45, 2157–2162.
- Cavaglieri, L., Orlando, J., Rodriguez, M., Chulze, S., and Etcheverry, M. (2005). Biocontrol of *Bacillus subtilis* against *Fusarium verticillioides* in vitro and at the maize root level. *Res. Microbiol.* 156, 748–754. doi: 10.1016/j.resmic.2005.03.001
- Cazorla, F., Romero, D., Pérez-García, A., Lugtenberg, B., Vicente, A. D., and Bloemberg, G. (2007). Isolation and characterization of antagonistic *Bacillus subtilis* strains from the avocado rhizosphere displaying biocontrol activity. *J. Appl. Microbiol.* 103, 1950–1959. doi: 10.1111/j.1365-2672.2007.03433.x
- Chandrasekaran, M., Senthilkumar, A., and Venkatesalu, V. (2011). Antibacterial and antifungal efficacy of fatty acid methyl esters from the leaves of *Sesuvium portulacastrum* L. *Eur. Rev. Med. Pharmacol. Sci.* 15, 775–780.
- Chen, Y., Gao, X., Chen, Y., Qin, H., Huang, L., and Han, Q. (2014). Inhibitory efficacy of endophytic *Bacillus subtilis* EDR4 against *Sclerotinia sclerotiorum* on rapeseed. *Biol. Control* 78, 67–76. doi: 10.1016/j.biocontrol.2014.07.012
- Chernin, L., and Chet, I. (2002). *Microbial Enzymes in Biocontrol of Plant Pathogens and Pests. Enzymes in the Environment: Activity, Ecology, and Applications*. New York, NY: Marcel Dekker, 171–225.
- Cho, K. M., Hong, S. Y., Lee, S. M., Kim, Y. H., Kahng, G. G., Lim, Y. P., et al. (2007). Endophytic bacterial communities in ginseng and their antifungal activity against pathogens. *Microb. Ecol.* 54, 341–351. doi: 10.1007/s00248-007-9208-3
- Cho, S. J., Park, S. R., Kim, M. K., Lim, W. J., Ryu, S. K., An, C. L., et al. (2002). Endophytic *Bacillus* sp. isolated from the interior of balloon flower root. *Biosci. Biotechnol. Biochem.* 66, 1270–1275.
- Courtois, S., Cappellano, C. M., Ball, M., Francou, F.-X., Normand, P., Helyncq, G., et al. (2003). Recombinant environmental libraries provide access to microbial diversity for drug discovery from natural products. *Appl. Environ. Microbiol.* 69, 49–55.
- de Souza, J. T., de Boer, M., de Waard, P., van Beek, T. A., and Raaijmakers, J. M. (2003). Biochemical, genetic, and zoosporicidal properties of cyclic lipopeptide surfactants produced by *Pseudomonas fluorescens*. *Appl. Environ. Microbiol.* 69, 1716–1717.
- Dos Santos, P. J. C., Savi, D. C., Gomes, R. R., Goulin, E. H., Senkiv, C. D. C., Tanaka, F. A. O., et al. (2016). Diaporthe endophytica and *D. terebinthifolii* from medicinal plants for biological control of *Phyllosticta citricarpa*. *Microbiol. Res.* 186, 153–160. doi: 10.1016/j.micres.2016.04.002
- Droby, S. (2005). Improving quality and safety of fresh fruits and vegetables after harvest by the use of biocontrol agents and natural materials. *Acta Hort.* 709, 45–52.
- Ebrahimi, A., Asgharian, S., and Habibian, S. (2010). Antimicrobial activities of isolated endophytes from some Iranian native medicinal plants. *Iran. J. Pharm. Sci.* 6, 217–222. doi: 10.17795/zjrms-2482
- Egamberdieva, D., Wirth, S., Behrendt, U., Ahmad, P., and Berg, G. (2017a). Antimicrobial activity of medicinal plants correlates with the proportion of antagonistic endophytes. *Front. Microbiol.* 8:199. doi: 10.3389/fmicb.2017.00199
- Egamberdieva, D., Wirth, S. J., Shurigin, V. V., Hashem, A., and Abd-Allah, E. F. (2017b). Endophytic bacteria improve plant growth, symbiotic performance of chickpea (*Cicer arietinum* L.) and induce suppression of root rot caused by *Fusarium solani* under Salt Stress. *Front. Microbiol.* 8:1887. doi: 10.3389/fmicb.2017.01887
- Erdogan, O., and Benlioglu, K. (2010). Biological control of *Verticillium wilt* on cotton by the use of fluorescent *Pseudomonas* spp. under field conditions. *Biol. Control* 53, 39–45. doi: 10.1016/j.biocontrol.2009.11.011
- Finking, R., and Marahiel, M. A. (2004). Biosynthesis of nonribosomal peptides. *Annu. Rev. Microbiol.* 58, 453–488. doi: 10.1146/annurev.micro.58.030603.123615
- Fisher, M. C., Henk, D. A., Briggs, C. J., Brownstein, J. S., Madoff, L. C., McCraw, S. L., et al. (2012). Emerging fungal threats to animal, plant and ecosystem health. *Nature* 484, 186–194. doi: 10.1038/nature10947
- Frankowski, J., Lorito, M., Scala, F., Schmid, R., Berg, G., and Bahl, H. (2001). Purification and properties of two chitinolytic enzymes of *Serratia plymuthica* HRO-C48. *Arch. Microbiol.* 176, 421–426. doi: 10.1007/s002030100347
- Gao, Z., Zhang, B., Liu, H., Han, J., and Zhang, Y. (2017). Identification of endophytic *Bacillus velezensis* ZSY-1 strain and antifungal activity of its volatile compounds against *Alternaria solani* and *Botrytis cinerea*. *Biol. Control* 105, 27–39. doi: 10.1016/j.biocontrol.2016.11.007
- Göre, M. E., Caner, Ö. K., Altın, N., Aydın, M. H., Erdoğan, O., Filizer, F., et al. (2009). Evaluation of cotton cultivars for resistance to pathotypes of *Verticillium dahliae*. *Crop Prot.* 28, 215–219. doi: 10.1016/j.cropro.2008.10.004
- Guo, J. W., Yang, L. F., Liu, Y. H., Yang, J., Wang, H. F., Li, L., et al. (2016). First report of pseudostem black spot caused by *Pestalotiopsis microspora* on Tsao-ko in Yunnan, China. *Plant Dis.* 100, 1021–1021. doi: 10.1094/PDIS-08-15-0920-PDN
- Guo, Y., Huang, E., Yang, X., Zhang, L., Yousef, A. E., and Zhong, J. (2016). Isolation and characterization of a *Bacillus atrophaeus* strain and its potential use in food preservation. *Food Control* 60, 511–518. doi: 10.1016/j.foodcont.2015.08.029
- Harwood, T., Tomlinson, I., Potter, C., and Knight, J. (2011). Dutch elm disease revisited: past, present and future management in Great Britain. *Plant Pathol.* 60, 545–555. doi: 10.1111/j.1365-3059.2010.02391.x

- Horikoshi, K. (2008). Past, present and future of extremophiles. *Extremophiles* 12, 1–2. doi: 10.1007/s00792-007-0127-5
- Hsouna, A. B., Trigui, M., Mansour, R. B., Jarraya, R. M., Damak, M., and Jaoua, S. (2011). Chemical composition, cytotoxicity effect and antimicrobial activity of *Ceratonia siliqua* essential oil with preservative effects against *Listeria* inoculated in minced beef meat. *Int. J. Food Microbiol.* 148, 66–72. doi: 10.1016/j.jfoodmicro.2011.04.028
- Janisiewicz, W. J., and Korsten, L. (2002). Biological control of postharvest diseases of fruits. *Annu. Rev. Phytopathol.* 40, 411–441.
- Jiang, C.-H., Wu, F., Yu, Z.-Y., Xie, P., Ke, H.-J., Li, H.-W., et al. (2015). Study on screening and antagonistic mechanisms of *Bacillus amyloliquefaciens* 54 against bacterial fruit blotch (BFB) caused by *Acidovorax avenae* subsp. *citruilli*. *Microbiol. Res.* 170, 95–104. doi: 10.1016/j.micres.2014.08.009
- Juzwik, J., Harrington, T. C., MacDonald, W. L., and Appel, D. N. (2008). The origin of *Ceratocystis fagacearum*, the oak wilt fungus. *Annu. Rev. Phytopathol.* 46, 13–26. doi: 10.1146/annurev.phyto.45.062806.094406
- Kandasamy, S., Sahu, S. K., and Kandasamy, K. (2012). In Silico studies on fungal metabolite against skin cancer protein (4, 5-Diarylisoazole HSP90 Chaperone). *ISRN Dermatol.* 2012, 626214. doi: 10.5402/2012/626214
- Kavitha, A., Prabhakar, P., Narasimhulu, M., Vijayalakshmi, M., Venkateswarlu, Y., Rao, K. V., et al. (2010). Isolation, characterization and biological evaluation of bioactive metabolites from *Nocardia levis* MK-VL_113. *Microbiol. Res.* 165, 199–210. doi: 10.1016/j.micres.2009.05.002
- Korsten, L., De Villiers, E., Wehner, F., and Kotzé, J. (1997). Field sprays of *Bacillus subtilis* and fungicides for control of preharvest fruit diseases of avocado in South Africa. *Plant Dis.* 81, 455–459. doi: 10.1094/PDIS.1997.81.5.455
- Kun, L., Zhixiang, Z., and Xiaofei, L. (2010). Screening for PKS gene from soil metagenomics library and identification of the actives against root-knot nematodes. *Plant Prot.* 36, 38–42.
- Kusari, S., Pandey, S. P., and Spittler, M. (2013). Untapped mutualistic paradigms linking host plant and endophytic fungal production of similar bioactive secondary metabolites. *Phytochemistry* 91, 81–87. doi: 10.1016/j.phytochem.2012.07.021
- Lacava, P. T., Li, W., Araujo, W. L., Azevedo, J. L. C., and Hartung, J. S. (2007). The endophyte *Curtobacterium flaccumfaciens* reduces symptoms caused by *Xylella fastidiosa* in *Catharanthus roseus*. *J. Microbiol.* 45, 388–393.
- Li, L., Mohamad, O. A. A., Ma, J., Friel, A. D., Su, Y., Wang, Y., et al. (2018). Synergistic plant-microbe interactions between endophytic bacterial communities and the medicinal plant *Glycyrrhiza uralensis* F. *Antonie Van Leeuwenhoek* 2018, 1–14. doi: 10.1007/s10482-018-1062-4
- Liao, W. C., Lin, Y. H., Chang, T. M., and Huang, W. Y. (2012). Identification of two licorice species, *Glycyrrhiza uralensis* and *Glycyrrhiza glabra*, based on separation and identification of their bioactive components. *Food Chem.* 132, 2188–2193. doi: 10.1016/j.foodchem.2011.12.051
- Lin, T., Zhao, L., Yang, Y., Guan, Q., and Gong, M. (2013). Potential of endophytic bacteria isolated from ‘*Sophora alopecuroides*’ nodule in biological control against *Verticillium wilt* disease. *Aust. J. Crop Sci.* 7:139.
- Liu, L., Salam, N., Jiao, J.-Y., Jiang, H.-C., Zhou, E.-M., Yin, Y.-R., et al. (2016). Diversity of culturable thermophilic Actinobacteria in hot springs in Tengchong, China and studies of their biosynthetic gene profiles. *Microb. Ecol.* 72, 150–162. doi: 10.1007/s00248-016-0756-2
- Liu, Y.-H., Guo, J.-W., Salam, N., Li, L., Zhang, Y.-G., Han, J., et al. (2016). Culturable endophytic bacteria associated with medicinal plant *Ferula songorica*: molecular phylogeny, distribution and screening for industrially important traits. *3 Biotech* 6:209. doi: 10.1007/s13205-016-0522-7
- Liu, Y., Guo, J., Li, L., Asem, M. D., Zhang, Y., Mohamad, O. A., et al. (2017). Endophytic bacteria associated with endangered plant *Ferula sinkiangensis* KM Shen in an arid land: diversity and plant growth-promoting traits. *J. Arid Land* 9, 432–445. doi: 10.1007/s40333-017-0015-5
- Lodewyckx, C., Vangronsveld, J., Porteous, F., Moore, E. R., Taghavi, S., Mezgeay, M., et al. (2002). Endophytic bacteria and their potential applications. *Crit. Rev. Plant Sci.* 21, 583–606. doi: 10.1080/0735-260291044377
- Loqman, S., Barka, E. A., Clément, C., and Ouhdouch, Y. (2009). Antagonistic actinomycetes from Moroccan soil to control the grapevine gray mold. *World J. Microbiol. Biotechnol.* 25, 81–91. doi: 10.1007/s11274-008-9864-6
- Maldonado-González, M. M., Bakker, P. A., Prieto, P., and Mercado-Blanco, J. (2015). *Arabidopsis thaliana* as a tool to identify traits involved in *Verticillium dahliae* biocontrol by the olive root endophyte *Pseudomonas fluorescens* PICF7. *Front. Microbiol.* 6:266. doi: 10.3389/fmicb.2015.00266
- Metsä-Ketelä, M., Salo, V., Halo, L., Hautala, A., Hakala, J., Mäntsälä, P., et al. (1999). An efficient approach for screening minimal PKS genes from *Streptomyces*. *FEMS Microbiol. Lett.* 180, 1–6. doi: 10.1016/S0378-1097(99)00453-X
- Nejatzadeh-Barandozi, F. (2013). Antibacterial activities and antioxidant capacity of *Aloe vera*. *Org. Med. Chem. Lett.* 3:5. doi: 10.1186/2191-2858-3-5
- Nie, Y., Zeng, X.-C., Yang, Y., Luo, F., Luo, X., Wu, S., et al. (2012). A novel class of antimicrobial peptides from the scorpion *Heterometrus spinifer*. *Peptides* 38, 389–394. doi: 10.1016/j.peptides.2012.09.012
- Peng, H., Thevs, N., Beckmann, V., and Abdusalih, N. (2016). Economic performance of cotton and fruit plantations in arid regions: observation from the Tarim River Basin, NW China. *Asian J. Agric. Ext. Econ. Sociol.* 8, 1–15. doi: 10.9734/AJAEES/2016/22254
- Qing, W., Zuo, J.-H., Qian, W., Yang, N., and Gao, L.-P. (2015). Inhibitory effect of chitosan on growth of the fungal phytopathogen, *Sclerotinia sclerotiorum*, and sclerotinia rot of carrot. *J. Integr. Agric.* 14, 691–697. doi: 10.1016/S2095-3119(14)60800-5
- Radhakrishnan, R., Hashem, A., and Abd-Allah, E. F. (2017). *Bacillus*: a biological tool for crop improvement through bio-molecular changes in adverse environments. *Front. Physiol.* 8:667. doi: 10.3389/fphys.2017.00667
- Rodríguez-Meizoso, I., Jaime, L., Santoyo, S., Señoráns, F., Cifuentes, A., and Ibáñez, E. (2010). Subcritical water extraction and characterization of bioactive compounds from *Haematococcus pluvialis* microalga. *J. Pharm. Biomed. Anal.* 51, 456–463. doi: 10.1016/j.jpba.2009.03.014
- Rokade, Y., and Sayyed, R. (2009). Naphthalene derivatives: a new range of antimicrobials with high therapeutic value. *Rasayan J. Chem.* 2, 972–980.
- Saikkonen, K., Wäli, P., Helander, M., and Faeth, S. H. (2004). Evolution of endophyte-plant symbioses. *Trends Plant Sci.* 9, 275–280. doi: 10.1016/j.tplants.2004.04.005
- Santhanam, R., Groten, K., Meldau, D. G., and Baldwin, I. T. (2014). Analysis of plant-bacteria interactions in their native habitat: bacterial communities associated with wild tobacco are independent of endogenous jasmonic acid levels and developmental stages. *PLoS One* 9:e94710. doi: 10.1371/journal.pone.0094710
- Schneemann, I., Nagel, K., Kajahn, I., Labes, A., Wiese, J., and Imhoff, J. F. (2010). Comprehensive investigation of marine Actinobacteria associated with the sponge *Halichondria panicea*. *Appl. Environ. Microbiol.* 76, 3702–3714. doi: 10.1128/AEM.00780-10
- Sermakkani, M., and Thangapandian, V. (2012). GC-MS analysis of *Cassia italica* leaf methanol extract. *Asian J. Pharm. Clin. Res.* 5, 90–94.
- Sharma, D., Pramanik, A., and Agrawal, P. K. (2016). Evaluation of bioactive secondary metabolites from endophytic fungus *Pestalotiopsis neglecta* BAB-5510 isolated from leaves of *Cupressus torulosa* D. Don. *3 Biotech* 6:210. doi: 10.1007/s13205-016-0518-3
- Slepecky, R., and He, H. (1992). “The genus *Bacillus*-Nonmedical,” in *The Prokaryotes*, eds A. Balows, H. G. Trüper, M. Dworkin, W. Harder, and K.-H. Schleifer (New York, NY: Springer-Verlag), 1663–1696.
- Strobel, G., and Daisy, B. (2003). Bioprospecting for microbial endophytes and their natural products. *Microbiol. Mol. Biol. Rev.* 67, 491–502. doi: 10.1128/MMBR.67.4.491-502.2003
- Sturz, A. V., Christie, B. R., and Nowak, J. (2000). Bacterial endophytes: potential role in developing sustainable systems of crop production. *Crit. Rev. Plant Sci.* 19, 1–30. doi: 10.1080/07352680091139169
- Sun, G., Yao, T., Feng, C., Chen, L., Li, J., and Wang, L. (2017). Identification and biocontrol potential of antagonistic bacteria strains against *Sclerotinia sclerotiorum* and their growth-promoting effects on *Brassica napus*. *Biol. Control* 104, 35–43. doi: 10.1016/j.biocontrol.2016.10.008
- Tambekar, D., Tiwari, A., and Tambekar, S. (2014). Studies on production of antimicrobial substances from *Bacillus* species isolated from Lonar Lake. *Indian J. Appl. Res.* 4, 502–506. doi: 10.15373/2249555X/August2014/131
- Ting, A. S. Y. (2014). “Biosourcing endophytes as biocontrol agents of wilt diseases,” in *Advances in Endophytic Research*, eds V. Verma and A. Gange (New Delhi: Springer), 283–300.

- Vieira, M. L., Hughes, A. F., Gil, V. B., Vaz, A. B., Alves, T. M., Zani, C. L., et al. (2011). Diversity and antimicrobial activities of the fungal endophyte community associated with the traditional Brazilian medicinal plant *Solanum cernuum* Vell. (Solanaceae). *Can. J. Microbiol.* 58, 54–66. doi: 10.1139/w11-105
- Walker, R., Powell, A. A., and Seddon, B. (1998). Bacillus isolates from the spermosphere of peas and dwarf French beans with antifungal activity against *Botrytis cinerea* and Pythium species. *J. Appl. Microbiol.* 84, 791–801. doi: 10.1046/j.1365-2672.1998.00411.x
- Weller, D. M. (1988). Biological control of soilborne plant pathogens in the rhizosphere with bacteria. *Annu. Rev. Phytopathol.* 26, 379–407. doi: 10.1146/annurev.py.26.090188.002115
- Yadeta, K. A., and Thomma, B. P. (2013). The xylem as battleground for plant hosts and vascular wilt pathogens. *Front. Plant Sci.* 4:97. doi: 10.3389/fpls.2013.00097
- Zhang, B., Yang, Y., Chen, T., Yu, W., Liu, T., Li, H., et al. (2012). Island cotton Gbve1 gene encoding a receptor-like protein confers resistance to both defoliating and non-defoliating isolates of *Verticillium dahliae*. *PLoS One* 7:e51091. doi: 10.1371/journal.pone.0051091
- Zhang, Q., and Ye, M. (2009). Chemical analysis of the Chinese herbal medicine Gan-Cao (licorice). *J. Chromatogr. A* 1216, 1954–1969. doi: 10.1016/j.chroma.2008.07.072
- Zheng, Y., Chen, F., and Wang, M. (2013). “Use of Bacillus-based biocontrol agents for promoting plant growth and health,” in *Bacteria in Agrobiolgy: Disease Management*, ed. D. K. Maheshwari (Berlin: Springer), 243–258. doi: 10.1016/j.chroma.2008.07.072

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2018 Mohamad, Li, Ma, Hatab, Xu, Guo, Rasulov, Liu, Hedlund and Li. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.