



Virginia Commonwealth University
VCU Scholars Compass

Hepatobiliary Cancers: Pathobiology and
Translational Advances

Dept. of Pathology

2017

Regulation of Biliary Progenitor Cell Proliferation by Hedgehog Signaling in In Vivo and Ex Vivo Organoid Models

Nataliya Razumilava M.D.

The University Of Michigan, razumila@med.umich.edu

Junya Shiota M.D.

The University Of Michigan, shiotaj@med.umich.edu

Ramon Ocadiz-Ruiz Ph.D.

The University Of Michigan, jocadizr@med.umich.edu

See next page for additional authors

Follow this and additional works at: http://scholarscompass.vcu.edu/hepa_cancers

 Part of the [Biological Phenomena, Cell Phenomena, and Immunity Commons](#), [Digestive System Diseases Commons](#), [Medical Cell Biology Commons](#), and the [Neoplasms Commons](#)

© The Author(s)

Downloaded from

http://scholarscompass.vcu.edu/hepa_cancers/11

This Abstract Accepted for Presentation is brought to you for free and open access by the Dept. of Pathology at VCU Scholars Compass. It has been accepted for inclusion in Hepatobiliary Cancers: Pathobiology and Translational Advances by an authorized administrator of VCU Scholars Compass. For more information, please contact libcompass@vcu.edu.

Authors

Nataliya Razumilava M.D.; Junya Shiota M.D.; Ramon Ocadiz-Ruiz Ph.D.; Nureen Hanisah Mohamad Zaki; Kais Zakharia M.D.; Michael Mathew Hayes; and Juanita Lynn Merchant M.D., Ph.D.

Regulation of Biliary Progenitor Cell Proliferation by Hedgehog Signaling in *In Vivo* and *Ex Vivo* Organoid Models

N Razumilava,¹ J Shiota,¹ R Ocadiz-Ruiz,¹ NH Mohamad Zaki,¹ K Zakharia,¹ MM Hayes,¹ JL Merchant¹

¹Division of Gastroenterology, University of Michigan, Ann Arbor, MI

Background: Cholangiopathies are a group of chronic progressive diseases with poor outcomes, and include primary sclerosing cholangitis and cholangiocarcinoma (CCA). Hedgehog (Hh) signaling mediates biliary repair in injury and is activated in CCA. The steps from chronic inflammation to CCA are not well defined, but are thought to include hyperplasia of the PBGs, the niche for biliary progenitor cells (BPCs). **Objectives:** To elucidate the effects of Hh signaling on the BPCs. **Methods:** We used wild type (WT) mice, and mice with Hh pathway activation (*pCMV-Shh^{HA}*) and inhibition (*Gli1^{-/-}*). We assessed the mouse bile ducts (BDs) for proliferation (BrdU) and expression of differentiated cell markers *in vivo*. We used the novel 3-D biliary organoid (BO) model and evaluated effects of genetic and pharmacological Hh signaling activation [recombinant Shh (rShh)] and inhibition (GANT-61) *ex vivo*. **Results:** We demonstrated that Shh expression in the *pCMV-Shh^{HA}* mice localized to the PBGs and Gli1 was expressed in the *Gli1^{-/-}* mice stroma in homeostasis. Hh signaling did not affect PBG cell proliferation but rather promoted PBG cell differentiation in homeostasis *in vivo*. However, the *pCMV-Shh^{HA}* mice-derived organoids and BOs treated with rShh demonstrated increased BO growth and proliferation *ex vivo*. Pharmacological Gli inhibition significantly decreased growth and proliferation in the BOs from the WT and *pCMV-Shh^{HA}* mice. **Conclusions:** Hh signaling regulates PBG proliferation, which can contribute to PBG hyperplasia, in a paracrine and autocrine manner. The cross-talk between the Shh ligand-producing PBGs and Gli1+ stromal cells is required to control BPC proliferation.