

## Virginia Commonwealth University VCU Scholars Compass

Biology and Medicine Through Mathematics Conference

2018

Jun 1st, 10:00 AM - 10:30 AM

## Transport Phenomena in Field Effect Transistors

Ryan M. Evans National Institute of Standards and Technology, ryan.evans@nist.gov

Arvind Balijepalli National Institute of Standards and Technology, arvind.balijepalli@nist.gov

Anthony Kearsley National Institute of Standards and Technology, anthony.kearsley@nist.gov

Follow this and additional works at: https://scholarscompass.vcu.edu/bamm Part of the <u>Medicine and Health Sciences Commons</u>, <u>Numerical Analysis and Computation</u> <u>Commons</u>, and the <u>Partial Differential Equations Commons</u>

https://scholarscompass.vcu.edu/bamm/2018/friday/9

This Event is brought to you for free and open access by the Dept. of Mathematics and Applied Mathematics at VCU Scholars Compass. It has been accepted for inclusion in Biology and Medicine Through Mathematics Conference by an authorized administrator of VCU Scholars Compass. For more information, please contact libcompass@vcu.edu.

## **Transport Phenomena in Field Effect Transistors**

<sup>1,\*</sup> Dr. Ryan Evans, <sup>2</sup>Dr. Arvind Balijepalli, <sup>1</sup>Dr. Anthony Kearsley

<sup>1</sup>Applied and Computational Mathematics Division <sup>2</sup>Engineering Physics Division National Institute of Standards and Technology 100 Bureau Drive Gaithersburg, MD 20899 \*ryan.evans@nist.gov

## Abstract

Tailoring therapies to individuals for personalized care can be safer and yield superior outcomes with lower doses for conditions such as diabetes, Alzheimers disease, or even certain cancers. However, widespread use of personalized care is currently limited by inability to measure pathology and detect biomarkers. Moreover, existing strategies require specialized facilities, can be slow to perform and can be expensive. This has led to the development of a new portable detection tool known as a field effect transistor (FET). Very well-suited for biomarker measurements due their high charge sensitivity and direct signal transduction, FETs allow label-free measurements at physiological concentrations. Chemical reactants are injected at the top of solution-well and diffuse through a well to bind with another chemical reactant immobilized to a narrow band on the well-floor. A resulting response curve allows for biomarker measurement and estimation of key parameters, such as binding affinities. A mathematical model for FET experiments will be presented that takes the form of a diffusion coupled to a nonlinear equation that describes the evolution of the reacting species concentration.