

Masthead Logo

Virginia Commonwealth University  
VCU Scholars Compass

---

Biology and Medicine Through Mathematics  
Conference

2019

---

May 15th, 3:30 PM

# Modeling and control of enzootic West Nile virus transmission: The role of the avian nesting curve

Suzanne Robertson

*Virginia Commonwealth University*, [srobertson7@vcu.edu](mailto:srobertson7@vcu.edu)

Kevin Caillouet

*St. Tammany Parish Mosquito Abatement District*, [caillouet@stpmd.org](mailto:caillouet@stpmd.org)

Elsa Schaefer

*Marymount University*, [schaefer@marymount.edu](mailto:schaefer@marymount.edu)

Follow this and additional works at: <https://scholarscompass.vcu.edu/bamm>

Part of the [Life Sciences Commons](#), and the [Physical Sciences and Mathematics Commons](#)

---

<https://scholarscompass.vcu.edu/bamm/2019/wed/10>

This Event is brought to you for free and open access by the Dept. of Mathematics and Applied Mathematics at VCU Scholars Compass. It has been accepted for inclusion in Biology and Medicine Through Mathematics Conference by an authorized administrator of VCU Scholars Compass. For more information, please contact [libcompass@vcu.edu](mailto:libcompass@vcu.edu).

**Title:** Modeling and control of enzootic *West Nile virus* transmission: The role of the avian nesting curve

**Abstract:** Introduced into the United States in 1999, *West Nile virus* (WNV) remains a major public health concern. The ecological mechanisms responsible for the timing and magnitude of seasonal transmission are not well understood, though outbreaks have been widely observed to be associated with the end of the avian nesting season. Newly hatched birds, or nestlings, have less feather coverage and fewer defense mechanisms than older birds, rendering them more vulnerable to mosquitoes. The rate at which new nestlings are produced is determined by the avian nesting curve, which can vary with avian species and is also influenced by climate. We use a mathematical model incorporating avian (host) stage-structure and within-species heterogeneity in the form of stage-specific mosquito (vector) biting rates to investigate the connection between properties of the avian nesting curve and enzootic WNV transmission, as well as implications for control of WNV.