

Masthead Logo

Virginia Commonwealth University
VCU Scholars Compass

Biology and Medicine Through Mathematics
Conference

2019

May 15th, 6:00 PM

Immunofluorescence Image Feature Analysis and Clustering Pipeline for Distinguishing Epithelial-Mesenchymal Transition

Shreyas Hirway

Virginia Commonwealth University, hirwaysu@vcu.edu

Nadiah Hassan

Virginia Commonwealth University, hassannt@vcu.edu

Dr. Christopher Lemmon

Virginia Commonwealth University, clemmon@vcu.edu

See next page for additional authors

Follow this and additional works at: <https://scholarscompass.vcu.edu/bamm>

Part of the [Biological Engineering Commons](#), [Life Sciences Commons](#), [Medicine and Health Sciences Commons](#), and the [Numerical Analysis and Computation Commons](#)

<https://scholarscompass.vcu.edu/bamm/2019/wed/19>

This Event is brought to you for free and open access by the Dept. of Mathematics and Applied Mathematics at VCU Scholars Compass. It has been accepted for inclusion in Biology and Medicine Through Mathematics Conference by an authorized administrator of VCU Scholars Compass. For more information, please contact libcompass@vcu.edu.

Presenter Information

Shreyas Hirway, Nadiah Hassan, Dr. Christopher Lemmon, and Dr. Seth Weinberg

Epithelial-Mesenchymal Transition (EMT) is a biological process essential for tissue regeneration and implicated in pathological process such as cancer metastasis, in which epithelial cells transdifferentiate into mesenchymal cells. We devised an image processing pipeline to distinguish between tissues comprised of epithelial and mesenchymal images based on extracted features, using immunofluorescence images which demonstrate differing cellular and tissue structure and biochemical markers. Mammary epithelial cells were incubated with 0ng/mL (control) or 10ng/mL TGF- β 1 and cultured for 48 hours. Cells were fixed, stained and imaged for E-cadherin, actin, fibronectin and nuclei via immunofluorescence microscopy. Feature selection was performed on both composites images and different combinations of individual channels using the MATLAB Bag-of-Features tool. Principal Component Analysis (PCA) was performed on the extracted data, and a normalized epithelial and mesenchymal score was defined for each image, based on Euclidean distance from cluster centers. Each image was also rated on a Likert (1-5) scale, by 4 lab members blinded to treatment conditions. PCA produced distinct clusters between epithelial and mesenchymal images. We found that increasing the number of extracted features required more components to explain a given variance. Surprisingly, fewer features produced greater cluster separation and more cohesive clusters. Amongst the analysis of individual channels, imaging of fibronectin yielded the best cluster separation. Lab member ratings agreed closely with epithelial and mesenchymal scores. Overall, the feature extraction and PCA pipeline distinguished epithelial and mesenchymal immunofluorescence images, and future work will focus on characterization of images in intermediate stages of the EMT process.