

Masthead Logo

Virginia Commonwealth University
VCU Scholars Compass

Biology and Medicine Through Mathematics
Conference

2019

May 15th, 6:00 PM

A pharmacokinetic model of lead absorption and calcium competitive dynamics

Anca R. Radulescu

State University of New York at New Paltz, radulesa@newpaltz.edu

Follow this and additional works at: <https://scholarscompass.vcu.edu/bamm>

Part of the [Medicine and Health Sciences Commons](#), [Ordinary Differential Equations and Applied Dynamics Commons](#), and the [Other Pharmacology, Toxicology and Environmental Health Commons](#)

<https://scholarscompass.vcu.edu/bamm/2019/wed/21>

This Event is brought to you for free and open access by the Dept. of Mathematics and Applied Mathematics at VCU Scholars Compass. It has been accepted for inclusion in Biology and Medicine Through Mathematics Conference by an authorized administrator of VCU Scholars Compass. For more information, please contact libcompass@vcu.edu.

A pharmacokinetic model of lead absorption and calcium competitive dynamics

Anca Rădulescu, SUNY New Paltz, radulesa@newpaltz.edu
Tucker Lundgren, SUNY New Paltz, lundgres2@hawkmil.newpaltz.edu

Lead is a naturally-occurring element. It has been known to man for a long time, and it is one of the longest established poisons. The current consensus is that no level of lead exposure should be deemed “safe.” New evidence regarding the blood levels at which morbidities occur has prompted the CDC to reduce the screening guideline of 10 $\mu\text{g}/\text{dl}$ to 2 $\mu\text{g}/\text{dl}$. Measurable cognitive decline (reduced IQ, academic deficits) have been found to occur at levels below 10mg/dl.

Knowledge of lead pharmacology allows us to better understand its absorption and metabolization, mechanisms that produce its medical consequences. Based upon an original and very simplified compartmental model of Rabinowitz (1973) with only three major compartments (blood, bone and soft tissue), extensive biophysical models sprouted over the following two decades. However, none of these models have been specifically designed to use new knowledge of lead molecular dynamics to understand its deleterious effects on the brain. We build and analyze a compartmental model of lead pharmacokinetics, focused specifically on addressing neurotoxicity. We use traditional phase space methods, parameter sensitivity analysis and bifurcation theory to study the transitions in the system’s behavior in response to various physiological parameters.

We conclude that modeling the complex interaction of lead and calcium along their dynamic trajectory may successfully explain counter-intuitive effects on systemic function and neural behavior which could not be addressed by existing linear models. Our results encourage further efforts towards using nonlinear phenomenology in conjunction with empirically driven system parameters, to obtain a biophysical model able to provide clinical assessments and predictions.