

Masthead Logo

Virginia Commonwealth University
VCU Scholars Compass

Biology and Medicine Through Mathematics
Conference

2019

May 15th, 6:30 PM

Ebola: Impact of hospital's admission policy in an overwhelmed scenario

Mondal H. Zahid

University of Texas at Arlington, mdmondal.zahid@mavs.uta.edu

Follow this and additional works at: <https://scholarscompass.vcu.edu/bamm>

Part of the [Disease Modeling Commons](#), [Epidemiology Commons](#), and the [Physical Sciences and Mathematics Commons](#)

<https://scholarscompass.vcu.edu/bamm/2019/wed/32>

This Event is brought to you for free and open access by the Dept. of Mathematics and Applied Mathematics at VCU Scholars Compass. It has been accepted for inclusion in Biology and Medicine Through Mathematics Conference by an authorized administrator of VCU Scholars Compass. For more information, please contact libcompass@vcu.edu.

Ebola: Impact of hospital's admission policy in an overwhelmed scenario

Mondal Hasan Zahid and Christopher M. Kribs

January 31, 2019

Infectious disease outbreaks sometimes overwhelm healthcare facilities. A recent case occurred in West Africa in 2014 when an Ebola virus outbreak overwhelmed facilities in Sierra Leone, Guinea and Liberia. In such scenarios, how many patients can hospitals admit to minimize disease burden? This study considers what type of hospital admission policy during a hypothetical Ebola outbreak can better serve the community, if overcrowding degrades the hospital setting. Our result shows that which policy minimizes loss to the community depends on the initial estimation of the control reproduction number, R_0 . When the outbreak grows extremely fast ($R_0 \gg 1$) it is better (in terms of total disease burden) to stop admitting patients after reaching the carrying capacity because overcrowding in the hospital makes the hospital setting ineffective at containing infection, but when the outbreak grows only a little faster than the system's ability to contain it ($R_0 \gtrsim 1$), it is better to admit patients beyond the carrying capacity because limited overcrowding still reduces infection more in the community. However, when R_0 is no more than a little greater than 1 (for our parameter values, 1.012), both policies result the same because the number of patients never exceeds the maximum capacity.