

Stephen F. Austin State University
SFA ScholarWorks

Faculty Publications

Chemistry and Biochemistry

2010

Mechanistic and computational studies of oxidatively-induced aryl-CF₃ bond formation at palladium

J. Brannon Gary

Stephen F Austin State University Department of Chemistry & Biochemistry, garyjb@sfasu.edu

Follow this and additional works at: https://scholarworks.sfasu.edu/chemistry_facultypubs

 Part of the [Chemistry Commons](#)

Tell us how this article helped you.

Repository Citation

Gary, J. Brannon, "Mechanistic and computational studies of oxidatively-induced aryl-CF₃ bond formation at palladium" (2010). *Faculty Publications*. 65.

https://scholarworks.sfasu.edu/chemistry_facultypubs/65

This Article is brought to you for free and open access by the Chemistry and Biochemistry at SFA ScholarWorks. It has been accepted for inclusion in Faculty Publications by an authorized administrator of SFA ScholarWorks. For more information, please contact cdsscholarworks@sfasu.edu.



Published in final edited form as:

J Am Chem Soc. 2011 May 18; 133(19): 7577–7584. doi:10.1021/ja201726q.

Mechanistic and Computational Studies of Oxidatively-Induced Aryl–CF₃ Bond-Formation at Pd: Rational Design of Room Temperature Aryl Trifluoromethylation

Nicholas D. Ball, J. Brannon Gary, Yingda Ye, and Melanie S. Sanford*

Department of Chemistry, University of Michigan, 930 N. University Avenue, Ann Arbor, Michigan 48109

Abstract

This article describes the rational design of 1st generation systems for oxidatively-induced Aryl–CF₃ bond-forming reductive elimination from Pd^{II}. Treatment of (dtbpy)Pd^{II}(Aryl)(CF₃) (dtbpy = di-*tert*-butylbipyridine) with NFTPT (*N*-fluoro-1,3,5-trimethylpyridium triflate) afforded the isolable Pd^{IV} intermediate (dtbpy)Pd^{IV}(Aryl)(CF₃)(F)(OTf). Thermolysis of this complex at 80 °C resulted in Aryl–CF₃ bond-formation. Detailed experimental and computational mechanistic studies have been conducted to gain insights into the key reductive elimination step. Reductive elimination from this Pd^{IV} species proceeds via pre-equilibrium dissociation of TfO[−] followed by Aryl–CF₃ coupling. DFT calculations reveal that the transition state for Aryl–CF₃ bond formation involves the CF₃ acting as an electrophile with the Aryl ligand acting as a nucleophilic coupling partner. These mechanistic considerations along with DFT calculations have facilitated the design of a 2nd generation system utilizing the tmeda (*N,N,N',N'*-tetramethylethylenediamine) ligand in place of dtbpy. The tmeda complexes undergo oxidative trifluoromethylation at room temperature.

Introduction


The formation of C–CF₃ bonds is an important transformation for the construction of pharmaceuticals and agrochemicals.¹ Replacing a methyl group with a trifluoromethyl substituent can have a profound effect on the physical and biological properties of a molecule.² As a result, there is high demand for versatile synthetic methods for generating carbon–CF₃ bonds.³ While there has been significant progress in the construction of sp³-carbon–CF₃ linkages,³ there are comparatively fewer methods for Aryl–CF₃ bond-formation.^{4,5,6,7}

Transition metal-catalyzed cross-coupling between Aryl–X and CF₃–Y would serve as an attractive method for the synthesis of benzotrifluorides.^{8,9,10,11} The use of Pd-based catalysts for such transformations is of particular interest because they serve as versatile and widely used catalysts for a variety of other carbon–carbon bond-forming reactions.¹² However, developments in this area have been limited by the challenges associated with a key step of the cross-coupling catalytic cycle, namely Aryl–CF₃ bond-forming reductive elimination from Pd^{II} centers.^{4,13} The vast majority of known Pd^{II}(Aryl)(CF₃) complexes are inert to Aryl–CF₃ coupling at temperatures as high as 150 °C.^{24,21}


* mssanfor@umich.edu .

Supporting Information Available. Experimental and computational details and spectroscopic data for new compounds. This material is available free of charge via the internet at <http://pubs.acs.org>.

Two strategies have been utilized in the literature to address this challenge. The first has focused on achieving Aryl–CF₃ bond-forming reductive elimination from Pd^{II} via steric and electronic modification of the ancillary ligands (L) at (L)_nPd^{II}(Aryl)(CF₃). For example, pioneering work by Grushin demonstrated that the Xantphos ligand (Xantphos = 4,5-bis(diphenylphosphino)-9,9-dimethylxanthene) facilitates high yielding formation of trifluorotoluene from (Xantphos)Pd^{II}(Ph)(CF₃) at 80 °C (eq. 1).¹⁴ More recently, Buchwald has elegantly demonstrated that the sterically large monodentate phosphine ligand Brettphos (Brettphos = 2-(dicyclohexylphosphino)-3,6-dimethoxy-2',4',6'-triisopropyl-1,1'-biphenyl) promotes stoichiometric Aryl–CF₃ coupling from (Brettphos)Pd^{II}(Aryl)(CF₃) at 80 °C (eq. 2).¹⁵ Remarkably, this latter system was successfully applied to the catalytic trifluoromethylation of aryl chlorides with Et₃SiCF₃ at 130–140 °C.¹⁵



While this first approach has provided important breakthroughs, Pd^{II}-mediated Aryl–CF₃ bond-forming reactions remain limited by the requirement for specialized and expensive phosphine ligands^{16,17} relatively high reaction temperatures (80–140 °C), and the need for expensive Et₃SiCF₃¹⁸ in catalytic processes. As a result, several groups have focused on a second, complementary strategy for achieving Aryl–CF₃ bond-forming reductive elimination from Pd.^{19,20,21} This approach hinges on changing the oxidation state, rather than the ancillary ligand environment, at the metal center. As shown in eq. 3, it was expected that palladium(IV) complexes of general structure (L)₂(X)₂Pd^{IV}(Aryl)(CF₃) (**A**) would be highly kinetically and thermodynamically reactive towards Aryl–CF₃ coupling. This hypothesis is predicated on literature reports showing that Pd^{IV} complexes participate in numerous carbon-heteroatom bond-forming reductive elimination reactions that remain challenging at Pd^{II} centers.^{22,23} A key advantage of this approach would be that Pd^{IV} intermediate **A** could potentially be accessed using nucleophilic (CF₃[−])²⁴, electrophilic (CF₃⁺)²⁵, or free-radical (CF₃·)²⁶ based trifluoromethylating reagents.



Two preliminary examples have shown the viability of this approach. First, Yu and coworkers have elegantly demonstrated the Pd^{II/IV}-catalyzed ligand-directed C–H trifluoromethylation with electrophilic trifluoromethylating reagents (CF₃⁺ reagents).¹⁹ Subsequent stoichiometric studies implicated a mechanism involving: (a) C–H activation to form a cyclometalated Pd^{II}–Aryl intermediate, (b) oxidation with CF₃⁺ to generate

$\text{Pd}^{\text{IV}}(\text{Aryl})(\text{CF}_3)$, and (c) Aryl– CF_3 bond-forming reductive elimination from Pd^{IV} to release the product.²⁰


In a second example, our group has shown the viability of stoichiometric Aryl– CF_3 bond-forming reductive elimination from Pd^{IV} centers bearing σ -aryl ligands that do not contain chelate directing groups.²¹ In this system, the key Pd^{IV} intermediate is generated via oxidation of a pre-assembled $\text{Pd}^{\text{II}}(\text{Aryl})(\text{CF}_3)$ species with an *N*-fluoropyridinium reagent. We report herein a detailed mechanistic study of Aryl– CF_3 bond-forming reductive elimination from Pd^{IV} in this latter system. This work provides valuable insights into the influence of ancillary ligands, the role of each coupling partner, and the nature of the transition state for this transformation. These mechanistic studies have also allowed us to rationally design the first examples of *room temperature* Aryl– CF_3 bond-formation from palladium.

Results and Discussion


Development and Scope of Aryl– CF_3 Coupling at Pd^{IV} .

Our studies began with the synthesis of a series of $\text{Pd}^{\text{II}}\text{--CF}_3$ complexes of general structure $(\text{dtbpy})\text{Pd}^{\text{II}}(\text{Aryl})(\text{CF}_3)$ (**1a–i**, dtbpy = 4,4'-di-*tert*-butyl-2,2'-bipyridine). These compounds were prepared by the treatment of $(\text{dtbpy})\text{Pd}^{\text{II}}(\text{Aryl})(\text{I})$ with CsF followed by TMSCF_3 in THF at 23 °C (Table 1). The products were isolated as yellow solids in 32-70% yield.²¹ X-ray quality crystals of **1a** were obtained by vapor diffusion of pentanes into a dichloromethane solution of **1a**, and the X-ray crystal structure of this complex is shown in Figure 1.

The Pd^{II} complexes (**1a–i**) are inert towards direct Aryl– CF_3 bond-forming reductive elimination. For example, heating **1a** at 130 °C for 72 h produced <5% of 4-fluorobenzotrifluoride (**2a**), and the Pd^{II} starting material could be recovered in >80% yield (eq. 4). Importantly, similarly low reactivity has been reported in the literature for other $(\text{L}\sim\text{L})\text{Pd}^{\text{II}}(\text{Aryl})(\text{CF}_3)$ complexes ($\text{L}\sim\text{L}$ = diphenylphosphinoethane, diphenylphosphinopropane, and diphenylphosphinobenzene).^{24a,b} As discussed above, the only demonstrated examples of Aryl– CF_3 bond-forming reductive elimination from Pd^{II} require relatively high temperatures (80 °C) and specialized phosphine ligands.^{14,15a}




We reasoned that the $2e^-$ oxidation of $(\text{dtbpy})\text{Pd}^{\text{II}}(\text{Aryl})(\text{CF}_3)$ would yield a highly reactive Pd^{IV} adduct that might undergo more facile Aryl– CF_3 bond-forming reductive elimination (eq. 5, *i*). Thus, we examined the reaction of **1a** with *N*-bromosuccinimide (NBS), *N*-chlorosuccinimide (NCS), and $\text{PhI}(\text{OAc})_2$, which are all well-known to promote the oxidation of Pd^{II} to Pd^{IV} .²² As shown in Table 2, all three oxidants reacted rapidly with **1a** in nitrobenzene- d_5 at 80 °C. However, the desired trifluoromethylated product was not obtained; instead, the major organic product contained a nucleophile derived from the oxidant (Br, Cl, or OAc, respectively). These results are consistent with the formation of a Pd^{IV} intermediate, from which Aryl–X (X = OAc, Br, and Cl) bond-forming reductive elimination is significantly faster than Aryl– CF_3 coupling (eq. 5, *ii*).²¹



In an effort to avoid competing reductive elimination processes, we next examined the use of electrophilic fluorinating reagents (F^+ sources). These reagents were selected based on the hypothesis that fluoride (the X-type ligand introduced to Pd^{IV} by F^+ sources) might undergo slower reductive elimination than CF_3 .^{22d,27,28} Gratifyingly, a variety of different F^+ reagents reacted with **1a** to afford modest to excellent yields of the trifluoromethylated product **2a** after 3 h at 80 °C (Table 2, entries 4-10). The optimal electrophilic fluorinating reagent was *N*-fluoro-1,3,5-trimethylpyridium triflate (NFTPT), which provided **2a** in 70% yield as determined by ^{19}F NMR spectroscopy.²⁹ Importantly, <5% of products derived from Aryl-F or Aryl-OTf coupling were observed under these conditions.

This transformation was next applied to complexes **1b-i**, which contain sterically and electronically diverse aryl groups. As shown in Table 3, the yield of trifluoromethylated product was relatively insensitive to the electronic properties of the arene, and the reaction proceeded with good results in systems containing both electron withdrawing [*e.g.*, CN, C(O)Ph] and electron donating (*e.g.*, CH_3 , OCH_3) *para*- and *meta*-substituents.³⁰

At room temperature, the oxidation of **1a** by NFTPT produced a single major intermediate. This species (**4**) was isolated from dichloroethane as a yellow solid in 53% yield (eq 6). Analysis of **4** by ^{19}F NMR spectroscopy in $MeCN-d_3$ showed four characteristic resonances: a doublet at -30.9 ppm (Pd- CF_3), a singlet at -79.4 ppm (Pd-OTf), a multiplet at -117.1 ppm (Pd-ArF), and a quartet at -256.5 ppm (Pd-F) in a 3: 3: 1: 1 ratio. In nitrobenzene- d_5 , broad resonances were observed with similar chemical shifts in the same 3: 3: 1: 1 ratio.³¹ The 1H NMR spectrum of **4** showed two singlets at 8.47 and 8.40 ppm (5,5' protons of dtbpy) as well as two singlets at 1.49 and 1.41 ppm (*t*Bu groups of dtbpy).




X-ray quality crystals were obtained via vapor diffusion of pentanes into a DCE solution of **4**. As shown in Figure 2, the solid state structure of **4** shows the octahedral Pd^{IV} species (dtbpy) $Pd(p-FC_6H_4)(CF_3)(F)(OTf)$. Interestingly, the Pd- CF_3 bond distance in **4** (2.009(4) Å) is nearly identical to that of the Pd^{II} - CF_3 starting material **1a** (2.005(3) Å). However, the Pd-N bond lengths of **4** (2.038(4) and 2.082(4) Å) are significantly shorter than those in **1a** (2.107(2) and 2.143(3) Å).

Mechanistic Study of Aryl- CF_3 Bond-Forming Reductive Elimination from Pd^{IV} .

There are at least three possible pathways for Aryl- CF_3 bond-forming reductive elimination from **4**, mechanisms **A**, **B**, and **C** (Figure 3). Mechanism **A** involves initial triflate dissociation to form a cationic five-coordinate Pd^{IV} intermediate **I** and subsequent Aryl- CF_3 bond formation. Mechanism **B** involves fluoride dissociation, followed by Aryl- CF_3 coupling from intermediate **II**. Mechanism **C** proceeds via concerted C- CF_3 bond-forming reductive elimination. Notably, there is significant literature precedent for both ionic^{22e, Error! Bookmark not defined.,32,33} and concerted^{22e,34} reductive elimination mechanisms from octahedral group 10 metal complexes.

Order in triflate.—The addition of 1 equiv of NBu₄OTf to the thermolysis of **4** in NO₂Ph-*d*₅ at 50 °C significantly slowed the initial rate of formation of **2a** (from $2.21 \times 10^{-5} \text{ M s}^{-1}$ to $1.35 \times 10^{-5} \text{ M s}^{-1}$).³⁵ Furthermore, an excellent linear fit was observed for a plot of initial rate versus 1/[NBu₄OTf] (Figure 4).

In order to confirm that the presence of TfO[−] was responsible for the observed effect, this reaction was next examined in the presence of NBu₄PF₆, which contains a non-coordinating anion. The use of 1 equiv of NBu₄PF₆ under otherwise identical conditions resulted in a >2-fold increase in the initial rate of reductive elimination to $5.57 \times 10^{-5} \text{ M s}^{-1}$. This result indicates that inhibition by NBu₄OTf is specifically due to the TfO[−] anion. Furthermore, it suggests that enhancing the polarity of the medium (by adding non-coordinating ions) accelerates Aryl–CF₃ reductive elimination. Both pieces of data are consistent with ionic mechanism **A** operating in this system, as reflected by the rate expression in eq. 7.³⁶




Activation parameters.—The initial rate of C–CF₃ bond-forming reductive elimination from **4** in NO₂Ph-*d*₅ was next examined as a function of temperature. An Eyring plot of the data showed that ΔH^\ddagger is $+29.1 \pm 0.2 \text{ kcal/mol}$, while $\Delta S^\ddagger = +9.48 \pm 0.8 \text{ eu}$. The observed entropy of activation indicates a significant increase in disorder in the transition state for this reductive elimination reaction. Similar values have been observed for carbon-heteroatom bond-forming reductive elimination reactions from Pd^{IV} that proceed by ionic mechanisms.^{22e,23e,37} For example, C–OAc reductive elimination from Pd^{IV} complex (phpy)₂Pd^{IV}(OAc)₂ (phpy = 2-phenylpyridine), which is proposed to involve pre-equilibrium dissociation of an acetate ligand, showed ΔS^\ddagger of $+4.2 \pm 1.2 \text{ eu}$.^{22e}

Electronic effects.—Finally, electronic effects on reductive elimination were evaluated using Pd^{IV} complexes containing *p*-fluoro, *p*-methyl, and *p*-trifluoromethyl-substituted aryl groups (complexes **4–6**). Thermolysis of **4**, **5**, and **6** at 80 °C in NO₂Ph-*d*₅ for 3 h afforded the corresponding benzotrifluorides in 77%, 93%, and 65% yield, respectively (Table 4). The initial rates of reductive elimination from these complexes at 50 °C in NO₂Ph-*d*₅ show that electron donating substituents accelerate the reaction. For example, with X = CH₃ (**5**), the initial rate is more than 20 times greater than with X = F (**4**). Additionally, when X = CF₃ (**6**), the initial rate is nearly two times slower than from **4** (Table 4).

While Table 4 clearly shows that electron-donating aryl substituents accelerate this reaction, it is challenging to definitively interpret this data in the context of the C–CF₃ bond-forming event. Mechanism **A** is a two-step process involving triflate dissociation followed by Aryl–CF₃ coupling. Thus, the faster rate with complex **5** may be due to the stronger *trans* effect of a more electron donating σ -aryl ligand and/or from a partial positive charge buildup on the aromatic ring in the transition state for Aryl–CF₃ coupling.

DFT calculations.—We next turned to DFT calculations to more fully explore the mechanism of Aryl–CF₃ bond formation. The complex [(bpy)Pd^{IV}(Ph)(CF₃)(F)(OTf)] (**7**)

(bpy = 2,2'-bipyridyl, eq. 8) was employed as a model for **5** (Table 4), and the CEP-31G(d)^{38,39} basis set and M06^{40,41} functional were used along with a single point solvent correction in nitrobenzene (SMD solvation model).⁴² As shown in eq. 8, the calculated ΔH_{298} for loss of TfO⁻ from **7** in nitrobenzene to form cationic intermediate $[(\text{bpy})\text{Pd}^{\text{IV}}(\text{Ph})(\text{CF}_3)(\text{F})]^+$ (**8**) is 15.0 kcal/mol. Furthermore, the activation enthalpy (ΔH_{298}^\ddagger) for Ph–CF₃ bond-forming reductive elimination from **8** is 13.7 kcal/mol. Thus, we calculate that mechanism **A** has an overall ΔH_{298}^\ddagger of 28.7 kcal/mol, which is in excellent agreement with the experimental value ($+29.1 \pm 0.2$ kcal/mol, *vide supra*).




As discussed above, we observe experimentally that trifluoromethylated products are formed selectively over the corresponding fluorinated compounds. To gain further insights into this selectivity, we used DFT to examine the transition state for Ph–F bond-forming reductive elimination from intermediate **8**. As shown in eq. 8, ΔH_{298}^\ddagger (DFT) from Ph–F coupling is 14.7 kcal/mol. As such, this is a higher energy pathway than Ph–CF₃ bond-formation (ΔH_{298}^\ddagger (DFT) = 1 kcal/mol), consistent with the experimental results.

The calculated charge distribution of intermediate **8** using Natural Bond Order (NBO) analysis^{43,44} indicates that the CF₃ carbon carries a significant positive charge (+1.18), while the α -carbon of the phenyl ligand bears a charge of +0.07.⁴⁵ This charge difference is amplified in the transition state for C–CF₃ bond-forming reductive elimination, where the CF₃ carbon carries an enhanced positive charge of +1.24, and the charge on the Ph carbon decreases to –0.11. These data imply that the Ph group is acting as the nucleophile during the bond-forming event. This is in sharp contrast to other reports of reductive elimination from Pd^{IV} in which the aryl or alkyl ligand typically serves as the electrophilic coupling partner.^{22,23}

We next used DFT to determine the transition state enthalpies (ΔH_{298}^\ddagger) for Aryl–CF₃ coupling from complexes of general structure $[(\text{bpy})\text{Pd}^{\text{IV}}(p\text{-XC}_6\text{H}_4)(\text{CF}_3)(\text{F})]^+$. As expected based on the NBO analysis, ΔH_{298}^\ddagger was smallest with electron donating *para* substituents (X) on the aromatic ring, consistent with a transition state involving nucleophilic attack by σ -aryl ligand on the CF₃ moiety (Table 5). Furthermore, the ΔH_{298}^\ddagger values showed better correlation with Hammett σ^+ values for X than the corresponding σ or σ^- parameters. This implicates significant resonance effects in the transition state for Aryl–CF₃ coupling.⁴⁶

Finally, we sought to identify supporting ligands that would lower the energy barrier for Ph–CF₃ bond-forming reductive elimination in this system. Literature studies have shown that (tmeda)Pd^{IV}(CH₃)₂(Ph)(I) (tmeda = tetramethylethylenediamine) is significantly more reactive towards C–C bond-forming reductive elimination than its bipyridine analogue (bpy)Pd^{IV}(CH₃)₂(Ph)(I)⁴⁷. Thus, we hypothesized that using tmeda in place of dtbpy in our system might impart a similar effect. Consistent with this hypothesis, DFT calculations show that the nitrobenzene solvent-corrected transition state enthalpy (ΔH_{298}^\ddagger) for formation of trifluorotoluene from $[(\text{tmeda})\text{Pd}^{\text{IV}}(\text{Ph})(\text{CF}_3)(\text{F})]^+$ is 0.8 kcal/mol lower than that for the analogous bpy complex (12.9 versus 13.7 kcal/mol) (eq. 9). Furthermore, ΔH_{298} for the loss of OTf⁻ from $[(\text{tmeda})\text{Pd}^{\text{IV}}(\text{Ph})(\text{CF}_3)(\text{F})(\text{OTf})]$ is >10 kcal/mol lower than that from **7** (4.2 versus 15 kcal/mol). Thus, the calculated overall ΔH_{298}^\ddagger for reductive

elimination from the tmeda complex is 17.1 kcal/mol, suggesting that Aryl-CF₃ coupling should proceed at significantly lower temperatures than from **7**.



Room Temperature Arene Trifluoromethylation.

To assess the effect of diamine ligands experimentally, a series of (N~N)Pd^{II}(Ph)(CF₃) complexes (**9**, **10**, and **11e**) were prepared by the reaction of (N~N)Pd^{II}(Ph)(I) with CsF followed by TMSCF₃ (see Supporting Information for full details).⁴⁸ Treatment of **9**, **10**, and **11e** with NFTPT under our standard conditions (80 °C for 3 h in NO₂Ph) resulted in clean formation of trifluorotoluene in modest to excellent yield (Table 6). The readily available and inexpensive tmeda ligand was particularly effective. Tmeda complex **11e** afforded 90% yield of trifluorotoluene at 80 °C (entry 4), and, most remarkably, provided 83% yield *at room temperature* (entry 5). To our knowledge, this is first example of room temperature arene trifluoromethylation at a Pd center.⁴⁹

The scope of room temperature oxidative trifluoromethylation from (tmeda)Pd^{II}(Aryl)(CF₃) was found to be quite broad.⁵⁰ These reactions proceeded efficiently and in high yield with electron donating and electron neutral substituents on the σ-aryl ligand (for example, Table 7, entries 4-7). The room temperature reactions were lower yielding with arenes bearing highly electron-withdrawing substituents like CF₃ and CN (entries 2 and 3), which is consistent with the proposed transition state (*vide supra*). Finally, tmeda complexes containing *ortho*-substituted aryl groups underwent high yielding room temperature oxidative trifluoromethylation (entries 8 and 9). In contrast, these were poorly effective substrates in the dtbpa system, even at 80 °C.³⁰

Conclusion.

This article describes the rational design of 1st generation systems for oxidatively-induced Aryl-CF₃ bond forming reductive elimination from Pd. Experimental mechanistic studies implicate Aryl-CF₃ coupling from a cationic five-coordinate intermediate, and DFT suggests that the CF₃ ligand serves as the electrophilic partner during bond formation. Our investigations into the scope and mechanism of this reaction have facilitated the development of a 2nd generation ligand system that enables Aryl-CF₃ coupling *at room temperature*. This work provides the basis for the design of novel Pd^{II/IV}-catalyzed trifluoromethylation reactions of aryl metal species (metal = B, Sn, Si) or simple arene C-H bonds. Efforts in this area are currently underway in our group and will be reported in due course.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank the NIH [GM073836 and F31GM089141 (fellowship to NDB)], the National Science Foundation Graduate Research Fellowship and Murrill Memorial Scholarship (fellowship to JBG) and the Research Corporation Cottrell Scholar Program for support of this research. Additional unrestricted support from Dupont is also gratefully acknowledged. We also thank Paul Lennon and Jim Windak for assistance with mass spectroscopy as well as Prof. Tom Cundari and Prof. Brian Yates for valuable discussions on DFT calculations.


References

1. Kirk KL. *Org. Process Res. Dev.* 2008; 12:305–321.
2. Purser S, Moore PR, Swallow S, Gouverneur V. *Chem. Soc. Rev.* 2008; 37:320–330. [PubMed: 18197348]
3. (a) Ma JA, Cahard D. *Chem. Rev.* 2004; 104:6119–6146. [PubMed: 15584697] (b) Ma JA, Cahard D. *Chem. Rev.* 2008; 108:PR1–PR43. [PubMed: 18798358] (c) Prakash GKS, Chacko S. *Curr. Opin. Drug Disc. Dev.* 2008; 11:793–802. (d) Shibata N, Mizuta S, Kawai H. *Tetrahedron: Asymmetry.* 2008; 19:2633–2644.
4. Grushin VV. *Acc. Chem. Res.* 2010; 43:160–171. [PubMed: 19788304]
5. Swarts F. *Bull. Acad. R. Belg.* 1892; 24:415–429.
6. (a) Henne AL, Whaley AM, Stevenson JK. *J. Am. Chem. Soc.* 1941; 63:3478–3479. (b) Furuta S, Kuroboshi M, Hiyama T. *Bull. Chem. Soc. Jpn.* 1999; 72:805–819.
7. Yang JJ, Kirchmeier RL, Shreeve JM. *J. Org. Chem.* 1998; 63:2656–2660. [PubMed: 11672133]
8. For examples of Cu-mediated reaction of TMSCF_3 with aryl halides, see: (a) Kobayashi Y, Kumadaki I. *Tetrahedron Lett.* 1969; 10:4095–4096. (b) Konderatenko NV, Vechirko EP, Yagupolskii LM. *Synthesis.* 1980:932–933. (c) Matsui K, Tobita E, Ando M, Kondo K. *Chem. Lett.* 1981; 10:1719–1720. (d) Suzuki H, Yoshida Y, Osuka A. *Chem. Lett.* 1982; 11:135–136. (e) Burton DJ, Wiemers DM. *J. Am. Chem. Soc.* 1985; 107:5014–5015. (f) Urata H, Fuchikami T. *Tetrahedron Lett.* 1991; 32:91–94. (g) Dubinina GG, Furutachi H, Vivic DA. *J. Am. Chem. Soc.* 2008; 130:8600–8601. [PubMed: 18543912] (h) Dubinina GG, Ogikubo J, Vivic DA. *Organometallics.* 2008; 27:6233–6235.
9. For the Cu-mediated reaction of TMSCF_3 with aryl boronic acids, see (a) Chu L, Qing F-L. *Org. Lett.* 2010; 12:5060–5063. [PubMed: 20923196] (b) Senecal TD, Parsons AT, Buchwald SL. *J. Org. Chem.* 2011; 76:1174–1176.
10. Zhang C-P, Wang Z-L, Chen Q-Y, Zhang C-T, Gu Y-C, Xiao J-C. *Angew. Chem. Int. Ed.* 2011; 50:1896–1900.
11. (a) Oishi M, Kondo H, Amii H. *Chem. Commun.* 2009:1909–1911. (b) Knauber T, Arikan F, Röschenhaler G, Gooßen LJ. *Chem. Eur. J.* 2011; 17:2689–2697.
12. For recent reviews see: (a) Hassan J, Sévignon M, Gozzi C, Schulz E, Lemaire M. *Chem. Rev.* 2002; 102:1359–1470. [PubMed: 11996540] (b) Beccalli EM, Broggin G, Martinelli M, Sottocornola S. *Chem. Rev.* 2007; 107:5318–5365. [PubMed: 17973536] (c) Chen X, Engle KM, Wang D-H, Yu J-Q. *Angew. Chem., Int. Ed.* 2009; 48:5094–5115. (d) McGlacken GP, Bateman LM. *Chem. Soc. Rev.* 2009; 38:2447–2464. [PubMed: 19623360] (e) Daugulis O, Do H-Q, Shabashov D. *Acc. Chem. Res.* 2009; 42:1074–1086. [PubMed: 19552413]
13. For reviews on $[\text{M}]-\text{CF}_3$ and $[\text{M}]-\text{R}_f$ (R_f = perfluoroalkyl) complexes, see: (a) Hughes RP. *Adv. Organomet. Chem.* 1990; 31:183–267. (b) Morrison JA. *Adv. Organomet. Chem.* 1993; 35:211–239.
14. Grushin VV, Marshall WJ. *J. Am. Chem. Soc.* 2006; 128:12644–12645. [PubMed: 17002347]
15. Cho EJ, Senecal TD, Kinzel T, Zhang Y, Watson DA, Buchwald SL. *Science.* 2010; 328:1679–1681. [PubMed: 20576888] Pd-catalyzed coupling between ArI and $[\text{Zn}-\text{CF}_3]$: Kitazume T, Ishikawa N. *Chem. Lett.* 1982:137–140.
16. Current price for Xantphos = \$18,122/mol. Determined based on the largest quantity of Xantphos available from Sigma-Aldrich on March 17, 2011 (25 g/\$783.00).
17. Current price for Brettphos = \$99,195/mol. Determined based on the largest quantity of Brettphos available from Strem Chemicals on March 17, 2011 (5 g/\$924.00).

18. Price for the largest quantity of TESC_F₃ available from Sigma-Aldrich on March 17, 2011: 1 g/\$72.90, \$13,433/mol. Price for the largest quantity of TMSC_F₃ available from Sigma-Aldrich on March 17, 2011: 25 mL/\$397.00, \$2347/mol.
19. Wang X, Truesdale L, Yu JQ. *J. Am. Chem. Soc.* 2010; 132:3648–3649. [PubMed: 20184319]
20. Ye Y, Ball ND, Kampf JW, Sanford MS. *J. Am. Chem. Soc.* 2010; 132:14682–14687. [PubMed: 20866077]
21. Ball ND, Kampf JW, Sanford MS. *J. Am. Chem. Soc.* 2010; 132:2878–2879. [PubMed: 20155920]
22. (a) Dick AR, Kampf JW, Sanford MS. *J. Am. Chem. Soc.* 2005; 127:12790–12791. [PubMed: 16159259] (b) Whitfield SR, Sanford MS. *J. Am. Chem. Soc.* 2007; 129:15142–15143. [PubMed: 18004863] (c) Dick AR, Remy MS, Kampf JW, Sanford MS. *Organometallics.* 2007; 26:1365–1370. (d) Ball ND, Sanford MS. *J. Am. Chem. Soc.* 2009; 131:3796–3797. [PubMed: 19249867] (e) Racowski JM, Dick AR, Sanford MS. *J. Am. Chem. Soc.* 2009; 131:10974–10983. [PubMed: 19459631] (f) Arnold PL, Sanford MS, Pearson SM. *J. Am. Chem. Soc.* 2009; 131:13912–13913. [PubMed: 19788324]
23. For examples, see: (a) Alsters PL, Engel PF, Hogerheide MP, Copijn M, Spek AL, van Koten G. *Organometallics.* 1993; 12:1831–1844. (b) Lagunas MC, Gossage RA, Spek AL, van Koten G. *Organometallics.* 1998; 17:731–741. (c) van Belzen R, Elsevier CJ, Dedieu A, Veldman N, Spek AL. *Organometallics.* 2003; 22:722–736. (d) Canty AJ, Denny MC, Patel J, Sun H, Skelton BW, White AH. *J. Organomet. Chem.* 2004; 689:672–677. (e) Canty AJ, Denney MC, Skelton BW, White AH. *Organometallics.* 2004; 23:1122–1131. (f) Yamamoto Y, Kuwabara S, Matsuo S, Ohno T, Nishiyama H, Itoh K. *Organometallics.* 2004; 23:3898–3906. (g) Kaspi AW, Yahav-Levi A, Goldberg I, Vigalok A. *Inorg. Chem.* 2008; 47:5–7. [PubMed: 18052157] (h) Furuya T, Benitez D, Tkatchouk E, Strom AE, Tang P, Goddard WA III, Ritter T. *J. Am. Chem. Soc.* 2010; 132:3793–3807. [PubMed: 20196595] (i) Vicente J, Arcas A, Julia-Hernandez F, Bautista D. *Chem. Comm.* 2010; 46:7253–7255. [PubMed: 20740248] (j) Williamson O, Zavalij PY, Zhang J, Khaskin E, Vedernikov AN. *J. Am. Chem. Soc.* 2010; 132:14400–14402. [PubMed: 20866056] (k) Zhao X, Dong VM. *Angew. Chem. Int. Ed.* 2011; 50:932–934.
24. (a) Culkin DA, Hartwig JF. *Organometallics.* 2004; 23:3398–3416. (b) Grushin VV, Marshall WJ. *J. Am. Chem. Soc.* 2006; 128:4632–4641. [PubMed: 16594700] (c) McReynolds KA, Lewis RS, Ackerman LKG, Dubinina GG, Brennessel WW, Vici DA. *J. Fluorine Chem.* 2010; 131:1108–1112.
25. (a) Umemoto T. *Chem. Rev.* 1996; 96:1757–1778. [PubMed: 11848810] (b) Eisenberger P, Gischig S, Togni A. *Chem. Eur. J.* 2006; 12:2579–2586. (c) Kieltch I, Eisenberger P, Stanek K, Togni A. *Chimia.* 2008; 62:260–263.
26. Stanek K, Koller R, Togni A. *J. Org. Chem.* 2008; 73:7678–7685. [PubMed: 18771328]
27. For similar strategies to generate C–N or C–C bonds using F⁺ sources see: (a) Mei T-S, Wang X, Yu J-Q. *J. Am. Chem. Soc.* 2009; 131:10806–10807. [PubMed: 19606861] (b) Sibbald PA, Rosewall CF, Swartz RD, Michael FE. *J. Am. Chem. Soc.* 2009; 131:15945–15951. [PubMed: 19824646]
28. Engle KM, Mei TS, Wang X, Yu J-Q. *Angew. Chem. Int. Ed.* 2011; 50:1478–1491.
29. The reported yields in Tables 1 and 2 are ~10% lower than reported in our original communication (ref. 21), due to a minor error associated with the quantity of internal standard used for ¹⁹F NMR analysis. The correct yields for these transformations are reported in Table 3.
30. (dtbpy)Pd^{II}(Aryl)(CF₃) complexes containing *ortho*-substituted Aryl ligands reacted with NFTPT to provide very low yields of the corresponding benzotrifluorides. For example (dtbpy)Pd(*o*-CH₃Ph)(CF₃) afforded 13% yield of 2-methylbenzotrifluoride. Interestingly, the major product of this reaction was the corresponding fluorinated product (2-fluorotoluene, formed in 62% yield).
31. In MeCN-*d*₃, it is likely that the triflate ligand is replaced by a solvent molecule.
32. For examples of mechanisms like **A** and **B** for reductive elimination from Pd^{IV}, see: (a) Byers PK, Canty AJ, Crespo M, Puddephatt RJ, Scott JD. *Organometallics.* 1988; 7:1363–1367. (b) Ducker-Benfer C, van Eldik R, Canty AJ. *Organometallics.* 1994; 13:2412–2414. (c) Canty AJ, Jin H, Skelton BW, White AH. *Inorg. Chem.* 1998; 37:3975–3981. [PubMed: 11670512]
33. For examples of mechanisms similar to **A** and **B** for reductive elimination from Pt^{IV}, see: (a) Goldberg KI, Yan J, Winter EL. *J. Am. Chem. Soc.* 1994; 116:1573–1574. (b) Goldberg KI, Yan J, Breitung EM. *J. Am. Chem. Soc.* 1995; 117:6889–6896. (c) Williams BS, Goldberg KI. *J. Am.*

- Chem. Soc. 2001; 123:2576–2587. [PubMed: 11456927] (d) Vedernikov AN, Binfield SA, Zavalij PY, Khusnutdinova JR. J. Am. Chem. Soc. 2006; 128:82–83. [PubMed: 16390132] (e) Khusnutdinova JR, Zavalij PY, Vedernikov AN. Organometallics. 2007; 26:3466–3483. (f) Pawlikowski AV, Getty AD, Goldberg KI. J. Am. Chem. Soc. 2007; 129:10382–10393. [PubMed: 17672451] (g) Khusnutdinova JR, Newman LL, Zavalij PY, Lam Y-F, Vedernikov AN. J. Am. Chem. Soc. 2008; 130:2174–2175. [PubMed: 18220400] (h) Smythe NA, Grice KA, Williams BS, Goldberg KI. Organometallics. 2009; 28:277–288.
34. For examples of mechanisms similar to **C** for reductive elimination from Pd^{IV} and Pt^{IV}, see: (a) Crumpton DM, Goldberg KI. J. Am. Chem. Soc. 2000; 122:962–963. (b) Crumpton-Bregel DM, Goldberg KI. J. Am. Chem. Soc. 2003; 125:9442–9456. [PubMed: 12889975] (c) Arthur KL, Wang QL, Bregel DM, Smythe NA, O'Neill BA, Goldberg KI, Moloy KG. Organometallics. 2005; 24:4624–4628. (d) Reference 22e.
35. We initially examined Aryl–CF₃ bond-forming reductive elimination by monitoring the disappearance of **4** and concomitant appearance of **2a** over three half lives. These studies revealed that the rate of product formation decreases upon higher conversion to product (see Supporting Information for full details). We hypothesize that this is due to a build-up of TfO[−] as the reaction progresses, which then inhibits the 1st step of Aryl–CF₃ bond-forming reductive elimination from **4**. To avoid this apparent product inhibition, we used the initial rates method (measuring the first 10% conversion) for both the order and activation parameter studies.
36. The addition of 1 equiv of NMe₄F to **4** at 23 °C in NO₂Ph-*d*₅ produced a complex mixture of Pd–F and Pd–CF₃ containing products (as determined by ¹⁹F NMR spectroscopy). As a result, it was not possible to definitively assess the effect of F[−] on the rate of reductive elimination from **4**.
37. The entropy of activation for reductive elimination reactions from M^{IV} can be highly dependent on both the structure of the M^{IV} complex and the solvent. This is particularly the case in ionic reductive elimination reactions, where solvent ordering around charged species can have a significant effect. See ref. 32a.
38. Stevens WJ, Basch H, Krauss MJ. J. Chem. Phys. 1984; 81:6026–6033.
39. Stevens WJ, Krauss M, Basch H, Jasien PG. Can. J. Chem. 1992; 70:612–630.
40. Zhao Y, Truhlar DG. Theor. Chem. Acc. 2008; 120:215–241.
41. The M06 functional was selected based its success in DFT calculations of related late transition metal organometallic complexes. For examples, see: (a) Sieffert N, Bühl M. Inorg. Chem. 2009; 48:4622–4624. [PubMed: 19382761] (b) Benitez D, Tkatchouk E, Goddard WA III. Organometallics. 2009; 28:2643–2645. (c) Benitez D, Shapiro ND, Tkatchouk E, Wang YM, Goddard WA III, Toste FD. Nat. Chem. 2009; 1:482–486. [PubMed: 20161015] (d) Ariafard A, Hyland CJT, Canty AJ, Sharma M, Brookes NJ, Yates BF. Inorg. Chem. 2010; 49:11249–11253. [PubMed: 21043467]
42. Marenich AV, Cramer CJ, Truhlar DG. J. Phys. Chem. B. 2009; 113:6378–6396. [PubMed: 19366259]
43. Glendening, ED.; Reed, AE.; Carpenter, JE.; Weinhold, F. NBO Version 3.1.
44. For original references of theory, see: (a) Foster JP, Weinhold F. J. Am. Chem. Soc. 1980; 102:7211–7218. (b) Reed AE, Weinhold F. J. Chem. Phys. 1983; 78:4066–4073. (c) Reed AE, Weinstock RB, Weinhold F. J. Chem. Phys. 1985; 83:735–746. (d) Reed AE, Weinhold F. J. Chem. Phys. 1985; 83:1736–1740. (e) Carpenter JE. Extension of Lewis structure concepts to open-shell and excited-state molecular species. PhD thesis. 1987 University of Wisconsin-Madison, WI (f) Carpenter JE, Weinhold F. J. Mol. Struct. (Theochem). 1988; 46:41–62. (g) Reed AE, Curtiss LA, Weinhold F. Chem. Rev. 1988; 88:899–926. (h) Weinhold F, Carpenter JE, Naaman R, Vager Z. The Structure of Small Molecules and Ions. 1988:227–236. Plenum New York
45. The ground state structure of the analogous CH₃ complex ([{(bpy)Pd^{IV}(Ph)(CH₃)(F)]⁺ has calculated charges of −0.58 and +0.06 on the CH₃ and Ph carbons, respectively.
46. Assuming a similar entropy component in each transition state, a Hammett plot of log(*k*_X/*k*_H) versus σ⁺ with rate constants calculated based on the electronic energies (Δ*H*[‡]₂₉₈), provided a reasonably good linear correlation (ρ⁺ = −0.79; R² = 0.84, Figure S8). Much poorer correlations were observed with σ and σ[−] (ρ value of −1.17 and R² = 0.57; ρ[−] value of −0.86 and R² = 0.52).
47. Markies BA, Canty AJ, Boersma J, van Koten G. Organometallics. 1994; 13:2053–2058.

48. Complex **11b** can also be prepared by reacting $(\text{tmeda})\text{Pd}^{\text{II}}(\text{p-CF}_3\text{C}_6\text{H}_4)(\text{X})$ ($\text{X} = \text{acetate}$ or trifluoroacetate) with $\text{CsF}/\text{TMSCF}_3$ in THF. This synthetic pathway is potentially useful and relevant for catalytic applications of these transformations. For example, C–H functionalization reactions often use $\text{Pd}(\text{OAc})_2$ or $\text{Pd}(\text{TFA})_2$ as catalysts. For examples, see: Lyons TW, Sanford MS. *Chem. Rev.* 2010; 110:1147–1169. [PubMed: 20078038]



49. Vivic has reported that the reaction of $(\text{NHC})\text{Cu}^{\text{I}}(\text{CF}_3)$ ($\text{NHC} = \text{heterocyclic carbene}$) with aryl iodides provides Aryl– CF_3 products at room temperature. The mechanism of the transformation is still under investigation. See refs 8g and h.
50. For the synthesis of these complexes, see Supporting Information.




Figure 1. ORTEP drawing of complex **1a**. Thermal ellipsoids are drawn at 50% probability, hydrogen atoms are omitted for clarity. Selected bond lengths (Å): Pd–C(1) 2.005(3), Pd–C(25) 2.007(4), Pd–N(1) 2.107(2), Pd–N(2) 2.143(3). Selected bond angles (°): C(1)–Pd–C(25) 89.99(13), C(1)–Pd–N(1) 95.58(11), C(1)–Pd–N(2) 169.99(11), C(25)–Pd–N(1) 173.92(12), C(25)–Pd–N(2) 96.67(11).




Figure 2.

ORTEP drawing of complex **4**. Thermal ellipsoids are drawn at 50% probability, hydrogen atoms are omitted for clarity. Selected bond lengths (Å): Pd–C(1) 2.009(5), Pd–C(25) 2.018(5), Pd–N(1) 2.038(4), Pd–N(15) 2.082(4), Pd–O(1) 2.226(3). Selected bond angles (°): C(19)–Pd–C(25) 91.09(15), C(19)–Pd–N(15) 92.18(16), C(25)–Pd–N(15) 175.68(17), C(19)–Pd–F(1) 91.09(15), C(25)–Pd–F(1) 83.31(16), C(19)–Pd(1)–O(1) 175.74 (16), C(25)–Pd(1)–O(1) 95.15 (16).




Figure 3. Three potential mechanisms for Aryl-CF₃ bond formation from **4**.





Figure 4. Plot of initial rate versus $1/[\text{OTf}]$ for reductive elimination from **4** to form **2a** in $\text{PhNO}_2\text{-}d_5$ at $50\text{ }^\circ\text{C}$. $y = 1.91 \times 10^{-7} + 9.86 \times 10^{-6}x$; $R^2 = 0.998$.


Table 1

Synthesis of Complexes **1a-i**





[L~L = dtbpy]

Entry	Aryl	Compound	Isolated Yield
1	<i>p</i> -FC ₆ H ₄	1a	70%
2	<i>p</i> -CNC ₆ H ₄	1b	54%
3	<i>p</i> -CF ₃ C ₆ H ₄	1c	63%
4	<i>p</i> -PhC(O)C ₆ H ₄	1d	51%
5	<i>p</i> -PhC ₆ H ₄	1e	58%
6	<i>p</i> -MeOC ₆ H ₄	1f	32%
7	<i>p</i> -MeC ₆ H ₄	1g	49%
8	C ₆ H ₅	1h	47%
9	<i>m</i> -MeC ₆ H ₄	1i	42%

Table 2

Reaction of **1a** with Diverse Oxidants (Oxidant-X)


Entry	Oxidant-X	Yield 2a	Yield 3a (X)
1		<5%	75% (Br)
2		<5%	70% (Cl)
3		<5%	20% (OAc)
4		<5%	nd (F) ^a
5		60%	<5%
6		53%	<5%

Entry	Oxidant-X	Yield 2a	Yield 3a (X)
7		69%	<5%
8		55%	<5%
9		68%	<5%
10		70%	<5%
11	XeF ₂	65%	5%

Yields were determined by ¹⁹F NMR spectroscopy and are an average of two runs. nd = not detected.


³**1a** accounted for the remaining mass balance.

Table 3

NFTPT-Promoted Aryl-CF₃ Coupling at Complexes **1a-i**

Entry	Compound	Aryl	Yield Aryl-CF ₃
1	1a	<i>p</i> -FC ₆ H ₄	70%
2	1b	<i>p</i> -CNC ₆ H ₄	25%
3	1c	<i>p</i> -CF ₃ C ₆ H ₄	55%
4	1d	<i>p</i> -PhC(O)C ₆ H ₄	56%
5	1e	<i>p</i> -PhC ₆ H ₄	70%
6	1f	<i>p</i> -MeOC ₆ H ₄	72%
7	1g	<i>p</i> -MeC ₆ H ₄	66%
8	1h	C ₆ H ₅	70%
9	1i	<i>m</i> -MeC ₆ H ₄	64%

Yields were determined by ¹⁹F NMR spectroscopy and are an average of two runs.

Table 4Initial Rates of Aryl-CF₃ Bond-Forming Reductive Elimination from Complexes **4-6**


Entry	X	Yield Aryl-CF ₃	Initial rate (M s ⁻¹) × 10 ⁵
1	CH ₃	93%	45.8
2	F	77%	2.21
3	CF ₃	65% ^a	1.43

Reactions were run in duplicate and yields were determined by ¹⁹F NMR spectroscopy.

^a1,4(bistrifluoromethyl)biphenyl formed in 21% yield.

Table 5


Gas Phase Values of ΔH_{298}^\ddagger for C–CF₃ Bond-Formation from [(bpy)Pd^{IV}(*p*-XC₆H₄)(CF₃)(F)]⁺^[a]



X	ΔH_{298}^\ddagger (kcal/mol)	σ^+
NMe ₂	6.89	-1.70
NH ₂	7.10	-1.30
OH	7.71	-0.92
OMe	7.86	-0.78
SMe	7.79	-0.60
Me	9.19	-0.30
F	8.41	-0.07
H	9.41	0
CF ₃	9.43	0.53
CN	9.23	0.71
NO ₂	9.23	0.78

^[a] Complexes are calculated using CEP-31G(d)/M06 level of theory.


Table 6

NFTPT-Promoted Aryl-CF₃ Coupling from Complexes **1f**, **9**, **10**, and **11e**

Entry	Compound	N-N	Temp	Yield Ph-CF ₃
1	1f	dtbpy	80 °C	66%
2	9		80 °C	99%
3	10		80 °C	40%
4	11e		80 °C	90%
5	11e		23 °C	83%

Reactions were run in duplicate and yields determined by ¹⁹F NMR spectroscopy.

Table 7

Synthesis and Reactivity of (tmeda)Pd(Aryl)(CF₃) Complexes

Entry	Compound	Aryl	Yield Aryl-CF ₃ (80 °C)	Yield Aryl-CF ₃ (23 °C)
1	11a	<i>p</i> -FC ₆ H ₄	81%	78%
2	11b	<i>p</i> -CF ₃ C ₆ H ₄	76% ^a	52% ^a
3	11c	<i>p</i> -CNC ₆ H ₄	60%	22%
4	11d	<i>p</i> -MeOC ₆ H ₄	92%	95%
5	11e	C ₆ H ₅	94%	88%
6	11f	<i>p</i> -MeC ₆ H ₄	90%	83%
7	11g	<i>m</i> -MeC ₆ H ₄	95%	95%
8	11h	<i>o</i> -MeC ₆ H ₄	85%	88%
9	11i	<i>o</i> -MeOC ₆ H ₄	90%	99%

These reactions were conducted at 80 °C for 3 h and at 23 °C for 1 h. Reactions were run in duplicate and all of the starting material was consumed. Yields were determined by ¹⁹F NMR spectroscopy.

^a At both temperatures, trifluorotoluene was formed in 13% yield.