

Spring 1968

PHYTOPLANKTON POPULATIONS IN RELATION TO TROPHIC LEVELS IN NEW HAMPSHIRE LAKES

GERHARD K. GRUENDLING

Follow this and additional works at: <https://scholars.unh.edu/dissertation>

Recommended Citation

GRUENDLING, GERHARD K., "PHYTOPLANKTON POPULATIONS IN RELATION TO TROPHIC LEVELS IN NEW HAMPSHIRE LAKES" (1968). *Doctoral Dissertations*. 877.
<https://scholars.unh.edu/dissertation/877>

This Dissertation is brought to you for free and open access by the Student Scholarship at University of New Hampshire Scholars' Repository. It has been accepted for inclusion in Doctoral Dissertations by an authorized administrator of University of New Hampshire Scholars' Repository. For more information, please contact nicole.hentz@unh.edu.

**This dissertation has been
microfilmed exactly as received**

69-762

**GRUENDLING, Gerhard K., 1939-
PHYTOPLANKTON POPULATIONS IN RELATION TO
TROPHIC LEVELS IN NEW HAMPSHIRE LAKES.**

**University of New Hampshire, Ph.D., 1968
Botany**

University Microfilms, Inc., Ann Arbor, Michigan

PHYTOPLANKTON POPULATIONS IN RELATION TO
TROPHIC LEVELS IN NEW HAMPSHIRE LAKES

by

GERHARD K. GRUENDLING

B. S., Springfield College, 1964

A THESIS

Submitted to the University of New Hampshire
In Partial Fulfillment of
The Requirements for the Degree of

Doctor of Philosophy

Graduate School

Department of Botany

June, 1968

This thesis has been examined and approved.

Arthur C. Mathieson

Thesis director, Arthur C. Mathieson,
Asst. Prof. of Botany

Albion R. Hodgdon

Albion R. Hodgdon, Prof. of Botany

Charlotte G. Nast

Charlotte G. Nast, Prof. of Botany

John J. Sasner

John J. Sasner, Asst. Prof. of Zoology

Philip J. Sawyer

Philip J. Sawyer, Assoc. Prof. of Zoology

Hannah T. Croasdale

Hannah T. Croasdale, Prof. of Biology
Dartmouth College

5-15-68

Date

ACKNOWLEDGEMENTS

I express my sincere gratitude to the following people at the University of New Hampshire who have made this study possible: Dr. A. C. Mathieson for his direction and valuable assistance during the study; Dr. P. J. Sawyer for the use of his limnological data and for valuable suggestions in reviewing the manuscript; Dr. A. R. Hodgdon, Dr. C. G. Nast, and Dr. J. J. Sasner, for aid in the preparation and reviewing of the manuscript; Mr. N. B. Reynolds for his untiring aid in the collection of field data; Mr. G. L. Byers, Director of the N. H. Water Resource Research Center. I also acknowledge the assistance of Dr. H. T. Croasdale of Dartmouth College for aid in the identification of desmids and for reviewing the manuscript; Dr. R. Patrick of the Academy of Natural Sciences of Philadelphia for aid in the identification of diatoms; and Mr. R. S. Kinniburgh of the N. H. Water Supply and Control Commission for aid in nutrient analyses and for the use of laboratory equipment. I am especially grateful to the faculty and staff of the Department of Botany of the University of New Hampshire for their encouragement and guidance and for the opportunity to use the facilities and equipment of the department.

The work upon which this thesis is based was supported in part by funds provided by the United States Department of Interior, Office of Water Resources Research, as authorized under the Water Resources Act of 1964, Public Law 88-379, through the Water Resources Research Center of the University of New Hampshire.

TABLE OF CONTENTS

LIST OF TABLES	vi
LIST OF FIGURES	vii
ABSTRACT	viii
INTRODUCTION	1
METHODS	3
ECOLOGY OF PHYTOPLANKTON AT NEWFOUND LAKE	6
Description of Area and Environmental Factors	6
Species Composition	10
Seasonal Periodicity of Standing Crop	11
Vertical Distribution of Standing Crop	16
Occurrence of Reproductive Spores	18
Calculation of Phytoplankton Indices	19
ECOLOGY OF PHYTOPLANKTON AT WINNISQUAM LAKE	21
Description of Area and Environmental Factors	21
Species Composition	24
Seasonal Periodicity of Standing Crop	25
Effects of Copper Sulfate Treatment on Standing Crop	30
Vertical Distribution of Standing Crop	32
Occurrence of Reproductive Spores	35
Calculation of Phytoplankton Indices	36
DISCUSSION	37
SUMMARY	52
BIBLIOGRAPHY	54
TABLES	59
FIGURES	77

LIST OF TABLES

1.	Seasonal Occurrence of Phytoplankton-Newfound Lake ...	59
2.	Summary of Major Phytoplankton Components - Newfound Lake	64
3.	Vertical Distribution of Phytoplankton - Percent of Total Standing Crop - Newfound Lake	65
4.	Summary of Reproductive Spore Periodicity	66
5.	Summary of Phytoplankton Indices and Proposed Trophic Status - Newfound Lake	67
6.	Seasonal Occurrence of Phytoplankton - Winnisquam Lake	68
7.	Summary of Major Phytoplankton Components - Winnisquam Lake	74
8.	Vertical Distribution of Phytoplankton - Percent of Total Standing Crop - Winnisquam Lake.....	75
9.	Summary of Phytoplankton Indices and Proposed Trophic Status - Winnisquam Lake	76

LIST OF FIGURES

1.	Map of Newfound Lake, New Hampshire	77
2.	Map of Winnisquam Lake, New Hampshire	78
3.	Seasonal Temperature Variations - Newfound Lake	79
4.	Seasonal Variations in Light Penetration - 1% Level ..	80
5.	Total Phytoplankton - Newfound Lake	81
6.	Orthophosphate - Newfound Lake	82
7.	Nitrate Nitrogen - Newfound Lake	83
8.	Silicon Dioxide - Newfound Lake	84
9.	Major Phytoplankton Components - Newfound Lake	85
10.	Seasonal Temperature Variations - Winnisquam Lake	86
11.	Orthophosphate - Winnisquam Lake	87
12.	Nitrate Nitrogen - Winnisquam Lake	88
13.	Silicon Dioxide - Winnisquam Lake	89
14.	Total Phytoplankton - Winnisquam Lake	90
15.	Major Phytoplankton Components - Winnisquam Lake	91

ABSTRACT

PHYTOPLANKTON POPULATIONS IN RELATION TO TROPHIC LEVELS IN NEW HAMPSHIRE LAKES

by

GERHARD K. GRUENDLING

The relationships between composition, periodicity, and abundance of phytoplankton species and the trophic status of two New Hampshire lakes (Newfound and Winnisquam) were investigated. The lakes had vastly different concentrations of nitrate and phosphate and provided an excellent opportunity for a comparative study of the basic ecology and nutrient requirements of phytoplankton. They also provided an opportunity to test the usefulness of algal indicators and quotients for determining trophic status.

Regular collections (approximately fortnightly) were made at each lake from June 25, 1966 to November 10, 1967. Samples of phytoplankton were taken from six depths at three stations with a 4-liter Van Dorn water sampler. Samples were also taken for nutrient analyses of orthophosphate, nitrate nitrogen, and silicon dioxide. Qualitative determinations were from living material. The total number of phytoplankton cells were enumerated from fixed material under an inverted microscope.

The differences in the total standing crop at the two lakes were correlated with their differences in nutrient concentrations. Orthophosphate appeared to be the primary limiting factor of algal growth at the oligotrophic lake (Newfound), while light, temperature, and possible extracellular products were the primary limiting factors at the eutrophic lake (Winnisquam). The vertical distribution of the standing crop was a function of light penetration and the types of algal populations. A discussion of the factors influencing the periodicity, abundance, and distribution of phytoplankton species at the two lakes is presented, and the results are compared with the findings of other workers.

The species composition at the lakes was dependent upon their trophic status. Dominant species were considered to be the best indicators of trophic levels and a list of possible indicator organisms is given. A summary of possible rare indicators is also included. It is suggested that the species composition, abundance, and periodicity of phytoplankton should be considered when determining indicator organisms.

The phytoplankton quotients proposed by various authors were applied to the lakes. Most of the indices correctly designated the eutrophic lake, but not the oligotrophic lake. An evaluation of several phytoplankton quotients is given and a few suggestions are presented for improving phytoplankton quotients.

INTRODUCTION

The present investigation was initiated in order to evaluate the relationships between composition, periodicity, and abundance of phytoplankton species and the trophic status of two New Hampshire lakes (Newfound Lake and Winnisquam Lake). The lakes are similar in origin, geology, size, depth, altitude, and certain of their chemical and physical characteristics. However, they have vastly different concentrations of nitrate and phosphate, for Newfound Lake is oligotrophic while Winnisquam Lake is in the early stages of eutrophication. Winnisquam Lake was considered to be oligotrophic in 1936 (Newell, 1963), but since then the process of eutrophication has been accelerated by the liberation of treated sewage effluent into the lake. The two lakes provided an excellent opportunity for a comparative study of the basic ecology and nutrient requirements of phytoplankton populations.

It was also felt that the two lakes provided an opportunity to test the usage of algal indicators and quotients for determining trophic status - especially of soft-water lakes. Few North American workers have attempted to use algal indicators and quotients to evaluate the trophic levels of lakes (Palmer, 1957; Rawson, 1956), although a number of studies have been done on the effects of pollutants on algae

in flowing waters (Lackey, 1938, 1941; Patrick, 1949, 1961, 1964). A detailed study of nutrient and phytoplankton relationships has never been undertaken in New Hampshire lakes.

METHODS

Regular collections (approximately fortnightly) were made at each lake from June 25, 1966 to November 10, 1967, except during December and January when ice conditions prevented access. Three stations were established at each lake (Fig. 1 & 2), and six samples (1, 3, 5, 10, 15, and 20 meters) were taken at each station with a 4-liter Van Dorn water sampler. Twenty-five ml subsamples were collected from each depths for phytoplankton enumeration. These were preserved with three drops of acid Lugol's solution. One liter subsamples were also taken from 1, 5, and 10 meter depths for nutrient analysis. The nutrient samples were placed in polypropylene bottles, refrigerated and later taken in the laboratory for analysis. Vertical hauls (20 meters in depth) were collected at each station with a # 25 plankton net, in order to concentrate living phytoplankton for identification. The collections were refrigerated and then taken to the laboratory for analysis. Light penetration was determined with a Whitney Model LMD-8A Underwater Light Meter. Temperatures were recorded with a standard mercury thermometer and a Whitney Model TC-5 Underwater Thermometer.

Living samples were used whenever possible for the identification of organisms. Nannoplankton were concentrated

with a .45 micron millipore filter (living samples) and by sedimentation (fixed samples). The generic designations listed by Smith (1950) were used whenever possible. A variety of sources were consulted for specific identifications: Ahlstrom (1937); Eddy (1930); Huber-Pestalozzi (1938-1961); Patrick and Reimer (1966); Prescott (1962); Smith (1920, 1924); Teiling (1967); West and West (1904-1912); and West, West, and Carter (1923). Voucher specimens have been deposited in the University of New Hampshire Herbarium.

The phytoplankton were enumerated according to the inverted microscope technique outlined by Lund, Kipling, and Le Cren (1958), using a Unitron Model Bi-4777 inverted microscope. A one ml unconcentrated sample was added to a sedimentation tube, it was allowed to settle for 18 hours, and then the phytoplankton were counted under 430X magnification. The total number of cells of each species per ml was counted.

Nutrient analyses were performed, whenever possible, immediately upon return to the laboratory. Otherwise the samples were immediately frozen and stored at -10 C until they were analyzed. The procedures outlined by the American Public Health Association (1965) were used, i.e., the stannous chloride method for orthophosphate, the phenoldisulfonic acid method for nitrate nitrogen, and the colorimetric heteropoly blue method for silica. All colorimetric analyses

were determined with a Beckman DU-2 spectrophotometer.

ECOLOGY OF PHYTOPLANKTON AT NEWFOUND LAKE

Description of Area and Environmental Factors

Newfound Lake is located at approximate $43^{\circ} 40'$ N latitude and $71^{\circ} 46'$ W longitude. It has an area of 1,661 hectares and a maximum depth of 55 meters (Newell, 1960). The lake is a preglacial land form which was slightly modified (mainly deepened) by Pleistocene glaciation. Most of the lake is underlain by Devonian bedrock of fine to coarse grained biotite-muscovite granite and quartz monzonite. In a few areas, at the extreme north and south ends, the bedrock is composed of micaceous quartzite and coarse grained mica schist (Billings, 1956).

Philip J. Sawyer (personal communication) studied the limnological characteristics of Newfound Lake from 1965-1967. His data have shown that it is an extremely soft-water lake. The pH ranged from 6.7-7.2; methyl orange alkalinity values ranged from 5.0-9.0 ppm; no phenolphthalein alkalinity was detected; and the total dissolved solids ranged from 36-45 ppm. The lake never had an oxygen deficiency in the hypolimnion and usually had over 90 per cent oxygen saturation throughout the water column.

Figure 3 shows the seasonal variation of water temperatures at Newfound Lake. Five depths are recorded, and each

point represents the mean temperature found at all stations for that depth. The temperatures at the three stations did not vary more than ± 0.3 C on any occasion. The maximum surface water temperatures (23.0 to 25.0 C) were recorded during July and August. The temperatures at the lower depths reached their maxima somewhat later. In August, 1966, the surface temperatures decreased rapidly and by September 19, the upper 10 meters of the lake were isothermal. The fall overturn was completed by October 17 and at this time the water column was isothermal at 11.8 C. During February and March, 1967, the temperatures under the ice ranged from 0.5 - 2.0 C; a slight inverse stratification was evident. The lake was free of ice in early April and the spring overturn then occurred. Thereafter, the temperature of the water column increased rapidly and thermal stratification was evident in early June. After June the surface temperatures increased rapidly until a maximum of 25.0 C was recorded on August 1. During the summer period the temperatures at 10 meters ranged from 8.5-11.5 C; those at 15 meters ranged from 8.5-10.0 C. During the later summer and fall the temperature again decreased rapidly, and isothermal conditions were recorded in early November.

Figure 4 shows the seasonal variation of light penetration (1% of the surface light). Each point depicts the mean value recorded for all stations. In general, there was a

correlation between the total standing crop and the depth of light penetration (compare Figures 4 & 5). Thus, as the total standing crop increased, the light penetration decreased, and vice versa. The light penetration was low during the spring of 1966, and a minimum transparency was recorded on August 24 (12 meters). The minimum corresponded with the maximum standing crop of the year. The transparency of the water then increased steadily until fall, at which time decreased light penetration was again recorded. During February, March, and early April of 1967, the ice and snow cover on the lake varied considerably. A maximum cover of 50 cm of ice and 45.7 cm of snow occurred on March 3. The maximum cover corresponded with the lowest light penetration (2.5 meters). During the ensuing two weeks, there was considerable melting, and the light penetration increased rapidly. The light penetration fluctuated throughout the summer and fall, in association with minor peaks of phytoplankton.

The seasonal levels of nutrients (orthophosphate, nitrate nitrogen, and silicon dioxide) at Newfound Lake are shown graphically in Figures 6, 7, 8, respectively. Each point represents the mean value of the three stations.

The yearly mean orthophosphate values recorded were .011 mg/l at 1 meter, .011 mg/l at 5 meters, and .010 mg/l at 10 meters. During 1966, the maximum orthophosphate concen-

tration (.020 mg/l) occurred on October 3; smaller peaks were recorded in June and September. A minimum concentration (.002 mg/l) was found on November 23; other low values were recorded in August and late September. During 1967 the orthophosphate concentrations ranged from .032 mg/l (February 20) to .002 mg/l (May 5). Two lesser peaks occurred during June and November. Low values occurred during late March, April, July, and August. In general, the seasonal patterns of orthophosphate concentrations were similar at each depth.

The yearly mean nitrate nitrogen concentrations recorded were .057 mg/l at 1 meter, .052 mg/l at 5 meters, and .055 mg/l at 10 meters. During 1966, peaks occurred on June 25 (.064 mg/l) and on November 23 (.082 mg/l). A low of .019 mg/l was recorded on October 3. During 1967, the highest concentration (.097 mg/l) was recorded on February 20. After February there was a rapid decline to a low of .046 mg/l on May 5. The summer concentrations fluctuated considerably and the lowest concentration (.044 mg/l) was recorded on November 10. The concentrations at the three depths showed similar seasonal patterns. Exceptions were recorded on November 23, 1966 and February 20, 1967.

The yearly mean silicon dioxide concentrations were .849 mg/l at 1 meter, .841 mg/l at 5 meters, and .770 mg/l at 10 meters. The concentrations during 1966 were lowest in

the spring (.687 mg/l on June 25) and fall (.210 mg/l on November 23). The maximum concentration (2.120 mg/l) was recorded on August 24. The silicon dioxide concentrations fluctuated extensively during 1967. The winter concentrations were generally low, and ranged between .181 and .820 mg/l. The concentrations rose steadily throughout the spring and summer and a maximum of 2.095 mg/l was recorded on August 11. During 1966, concentrations of silicon dioxide were similar at the three depths, but they differed considerably during 1967. This was especially true during July and August, when the concentrations at 1 meter were much higher than the concentrations at 5 and 10 meters.

Species Composition

A total of 105 species of phytoplankton were identified at Newfound Lake. A complete list of the species and their seasonal periodicity is presented in Table 1. No particular group of algae dominated the phytoplankton from Newfound Lake. The largest number of species belonged to the Chlorophyceae (a total of 43) and 26 of these were desmids. The other major components of the total phytoplankton flora were as follows: Chrysophyceae (19 species), Bacillariophyceae (13 species), Cyanophyceae (10 species), and Dinophyceae (10 species).

The phytoplankton flora of Newfound Lake was characterized by having a large number of species that were present during the entire year. Thirty-six species were found in every collection; these were primarily members of the Desmidiaceae, Chrysophyceae, and Bacillariophyceae. The ephemeral species were chiefly species of the Chroococcales, Chlorophyceae (exclusive of desmids), and Dinophyceae. In general, the summer and fall phytoplankton flora of Newfound Lake was composed of a large number of species of Cyanophyceae, Chrysophyceae, and Desmidiaceae. The Chrysophyceae and Bacillariophyceae contributed the largest number of species during the winter and spring.

Seasonal Periodicity of Standing Crop

Figure 5 shows the seasonal periodicity of the total standing crop, while Figure 9 shows the major groups contributing to the total standing crop. Each point in these figures represents the mean of five collections (1, 3, 5, 10, and 15 meters). Thus, a general indication of the average standing crop within the photic zone is given. The mean monthly cell counts for the major species are depicted in Table 2.

The standing crop was higher during the summer of 1966 than in 1967. The total number of phytoplankton was low during June, 1966 (3.2×10^5 cells/l), but increased steadily

to a peak of 12.7×10^5 cells/l on August 24. It then decreased slowly to 3.9×10^5 cells/l on October 3. After the fall overturn, the phytoplankton numbers then increased to a fall maximum of 5.7×10^5 cells/l on November 23. The lowest standing crop (1.2-2.0 cells/l) occurred during the winter of 1967. No significant increase was detected until early June. At this time, the cell numbers began to increase and a maximum of 6.9×10^5 cells/l was recorded on July 6; this was the highest value of 1967. The total population then fluctuated slightly during the summer, and reached a low of 2.9×10^5 cells/l on October 16. The population then increased again during November.

Members of the Cyanophyceae and Chrysophyceae were the major contributors to the standing crop at Newfound Lake (Fig. 9). In June, 1966, the numbers of Cyanophyceae increased steadily until a peak of 8.0×10^5 cells/l was reached on September 19. Gomphosphaeria lacustris and Aphanothece clathrata were the major contributors on this date. The cell numbers then decreased rapidly throughout the fall, but increased again in November, reaching a peak on November 23. The standing crop of Cyanophyceae was at a low level (.35-.53 cells/l) during the winter of 1967. The population started to increase during June and reached a peak of 4.0×10^5 cells/l on August 1. Gomphosphaeria lacustris was again the dominant

species. The population then decreased throughout the fall and a final pulse occurred on November 10.

The Chrysophycean population at Newfound Lake can be characterized as having two pulses a year. One occurred in the summer, and the other occurred in late fall. Beginning in June, 1966, a rapid increase of Chrysosphaerella longispina, Dinobryon cylindricum, and D. bavaricum took place. A maximum of 3.7×10^5 cells/l of Chrysosphaerella longispina occurred on August 8. The seasonal maximum recorded for the Chrysophyceae (12.0×10^5 cells/l) occurred on August 24, when a sudden pulse of Uroglenopsis americana developed. After August 24, the standing crop of Chrysophyceae declined rapidly to the lowest point recorded of $.03 \times 10^5$ cells/l on September 19. There was then a gradual increase in numbers, especially of Dinobryon cylindricum and D. bavaricum, until a maximum of 1.7×10^5 cells/l was reached on November 23. The standing crop during the winter of 1967 was dominated by Synura uvella. Population levels were very low from February through early June (ranging between $.05$ and $.35 \times 10^5$ cells/l), but beginning in early June, 1967 there was a rapid increase. Uroglenopsis americana, Dinobryon bavaricum, and D. cylindricum again contributed to the maximum (4.8×10^5 cells/l) which occurred on July 6. Chrysosphaerella longispina was not as abundant in 1967 as in 1966. After July, there was a

rapid decline in the standing crop until early fall; however, during this decline period there was a substantial increase in Dinobryon vanohoffenii and Synura uvella. The fall pulse of 1967 was not as great as that during 1966. A maximum of only $.48 \times 10^5$ cells/l was recorded on October 16. The main contributors were Synura uvella, Dinobryon cylindricum, and D. divergens. In general, the populations of Chrysophyceae followed the same yearly trends throughout the study.

Members of the Bacillariophyceae were the third largest contributors to the total standing crop. In 1966, the standing crop increased through June and reached a peak of 2.3×10^5 cells/l on July 25. The major components were Asterionella formosa var. formosa and Rhizosolenia eriensis. The population then decreased steadily through September. Two pulses were recorded during the fall; one occurred during the breakdown of thermal stratification and the other followed the complete overturn. The first pulse ($.60 \times 10^5$ cells/l) occurred on October 3 and it was chiefly composed of Melosira islandica and Asterionella formosa var. formosa. The second maxima ($.75 \times 10^5$ cells/l) of the same species occurred on November 23. The Bacillariophyceae were the major components of the winter population. The maximum standing crop (1.10×10^5 cells/l) occurred on April 21, 1967, immediately after the spring overturn. The population was dominated by

Asterionella formosa var. formosa, A. formosa var. gracillima, and Melosira islandica. The numbers of diatoms decreased steadily through June, but thereafter increased. A maximum population of $.91 \times 10^5$ cells/l was recorded on July 21, and was primarily composed of Rhizosolenia eriensis. Instead of decreasing to a summer low, as during 1966, the standing crop of diatoms remained relatively high during the summer and fall. The high standing crop during the summer was primarily due to large numbers of Rhizosolenia eriensis and Asterionella formosa var. formosa. The only substantial increase of diatoms during the fall occurred on October 1, when Tabellaria fenestrata reached $.25 \times 10^5$ cells/l.

Members of the Chlorophyceae did not contribute significantly to the total standing crop. Their standing crop remained relatively low throughout the year except for small pulses during June and July, 1966 (1.0×10^5 cells/l) and during June ($.50 \times 10^5$ cells/l) and September, 1967 ($.70 \times 10^5$ cells/l). The major contributors were Botryococcus braunii, Gloeocystis gigas, and Crucigenia rectangularis.

Members of the Cryptophyceae contributed less than the Chlorophyceae to the standing crop. The populations were low during the summer and reached maximum levels in the fall - 1.1×10^5 cells/l on October 3, 1966 and 1.0×10^5 cells/l on November 10, 1967. Rhodomonas lacustris was the major

component of the fall blooms. Cryptomonas ovata and C. marsonnii tended to reach smaller peaks during late summer.

The Dinophyceae, although conspicuous at times due to their large size, contributed very little to the total standing crop. Peridinium limbatum was the major contributor and it was most abundant in the early spring and late fall, with maximum concentrations of $.03 \times 10^5$ cells/l. During the winter, substantial populations of dinoflagellates also occurred. Peridinium willei was the major contributor. Although the dinoflagellate populations were generally abundant during the spring and fall, the largest pulse occurred during the summer of 1967. On August 1 and August 11, Peridinium inconspicuum was found in quantities of .07 and .06 cells/l, respectively.

Vertical Distribution of Standing Crop

Table 3 summarizes the vertical distribution of the standing crop at Newfound Lake. The bulk of the phytoplankton occurred at 3, 5, and 10 meters. All of the Cyanophyceae, except Anabaena flos-aquae, were most abundant between 5 and 10 meters. Anabaena flos-aquae the only filamentous blue-green alga of any significance at Newfound Lake, occurred throughout the summer in largest numbers at the surface.

The Chrysophyceae generally reached maximum standing crops at 5 and 10 meters. An exception to this was the

occurrence of Dinobryon spp. in large numbers at the surface during June and July, 1966. However, during other peaks of Dinobryon spp. the algae did not occur at the surface, but they reached their maximum numbers at 3, 5, and 10 meters. An interesting vertical distribution was noted in Mallomonas tonsurata var. alpina. During the period of ice cover, when light and temperatures were low, it reached maximum abundance at one meter. During the summer, it was found in maximum numbers at 10 and 15 meters, where the light and temperature regimes were lower than at the surface. During the late fall, it was again higher in the water column (3-5 meters). Such observations suggest that the vertical distribution is dependent upon certain critical temperature and/or light relationships.

The Bacillariophycean populations were usually most abundant at 5 and 10 meters - even during winter when the ice and snow cover was substantial. It was also noted, that as diatom populations declined, they tended to occur progressively deeper in the water column. This sinking was especially evident after peaks of Melosira islandica, Asterionella formosa var. formosa and Rhizosolenia eriensis. Rhizosolenia eriensis reached maximum numbers at one meter during the early summer. As the population began to decline the depth at which the maximum number of organisms was found increased, until the

alga was almost completely absent from the photic zone. The same sequence occurred again during the summer of 1967.

Melosira islandica was abundant near the surface only after the spring and fall overturns. Following the overturns, it sank quickly to greater depths. Asterionella formosa var. formosa, Cyclotella bodanica, and C. compta never developed maximum numbers near the surface, but usually reached a maximum at 5 and 10 meters.

During the ice free period the members of the Cryptophyceae were evenly distributed throughout the photic zone. However, during the winter they were generally found closer to the surface, although large number of Rhodomonas lacustris and Cryptomonas spp. were found still well below the point of light extinction. Most of the Dinophyceae were evenly distributed throughout the water, except for Peridinium inconspicuum which was most abundant at 1 and 3 meters. Peridinium willei reached maximum numbers at 1 meter under the ice but after the ice had melted it tended to become more evenly distributed. The Chlorophyceae generally occurred in the surface waters and reached maximum quantities at 1 and 3 meters.

Occurrence of Reproductive Spores

Table 4 summarizes the reproductive spore periodicity of phytoplankton observed during the study. Spores were only

found in three species. Akinetes were present on Anabaena flos-aquae throughout its growing season, except for November. No obvious correlations were evident between akinete production and environmental factors. Statospores were found on Dinobryon bavaricum during June and July, 1967 and on Dinobryon cylindricum during October and November, 1966 and June and July, 1967. In all cases, the formation of statospores occurred when the standing crops of Dinobryon spp. were at their peaks and when the orthophosphate concentrations were low (between .008 and .015 mg/l). No other correlations were evident between spore formation and other environmental factors.

Calculation of Phytoplankton Indices

An attempt was made to assess the trophic level of Newfound Lake by applying the phytoplankton quotients proposed by various authors. The Chlorophycean quotient suggested by Thunmark (1945), and the Myxophycean, Diatom, Euglenine, and Compound quotients developed by Nygaard (1949) were used. A summary of the index values and the suggested trophic status for Newfound Lake is found in Table 5. Calculations were based on yearly records of species, as well as distinct seasonal periods (i.e. summer and winter). Values for indices other than the Diatom index were not calculated for the winter, since Nygaard considered them applicable only to summer collec-

tions. A summary of the quotients and their interpretations is as follows:

QUOTIENT	OLIGOTROPHIC	EUTROPHIC
Chlorophycean =		
$\frac{\text{Chlorococcales}}{\text{Desmidiace}}$	< 1.0	> 1.0
Myxophycean =		
$\frac{\text{Myxophyceae}}{\text{Desmidiace}}$	< 0.4	0.1 - 3.0
Diatom =		
$\frac{\text{Centrales}}{\text{Pennales}}$	< 0.3	> 0.3
Euglenine =		
$\frac{\text{Euglenine}}{\text{Myxo. + Chlor.}}$	< 0.2	0.0 - 1.0
Compound =		
$\frac{\text{Myxo. + Chlor. + Cent. + Euglen.}}{\text{Desmidiace}}$	< 1.0	> 2.5

The results shown in Table 5 are conflicting and the designation of the trophic status of Newfound Lake depends upon the type of index used and the time for which it is calculated. The simple Chlorophycean index and the Euglenine index tend to classify the lake as oligotrophic. However, the Myxophycean, Diatom, and Compound indices place it as a mesotrophic or eutrophic lake. An interpretation of the significance of these results can be found in the discussion.

ECOLOGY OF PHYTOPLANKTON AT WINNISQUAM LAKE

Description of Area and Environmental Factors

Winnisquam Lake is located at approximate 43° 32' N latitude and 71° 31' W longitude. It is 1,725 hectares in area and has a maximum depth of 50 meters (Newell, 1963). The origin and basic morphometry are similar to that of Newfound Lake. The northern portion of the lake is underlain by medium to coarse grained biotite-quartz monzonite, while the bedrock in the southern portion is composed of micaceous quartzite and coarse grained mica schist (Billings, 1956).

Philip J. Sawyer (personal communication) has also studied the limnological characteristics of Winnisquam Lake. His data have shown that it is a soft-water lake. The pH ranged from 6.2-7.2, except during an extreme bloom of Aphanizomenon flos-aquae, when a value of 10.2 was recorded. Methyl orange alkalinity values ranged from 5.0 - 11.0 ppm. No phenolphthalein hardness was recorded except during bloom conditions. The oxygen values ranged from 100% saturation at the surface to complete oxygen depletion in the hypolimnion during the summer months. Total dissolved solids ranged from 55 - 75 ppm.

The seasonal cycle of temperature at Winnisquam Lake was similar to that at Newfound Lake (compare Figs. 3 & 10).

The maximum summer surface temperatures at Winnisquam Lake ranged between 23.0 and 25.0 C, and after the summer there was a steady decrease in temperatures until the fall overturn. During the winter of 1967 the temperatures under the ice showed an inverse stratification similar to that in Newfound Lake. The spring overturn occurred during April, and relatively isothermal conditions existed until early June. Thereafter, the summer stratification began to develop. The maximum summer surface temperatures were reached later in 1967 than in 1966. In addition, the complete overturn in the fall of 1967 did not occur until much later.

Figure 4 illustrates the seasonal variation of light penetration at Winnisquam Lake. Because the total standing crop at Winnisquam Lake was much greater than at Newfound Lake, the light penetration at Winnisquam Lake was reduced. During 1966, the light penetration ranged from 7.0 to 10.0 meters, depending on seasonal variations of the standing crop. The maximum ice (60.0 cm) and snow cover (22.8 cm) at Winnisquam Lake was recorded on March 3, and at this time the 1% light level ranged between 4.2 and 5.8 meters. During April, the ice cover had just disappeared and the maximum spring standing crop was recorded. The light penetration reached its lowest value (3.2 meters) at this time. The occurrence of a large amount of particulate matter also contributed to the low

transparency. Summer transparencies were comparable to the previous summer. During the fall, the light penetration decreased drastically, corresponding to the time of the fall phytoplankton bloom.

Figures 11 - 13 summarize the seasonal values of nutrients (orthophosphate, nitrate nitrogen, and silicon dioxide) at Winnisquam Lake. The mean orthophosphate values were .035 mg/l at 1 meter, .036 mg/l at 5 meters, and .051 mg/l at 10 meters. During 1966 the orthophosphate concentrations ranged from .124 mg/l on September 12 to .019 mg/l on September 28. In 1967 the values ranged from maxima of .079 mg/l (September 6) and .076 mg/l (April 17) to minima of .012 mg/l (August 1) and .015 mg/l (March 3). At 1, 5, and 10 meters there were similar seasonal trends and only slight differences were evident at the depths sampled.

The mean nitrate nitrogen concentrations were .087 mg/l at 1 meter, .096 mg/l at 5 meters, and .122 mg/l at 10 meters. During 1966 they ranged from .201 mg/l (June 25) to .012 mg/l (September 12). In 1967, maxima of approximately .210 mg/l occurred from March through early April and in September. The minimum values (approximately .030 mg/l) occurred from late June through August. The nitrate concentrations at the three sampling depths showed a similar pattern, except during September, 1967, when the amount at 10 meters rose to .205 mg/l.

The concentrations at 10 meters were very high throughout September, but decreased again by October 16.

The mean silicon dioxide concentrations recorded during the study were .286 mg/l at 1 meter, .274 mg/l at 5 meters, and .266 mg/l at 10 meters. During June, July, and August, 1966 the values ranged from .160 to .220 mg/l. They decreased to a low of .030 mg/l on September 28, but they rose again during late fall. The maximum silicon dioxide concentration recorded during 1967 was .520 mg/l on March 3. After March the concentrations decreased and eventually leveled off during June, July, and August. A late fall peak (.400 mg/l) was recorded on November 3. The silicon dioxide concentrations recorded at different depths were quite variable, probably reflecting differences in utilization at the particular depths.

Species Composition

One hundred and forty-three species of algae were identified from the plankton at Winnisquam Lake. Several species, however, were considered transitory members in the open water - e.g. some of the filamentous green algae (Mougeotia, Oedogonium, and Spirogyra), certain species of Oscillatoria, and the diatom Cymbella. Table 6 summarizes all of the species found and their periodicity of occurrence.

The green algae contributed the most species to the

phytoplankton at Winnisquam Lake - e.g. ninety-three species were members of the Chlorophyceae. There were 32 desmid species found. Other major contributing groups were the Bacillariophyceae (14 species), Chrysophyceae (13 species), and Cyanophyceae (10 species). The lake also had a large number of ephemeral species; they contributed to the marked fluctuations in the species composition throughout the year. Only 21 species were present in all of the 29 collections made during the study. These persistent species included a few members from all of the algal classes present. The majority of the ephemeral species were members of the Chlorophyceae. In general, the summer phytoplankton flora of Winnisquam Lake was composed of a large number of species of Chlorophyceae and filamentous Cyanophyceae. In the spring and fall, the Bacillariophyceae, Chrysophyceae, and Chlorophyceae (mainly desmids), contributed the largest number of species. Diatoms were the major components of the winter flora.

Seasonal Periodicity of Standing Crop

Figure 14 summarizes the total standing crop at Winnisquam Lake, while Figure 15 shows the major components of the total phytoplankton. Each point on Figures 14 and 15 represents the mean value of four depths (1, 3, 5, and 10 meters) and three stations, i.e. 12 collections. The mean

monthly cell counts for the major species contributing to the standing crop are also depicted in Table 7.

The total standing crop was much higher during 1967 than in 1966; in some cases the difference was as great as 1.0×10^7 cells/l. Large populations of algae, chiefly blue-green, occurred during July and August, 1966. A maximum of 2.2×10^6 cells/l was found on August 8. The populations then decreased to their lowest points ($.7 \times 10^6$ cells/l) during August and September. A second pulse (1.7×10^6 cells/l) was recorded after the fall overturn; it was primarily composed of blue-green algae and diatoms. The fall pulse was then followed by a gradual decline in the standing crop.

During February and March of 1967, the standing crop (mainly diatoms) was at its lowest point (between $.45$ and $.55 \times 10^6$ cells/l). A rapid increase in the phytoplankton numbers of all algal groups occurred after the spring overturn. By April 29, a maximum of 9.0×10^6 cells/l was found. A rapid decline in numbers then ensued and a summer low of 1.3×10^6 cells/l was recorded on June 2. Although the cell counts recorded on June 2 were relatively low for 1967, they were still higher than most of the 1966 records. The standing crop increased rapidly throughout the summer and fall and reached a maximum of 1.7×10^8 cells/l on September 29. At this date, most of the standing crop consisted of Aphanizomenon

flos-aquae and it remained very high until the study was terminated in November.

The Cyanophyceae and Chlorophyceae were the major phytoplankton components at Winnisquam Lake. A blue-green maximum of 1.5×10^6 cells/l (chiefly Anabaena flos-aquae) was recorded on August 8, 1966, and a smaller peak of $.68 \times 10^6$ cells/l (mostly Aphanizomenon flos-aquae) occurred in October 12. During February and March of 1967 a substantial population of Oscillatoria limnetica developed under the ice. Their cell numbers increased rapidly after the spring overturn and a peak of 3.6×10^6 cells/l was recorded on April 29. A peak of 3.2×10^6 cells/l occurred on July 6, 1967; it consisted mostly of Anabaena flos-aquae and Aphanizomenon flos-aquae. The final pulse began in early August and a maximum of 1.7×10^6 cells/l was recorded on September 29 (almost entirely Aphanizomenon flos-aquae).

The Chlorophyceae were another major group contributing to the total standing crop at Winnisquam Lake. One major peak (1.5×10^6 cells/l) was observed on July 18, 1966; it was composed chiefly of Gloeocystis vesiculosa and Eudorina elegans. The populations then leveled off during the rest of the summer and fall, and cell counts of $.3 - .5 \times 10^6$ cells/l were recorded. This late summer and fall population also had Dictyosphaerium pulchellum, Staurastrum pingue, and

Schroederia judayi as important contributors. The green algal populations were at their lowest point during February and March, 1967. After the spring overturn their cell numbers increased drastically and a maximum of 2.1×10^6 cells/l was found on April 29. Almost all of the material at this time was Ankistrodesmus falcatus var. mirabilis. The standing crop then gradually decreased until a second pulse occurred during July and August. The summer maximum of 1.6×10^6 cells/l was observed on August 1; it was primarily composed of Dictyosphaerium pulchellum and Gloeocystis vesiculosa. The population of green algae then decreased rapidly to a low level which was maintained until the termination of the study.

The members of the Bacillariophyceae contributed significantly to the winter, spring and late fall standing crop. During the summer of 1966, the populations were generally low until August. At this time, there was an increase in Tabellaria fenestrata var. fenestrata and a peak of approximately $.13 \times 10^6$ cells/l was found from September 12 to October 12. The maximum diatom standing crop ($.64 \times 10^6$ cells/l) occurred on November 23, and it consisted chiefly of Fragilaria crotonensis, Asterionella formosa var. formosa, and Melosira italica.

The winter diatom populations were somewhat lower than those recorded during the fall and they were primarily composed

of Melosira italica, Asterionella formosa var. formosa, and A. formosa var. gracillima. The diatom population increased during April and it reached a peak of 2.3×10^6 cells/l on April 29. The total population then decreased to an early summer low; however, Synedra ulna reached its maximum standing crop ($.18 \times 10^6$ cells/l) on May 11. The standing crop recorded during the summer of 1967 was different from that of 1966. Peaks of Fragilaria crotonensis were observed on July 21 ($.35 \times 10^6$ cells/l) and on August 1 ($.22 \times 10^6$ cells/l). There was then a steady decline in the diatom population and no significant fall diatom pulse occurred.

The Chrysophyceae contributed insignificantly to the total standing crop at Winnisquam Lake. Synura uvella and Dinobryon spp. were the most important individuals of the class. Small peaks were recorded in June and July of 1966, and a high of $.22 \times 10^6$ cells/l was found on October 12. Synura uvella contributed $.19 \times 10^6$ cells/l to this peak. During 1967, the populations were even lower than in 1966. Peaks of Synura uvella were recorded on August 1 ($.10 \times 10^6$ cells/l) and on August 11 ($.12 \times 10^6$ cells/l). The populations of Chrysophyceae were scarcely detectable during the fall of 1967.

The Cryptophyceae were at their peak during the spring overturn. The major contributors were Rhodomonas lacustris

and Cryptomonas ovata; although other species of Cryptomonas were evident at the time. Their standing crop was fairly uniform throughout 1966 except for a small peak of $.24 \times 10^6$ cells/l which was recorded on October 12. The population was very low during the winter of 1967, but a rapid increase of Rhodomonas lacustris occurred immediately after the ice melted. The population reached a plateau of about $.80 \times 10^6$ cells/l on April 29, and it remained at this level throughout May. There was then a rapid decline in the Cryptophycean population and it remained at a low level until the end of the study.

Only one species of the Dinophyceae, Ceratium hirundinella, was consistently found in the cell counts, but several other ephemeral were also recorded. The largest population of Dinophyceae was found on September 12 ($.024 \times 10^6$ cells/l) and on September 28 ($.048 \times 10^6$ cells/l). The standing crop recorded during 1967 was never as high as that observed in 1966.

Effects of Copper Sulphate Treatment on The Standing Crop

The natural sequence of development at Winnisquam Lake was altered on July 21, 1967 when 0.5 ppm of copper sulphate were added by the New Hampshire Water Supply and Pollution Control Commission, to control a bloom of Anabaena flos-aquae and Aphanizomenon flos-aquae. Collections made before and on

the day of application revealed that the blue-green algal populations had reached their peak of 3.2×10^6 cells/l on July 6, and they were decreasing at the time of application. In contrast, the populations of Chlorophyceae, Bacillariophyceae, and Cryptophyceae were increasing before the application. Our records indicated that the total standing crop increased from 3.5×10^6 cells/l on July 21 to 3.8×10^6 cells/l on August 1. Several species decreased in numbers, but certain green algal populations continued to increase. The most striking example was Distyosphaerium pulchellum which increased from $.27 \times 10^6$ cells/l to 1.03×10^6 cells/l. Gloeocystis vesiculosa also increased (from $.39 \times 10^6$ to $.51 \times 10^6$ cells/l). The diatom and cryptomonad populations decreased. The blue-green populations decreased slightly and 1.8×10^6 cells/l were recorded on August 1. However, the standing crop of Aphanizomenon flos-aquae then began to increase and a maximum of 1.7×10^8 cells/l was recorded on September 29. This was the highest value recorded for a phytoplankton population during the study. The population of Aphanizomenon flos-aquae remained very high until the termination of the study. During the same period all of the other algal groups eventually decreased in abundance, and the normal fall pulsation did not occur.

Vertical Distribution of Standing Crop

Table 8 summarizes the vertical distribution of the standing crop at Winnisquam Lake. In contrast to the vertical distribution at Newfound, the largest portion of the standing crop at Winnisquam Lake occurred in the upper five meters. A large percentage of the phytoplankton occurred below 5 meters when populations were declining and sinking out of the photic zone. The vertical distributions of the standing crop also depended upon the periodicity of major phytoplankton groups. When there was a large population of blue-green algae the largest number of organisms usually occurred near the surface, but when diatoms were dominant the maximum numbers occurred lower in the water column.

Although the largest numbers of Cyanophyceae usually occurred at 1, 3, and 5 meters, at times they were observed in abundance at lower depths. For example, Oscillatoria limnetica showed a conspicuous seasonal difference in its vertical distribution. During late February and early March large populations were present in the upper three meters. By the end of March it had settled to 10 and 15 meters - even though it was still increasing in numbers. With the onset of the spring overturn, a large portion of the population was brought to the surface where it increased rapidly. The standing crop of Oscillatoria limnetica then decreased, sank to lower depths and finally disappeared from the photic zone.

The large bloom of Aphanizomenon flos-aquae, which occurred during the fall of 1967, did not appear to react in the same manner. On September 6 it was found in very large numbers at 1, 3, and 5 meters. During late September and October some of the population moved to 10 meters, with the decline and eventual decomposition of the alga occurring in the upper 10 meters.

The members of the Chlorophyceae were more evenly distributed within the upper ten meters than the Cyanophyceae. However, when large blooms of blue-green algae were found at the surface, the maximum vertical distribution of the green algae was reduced. An interesting vertical distribution was found in Closteriopsis longissima. During the ice-free periods, it was always confined to the 15 and 20 meter samples, and it was often found in relatively large numbers well below the 1% surface light level. The only time it was found higher in the water column (5 meters) was during February under the ice. Such observations imply that low light and/or low temperature are critical in determining its distribution. Staurostrum pingue also appeared to reach maximum numbers at greater depths than most green algae. Although it was found near the surface, its maximum population densities usually occurred at 10 meters.

During times of peak numbers the members of the Bacillariophyceae showed a maximum numbers of cells at 5 and 10 meters. When they began to decline they slowly sank. A

striking example of this was Melosira italica. During September, 1966 it was present in substantial quantities and at lower depths (15 and 20 meters). The alga was evenly distributed throughout the water column after the fall overturn; it was also at this point that peak numbers were reached. The population began to sink after the ice cover had formed and by March 20, it was confined again to 15 and 20 meters. With the onset of the spring overturn the alga again appeared in large numbers throughout the water column, and it was at this time that the maximum densities were reached. The population steadily settled out of the photic zone during the spring and summer. It was difficult to access whether other diatoms showed this type of movement.

Members of the Cryptophyceae occurred mainly in the surface waters. The largest portions of their standing crop usually occurred above 5 meters. However, during the spring and fall overturns they tended to become more uniformly distributed.

It was difficult to distinguish any distributional pattern for members of the Chrysophyceae and Dinophyceae since their standing crop was very low and they were very sporadic in occurrence. During peak populations, Synura uvella tended to reach maximum numbers at 5 and 10 meters.

Occurrence of Reproductive Spores

Table 4 summarizes the reproductive periodicity of six species of phytoplankton at Winnisquam Lake. Akinetes were almost always found during the summer on Anabaena flos-aquae, Anabaena Scheremetievi, and Gloeotrichia echinulata. No correlation was evident between akinete formation and environmental factors. Akinetes were found on Aphanizomenon flos-aquae from September through November, 1967. The akinetes first appeared when the standing crop of this alga had reached its highest level during the study. Dinobryon cylindricum formed statospores in April, May and early June of 1967. These dates were somewhat earlier than those recorded for this species at Newfound Lake. However, as at Newfound Lake, the statospores were produced during the peak of the standing crop. The nutrient levels were also decreasing, but they were not as low as those recorded at Newfound Lake. There was another peak of Dinobryon cylindricum at Winnisquam Lake during June and July, 1966, but there was no formation of statospores. Auxospores were found on Melosira italica during the peak of the standing crop (April 29, 1967), and they were present until early June when many of the spores were germinating. Auxospore formation occurred only on the smallest filaments, i.e. those ranging from 3.2 to 5.6 microns in diameter.

Calculation of Phytoplankton Quotients

The phytoplankton quotients proposed by Thunmark (1945) and Nygaard (1949) were also applied to Winnisquam Lake. A summary of the values and the proposed trophic status is given in Table 9. The Myxophycean and Euglenine indices classified the lake as oligotrophic. However, the Chlorophycean, Compound, and Diatom indices placed it as a mesotrophic or eutrophic lake. The index values were quite variable and depended on the algal groups employed and the time of year the data were obtained.

DISCUSSION

The differences in the total standing crop at the two lakes can be correlated with their differences in nutrient concentrations. Phosphate appears to be the most important limiting factor for algal growth at Newfound Lake. Hutchinson (1967) stated that phosphate is the limiting factor in many unproductive lakes. The primary factors limiting growth at Winnisquam Lake are probably light, temperature, and possibly extracellular substances; nutrients are usually not limiting. According to Lund (1965), the phosphate concentration in artificially enriched lakes is so high, that it may play a lesser role in controlling phytoplankton populations. Steeman-Nielsen (1955) has proposed that if a lake is very rich in nutrient, light may be the main limiting factor, especially in the fall. In New Hampshire lakes, temperature appears to control the initiation and cessation of growth of certain algal populations.

The total standing crop at Newfound Lake reached higher levels during the summer of 1966 than in 1967. The higher standing crop was correlated with slightly higher levels of phosphate in the surface waters during 1966. In contrast, the standing crop at Winnisquam Lake was higher during the summer and fall of 1967 than in 1966. The higher

standing crop was correlated with increased nitrate and phosphate concentrations. Apparently the nutrients were liberated by many of the algae after the lake had been treated with copper sulfate.

A large standing crop (mainly diatoms) was developed during the spring at Winnisquam Lake; this is typical of temperate lakes. High concentrations of nutrients were present during the winter, and after the light increased, the algal populations increased rapidly. The maximum standing crop (mainly Chrysophyceae) at Newfound Lake did not occur during the spring, but developed during early summer. There was a definite uptake of phosphate and nitrate by the phytoplankton during the winter and spring. However, a long lag period then occurred until the proper light and/or temperature requirements were reached for these organisms. These differing maxima may, perhaps, be explained in terms of differing species compositions at the two lakes.

The vertical distribution of the standing crop was chiefly a function of light penetration and the types of algal populations present. Since the standing crop at Winnisquam Lake was large and primarily composed of blue-green algae, the light penetration was reduced. Thus, the largest percentage of organisms occurred in the upper five meters. There also appeared to be little light inhibition of algal growth at the lake surface. At Newfound Lake, light inhibition appeared to

occur near the surface, since very low cell numbers were recorded from the one meter samples. The transparency at this lake was much greater and a large percentage of organisms occurred lower in the water column. Suppression of the actively growing zone could have had an affect on the development of the standing crop, since many of the algae had to grow at lower temperatures.

Pearsall (1932) observed that diatom populations in English Lakes were limited by silicon concentrations below .500 mg/l. In contrast, the silicon concentrations at Winnisquam Lake were seldom more than .500 mg/l, but diatom populations were extensive and increased even when the silicon levels were below .100 mg/l. Lund (1949, 1950) found that if other nutrients were not limiting, diatom populations would continue to increase. Such may be the case at Winnisquam Lake where phosphate and nitrate levels were very high. Jorgensen (1957) found that the lower limits for diatom growth were between .03 - .04 mg/l silicon; these are similar to the lower limits observed at Winnisquam Lake. The silicon concentrations at Newfound Lake were very high, and except for brief periods during the winter, they were always well above .500 mg/l. However, large diatom populations were never recorded and this was probably due to the low phosphate concentrations.

Specific diatom populations had different factors controlling their growth. Peaks of Asterionella formosa occurred at Winnisquam Lake shortly after maximum nitrate and silicon concentrations. However, the phosphate concentrations preceding the population peaks were generally low. A possible explanation has been presented by Mackereth (1953). He found in culture that there was a rapid uptake of phosphate by Asterionella formosa before any appreciable growth in the population was detected. At Newfound Lake, this same alga showed a similar pattern. Decreases in populations at both lakes were associated with decreases in nutrients. In addition, parasitism by chytrid fungi, as suggested by Canter and Lund (1948), possibly contributed to the decreases in the populations.

Maxima of Tabellaria fenestrata were usually reached later in the spring and earlier in the fall than most of the other diatoms. The populations at both lakes initiated growth when the phosphate and nitrate concentrations were low. In the two New Hampshire lakes, it appears that Tabellaria fenestrata had a preference for lower nutrient concentrations than other diatoms.

Another factor that contributed to the seasonal diatom fluctuations was the sinking phenomenon studied extensively by Lund (1954, 1955). He found that Melosira italica subsp.

subartica was brought to the surface during the spring and fall overturns, reached peak development, and then quickly sank out of the photic zone when less turbulent conditions occurred. Melosira italica at Winnisquam Lake and Melosira islandica at Newfound Lake also followed this type of pattern. Fragilaria crotonensis did not show the same development, since large numbers occurred during the summer stratification period.

Lund (1965) stated that lakes with low phosphorus content should have a large Chrysophyte population. At Newfound Lake, Dinobryon cylindricum, D. bavaricum, D. vanhoeffenii, Chryso-sphaerella longispina, and Uroglenopsis americana began development in early summer, when the phosphate concentrations were low. At Winnisquam Lake, Synura uvella increased when the phosphate and nitrate concentrations were low, while Dinobryon cylindricum did not increase substantially until after the vernal maximum. Hutchinson (1944) found similar results for Dinobryon divergens in Linsey Pond, Connecticut. He and other works (Bamforth, 1958; Rodhe, 1948; and Lund, 1965) have shown that the increase in Chrysophyceae standing crop was associated with low phosphate concentrations.

The Cyanophyceae populations at the two lakes were floristically distinct. Members of the Chroococcales were abundant at Newfound Lake and reached peak numbers during

the summer. They initiated growth when the surface temperatures increased and their maximum standing crop was recorded when the temperatures were between 20.0 and 25.0 C. No apparent correlations were noted between nutrient concentrations and the initiation of peak numbers. The Cyanophyceae populations at Winnisquam Lake also reached maximum numbers in the summer, although Oscillatoria limnetica increased greatly under the ice. A number of workers, including Eberly (1967) have found large quantities of Oscillatoria spp. at low light intensities and temperatures. The larger standing crop of Oscillatoria limnetica in April, October, and November indicates that optimum growth conditions are not restricted to the winter months.

Hammer (1964) studied lakes in Saskatchewan, Canada and found that blooms of Anabaena flos-aquae usually appeared one or two weeks after phosphate peaks and when the water temperatures were between 15.0 and 20.0 C. Similar results were recorded at Winnisquam Lake, except that temperatures were higher (22.0 - 25.0 C) and the blooms occurred later in the season. Hammer (1964) also found that blooms of Aphanizomenon flos-aquae developed after the Anabaena flos-aquae blooms. Aphanizomenon was rarely present until the temperatures reached 20.0 C and most blooms appeared at temperatures of 22.5 - 26.5 C. The alga appeared also to be influenced

directly by the phosphate concentrations. During the fall of 1966, Aphanizomenon flos-aquae initiated its growth at Winnisquam Lake approximately two weeks after high phosphate concentrations were recorded. The surface water temperatures ranged between 12.0 and 21.0 C. The population decline was correlated with a decrease in temperature. The alga was present throughout the winter and began to increase again during the early spring. However, substantial numbers were not recorded until the surface water temperatures had reached 20.0 C. Peak numbers were recorded when the temperatures ranged between 21.0 and 22.0 C. The initial growth was again associated with high concentrations of phosphate.

Aphanizomenon flos-aquae decreased slightly after the lake was treated with copper sulphate and then it increased to very high numbers. Copper sulphate treatment killed most of the Anabaena flos-aquae population and certain of the green algal populations. As a result, large amounts of nutrients were released into the surface waters. A very large peak of Aphanizomenon flos-aquae occurred after five to six weeks and few other algae were present. The low standing crop of other organisms could have been due to the shading effect of the large blue-green algal population. However, more probably it was due to an antagonistic substance released by Aphanizomenon. The abundance of Aphanizomenon appeared to

restrict the occurrence of a diatom pulse in the fall of 1967. Lefevre (1964) has reviewed the subject of extracellular products of algae and he has stated that Aphanizomenon flos-aquae releases substances which inhibit the growth of other algae. Shortly after Aphanizomenon began to decrease at Winnisquam Lake, other algae again began to increase.

The Chlorophyceae were more abundant at Winnisquam Lake than at Newfound Lake. Peaks of Ankistrodesmus falcatus var. mirabilis, Distyosphaerium pulchellum, and Eudorina elegans occurred shortly after high phosphate and nitrate concentrations were recorded. The very large populations of Dictyosphaerium pulchellum occurred immediately after the copper sulphate treatment. The highest nitrate concentrations were recorded during the winter at Winnisquam Lake. As the ice disappeared and light intensity increased, Ankistrodesmus falcatus var. mirabilis reached its maximum standing crop.

Lackey (1938, 1941) found that Cryptomonas ovata, Cryptomonas erosa, and Rhodomonas lacustris were inhibited by high concentrations of sewage. All three species were found in substantial quantities at both lakes. The initiation of their peaks could not be correlated with low nutrient levels. During the summer, Cryptomonas ovata and Rhodomonas lacustris developed peak numbers when the phosphate levels were low; but in the spring and fall, the initiation of pulses were associat-

ed with high nutrients. It is not possible to make exact comparisons with those of Lackey, since he did not include quantitative nutrient data.

The species composition of the two lakes was quite different and it is presumed to be dependant upon the trophic levels of the lakes. Seventy-three species were found exclusively at Winnisquam Lake. Of these, the dominant species were Aphanizomenon flos-aquae, Gleotrichia echinulata, Oscillatoria limnetica, Ankistrodesmus falcatus var. mirabilis, Eudorina elegans, Gloeocystis vesiculosa, Scenedesmus spp., Schroederia judayi, Staurastrum pingue, Staurastrum johnsonii, and Melosira italica. Forty-two species were found exclusively at Newfound Lake, of which the dominants were Aphanothece clathrata, Gloeothece linearis, Merismopedia punctata, Merismopedia tenuissima, Cosmarium bioculatum, Staurastrum pentacerum, Staurodesmus bulnheimii, Staurodesmus cuspidatus var. curvatus, Dinobryon vanhoeffenii, Mallomonas tonsurata var. alpina, Cyclotella bodanica, Melosira islandica, Peridinium inconspicuum, and Peridinium willei. One-hundred and thirty-three species were common to both lakes. Of these, Anabaena flos-aquae and Dictyosphaerium pulchellum are generally considered to be eutrophic species, but they were also found in very small quantities at Newfound Lake. The following species were abundant at Newfound Lake, but were found in

abundance at Winnisquam Lake only during periods of low nutrients: Dinobryon bavaricum, D. cylindricum, Chryso-sphaerella longispina, Synura uvella, Uroglenopsis americana, Gloeobotrys limnetica, and Rhizosolenia eriensis.

Two points must be considered when indicator organisms are selected. First, species which are found exclusively in a particular environment should be considered. Secondly, species that are dominant in a particular trophic situation should also be considered - even though they are found in other situations. For example, Hutchinson (1967) has reviewed most of the literature on indicator organisms and has pointed out that a number of workers have considered Dinobryon cylindricum var. palustre and Dinobryon bavaricum var. vanhoeffenii (D. vanhoeffenii in this study) as oligotrophic species. Dinobryon spp. can be considered oligotrophic indicators in New Hampshire lakes, even though Dinobryon cylindricum was found in the eutrophic situation. More detailed information about other New Hampshire lakes would be required to verify the possible indicators mentioned above, although most of the organisms suggested have also been proposed by several European workers (Brook, 1968; Jarnefelt, 1952; Nygaard, 1949; Round and Brook, 1959; Teiling, 1955; and Thunmark, 1945). However, there was some disagreement with the findings of other authors. Jarnefelt (1956) found

one eutrophic lake in Finland which was dominated by Cosmarium bioculatum. In New Hampshire the alga was found only in the oligotrophic lake. Rawson (1956) considered Asterionella formosa as an oligotrophic species in Saskatchewan lakes, although most European workers consider it to be associated with higher trophic levels. In both New Hampshire lakes it was relatively abundant at all times of the year, but it was usually associated with high nutrient levels. Lackey (1941) considered Cryptomonas ovata and Rhodomonas lacustris as oligotrophic indicators in rivers. However, these organisms showed a preference for higher nutrient levels in the lakes studied in New Hampshire. Round and Brook (1959) found that Dictyosphaerium pulchellum showed a preference for oligotrophic lakes in Ireland. In New Hampshire the alga was most abundant in the eutrophic lake.

Rawson (1956) has suggested that the rarer species would be more indicative of trophic status than the dominant species. If Rawson's assumption is correct, the species that are present for extended periods would probably be the best ones to use as indicators. Taking this into account, possible rare indicator organisms at Newfound Lake would then be Dinobryon crenulatum, D. suecium, Mallomonas pseudocoronata, Diceras phaseolus, and Peridinium wisconsinense, while at Winnisquam Lake they could be Closteriopsis longissima,

Kirchneriella lunaris, Kirchneriella obesa var. aperta, Schroederia setigera, and Synura adamsii. However, the designation of these species as indicators is highly speculative, and many more lakes would have to be investigated in order to obtain an accurate assessment of their usefulness.

The phytoplankton quotients proposed by Thunmark (1945) and Nygaard (1949), which give equal value to dominant and rare organisms, were applied to the New Hampshire lakes. Most of the indices calculated, correctly designated Winnisquam Lake as eutrophic. However, they gave a very poor characterization of Newfound Lake. It has been shown (Tables 5 and 9) that the simple Chlorophycean quotient of Thunmark was the only one that adequately described the trophic status of both lakes. It was reliable because the Chlorococcalean populations had many more species at Winnisquam Lake than at Newfound Lake. The Diatom Index did not give an accurate determination of trophic status because the species of diatoms were similar at both lakes. The Myxophycean Index was not accurate because the numbers of species of blue-green algae at the two lakes was comparable. The Euglenine Index was not applicable, since there were very few members of this group in the eutrophic lake (two species of Colacium and one Euglena.) The Compound Index was also erroneous, for many of the above reasons. In addition, the emphasis upon desmids in

the Compound Quotient gave an inaccurate determination of the trophic status of Newfound Lake. Nygaard assumed that many species of desmids were restricted to oligotrophic waters, but in New Hampshire, a larger number of species of desmids were found in the eutrophic situation. Hutchinson (1967) has stated that "the prevalence of a large number of desmids is probably directly determined by low calcium rather than by deficiency of nitrogen and phosphorus"; this might explain the similarity in the number of desmid species at the two lakes. Calcium was present only as calcium bicarbonate in both lakes except during peak blooms at Winnisquam.

Data collected in this study suggests that species composition, abundance, and periodicity should be taken into account when determining the usefulness of indicators. Species of diatoms could not be used as indicators in the two New Hampshire lakes (with the possible exception of Melosira italica and M. islandica) because most of the same species occur at both lakes. However, with information about their abundance and periodicity, some indication of trophic status at the lakes can be determined. Dinobryon spp. and other species of Chrysophyceae appear to be good oligotrophic indicators, since they occurred in abundance at Newfound Lake and were rare at Winnisquam Lake. The Cyanophyceae populations were distinct at the two lakes and in this case could be used

as indicators. Winnisquam Lake was dominated by filamentous blue-green algae, which are generally considered to be eutrophic indicators. Even though Anabaena flos-aquae occurred in small quantities at Newfound Lake, this should not eliminate it as a eutrophic indicator. The blue-green population at Newfound Lake was primarily composed of members of the Chroococcales. Certain members of the Chroococcales might be used as oligotrophic indicators, but more work must be done to determine their trophic requirements. The Chlorococcales as a whole, appear to be good indicators of eutrophic situations, because a large number of species occurred at Winnisquam Lake while very few species were found at Newfound Lake. The present study concludes that members of the Cryptophyceae can not be used satisfactorily as indicators.

Although it is difficult to draw definite conclusions from a study of two lakes, the detailed nutrient and phytoplankton data has led to the following suggestions. It is felt that Nygaard definitely improved the quotient concept by developing the Compound Quotient. However, as Hutchinson (1957) and Rawson (1956) have suggested, it is unfortunate that he based the entire denominator on one group of organisms (desmids). If other groups could be added to the denominator, the index would probably become more reliable for New Hampshire lakes. In particular, the number of species of Chrysophyceae could

be added. Also, the separation of the Centrales into its eutrophic (Stephanodiscus, Melosira, etc.) and oligotrophic (Cyclotella, etc.) components may be advantageous. Another suggestion would be to integrate the abundance of organisms with the number of species in determining the phytoplankton index. Quantitative measurement could be calculated as relative abundance of species in the particular lake, or as absolute numbers of organisms. Thus, differences in standing crop would be considered. Jarnefelt (1956) has developed a quotient based on a quantitative approach and it worked well in Finnish Lakes. However, his quotient can not be used until a large number of lakes from a particular area have been investigated, and thus it was not applied in New Hampshire.

SUMMARY

The relationships between composition, periodicity, and abundance of phytoplankton species and the trophic status of two lakes in New Hampshire were investigated. Newfound Lake is oligotrophic, while Winnisquam Lake is eutrophic. During the study the following conclusions were reached:

1) The differences in the total standing crop at the two lakes were correlated with their differences in nutrient concentrations. Orthophosphate appeared to be the primary limiting factor of algal growth at Newfound Lake, while light, temperature, and possibly extracellular products were the primary limiting factors at Winnisquam Lake.

2) A large vernal maximum, typical for temperate lakes, was recorded at Winnisquam Lake. Newfound Lake has no vernal peak and the maximum phytoplankton development was delayed until early summer. The differing maxima may be explained in terms of differing species composition at the two lakes.

3) The vertical distribution of the standing crop was a function of light penetration and the types of algal populations. The maximum standing crop at Winnisquam Lake occurred higher in the water column than at Newfound Lake.

Light inhibition of algal growth appeared to occur near the surface at Newfound Lake.

4) A discussion of the factors influencing the periodicity, abundance, and distribution of phytoplankton species at the two lakes is presented, and the results are compared with findings of other authors.

5) The species composition appeared to be dependent upon the trophic levels of the lakes. Dominant species were considered to be the best indicators of trophic status and a list of possible indicator organisms is given. A summary of possible rare indicators is also included.

6) The phytoplankton quotients proposed by Thunmark and Nygaard were applied to the lakes. Most of the indices correctly designated Winnisquam Lake as eutrophic, but incorrectly characterized Newfound Lake. Thunmark's Chlorophycean Quotient was the only quotient that adequately described both lakes. An evaluation of the phytoplankton quotients is given.

7) It is suggested that the species composition, abundance, and periodicity of phytoplankton should be considered when determining indicator organisms. A few other suggestions are given for improving phytoplankton quotients.

REFERENCES

- Ahlstrom, E. H. 1937. Studies on variability in the genus Dinobryon (Mastigophora). Trans. Amer. Microsc. Soc. 56 (2):139-159.
- American Public Health Association, et. al. 1965. Standard methods for the examination of water and waste water including bottom sediments and sludges. Twelfth ed. Amer. Pub. Health Assoc., Inc. New York. 769 p.
- Bamforth, S. S. 1958. Ecological studies on the planktonic protozoa of a small artificial pond. Limmol. Oceanogr. 3 (4):398-412.
- Billings, M. P. 1956. The geology of New Hampshire part II Bedrock geology. New Hampshire State Planning and Development Commission. Concord. 203 p.
- Brook, A. J. 1958. Changes in the phytoplankton of some Scottish hill lochs resulting from their artificial enrichment. Verh. Internat. Ver. Limmol. 13:298-305.
- Brook, A. J. 1959. The status of desmids in the plankton and the determination of phytoplankton quotients. J. Ecol. 47:429-445.
- Canter, H. M., and J. W. G. Lund. 1948. Studies on plankton parasites. I. Fluctuations in the numbers of Asterionella formosa Hass. in relation to fungal epidemics. New Phytol. 47(2):238-261.
- Eberly, W. R. 1967. Problems in the laboratory culture of planktonic blue-green algae. p. 7-34. In Environmental requirements of blue-green algae (Proceedings of a Symposium). U. S. Dept. Interior. Corvallis.
- Eddy, S. 1930. The fresh-water armored or thecate dinoflagellates. Trans. Amer. Microsc. Soc. 49(4):277-321.
- Hammer, U. T. 1964. The succession of "bloom" species of blue-green algae and some causal factors. Verh. Internat. Ver. Limmol. 15:829-836.

- Huber-Pestalozzi, G. 1938-1961. Das Phytoplankton des Susswassers. Vol. 16 Pt. 1-5. In Thienemann, A. (ed.) Die Binnengewasser. E. Schweizerbart'sche Verlagsbuchhandlung, Stuttgart.
- Hutchinson, G. E. 1944. Limnological studies in Connecticut. VII. A critical examination of the supposed relationship between phytoplankton periodicity and chemical changes in lake waters. *Ecology* 25(1):3-26.
- Hutchinson, G. E. 1967. A treatise on limnology. Vol. II Introduction to lake biology and the limnoplankton. John Wiley & Sons., Inc., New York. 1115 p.
- Jarnefelt, H. 1952. Plankton als Indikator der Trophiegruppen der Seen. *Suomal. Tiedeakat. Toim. (Ann. Acad. Sci. Fenn.) Ser. A. IV. Biol.* 18:2-29.
- Jarnefelt, H. 1956. Zur Limnologie einiger Gewässer Finnlands. XVI. Mit besonderer Berücksichtigung des Planktons. *Suomal. elain-ja Karsirt. Sevr. van. elain. Julk. (Annls Zool. Soc. Vanamo)*, 17, No. 7: 201 p (not seen)
- Jorgensen, E. G. 1957. Diatom periodicity and silicon assimilation. *Dansk. bot. Ark.* 18(1):6-54.
- Lackey, J. B. 1939. Protozoan plankton as indicators of pollution in a flowing stream. U. S. Treasury Dept. Public Health Reports. 53:2037-2058.
- Lackey, J. B. 1941. Two groups of flagellated algae serving as indicators of clean water. *J. Amer. Water Works Assoc.* 33:1099-1110.
- Lefevre, M. 1964. Extracellular products of algae, p. 337-367. In Jackson, D. F. (ed.) *Algae and man*. Plenum Press, New York.
- Lund, J. W. G. 1949. Studies on Asterionella I. The origin and nature of the cells producing seasonal maxima. *J. Ecol.* 37:389-419.
- Lund, J. W. G. 1950. Studies on Asterionella formosa Hass. II. Nutrient depletion and the spring maximum. *J. Ecol.* 38:1-35.

- Lund, J. W. G. 1954. The seasonal cycle of the plankton diatom Melosira italica (Ehr.) Kutz, subsp. subarctica O. Mull. J. Ecol. 42:151-179.
- Lund, J. W. G. 1955. Further observations on the seasonal cycle of Melosira italica (Ehr.) Kutz. subsp. subarctica O. Mull. J. Ecol. 43:91-102.
- Lund, J. W. G. 1965. The ecology of the freshwater phytoplankton. Biol. Rev. 40(2):231-293.
- Lund, J. W. G., C. Kipling, and E. D. Le Cren. 1958. The inverted microscope method of estimating algal numbers and the statistical basis of estimation by counting. Hydrobiologia 11(2):143-170.
- Mackereth, F. J. H. 1953. Phosphorus utilization by Asterionella formosa Hass. J. Exp. Bot. 4(12):296-313.
- Newell, A. E. 1960. Biological survey of the lakes and ponds in Coos, Grafton, and Carroll counties. New Hampshire Fish and Game Depart. Survey Report No. 8a, Concord, 297 p.
- Newell, A. E. 1963. Biological survey of the lakes and ponds in Sullivan, Merrimack, Belknap, and Strafford counties. New Hampshire Fish and Game Depart. Survey Report No. 8b. Concord. 276 p.
- Nygaard, G. 1949. Hydrobiological studies of some Danish ponds and lakes. II. The quotient hypothesis and some new or little known phytoplankton organisms. Kongl. Danske Vidensk. Selsk. Biol. Skr. 7(1):1-293.
- Palmer, C. M. 1959. Algae in water supplies, an illustrated manual on the identification, significance, and control of algae in water supplies. Public Health Serv. Publ. No. 657. R. A. Taft Sanitary Engin. Cen, Cincinnati. 88 p.
- Patrick, R. 1949. A proposed biological measure of stream conditions, based on a survey of the Conestoga Basin, Lancaster County, Pennsylvania. Proc. Acad. Nat. Sci. Philad. 101:277-341.

- Patrick, R. 1961. A study of the numbers and kinds of species found in rivers in eastern United States. Proc. Acad. Nat. Sci. Philad. 113(10):215-258.
- Patrick, R. 1964. A discussion of natural and abnormal diatom communities, p. 185-204. In Jackson, D. F. (ed.) Algae and man. Plenum Press, New York.
- Patrick, R. and C. W. Reimer. 1966. The diatoms of the United States exclusive of Alaska and Hawaii. Vol. 1 Monograph No. 13, Acad. Nat. Sci. Philad., Philadelphia. 688 p.
- Pearsall, W. H. 1932. Phytoplankton in English lakes II. The composition of the phytoplankton in relation to dissolved substances. J. Ecol. 20:241-262.
- Prescott, G. W. 1962. Algae of the western great lakes area. Rev. Ed. Wm. C. Brown Co., Publishers, Dubuque. 977 p.
- Rawson, D. S. 1956. Algal indicators of trophic lake types Limmol. Oceanogr. 1(1):18-25.
- Rodhe, W. 1948. Environmental requirements of fresh-water plankton algae. Symb. Bot. Upsaliens. 10(1):5-149.
- Round, F. E. and A. J. Brook. 1959. The phytoplankton of some Irish loughs and an assessment of their trophic status. Proc. Roy. Irish Acad. Sect. B. 60:167-191.
- Smith, G. M. 1920. Phytoplankton of the inland lakes of Wisconsin. Part I. Wisconsin Geol. Nat. Hist. Surv. Bull. No. 57. Madison. 243 p.
- Smith, G. M. 1924. Phytoplankton of the inland lakes of Wisconsin. Part II. Wisconsin Geol. Nat. Hist. Surv. Bull. No. 57. Madison. 227 p.
- Smith, G. M. 1950. The fresh-water algae of the United States. 2nd ed. McGraw-Hill Book Co., Inc., New York. 719 p.
- Steemann-Nielsen, E. 1955. The production of organic matter by the phytoplankton in a Danish lake receiving extraordinarily great amounts of nutrients. Hydrobiologia 7 (1-2):68-74.

- Teiling, E. 1955. Algae - Some mesotrophic phytoplankton indicators. Verh. Internat. Ver. Limnol. 12:212-215.
- Teiling, E. 1967. The desmid genus *Staurodesmus*: a taxonomic study. Arkiv. For. Botanik. 6(11):467-629.
- Thunmark, S. 1945. Zur Soziologie des Susswasserplanktons. Eine methodologisch-okologische Studie. Folia. Limnol. Scandin. 3:1-66.
- West, G. S. and W. West. 1904-1912. A monograph of the British Desmidiaceae. Vols. 1-4. The Ray Soc., London.
- West, G. S., W. West, and N. Carter. 1923. A monograph of the British Desmidiaceae. Vol. 5. The Ray Soc., London.

Table 1

Seasonal Occurrence of Phytoplankton - Newfoundland Lake

	1966						1967									
	J	J	A	S	O	N	F	M	A	M	J	J	A	S	O	N
CYANOPHYCEAE																
<u>Anabaena flos-aquae</u> (Lyngb.) De Breb.		X	X	X	X	X						X	X	X	X	X
<u>Aphanothece clathrata</u> G.S. West	X	X	X	X	X	X						X	X	X	X	X
<u>Chroococcus limneticus</u> Lemm.	X	X	X	X	X	X						X	X	X	X	X
<u>Chroococcus minimus</u> (Keissl.) Lemm.			X	X	X	X						X	X	X	X	X
<u>Gloethece linearis</u> Naegeli	X	X	X	X	X	X						X	X	X	X	X
<u>Gomphosphaeria aponina</u>																
var. <u>delicatula</u> Virieux	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<u>Gomphosphaeria lacustris</u> Chodat	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<u>Gomphosphaeria lacustris</u>																
var. <u>compacta</u> Lemm.	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<u>Merismopedia punctata</u> Meyen	X	X	X	X	X	X					X	X	X	X	X	X
<u>Merismopedia tenuissima</u> Lemm.	X	X	X	X	X	X						X	X	X	X	X
CHLOROPHYCEAE																
<u>Ankistrodesmus falcatus</u>																
var. <u>acicularis</u> (Braun) G.S. West		X	X	X	X							X	X	X		
<u>Botryococcus braunii</u> Kuetzing	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<u>Botryococcus protuberans</u>																
var. <u>minor</u> G.M. Smith	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<u>Closterium parvulum</u> Naegeli											X					
<u>Closterium kuetzingii</u> De Breb.		X				X	X				X					
<u>Coelastrum microporum</u> Naegeli	X	X						X	X	X	X					
<u>Cosmarium bioculatum</u> De Breb.	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<u>Cosmarium contractum</u>																
var. <u>papillatum</u> W. & G.S. West	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<u>Cosmarium punctulatum</u> De Breb.		X	X	X	X	X						X	X	X	X	X
<u>Crucigenia rectangularis</u> (Braun) Gay	X	X	X	X	X	X	X	X				X	X	X	X	X
<u>Desmidium swartzii</u> C.A. Agardh.											X					

Table 1 - continued

	1966						1967									
	J	J	A	S	O	N	F	M	A	M	J	J	A	S	O	N
<u>Dictyosphaerium pulchellum</u> Wood	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<u>Elakatothrix viridis</u> (Snow) Printz	X	X	X	X	X	X	X	X			X	X	X	X	X	
<u>Euastrum didelta</u> (Turp.) Ralfs.															X	X
<u>Euastrum pulchellum</u> De Breb.	X	X	X	X	X	X					X	X	X	X	X	X
<u>Gloeocystis gigas</u> (Kuetz.) Lagerheim	X	X	X	X	X							X	X	X	X	
<u>Gonatozygon monotaenium</u> De Bary					X											
<u>Gonatozygon pilosum</u> Wolle				X	X											
<u>Gymnozyga moniliformis</u> Ehr.														X		
<u>Hyalotheca dissiliens</u> (Smith) De Breb.		X	X		X	X					X	X	X		X	
<u>Nephrocytium limneticum</u> (Smith) Smith			X	X	X									X	X	
<u>Oocystis borgei</u> Snow	X	X	X	X			X		X	X	X	X	X	X	X	X
<u>Pediastrum boryanum</u> (Turp.) Meneghini	X	X								X	X					
<u>Pediastrum boryanum</u> var. <u>longicorne</u> Raciborski						X										
<u>Pediastrum simplex</u> var. <u>duodenarium</u> (Bailey) Rabenh.											X					
<u>Quadrigula closterioides</u> (Bohlin) Printz	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<u>Scenedesmus serratus</u> (Corda) Bohlin	X	X	X	X	X	X					X	X	X	X	X	X
<u>Sphaerocystis schroeteri</u> Chodat		X	X									X	X			
<u>Spondylosium planum</u> (Wolle) W.& G.S. West					X										X	X
<u>Staurastrum ankyroides</u> Wolle					X											
<u>Staurastrum arctiscon</u> (Ehr.) Lund.					X											
<u>Staurastrum limneticum</u> var. <u>cornutum</u> G.M. Smith				X	X	X							X	X	X	X
<u>Staurastrum pentacerum</u> (Wolle) Smith	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<u>Staurodesmus bulnheimii</u> (Racib.) Brook														X		
<u>Staurodesmus cuspidatus</u> (West) Teiling	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<u>Staurodesmus extensus</u> (Borge) Teiling	X	X	X	X	X	X	X				X	X	X	X	X	X
<u>Staurodesmus incus</u> var. <u>ralfsii</u> (West) Teiling							X	X	X	X	X					

Table 1 - continued

	1966						1967									
	J	J	A	S	O	N	F	M	A	M	J	J	A	S	O	N
<u>Staurodesmus mamillatus</u> (Nordst.) Teil.	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<u>Staurodesmus megacanthus</u> (Lund.) Thunm.	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<u>Staurodesmus subtriangularis</u> (Borge) Teil.			X	X	X	X	X	X			X	X	X	X	X	X
<u>Stylosphaeridium stipitatum</u> (Bachm.) Geitler & Gimesi			X	X	X	X						X	X	X		
<u>Tetraedron limneticum</u> Borge													X			
<u>Xanthidium antilopaeum</u> var. <u>polymazum</u> Nordst.						X										
EUGLENOPHYCEAE																
<u>Colacium arbuscula</u> Stein	X	X	X	X	X	X			X	X	X	X	X	X	X	X
<u>Colacium vesiculosum</u> Ehrenberg		X	X	X	X	X	X	X		X	X	X	X	X	X	X
XANTHOPHYCEAE																
<u>Gloeobotrys limneticus</u> (Smith) Pascher					X	X	X						X	X	X	X
<u>Ophiocytium capitatum</u> Wolle					X	X							X	X	X	
CHRYSOPHYCEAE																
<u>Chryso-sphaerella longispina</u> Lauterborn	X	X	X	X	X	X					X	X	X	X	X	X
<u>Diceras phaseolus</u> Fott	X	X			X	X					X			X	X	
<u>Dinobryon bavaricum</u> Imhof	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<u>Dinobryon crenulatum</u> W. & G.S. West	X	X			X	X				X	X			X	X	X
<u>Dinobryon cylindricum</u> Imhof	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<u>Dinobryon divergens</u> Imhof	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<u>Dinobryon sertularia</u> Ehrenberg	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<u>Dinobryon suecicum</u> Lemm.	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<u>Dinobryon vanhoeffenii</u> (Krieg.) Bachm.		X	X	X	X	X					X	X	X	X	X	X
<u>Epipyxis tabellariae</u> (Lemm.) Smith						X								X	X	X
<u>Epipyxis utriculus</u> Ehrenberg			X	X	X	X							X	X	X	

Table 1 - continued

	1966						1967									
	J	J	A	S	O	N	F	M	A	M	J	J	A	S	O	N
<u>Mallomonas acaroides</u> Perty					X	X									X	X
<u>Mallomonas fastigata</u> Zach.			X	X	X	X					X	X	X	X	X	X
<u>Mallomonas producta</u> (Zach.) Iwanoff		X	X	X							X	X	X	X		
<u>Mallomonas pseudocoronata</u> Prescott	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<u>Mallomonas tonsurata</u>																
var. <u>alpina</u> (Pasch. & Ruttn) Krieger	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<u>Mallomonas</u> sp.					X	X									X	X
<u>Synura uvella</u> Ehrenberg	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<u>Uroglenopsis americana</u> (Calkins) Lemm.	X	X	X	X							X	X	X	X	X	X
BACILLARIOPHYCEAE																
<u>Asterionella formosa</u> Hass.	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<u>Asterionella formosa</u>																
var. <u>gracillima</u> (Hantz.) Grun.							X	X	X	X						
<u>Cyclotella bodanica</u> Eulenz.	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<u>Cyclotella comta</u> (E.) Kg.	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<u>Cyclotella glomerata</u> Bachm.					X		X	X	X	X			X	X	X	X
<u>Fragilaria capucina</u> Desm.		X	X	X	X	X	X			X	X	X	X	X	X	X
<u>Fragilaria crotonensis</u> Kitton	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<u>Melosira distans</u> (E.) Kg.	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<u>Melosira islandica</u> O. Muller	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<u>Rhizosolenia eriensis</u> H.L. Smith	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<u>Synedra ulna</u> (Nitz.) Ehr.	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<u>Tabellaria fenestrata</u> (Lyngb.) Kuetz.	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<u>Tabellaria flocculosa</u> (Roth) Kuetz.	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
DINOPHYCEAE																
<u>Ceratium hirundinella</u> (Muell.) Dujardin	X	X	X	X	X	X					X	X	X	X	X	X
<u>Gonyaulax palustris</u> Lemm.							X	X	X	X	X					
<u>Gymnodinium fuscum</u> (Ehrenb.) Stein	X				X	X	X	X	X	X	X	X		X	X	X
<u>Gymnodinium palustre</u> Schilling							X	X	X							

Table 1 - continued

	1966						1967									
	J	J	A	S	O	N	F	M	A	M	J	J	A	S	O	N
<u>Gymnodinium</u> sp.							X	X								
<u>Peridinium</u> <u>cinctum</u> (Muell.) Ehren.					X	X							X	X	X	X
<u>Peridinium</u> <u>inconspicuum</u> Lemm.		X	X	X	X							X	X	X	X	
<u>Peridinium</u> <u>limbatum</u> (Stokes) Lemm.	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<u>Peridinium</u> <u>willei</u> Huitfeld-Kaas	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<u>Peridinium</u> <u>wisconsinense</u> Eddy		X	X	X	X							X	X	X	X	
CRYPTOPHYCEAE																
<u>Cryptomonas</u> <u>erosa</u> Ehren.	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<u>Cryptomonas</u> <u>erosa</u> var. <u>reflexa</u> Marsson							X	X	X		X					
<u>Cryptomonas</u> <u>marssonii</u> Skuja				X	X	X							X	X		X
<u>Cryptomonas</u> <u>ovata</u> Ehren.	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<u>Rhodomonas</u> <u>lacustris</u> Pascher & Ruttner	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X

Table 2

Summary of Major Phytoplankton Components (1×10^5 cells/l)

Newfound Lake

	1966						1967									
	J	J	A	S	O	N	F	M	A	M	J	J	A	S	O	N
<u>Aphanothece clathrata</u>	.15	.07	1.65	2.32	.55	.05	-	-	-	-	-	-	.05	.08	.17	-
<u>Gloeothece linearis</u>	.30	.18	.41	.11	.26	.04	-	-	-	-	.04	.06	.05	.06	.31	.60
<u>Gomphosphaeria lacustris</u>	.97	1.60	2.80	3.35	1.62	1.15	.40	.32	.37	.37	.49	1.55	3.27	2.32	1.35	1.95
<u>Merismopedia tenuissima</u>	-	1.13	.26	.70	.11	.03	-	-	-	-	-	.07	.05	.11	.28	.12
<u>Botryococcus braunii</u>	.12	.14	.09	.08	.04	.21	.16	.20	.10	.08	.07	.25	.09	.04	.03	.01
<u>Crucigenia rectangularis</u>	.02	.05	.05	.04	.03	.02	-	-	-	.01	.03	.04	.01	.03	.05	.03
<u>Gloeocystis gigas</u>	.14	.20	.09	.07	.01	-	-	-	-	.05	.12	.03	.23	.02	-	-
<u>Chryso-sphaerella longispina</u>	.06	1.15	2.56	.07	.01	.21	-	-	-	-	.02	.11	.04	.01	.26	-
<u>Dinobryon bavaricum</u>	.02	.47	.11	.07	.27	.25	.04	.01	.01	.04	1.47	1.24	.19	.01	.01	.01
<u>Dinobryon cylindricum</u>	.10	.26	.02	.02	.15	.79	.04	.01	.01	.01	.23	.59	.19	.02	.06	.04
<u>Dinobryon vanhoeffenii</u>	-	.11	.09	.01	-	-	-	-	-	-	-	.19	.32	.02	-	-
<u>Synura uvella</u>	-	-	-	-	-	.06	.15	.08	.11	.12	.03	-	.13	.04	.13	.02
<u>Uroglenopsis americana</u>	-	-	7.12	1.00	.08	-	-	-	-	-	-	1.15	.16	.08	-	-
<u>Asterionella formosa</u>	.16	.95	.32	.03	.14	.11	.45	.33	.46	.29	.04	.03	.22	.72	.08	.01
<u>Cyclotella bodanica</u>	.19	.11	.12	.04	.02	.06	.01	.01	.01	-	-	.03	.06	.09	.02	.02
<u>Melosira</u>	.02	.05	.04	.04	.24	.20	.20	.11	.46	.39	.05	.05	.04	.03	.02	.08
<u>Rhizosolenia eriensis</u>	.03	.58	.38	.06	.02	.01	.01	.01	.02	.04	.02	.53	.40	.06	.01	.01
<u>Tabellaria fenestrata</u>	.01	.01	.11	.12	.07	.03	-	-	.01	.08	.02	.01	.03	.02	.20	.09
<u>Cryptomonas ovata</u>	.02	.09	.23	.19	.14	.09	.07	.07	.04	.04	.02	.15	.09	.11	.06	.11
<u>Rhodomonas lacustris</u>	.01	.02	.02	.08	.57	.91	.21	.10	.26	.16	.02	.02	.06	.19	.34	.89

Table 3

Vertical Distribution of Phytoplankton - Percent of Total Standing Crop
Newfound Lake

	1966						1967									
	J	J	A	S	O	N	F	M	A	M	J	J	A	S	O	N
<u>1 meter</u>	19	15	14	19	11	13	11	11	17	13	48	21	6	13	17	11
<u>3 meters</u>	26	19	25	20	12	17	15	23	27	16	25	21	8	12	22	19
<u>5 meters</u>	27	34	28	22	16	17	32	31	12	23	16	25	15	19	20	27
<u>10 meters</u>	18	26	27	20	25	18	32	22	14	17	7	22	38	34	19	22
<u>15 meters</u>	9	5	4	9	23	20	10	13	15	16	3	10	21	21	21	13
<u>20 meters</u>	1	1	2	10	13	15	-	-	15	15	1	1	12	1	2	8

Table 4
 Summary of Reproductive Spore Periodicity

Organism	Reproductive Spore	1966					1967										
		J	J	A	S	O	N	F	M	A	M	J	J	A	S	O	N
NEWFOUND LAKE																	
<u>Anabaena flos-aquae</u>	akinetete		X	X	X	X							X	X	X	X	
<u>Dinobryon bavaricum</u>	stato-spore											X	X				
<u>Dinobryon cylindricum</u>	stato-spore					X	X					X	X				
WINNISQUAM LAKE																	
<u>Anabaena flos-aquae</u>	akinetete		X	X	X	X	X					X	X	X	X		
<u>Anabaena scheremetievi</u>	akinetete					X	X							X	X		
<u>Aphanizomenon flos-aquae</u>	akinetete														X	X	X
<u>Gloetrichia echinulata</u>	akinetete		X	X	X							X	X				
<u>Dinobryon cylindricum</u>	stato-spore									X	X	X					
<u>Melosira</u>	auxospore									X	X	X					

Table 5
Summary of Phytoplankton Indices and Proposed Trophic Status
Newfound Lake

	Yearly	Summer	Winter	Trophic Status
Chlorophycean Index	.46	.24 - .94	-	oligotrophic
Myxophycean Index	.46	.46 - 1.00	-	eutrophic
Diatom Index	.85	.85 - 1.20	1.00	eutrophic
Euglenophyta Index	.03	.03 - .05	-	oligotrophic
Compound Index	1.20	.96 - 2.55	-	meso-eutrophic

Table 6

Seasonal Occurrence of Phytoplankton - Winnisquam Lake

	1966						1967									
	J	J	A	S	O	N	F	M	A	M	J	J	A	S	O	N
CYANOPHYCEAE																
<u>Anabaena flos-aquae</u> (Lyngb.) De Breb.	X	X	X	X	X	X					X	X	X	X	X	
<u>Anabaena scheremetievi</u> Elenkin					X	X							X	X		
<u>Aphanizomenon flos-aquae</u> (L.) Ralfs	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<u>Gloeotrichia echinulata</u> (J.E. Smith) Rich.		X	X	X								X	X			
<u>Gomphosphaeria aponina</u> var. <u>delicatula</u> Virieux	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<u>Gomphosphaeria lacustris</u> var. <u>compacta</u> Lemm.				X	X	X							X	X	X	X
<u>Oscillatoria granulata</u> Gardner	X	X	X	X	X					X	X	X	X	X		
<u>Oscillatoria limnetica</u> Lemm.	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<u>Spirulina laxa</u> G.M. Smith		X	X									X	X	X		
<u>Tolypothrix lanata</u> Wartmann			X	X	X	X							X	X		
CHLOROPHYCEAE																
<u>Ankistrodesmus falcatus</u> (Corda) Ralfs		X	X	X							X	X	X	X		
<u>Ankistrodesmus falcatus</u> var. <u>acicularis</u> (A. Braun) G.S. West			X	X							X	X	X			
<u>Ankistrodesmus falcatus</u> var. <u>mirabilis</u> (West & West) G.S. West	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<u>Botryococcus braunii</u> Keutzing	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<u>Botryococcus protuberans</u> var. <u>minor</u> G.M. Smith							X	X	X	X	X					
<u>Characium curvatum</u> G.M. Smith		X	X	X									X	X		
<u>Characium gracilipes</u> Lambert			X	X									X	X		
<u>Characium limneticum</u> Lemm.		X	X	X								X	X	X		
<u>Chlorangium stentorinum</u> (Ehren.) Stein		X	X										X			
<u>Closteriopsis longissima</u> Lemm.	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X

Table 6 - continued

	1966						1967									
	J	J	A	S	O	N	F	M	A	M	J	J	A	S	O	N
<u>Closterium acerosum</u> (Shrank) Ehren.				X		X										
<u>Closterium diana</u> e Ehren.											X					
<u>Closterium kuetzingii</u> De Breb.	X	X					X	X	X	X	X	X				
<u>Closterium moniliferum</u> (Bory) Ehren.						X										
<u>Closterium parvulum</u> Naegeli								X	X							
<u>Coelastrum microporum</u> Naegeli	X	X	X	X	X	X					X	X	X	X	X	X
<u>Cosmarium contractum</u>																
<u>Var. papillatum</u> W. & G.S. West	X	X	X	X	X	X					X	X	X	X	X	X
<u>Cosmarium punctulatum</u> De Breb.				X	X	X							X	X	X	X
<u>Crucigenia rectangularis</u> (Braun) Gay					X	X						X	X			
<u>Desmidium baileyi</u> (Ralfs) Nordst.		X	X											X	X	
<u>Desmidium swartzii</u> C. A. Agardh.				X												X
<u>Dictyosphaerium ehrenbergianum</u> Naegeli					X	X								X	X	X
<u>Dictyosphaerium pulchellum</u> Wood	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<u>Dimorphococcus lunatus</u> A. Braun					X	X								X	X	X
<u>Elakatothrix gelatinosa</u> Wille	X	X	X	X	X	X					X	X	X	X		
<u>Elakatothrix viridis</u> (Snow) Printz			X	X	X	X	X	X				X	X	X	X	X
<u>Eudorina elegans</u> Ehren.	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<u>Gloeocystis ampla</u> (Kuetz.) Lagerheim				X	X	X						X	X			
<u>Gloeocystis gigas</u> (Kuetz.) Lagerheim	X	X	X	X	X	X					X	X	X	X	X	X
<u>Gloeocystis vesiculosa</u> Naegeli		X	X	X	X	X					X	X	X	X		
<u>Gonatozygon aculeatum</u> Hastings				X	X											
<u>Gonatozygon monotaenium</u> De Bary													X			
<u>Hyalotheca dissiliens</u> (Smith) De Breb.					X	X	X							X		
<u>Kirchneriella contorta</u> (Schmidle) Bohl.																X
<u>Kirchneriella lunaris</u> (Kirch.) Moebius		X	X	X	X	X						X	X	X	X	X
<u>Kirchneriella lunaris</u>																
<u>var. diana</u> e Bohl.				X									X			
<u>Kirchneriella obesa</u>																
<u>var. aperta</u> (Tell.) Brunnthaler	X	X	X	X	X	X	X	X			X	X	X	X	X	X

Table 6 - continued

	1966						1967									
	J	J	A	S	O	N	F	M	A	M	J	J	A	S	O	N
<u>Microsterias radiata</u> Hass.					X											
<u>Microsterias radiosa</u> (Lyngbye) Agardh														X		
<u>Mougeotia</u> spp. (sterile)	X	X							X	X	X	X	X			
<u>Nephrocytium ecdysiscepanum</u> W. West											X	X	X	X	X	X
<u>Nephrocytium limneticum</u> (Smith) Smith	X	X	X	X	X	X						X	X	X		
<u>Nephrocytium lunatum</u> W. West			X	X	X							X	X	X		
<u>Oedogonium</u> spp. (sterile)	X	X	X	X	X					X	X	X	X	X		
<u>Oocystis borgei</u> Snow	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<u>Oocystis elliptica</u> W. West			X	X												
<u>Oocystis lacustris</u> Chodat					X	X										
<u>Oocystis parva</u> West & West		X	X	X								X	X			
<u>Pediastrum araneosum</u> (Racib.) Smith				X										X		
<u>Pediastrum boryanum</u> (Turp.) Meneghini	X	X	X	X	X	X			X	X	X	X	X	X	X	X
<u>Pediastrum duplex</u> Meyen	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<u>Pediastrum duplex</u> var. <u>clathratum</u> (Braun) Lagerheim	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<u>Pediastrum duplex</u> f. <u>gracillimum</u> West & West		X	X	X	X	X						X	X	X	X	X
<u>Pediastrum tetras</u> (Ehrenb.) Ralfs												X	X			
<u>Pediastrum tetras</u> var. <u>tetraodon</u> (Corda) Rabenhorst		X	X	X								X	X	X	X	X
<u>Pleurotaenium trabecula</u> (Ehrenb.) Naeg.				X		X										
<u>Quadrigula closterioides</u> (Bohlin) Printz	X	X	X	X	X	X					X	X	X	X	X	X
<u>Quadrigula lacustris</u> (Chodat) Smith			X	X	X							X	X	X		
<u>Scenedesmus abundans</u> (Kirch.) Chodat		X	X										X			
<u>Scenedesmus acutiformis</u> Schroeder												X				
<u>Scenedesmus arcuatus</u> var. <u>platydisca</u> Smith	X	X	X	X	X	X					X	X	X	X	X	X
<u>Scenedesmus bijuga</u> (Turp.) Lagerheim			X	X								X				
<u>Scenedesmus brasiliensis</u> Bohlin	X	X	X	X	X						X	X	X	X	X	X

Table 6 - continued

	1966						1967									
	J	J	A	S	O	N	F	M	A	M	J	J	A	S	O	N
<u>Scenedesmus dimorphus</u> (Turp.) Kuetz.	X	X	X	X	X	X					X	X	X	X	X	X
<u>Scenedesmus opoliensis</u> Richter															X	X
<u>Scenedesmus quadricauda</u> (Turp.) De Breb.	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<u>Scenedesmus quadricauda</u> var. <u>parvus</u> Smith			X	X									X			
<u>Scenedesmus quadricauda</u> var. <u>westii</u> Smith			X	X									X	X	X	
<u>Schroederia judayi</u> Smith	X	X	X	X	X	X					X	X	X	X	X	
<u>Schroederia setigera</u> (Schroed.) Lemm.	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<u>Sorastrum spinulosum</u> Naegeli				X									X	X	X	
<u>Sphaerocystis schroeteri</u> Chodat	X	X	X	X	X			X	X	X	X	X	X			
<u>Spirogyra</u> spp. (sterile)	X	X	X	X	X	X		X	X	X	X	X	X			
<u>Spondylosium planum</u> (Wolle) W. & G.S. West	X	X	X	X							X	X	X	X	X	X
<u>Staurastrum anatinum</u> var. <u>longibrachiatum</u> W. & G.S. West			X	X	X	X	X	X	X	X	X	X	X	X	X	X
<u>Staurastrum arctiscon</u> (Ehren.) Lund											X					
<u>Staurastrum cingulum</u> (W. & G.S. West) Smith				X	X	X	X	X			X	X	X			
<u>Staurastrum johnsoni</u> W. & G.S. West	X	X	X	X	X	X	X	X			X	X	X	X	X	X
<u>Staurastrum limneticum</u> var. <u>cornutum</u> Smith			X	X	X	X							X	X		
<u>Staurastrum manfeldtii</u> Delp.			X	X	X	X										
<u>Staurastrum pilosum</u> Archer					X	X					X	X	X	X	X	
<u>Staurastrum pingue</u> Telling	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<u>Staurastrum pseudopelagicum</u> W. & G.S. West	X	X	X	X	X	X					X	X	X	X		
<u>Staurastrum vestitum</u> Ralfs				X	X									X		
<u>Staurodesmus dickiei</u> (Ralfs) Lillier.									X	X						
<u>Staurodesmus extensus</u> (Borge) Telling			X	X	X										X	X
<u>Staurodesmus incus</u> var. <u>ralfsii</u> (West) Telling							X	X	X	X	X	X				
<u>Staurodesmus mamillatus</u> (Nordst.) Teil.	X	X	X	X								X	X	X	X	X

Table 6 - continued

	1966						1967									
	J	J	A	S	O	N	F	M	A	M	J	J	A	S	O	N
<u>Staurodesmus megacanthus</u> (Lund.) Thunm.	X	X	X	X					X	X	X	X	X	X	X	X
<u>Staurodesmus subtriangularis</u> (Borge) Teil.	X	X	X	X	X	X	X			X	X	X	X	X	X	X
<u>Ulothrix subconstricta</u> G.S. West	X	X	X	X	X	X					X	X	X	X	X	
<u>Ulothrix tenerrima</u> Kuetz.	X	X	X	X	X	X			X	X	X	X	X	X		
<u>Ulothrix variabilis</u> Kuetz.		X	X								X	X	X			
<u>Xanthidium antilopaeum</u> (De Breb) Kuetz.		X			X	X								X	X	X
EUGLENOPHYCEAE																
<u>Colacium arbuscula</u> Stein		X	X								X	X	X			
<u>Colacium vesiculosum</u> Ehren.			X	X	X	X	X	X	X	X	X	X	X			
<u>Euglena spirogyra</u> Ehren.			X													
XANTHOPHYCEAE																
<u>Gloeobotrys limneticus</u> (Smith) Pascher				X	X							X	X			
<u>Stipitococcus capense</u> Prescott	X	X	X									X	X			
CHRYSOPHYCEAE																
<u>Chrysococcus</u> sp.							X	X	X							
<u>Chryso-sphaerella longispina</u> Lauterborn				X	X	X								X	X	X
<u>Dinobryon bavaricum</u> Imhof	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<u>Dinobryon cylindricum</u> Imhof	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<u>Dinobryon divergens</u> Imhof	X	X	X	X	X	X				X	X	X	X	X	X	X
<u>Dinobryon sertularia</u> Ehren.	X	X	X	X	X		X	X	X	X	X	X	X	X	X	X
<u>Epipyxis tabellariae</u> (Lemm.) Smith							X	X	X							
<u>Mallomonas elegans</u> Lemm.							X	X	X							
<u>Mallomonas fastigata</u> Zach.		X	X	X	X	X				X	X	X	X			
<u>Mallomonas producta</u> (Zach.) Iwanoff	X	X	X				X	X	X	X	X	X	X			
<u>Synura adamsii</u> Smith	X	X	X	X	X	X				X	X	X	X	X	X	X
<u>Synura uvella</u> Ehren.	X	X	X	X	X	X			X	X	X	X	X	X	X	X
<u>Uroglenopsis americana</u> (Calkins) Lemm.										X	X	X	X			

Table 6 - continued

	1966						1967									
	J	J	A	S	O	N	F	M	A	M	J	J	A	S	O	N
BACILLARIOPHYCEAE																
<u>Asterionella formosa</u> Hass.	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<u>Asterionella formosa</u> var. <u>gracillima</u> (Hantz.) Grun.							X	X	X							
<u>Cyclotella comta</u> (E.) Kg.	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<u>Cyclotella stelligera</u> Cleve & Grun.				X	X	X							X	X		
<u>Cymbella ventricosa</u>									X	X						
<u>Fragilaria capucina</u> Desm.								X	X	X	X					
<u>Fragilaria crotonensis</u> Kitton	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<u>Melosira distans</u> (E.) Kg.	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<u>Melosira italica</u> (E.) Kg.	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<u>Meridion circulare</u> var. <u>constrictum</u> (Ralfs) V.H.									X	X						
<u>Rhizosolenia eriensis</u> H.L. Smith	X						X	X	X	X	X					
<u>Synedra ulna</u> (Nitz.) Ehr.	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<u>Tabellaria fenestrata</u> (Lyngb.) Kuetz.	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<u>Tabellaria flocculosa</u> (Roth) Kuetz.									X	X	X					
DINOPHYCEAE																
<u>Ceratium hirundinella</u> (Muell.) Dujardin	X	X	X	X	X	X				X	X	X	X	X	X	X
<u>Glenodinium aciculiferum</u> (Lemm.) Lindem.								X	X	X	X					
<u>Gymnodinium</u> sp.								X	X	X	X					
<u>Peridinium cinctum</u> (Muell.) Ehren.	X	X			X	X	X	X	X	X	X		X	X	X	X
<u>Peridinium limbatum</u> (Stokes) Lemm.	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
CRYPTOPHYCEAE																
<u>Cryptomonas marssonii</u> Skuja			X	X	X	X						X	X	X	X	X
<u>Cryptomonas ovata</u> Ehren.	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<u>Rhodomonas lacustris</u> Pascher & Ruttner	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X

Table 7

Summary of Major Phytoplankton Components (1×10^5 cells/l)

Winnisquam Lake

	1966						1967									
	J	J	A	S	O	N	F	M	A	M	J	J	A	S	O	N
<u>Anabaena flos-aquae</u>	.03	1.92	5.33	.10	.01	-	-	-	-	-	.02	7.41	.16	.01	-	-
<u>Aphanizomenon flos-aquae</u>	-	-	.02	.58	5.03	.70	.17	.10	3.10	4.08	9.88	18.20	28.50	*	**	49.10
<u>Oscillatoria limnetica</u>	-	-	-	-	-	.01	.46	2.19	25.30	20.04	19.30	1.69	3.13	1.19	22.70	59.00
<u>Ankistrodesmus falcatus</u> var. <u>mirabilis</u>	.68	.02	.01	.01	.01	-	.18	.22	15.17	12.09	1.01	.21	.14	.03	.01	.01
<u>Dictyosphaerium pulchellum</u>	.42	.13	.49	2.30	.31	.02	-	-	-	.87	1.02	1.46	8.75	1.52	.22	.75
<u>Eudorina elegans</u>	.13	.52	.60	.29	.68	.22	-	-	.03	.05	.09	.15	.12	.07	.01	-
<u>Gloeoecystis vesiculosa</u>	3.77	12.50	1.32	.53	1.58	.78	-	-	-	-	.18	2.01	3.06	1.25	.18	.18
<u>Schroederia judayi</u>	.03	.03	.10	.46	.31	.28	-	-	.01	.12	.11	.05	.01	.01	-	-
<u>Staurastrum pingue</u>	.02	.09	.12	.04	.43	.76	-	-	-	-	.05	.03	.02	.02	.01	-
<u>Dinobryon cylindricum</u>	.25	.38	.05	.02	.02	.03	-	-	.12	.12	.18	.02	.02	-	-	-
<u>Synura uvella</u>	.71	.66	.09	.02	1.02	.04	-	-	-	-	.02	.19	1.08	.14	-	-
<u>Asterionella formosa</u>	.06	.07	.21	.20	.38	1.72	.44	.24	3.50	.72	.76	.05	.04	.02	.01	.04
<u>Fragilaria crotonensis</u>	.01	.01	.11	.15	.76	2.74	.06	.05	.37	.29	.13	3.50	1.47	.06	.01	.08
<u>Melosira</u>	.05	.05	.16	.61	.92	.98	3.24	2.10	10.05	8.90	.21	.05	.05	.03	.02	.17
<u>Synedra ulna</u>	.01	.02	.01	.01	.02	.09	.15	.12	.33	1.85	1.54	.47	.04	.01	.01	.02
<u>Tabellaria fenestrata</u>	.01	.01	.27	1.18	.64	.31	.01	.01	.10	.40	.80	.12	.11	.08	.08	.30
<u>Cryptomonas ovata</u>	.39	.32	.35	.38	.97	.41	.08	.02	.08	.89	.42	.47	.13	.12	.05	.28
<u>Rhodomonas lacustris</u>	1.17	.42	.54	.47	.77	.32	.04	.03	4.81	6.28	6.54	.43	.08	.04	.01	.02

* 1.37×10^8 cells/l** 1.05×10^8 cells/l


Table 8


Vertical Distribution of Phytoplankton - Percent of Total Standing Crop
Winnisquam Lake

	1966						1967									
	J	J	A	S	O	N	F	M	A	M	J	J	A	S	O	N
<u>1 meter</u>	26	27	47	26	25	16	5	20	21	23	39	45	28	31	28	38
<u>3 meters</u>	24	32	23	21	20	18	11	16	19	20	26	37	33	30	28	26
<u>5 meters</u>	23	28	18	18	21	15	37	17	18	19	19	12	19	32	23	18
<u>10 meters</u>	15	10	7	17	21	23	20	16	16	16	9	4	15	6	19	15
<u>15 meters</u>	8	2	3	9	7	15	18	16	15	11	5	1	4	1	1	2
<u>20 meters</u>	4	1	2	9	6	12	9	15	11	11	2	1	1	1	1	1

Table 9
 Summary of Phytoplankton Indices and Proposed Trophic Status
 Winnisquam Lake

	Yearly	Summer	Winter	Trophic Status
Chlorophycean Index	1.42	1.24 - 2.13	-	eutrophic
Myxophycean Index	.33	.27 - .50	-	oligo-mesotrophic
Diatom Index	.55	.44 - 1.00	.80 - 1.00	eutrophic
Euglenophyta Index	.03	.03 - .04	-	oligotrophic
Compound Index	2.00	1.60 - 2.90	-	meso-eutrophic





SEASONAL TEMPERATURE VARIATIONS - NEWFOUND LAKE




Figure 3

SEASONAL VARIATIONS IN LIGHT PENETRATION - 1% LEVEL




Figure 4




Figure 5

ORTHOPHOSPHATE - NEWFOUND LAKE




Figure 6

NITRATE NITROGEN - NEWFOUND LAKE




Figure 7

SILICON DIOXIDE - NEWFOUND LAKE




Figure 8

MAJOR PHYTOPLANKTON COMPONENTS - NEWFOUND LAKE




Figure 9

SEASONAL TEMPERATURE VARIATIONS - WINNISQUAM LAKE




Figure 10

ORTHOPHOSPHATE - WINNISQUAM LAKE




Figure 11

NITRATE NITROGEN - WINNISQUAM LAKE




Figure 12

SILICON DIOXIDE - WINNISQUAM LAKE




Figure 13




Figure 14




Figure 15