

University of New Hampshire  
University of New Hampshire Scholars' Repository

---

Life Sciences Faculty Scholarship

Life Sciences

---

5-25-2017

# Draft Genome Sequence of the Salt Water Bacterium *Oceanospirillum linum* ATCC 11336T

Ariel M. Trachtenberg  
*University of New Hampshire, Manchester*

Joshua G. Carney  
*University of New Hampshire, Manchester*

Joshua D. Linnane  
*University of New Hampshire, Manchester*

Bruce A. Rheaume  
*University of New Hampshire, Manchester*

Natalie L. Pitts  
*Colorado State University*

*See next page for additional authors*

Follow this and additional works at: [https://scholars.unh.edu/unhmbiology\\_facpub](https://scholars.unh.edu/unhmbiology_facpub)

---

## Recommended Citation

Trachtenberg, A.M., Carney, J.G., Linnane, J.D., Rheaume, B.A., Pitts, N.L., Mykles, D.L., and MacLea, K.S. Draft Genome Sequence of the Salt Water Bacterium *Oceanospirillum linum* ATCC 11336T. *Genome Announc*, 5(21):e00395-17. doi: 10.1128/genomeA.00395-17, 2017.

This Article is brought to you for free and open access by the Life Sciences at University of New Hampshire Scholars' Repository. It has been accepted for inclusion in Life Sciences Faculty Scholarship by an authorized administrator of University of New Hampshire Scholars' Repository. For more information, please contact [nicole.hentz@unh.edu](mailto:nicole.hentz@unh.edu).

---


**Authors**

Ariel M. Trachtenberg, Joshua G. Carney, Joshua D. Linnane, Bruce A. Rheaume, Natalie L. Pitts, Donald L. Mykles, and Kyle S. MacLea



# Draft Genome Sequence of the Salt Water Bacterium *Oceanospirillum linum* ATCC 11336<sup>T</sup>

Ariel M. Trachtenberg,<sup>a</sup> Joshua G. Carney,<sup>a</sup> Joshua D. Linnane,<sup>a</sup>

Bruce A. Rheume,<sup>a</sup> Natalie L. Pitts,<sup>b</sup> Donald L. Mykles,<sup>b</sup>  Kyle S. MacLea<sup>a,c,d</sup>

Biology Program, University of New Hampshire, Manchester, New Hampshire, USA<sup>a</sup>; Department of Biology, Colorado State University, Fort Collins, Colorado, USA<sup>b</sup>; Biotechnology Program, University of New Hampshire, Manchester, New Hampshire, USA<sup>c</sup>; Department of Life Sciences, University of New Hampshire, Manchester, New Hampshire, USA<sup>d</sup>

**ABSTRACT** *Oceanospirillum linum* ATCC 11336<sup>T</sup> is an aerobic, bipolar-tufted gammaproteobacterium first isolated in the Long Island Sound in the 1950s. This announcement offers a genome sequence for *O. linum* ATCC 11336<sup>T</sup>, which has a predicted genome size of 3,782,189 bp (49.13% G+C content) containing 3,540 genes and 3,361 coding sequences.

*Oceanospirillum linum* ATCC 11336<sup>T</sup> is an obligately aerobic, bipolar-tufted gamma-proteobacterium taken from the tidal estuary waters of the Long Island Sound in the United States in 1957 by Williams and Rittenberg (1). *O. linum* was combined with *Spirillum atlanticum* (ATCC 12753) and placed in *Oceanospirillum* after the split of the 1832 genus *Spirillum* Ehrenberg, and it was designated the type species for the genus (1–6). The genus currently contains four other named species: *O. multiglobuliferum*, *O. maris*, *O. beijerinckii*, and *O. niense*. Our research group sequenced *O. linum* (this paper) and *O. multiglobuliferum* (7). Putative *Oceanospirillum* strain MED92 was determined to have <93% *Oceanospirillaceae* sequence similarity at the 16S rRNA gene and was instead assigned to the new genus *Neptuniibacter* as *N. caesariensis* (8).

Unlike freshwater spirilla, *O. linum* is a halophile, capable of growing under conditions of up to 9.75% (wt/vol) (3) or 12% (8) NaCl. It is a strict aerobe, producing polyhydroxybutyrate (PHB) intracellularly, is oxidase and catalase positive, and cannot oxidize or ferment carbohydrates or break down starches, casein, or hippurate (3). *O. linum* is unique in its genus for the ability to use as a sole nitrogen source L-methionine when the cells are provided succinate plus malate as carbon sources (3). Few carbon sources are used (3, 8), although acetate may be used as a sole carbon source when ammonium ions are used as a sole nitrogen source (3).

*O. linum* ATCC 11336<sup>T</sup> was purchased from ATCC (Manassas, VA, USA) in lyophilized form, rehydrated, and cultured in marine broth or agar (ATCC medium 2216) at 28°C and atmospheric pressure for 48 h. A single colony was grown in log phase, and genomic DNA (gDNA) isolation from these bacteria was achieved using the Genomic-tip 500/G kit (Qiagen, Valencia, CA, USA). The gDNA was fragmented, tagged with adapters using the Nextera DNA library prep kit (Illumina, San Diego, CA, USA), and sequenced with an Illumina HiSeq 2500 sequencer. Two hundred fifty base pair paired-end reads were generated at the Hubbard Center for Genome Studies at the University of New Hampshire (Durham, NH, USA), and Trimmomatic was used for bioinformatic removal of adapter sequences and trimming prior to gene analysis (9).

The genome of *Oceanospirillum linum* was assembled from 12,571,740 reads into 289 contigs using SPAdes version 3.8.0 (10). These contigs were interpreted with QUAST version 4.1 to have a total length of 3,782,189 bp, a G+C% of 49.13%, and an average

Received 31 March 2017 Accepted 4 April 2017 Published 25 May 2017

**Citation** Trachtenberg AM, Carney JG, Linnane JD, Rheume BA, Pitts NL, Mykles DL, MacLea KS. 2017. Draft genome sequence of the salt water bacterium *Oceanospirillum linum* ATCC 11336<sup>T</sup>. *Genome Announc* 5:e00395-17. <https://doi.org/10.1128/genomeA.00395-17>.

**Copyright** © 2017 Trachtenberg et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Kyle S. MacLea, [Kyle.MacLea@UNH.edu](mailto:Kyle.MacLea@UNH.edu).

coverage of  $1,702\times$  (11). The largest contig found was 1,127,340 bp, with an  $N_{50}$  value of 573,653 bp. The G+C% results are in strong agreement with previous reports of G+C content of 48% (3, 12, 13) and 49% (14).

The National Center for Biotechnology Information (NCBI) automatic annotation pipeline (PGAP) was used for genome annotation (15). A total of 3,540 genes, 3,361 coding sequences (CDSs), 96 RNA genes, 83 pseudogenes, and 3 clustered regularly interspaced short palindromic repeat (CRISPR) arrays were discovered by PGAP.

**Accession number(s).** This whole-genome shotgun project has been deposited at DDBJ/EMBL/GenBank under the accession no. [MTSD00000000](https://doi.org/10.1093/genomea.00396-17). The version described in this paper is version MTSD02000000.

## ACKNOWLEDGMENTS

Sequencing was undertaken at the Hubbard Center for Genome Studies at UNH with the kind assistance of Kelley Thomas, Jordan Ramsdell, and Stephen Simpson. The following underwriters of SciFund Challenge 2 graciously and generously supported this work: Robert and Lorraine MacLea, Edwin Anderson, Jennifer MacLea, Peter Harmon, Annuka Pasi, Francis Portland, Gilbert and Jeanne Slater, and the poet K. P. Anderson. This work was a project of the Microbiology Education through Genome Annotation-New Hampshire (MEGA-NH) program.

The Biology Program at UNH Manchester provided funds for sequencing. Funding awarded to K.S.M. from SciFund Challenge 2 (2012) also contributed to the supply and sequencing costs for this project. The funders had no role in the study design, data collection and interpretation, or the decision to submit the work for publication.

## REFERENCES

- Williams MA, Rittenberg SC. 1957. A taxonomic study of the genus *Spirillum* Ehrenberg. *Int Bull Bacteriol Nomencl Taxon* 7:49–111. <https://doi.org/10.1099/0096266X-7-2-49>.
- Ehrenberg CG. 1832. Beiträge zur Kenntnis der Organization der Infusorien und ihrer geographischen Verbreitung besonders in Sibirien. *Abh K Akad Wiss Berl* 1830:1–88.
- Hylemon PB, Wells JS, Krieg NR, Jannasch HW. 1973. The genus *Spirillum*: a taxonomic study. *Int J Syst Bacteriol* 23:340–380. <https://doi.org/10.1099/00207713-23-4-340>.
- Sneath PHA, McGowan V, Skerman VBD. 1980. Approved lists of bacterial names. *Int J Syst Evol Microbiol* 30:225–420. <https://doi.org/10.1099/00207713-30-1-225>.
- Terasaki Y. 1979. Transfer of five species and two subspecies of *Spirillum* to other genera (*Aquaspirillum* and *Oceanospirillum*), with emended descriptions of the species and subspecies. *Int J Syst Bacteriol* 29:130–144. <https://doi.org/10.1099/00207713-29-2-130>.
- Pot B, Gillis M, De Ley J. 1992. The genus *Oceanospirillum*, p 3230–3236. *In* *The prokaryotes*. Springer, New York, NY.
- Carney JG, Trachtenberg AM, Rheume BA, Linnane JD, Pitts NL, Mykles DL, MacLea KS. 2017. Genome sequence of amarine spirillum, *Oceanospirillum multiglobuliferum* ATCC 33336<sup>T</sup>, isolated from Japan. *Genome Announc* 5:e00396-17. <https://doi.org/10.1128/genomeA.00396-17>.
- Arahal DR, Lekunberri I, González JM, Pascual J, Pujalte MJ, Pedrós-Alió C, Pinhassi J. 2007. *Neptuniibacter caesariensis* gen. nov., sp. nov., a novel marine genome-sequenced gammaproteobacterium. *Int J Syst Evol Microbiol* 57:1000–1006. <https://doi.org/10.1099/ijs.0.64524-0>.
- Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120. <https://doi.org/10.1093/bioinformatics/btu170>.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Pribelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
- Gurevich A, Saveliev V, Vyahhi N, Tesler G. 2013. QUASt: quality assessment tool for genome assemblies. *Bioinformatics* 29:1072–1075. <https://doi.org/10.1093/bioinformatics/btt086>.
- Satomi M, Kimura B, Hayashi M, Shouzen Y, Okuzumi M, Fujii T. 1998. *Marinospirillum* gen. nov., with descriptions of *Marinospirillum megaterium* sp. nov., isolated from kusaya gravy, and transfer of *Oceanospirillum minutulum* to *Marinospirillum minutulum* comb. nov. *Int J Syst Bacteriol* 48:1341–1348. <https://doi.org/10.1099/00207713-48-4-1341>.
- Satomi M, Kimura B, Hamada T, Harayama S, Fujii T. 2002. Phylogenetic study of the genus *Oceanospirillum* based on 16S rRNA and *gyrB* genes: emended description of the genus *Oceanospirillum*, description of *Pseudospirillum* gen. nov., *Oceanobacter* gen. nov. and *Terasakiella* gen. nov. and transfer of *Oceanospirillum jannaschii* and *Pseudomonas stanieri* to *Marinobacterium* as *Marinobacterium jannaschii* comb. nov. and *Marinobacterium stanieri* comb. nov. *Int J Syst Evol Microbiol* 52:739–747. <https://doi.org/10.1099/00207713-52-3-739>.
- Bowditch RD, Baumann L, Baumann P. 1984. Description of *Oceanospirillum kriegii* sp. nov. and *O. jannaschii* sp. nov. and assignment of two species of *Alteromonas* to this genus as *O. commune* comb. nov. and *O. vagum* comb. nov. *Curr Microbiol* 10:221–229. <https://doi.org/10.1007/BF01627259>.
- Tatusova T, DiCuccio M, Badretdin A, Chetvermin V, Ciufu S, Li W. 2013. Prokaryotic genome annotation pipeline. National Center for Biotechnology Information, Bethesda, MD.