

University of New Hampshire  
University of New Hampshire Scholars' Repository

---

Life Sciences Faculty Scholarship

Life Sciences

---

10-29-2015

# Draft Whole-Genome Sequence of the Marine Bacterium *Idiomarina zobellii* KMM 231T

Scott Mithoefer

*University of New Hampshire, Manchester*

Bruce A. Rheaume

*University of New Hampshire, Manchester*

Kyle S. MacLea

*University of New Hampshire, Manchester, [kyle.maclea@unh.edu](mailto:kyle.maclea@unh.edu)*

Follow this and additional works at: [https://scholars.unh.edu/unhmbiology\\_facpub](https://scholars.unh.edu/unhmbiology_facpub)

---

## Recommended Citation

Mithoefer, S., Rheaume, B.A., and MacLea, K.S. Draft Whole-Genome Sequence of the Marine Bacterium *Idiomarina zobellii* KMM 231T. *Genome Announc*, 3(5):e01257-15. doi: 10.1128/genomeA.01257-15, 2015.?

This Article is brought to you for free and open access by the Life Sciences at University of New Hampshire Scholars' Repository. It has been accepted for inclusion in Life Sciences Faculty Scholarship by an authorized administrator of University of New Hampshire Scholars' Repository. For more information, please contact [nicole.hentz@unh.edu](mailto:nicole.hentz@unh.edu).

# Draft Whole-Genome Sequence of the Marine Bacterium *Idiomarina zobellii* KMM 231<sup>T</sup>

Scott Mithoefer,<sup>a</sup> Bruce A. Rheaume,<sup>a</sup>  Kyle S. MacLea<sup>a,b</sup>

Biology Program, University of New Hampshire, Manchester, New Hampshire, USA<sup>a</sup>; Biotechnology Program, University of New Hampshire, Manchester, New Hampshire, USA<sup>b</sup>

***Idiomarina zobellii* was isolated from the northwest Pacific Ocean at a depth of 4,000 to 5,000 m in 1985. The draft whole-genome shotgun sequence of *I. zobellii* KMM 231<sup>T</sup> described in this paper has a predicted length of 2,602,160 bp, containing 2,570 total genes, 52 tRNAs, and a G+C content of 47.10%.**

Received 12 September 2015 Accepted 23 September 2015 Published 29 October 2015

Citation Mithoefer S, Rheaume BA, MacLea KS. 2015. Draft whole-genome sequence of the marine bacterium *Idiomarina zobellii* KMM 231<sup>T</sup>. *Genome Announc* 3(5):e01257-15. doi:10.1128/genomeA.01257-15.

Copyright © 2015 Mithoefer et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Kyle S. MacLea, [kyle.maclea@unh.edu](mailto:kyle.maclea@unh.edu).

The bacterium *Idiomarina zobellii* strain KMM 231<sup>T</sup> was originally collected in 1985 from the northwest Pacific Ocean at a depth of 4,000 to 5,000 m by Ivanova et al. (1). *I. zobellii* is a Gram-negative gammaproteobacterium bacillus, with fimbriae and a single long flagellum. It is heterotrophic and grows under strict aerobic conditions in extreme oceanic environments of low temperature, high salinity, and pressures between 397 and 497 atm (1).

*Idiomarina zobellii* strain KMM 231<sup>T</sup> was obtained from ATCC (BAA-313) as a purified, freeze-dried sample and cultured in marine broth at a temperature of 30°C for 65 h at 1 atm. Genomic DNA extraction and isolation was performed using the Qiagen Genomic-tip 500/g kit (Valencia, CA, USA). The purified genomic DNA was fragmented and marked with adapters with the Nextera DNA Library prep kit (Illumina, San Diego, CA, USA), and the sequence library of 250-bp paired-end reads was generated on a HiSeq 2500 sequencer (Illumina) at the Hubbard Center for Genome Studies at the University of New Hampshire; 44,672,634 reads were produced and trimmed using Trimmomatic (2), and sequence assembly was achieved using the SPAdes genome assembler version 3.5.0 (3), which represented a 1,212.74-fold genome coverage when the contigs were analyzed with QUAST version 2.3 (4). Analysis of the genome assembly showed a length of 2,602,160 bp in 86 contiguous segments with a G+C content of 47.10% (4). Additionally, the  $N_{50}$  for the sequence was 248,687 bp, with the largest contiguous segment being 425,886 bp (4). Annotation of the sequence was accomplished through the NCBI Prokaryotic Genome Annotation Pipeline (5), which found 2,570 total genes of which 2,440 were coding sequences, 52 were tRNAs, and the balance were pseudogenes, rRNAs, and ncRNAs. A partial gene sequence for the 16S rRNA was noted, and comparison to the NCBI database through BLAST (6) revealed 100% similarity to previously deposited 16S rRNA sequences (e.g., DDBJ/ENA/GenBank accession number NR\_024892) from *I. zobellii* strains, as well as close matches to other *Idiomarina* spp.

**Nucleotide sequence accession numbers.** This whole-genome shotgun project has been deposited in DDBJ/ENA/GenBank under the accession number [LHSG00000000](https://www.ncbi.nlm.nih.gov/nuclink/LHSG00000000). The version described in this paper is the first version, LHSG01000000.

## ACKNOWLEDGMENTS

Sequencing was undertaken at the Hubbard Center for Genome Studies at UNH with the kind assistance of Kelley Thomas, Jordan Ramsdell, and Stephen Simpson. Joseph Eager, Naida Idrizovic, Timothy Labonville, Jesse Pinard, Jessie Schofield, and Zachary Window were summer 2015 students in BSCI 737 Microbial Genomics who contributed helpful discussions to the completion of this work. This work was a project of the Microbiology Education through Genome Annotation–New Hampshire (MEGA-NH) program.

The Biological Sciences program at UNH Manchester provided funds for sequencing. The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication.

## REFERENCES

- Ivanova EP, Romanenko LA, Chun J, Matte MH, Matte GR, Mikhailov VV, Svetashev VI, Huq A, Mangel T, Colwell RR. 2000. *Idiomarina* gen. nov., comprising novel indigenous deep-sea bacteria from the Pacific Ocean, including descriptions of two species, *Idiomarina abyssalis* sp. nov. and *Idiomarina zobellii* sp. nov. *Int J Syst Evol Microbiol* 50:901–907. <http://dx.doi.org/10.1099/00207713-50-2-901>.
- Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120. <http://dx.doi.org/10.1093/bioinformatics/btu170>.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Pribelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <http://dx.doi.org/10.1089/cmb.2012.0021>.
- Gurevich A, Saveliev V, Vyahhi N, Tesler G. 2013. QUAST: quality assessment tool for genome assemblies. *Bioinformatics* 29:1072–1075. <http://dx.doi.org/10.1093/bioinformatics/btt086>.
- Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Ciufu S, Li W. 2013. Prokaryotic genome annotation pipeline. National Center for Biotechnology Information, Bethesda, MD.
- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search tool. *J Mol Biol* 215:403–410. [http://dx.doi.org/10.1016/S0022-2836\(05\)80360-2](http://dx.doi.org/10.1016/S0022-2836(05)80360-2).