

South Dakota State University
**Open PRAIRIE: Open Public Research Access Institutional
Repository and Information Exchange**

South Dakota Beef Report, 2003

Animal Science Reports

2003

Increasing the Value of the Round and Sirloin Through Pre-Rigo Skeletal Separations

Bruce C. Shanks

South Dakota State University

Duane M. Wulf

South Dakota State University

Brian J. Reuter

South Dakota State University

Robert J. Maddock

South Dakota State University

Follow this and additional works at: http://openprairie.sdstate.edu/sd_beefreport_2003


 Part of the [Animal Sciences Commons](#)

Recommended Citation

Shanks, Bruce C.; Wulf, Duane M.; Reuter, Brian J.; and Maddock, Robert J., "Increasing the Value of the Round and Sirloin Through Pre-Rigo Skeletal Separations" (2003). *South Dakota Beef Report, 2003*. Paper 8.

http://openprairie.sdstate.edu/sd_beefreport_2003/8

This Report is brought to you for free and open access by the Animal Science Reports at Open PRAIRIE: Open Public Research Access Institutional Repository and Information Exchange. It has been accepted for inclusion in South Dakota Beef Report, 2003 by an authorized administrator of Open PRAIRIE: Open Public Research Access Institutional Repository and Information Exchange. For more information, please contact michael.biondo@sdstate.edu.



Increasing the Value of the Round and Sirloin Through Pre-Rigor Skeletal Separations

Bruce C. Shanks¹, Duane M. Wulf², Brian J. Reuter¹, and Robert J. Maddock³
Department of Animal and Range Sciences

BEEF 2003 – 07

Summary

Thirty crossbred steers were utilized to explore and compare tenderness improvements in beef round and sirloin muscles resulting from various methods of pre-rigor skeletal separations. Animals were slaughtered according to industry procedures and at 60 min postmortem one of six treatments were randomly applied to each side: **A)** control, **B)** saw pelvis at the sirloin-round junction, **C)** separate the pelvic-femur joint, **D)** saw femur at mid-point, **E)** combination of B and C, and **F)** combination of B and D. After 48-h, the following muscles were excised from each side: semimembranosus (**SM**), biceps femoris (**BF-R**), semitendinosus (**ST**), and adductor (**AD**) from the round; vastus lateralis (**VL**) and rectus femoris (**RF**) from the knuckle; and gluteus medius (**GM**), biceps femoris (**BF-S**) and psoas major (**PM**) from the sirloin. Following a 10 d ageing period, samples were removed from each muscle to determine the effect of treatments on sarcomere length and Warner-Bratzler shear force. Sarcomere lengths differed between treatments for SM, AD, ST, GM, and PM. Treatment C resulted in longer sarcomeres than controls for SM, AD, and ST. All pre-rigor skeletal separation treatments yielded shorter sarcomeres for the PM as compared to controls. Warner-Bratzler shear force differed between treatments for RF, ST and PM. For RF, all treatments, except B, resulted in lower ($P < 0.05$) shear values than for controls. Treatment F resulted in higher shear force values for the PM than controls ($P < 0.05$). Also, treatments B, D, and F increased shear force of the ST relative to controls ($P < 0.05$). Correlations between sarcomere length and shear force were found to be low and quite variable between muscles. In general, treatments increased sarcomere length of several muscles from the sirloin/round region, but had mixed effects on shear force values.

Key Words: Beef, Sarcomeres, Skeletal Separation, Tenderness

Introduction

The National Beef Tenderness Survey (Morgan et al., 1991), conducted in 1990, identified problems with tenderness in beef rounds and top sirloin steaks. A follow-up study, the National Beef Tenderness Survey-1998 (Brooks et al., 2000), revealed that improvements in tenderness of retail cuts from the round were still needed. In an effort to improve tenderness, some researchers have centered on physically stretching or controlling the shortening of sarcomeres during rigor development.

Two methods that have been considered and extensively investigated include alternative suspension of carcasses (Herring et al., 1965; Hostetler et al., 1970a,b; Hostetler et al., 1971; Hostetler and Carpenter, 1972; Hostetler et al., 1972; Hostetler et al., 1973; Smith et al., 1979; Barnier and Smulders, 1994; Eikelenboom et al., 1998) and applying tension to muscles with weights or mechanical devices (Buege and Stouffer, 1974; Sonaiya and Stouffer, 1982). Even hind leg "twisting" (Oduanya and Okubanjo, 1983) has been attempted. However, these procedures have not been readily adopted by the industry.

More recently, researchers have studied pre-rigor skeletal cuts (separations) to improve beef tenderness (Cotroneo, 1992; Wang et al., 1994; Wang et al., 1996; Claus et al., 1997; Ludwig et al., 1997; Beaty et al., 1999). This procedure, sometimes referred to as the "Tendercut Process," has been tested on the longissimus muscle and on sirloin and round cuts. Researchers have found tenderness improvements in the longissimus muscle, round, and sirloin; but the greatest improvement has been shown in the longissimus muscle. Furthermore, these researchers have only reported results for one cut location in the round/sirloin region and tenderness improvements have not been reported on all of the major round and sirloin muscles. Therefore, this study was designed to explore and compare tenderness improvements in beef round and

¹ Graduate Student

² Associate Professor

³ Assistant Professor

sirloin muscles resulting from various methods of pre-rigor skeletal separations.

Materials & Methods

Carcass Treatment. Thirty crossbred steers were slaughtered in four groups (two groups per week) at the South Dakota State University Meat Laboratory according to industry procedures. Carcasses were suspended by the Achilles tendon in the common vertical position, split, and one of six treatments were randomly applied to each side: **A)** control, **B)** saw pelvis at the sirloin-round junction, **C)** separate the pelvic-femur joint, **D)** saw femur at mid-point, **E)** combination of B and C, and **F)** combination of B and D. For treatments C and E, connective tissue adjacent to the pelvic-femur joint was either left intact or completely severed for the first two and last two slaughter groups, respectively. Average time between stunning and treatment application was 60 min and ranged from 47 to 76 min.

Carcass Length Measurement. Carcass length was measured from top of the carcass rail to the most anterior point of the first cervical vertebrae prior to and immediately after treatment application and at 24-h after treatment. To compensate for the length of the trolley, 3.25 in was subtracted from each measurement. Average length of control carcasses at treatment time was 99.0 in and at 24-h was 100.7 in. Initial and total 24-h carcass length drops were calculated from carcass length measurements.

Sampling and Storage. Following a 48-h chill period in a 34°F cooler, carcasses were ribbed, and USDA yield and quality grade data were collected from right sides by experienced evaluators. At 48-h postmortem, the following muscles were excised from each side: semimembranosus (**SM**), biceps femoris (**BF-R**), semitendinosus (**ST**), and adductor (**AD**) from the round; vastus lateralis (**VL**) and rectus femoris (**RF**) from the knuckle; and gluteus medius (**GM**), biceps femoris (**BF-S**) and psoas major (**PM**) from the sirloin. Psoas major muscles were only obtained from the last two slaughter groups. Muscles were then vacuum-packaged and aged until 10 d postmortem in a 35°F cooler before being frozen and stored (-0.4°F). At a later date, whole muscles were removed from freezer storage and 1.0 in thick steaks were cut frozen on a bandsaw from similar locations within the muscles. For sarcomere length determination, a 3 to 5 g sample was removed

from frozen steaks adjacent to steaks designated for shear force. Shear force and sarcomere length samples were then individually vacuum-packaged or placed in Whirl-Pac's®, respectively, and stored (2°F) for later analysis.

Warner-Bratzler Shear Determinations. Steaks were thawed for 24-h in a 37°F cooler and then broiled on Farberware Open Hearth electrical broilers (Farberware, Bronx, NY). Steaks were turned every 4 min during broiling until an internal temperature of 160°F was reached. Internal temperature was monitored by a digital thermometer placed in the approximate geometric center of each steak. Cooked steaks were cooled to room temperature (≈72°F) before four to eight cores (0.5 in) were removed parallel to the longitudinal orientation of the muscle fibers. Individual cores were sheared once on a Warner-Bratzler shear machine. An average shear force was calculated and recorded for each steak.

Sarcomere Length Measurements. Sarcomere length was determined using a modified laser diffraction method (Cross et al., 1980). Approximately four g of tissue was cut from each frozen sample, placed into 15 to 20 ml of cold solution containing 0.25 M Sucrose and 0.002 M KCl, and homogenized until fiber separation was noted. A drop of homogenate was then placed on a slide and sarcomere lengths were measured with a He-Ne laser (Model 155A, Spectra-Physics, Inc., Mt. View, CA). Nine measurements were made per sample. Calculations were performed according to the formula by Cross et al. (1980).

Statistical Analysis. Simple descriptive statistics were computed for live weight and carcass traits to characterize the sample of animals obtained for the experiment.

Data were analyzed (SAS, 1994) as a randomized incomplete block design, with animal serving as the block (six treatments with two treatments per block). For those dependent variables where animal was not a significant ($P > 0.05$) source of variation, the animal effect was removed and data were analyzed as a completely randomized design. Least-squares means were calculated and separated for significant ($P < 0.05$) treatment effects using pair-wise *t*-tests. To examine relationships between sarcomere length and shear force, simple correlations were computed within muscles (SAS, 1994).

Results & Discussion

Mean carcass trait values (Table 1) were generally representative of the population sampled in the 1995 NBQA (Boleman et al., 1998). However, less variation existed among carcasses in this project than in the 1995 NBQA. Therefore, this group of carcasses was an excellent test sample because they were: a) representative of the industry average, and b) consistent.

Table 2 presents means for initial and total 24-h carcass length drop of treated and control sides. Treatment F resulted in the greatest initial carcass length drop (2.93 in); treatments B, D, and E were intermediate; and treatment C resulted in the least amount of initial carcass length drop (1.25 in). Subsequently, sides subjected to treatment F had the largest amount of total carcass length drop at 24-h (4.18 in). However, even though treatment D resulted in a moderate (1.60 in) amount of initial carcass length drop, total carcass length drop at 24-h was minimal (2.38 in) and not different from control sides ($P < 0.05$).

Sarcomere lengths differed between treatments for SM, AD, ST, GM, and PM (Table 3). In general, either treatments B and C individually or combined (treatment E) were the most effective at lengthening sarcomeres. For SM, treatments B, C, E, and F resulted in longer sarcomeres than controls. For AD, treatments B, C, D, and E resulted in longer sarcomeres than controls. For ST, treatment C resulted in longer sarcomeres than controls. For GM, only treatments B and E resulted in longer sarcomeres than control counterparts. Apparently, longer sarcomeres observed in the SM, AD, ST, and GM were due to stretching which resulted from the skeletal separations. Correspondingly, Beaty et al. (1999) found that the Tendercut process, which is analogous to treatment B in the current study, increased sarcomere length in the SM and ST.

All treatments yielded shorter sarcomeres in the PM muscle as compared to controls (Table 3). Herring and colleagues (1965) observed similar complexities; they discovered that horizontal placement versus conventional suspension of carcasses resulted in lengthened sarcomeres for several muscles, but considerably shortened sarcomeres for the PM. In the current study, control sides had an average sarcomere length

of 3.52 μm , versus 2.41 μm for the average of treatments B through F. Treatment D resulted in a lesser degree of sarcomere shortening as compared to the other treatments, which was likely due to the greater linear distance between the point of skeletal separation (mid-point of the femur) and the PM muscle. Thus, with treatment D, intact connective tissue and tendons associated with the PM muscle may have maintained adequate resistance, hence keeping sarcomeres from shortening as much as with other treatments. In contrast to treatment D, the posterior insertion of the PM muscle was in close proximity to the site of treatment application for B, C, E, and F. Therefore, shorter sarcomeres found in the PM for treatments B, C, E, and F were probably a result of tension release, which probably occurred when connective tissue and tendons associated with the PM muscle were severed during treatment application.

Treatments had no effect ($P > 0.05$) on sarcomere length for BF-R, VL, or RF. The lack of response observed in sarcomere length for BF-R may have reflected the anatomical location of the BF-R in relation to the treatment sites. For the VL and RF, one could speculate that substantial stretching already occurs with traditional carcass hanging procedures. Thus, the weight and angle of conventionally suspended carcasses may be more effective than pre-rigor cuts at increasing sarcomere length in these muscles. In contrast to our results, Beaty et al. (1999) found that the Tendercut process increased BF-R sarcomere length and Wang et al. (1994) found that the Tendercut process resulted in significantly longer sarcomeres for RF and VL compared to control samples. In a later study, Wang et al. (1996) also found longer sarcomeres for Tendercut treated RF and BF steaks.

For RF, all treatments, except B, resulted in lower ($P < 0.05$) shear values than for controls (Table 3). Inconsistent with our results, some researchers have indicated that a pre-rigor cut at the round/sirloin juncture, identical to our treatment B, enhanced tenderness in the RF muscle (Wang et al., 1994; Wang et al., 1996; Claus et al., 1997). Differences between treatments and controls for shear force values were not found in the present study for SM, AD, BF-R, VL, GM, or BF-S ($P > 0.05$). In agreement with our findings, Beaty et al. (1999) reported no difference in BF-R, ST, and SM

shear force between Tendercut treated and control sides. In contrast to our results, other studies have found that VL (Wang et al., 1994) and GM (Claus et al., 1997) from Tendercut-treated carcasses had lower shear force values when compared to controls. However, Wang et al. (1996) and Claus et al. (1997) discovered no improvement in Warner-Bratzler or Lee-Kramer shear values for Tendercut treated BF steaks. These authors suggested that the location of the BF relative to the treatment site was too far apart to sufficiently stretch the muscle. They also acknowledged that the amount of collagen in the BF could have masked the effect of the treatment.

In the present study, Treatment F resulted in higher shear force values for the PM than controls ($P < 0.05$). Also, treatments B, D, and F increased shear force of the ST relative to controls ($P < 0.05$). Hostetler and Carpenter (1972) showed tendencies for the PM and ST to decrease in tenderness with alternative versus conventional suspension treatments, while other muscles from the round/sirloin region remained unchanged or improved.

Locker (1960) demonstrated that as sarcomere length decreases, tenderness of muscles declines. Previous correlations between sarcomere length and shear force of several different muscles have ranged from -0.34 to -0.80 (Herring et al., 1965; Hostetler et al., 1972; Dutson et al., 1976; Wang et al., 1994). Therefore, one would have expected the muscles in this study with longer sarcomeres to have enhanced tenderness. However, only the RF, which had similar ($P > 0.05$) sarcomere lengths for control and treated sides, responded favorably in tenderness. Even more noteworthy, in the ST, treatments B, D, and F produced substantially longer sarcomeres than controls, but control muscles were more tender ($P < 0.05$). Correspondingly, Barnier and Smulders (1994) observed increases in sarcomere length for the SM, GM, ST, and BF as a result of alternative carcass positioning and Beaty et al. (1999) observed increases in sarcomere length for Tendercut treated BF, ST, and SM, but both studies reported negligible or adverse changes in tenderness.

Correlations between sarcomere length and shear force were found to be low and quite variable among muscles (Table 5). For AD, VL, RF, and PM muscles, significant ($P < 0.05$)

negative correlations (-0.26 to -0.36) were detected indicating that longer sarcomeres were associated with lower shear force values. Yet, for SM, BF-R, and GM correlations between sarcomere length and shear force were slight and not statistically different than zero ($P > 0.05$). Indeed, a positive correlation (0.26) was observed for ST indicating that longer sarcomeres were associated with higher shear force values. An earlier series of studies (Hostetler et al., 1970; Hostetler et al., 1972; Hostetler et al., 1973) established that increased sarcomere length in ST was not always associated with improved shear force and taste panel tenderness. Hostetler et al. (1973) also found that increased sarcomere length was accompanied by increased shear force in the BF, and considerable nonlinearity was found between change in sarcomere length and change in shear force for AD, GM, and PM muscles. These authors attributed the lack of tenderness improvement seen with longer sarcomeres to the amount of connective tissue present in the muscles. Another possible explanation for these findings was elucidated in a detailed experiment conducted by Marsh and Carse (1974). Marsh and Carse (1974) detected a "peak" of toughness in muscles which were held in a 25-30% extended state during rigor onset. Hostetler et al. (1972) concluded that sarcomere length is only one of many numerous factors associated with meat tenderness; our findings strongly support this presumption.

Implications

Most of the pre-rigor skeletal treatments studied increased sarcomere length of muscles from the sirloin/round region, but had mixed effects on shear force values, thus clearly demonstrating that meat tenderness is not always positively associated with sarcomere length. None of the five treatments studied appears to have practical application in their current form because they either: a) had only minimal effects on tenderness, or b) increased tenderness in some muscles while decreasing tenderness in other muscles. Because some treatments did improve tenderness in some muscles, it may be possible to modify one or more of the treatments studied in order to elicit only positive effects on tenderness and thereby increase the value of beef cuts from the round and sirloin.

Literature Cited

- Barnier, V. M. H. and F. J. M. Smulders. 1994. The effect of pelvic suspension on shear force values in various beef muscles. Proc. 40th Int. Congr. Meat Sci. Technol., The Hague, S-IVB, 5.
- Beaty, S. L., J. K. Apple, L. K. Rakes, and D. L. Kreider. 1999. Early postmortem skeletal alterations effect on sarcomere length, myofibrillar fragmentation, and muscle tenderness of beef from light-weight, Brangus heifers. *J. Muscle Foods*. 10:67-78.
- Boleman, S. L., S. J. Boleman, W. W. Morgan, D. S. Hale, D. B. Griffin, J. W. Savell, R. P. Ames, M. T. Smith, J. D. Tatum, T. G. Field, G. C. Smith, B. A. Gardner, J. B. Morgan, S. L. Northcutt, H. G. Dolezal, D. R. Gill, and F. K. Ray. 1998. National Beef Quality Audit-1995: Survey of producer-related defects and carcass quality and quantity attributes. *J. Anim. Sci.* 76:96-103.
- Brooks, J. C., J. B. Belew, D. B. Griffin, B. L. Gwartney, D. S. Hale, W. R. Henning, D. D. Johnson, J. B. Morgan, F. C. Parrish, Jr., J. O. Reagan, and J. W. Savell. 2000. National Beef Tenderness Survey-1998. *J. Anim. Sci.* 78:1852-1860.
- Buege, D. R. and J. R. Stouffer. 1974. Effects of prerigor tension on tenderness of intact bovine and ovine muscle. *J. Food Sci.* 39:396-401.
- Claus, J. R., H. Wang, and N. G. Marriott. 1997. Prerigor carcass muscle stretching effects on tenderness of grain-fed beef under commercial conditions. *J. Food Sci.* 62:1231-1243.
- Cotroneo, C. J. 1992. Selected skeletal alteration to improve beef tenderness. M. S. Thesis. Virginia Polytechnic and State University, Blacksburg, VA.
- Cross, H. R., R. L. West, and T. R. Dutson. 1980. Comparison of methods for measuring sarcomere length in beef semitendinosus. *Meat Sci.* 5:261-266.
- Dutson, T. R., R. L. Hostetler, and Z. L. Carpenter. 1976. Effect of collagen levels and sarcomere shortening on muscle tenderness. *J. Food Sci.* 41:863-866.
- Eikelenboom, G., V. M. H. Barnier, A. H. Hoving-Bolink, F. J. M. Smulders, and J. Culioli. 1998. Effect of pelvic suspension and cooking temperature on the tenderness of electrically stimulated and aged beef, assessed with shear and compression tests. *Meat Sci.* 49:89-99.
- Herring, H. K., R. G. Cassens, and E. J. Briskey. 1965. Further studies on bovine muscle tenderness as influenced by carcass position, sarcomere length, and fiber diameter. *J. Food Sci.* 30:1049-1054
- Hostetler, R. L. and Z. L. Carpenter. 1972. Temperature aging and the Texas A&M tenderstretch. *Tex. Agric. Exp. Sta. Prog. Rep. Consol. Prog. Rep.* 3111/3131. p. 69-72.
- Hostetler, R. L., B. A. Link, W. A. Landmann, H. A. Fitzhugh, Jr. 1973. Effect of carcass suspension method on sensory panel scores for some major bovine muscles. *J. Food Sci.* 38:264-267.
- Hostetler, R. L., B. A. Link, W. A. Landmann, H. A. Fitzhugh, Jr. 1972. Effect of carcass suspension on sarcomere length and shear force of some major bovine muscles. *J. Food Sci.* 37:132-135.
- Hostetler, R. L., B. A. Link, W. A. Landmann, H. A. Fitzhugh, Jr. 1971. Improving beef tenderness by preventing muscle shortening in the carcass. *Tex. Agric. Exp. Sta. Pr-2988* p. 80-83.
- Hostetler, R. L., B. A. Link, W. A. Landmann, H. A. Fitzhugh, Jr. 1970a. Influence of carcass position during rigor mortis on tenderness of beef muscles: Comparison of two treatments. *J. Anim. Sci.* 31:47-50.
- Hostetler, R. L., B. A. Link, W. A. Landmann, H. A. Fitzhugh, Jr., Z. L. Carpenter, and F. A. Orts. 1970b. Texas A&M tenderstretch beef; A method of carcass suspension for improving tenderness. *Tex. Agric. Exp. Sta. Mp 943*.
- Locker, R. H. 1960. Degree of muscular contraction as a factor in tenderness of beef. *Food Res.* 25:304-307.
- Ludwig, C. J., J. R. Claus, N. G. Marriott, J. Johnson, and H. Wang 1997. Skeletal alteration to improve beef longissimus muscle tenderness. *J. Anim. Sci.* 75:2404-2410.
- Marsh, B. B. and W. A. Carse. 1974. Meat tenderness and the sliding-filament hypothesis. *J. Fd. Technol.* 9:129-139.
- Morgan, J. B., J. W. Savell, D. S. Hale, R. K. Miller, D. B. Griffin, H. R. Cross, and S. D. Shackelford. 1991. National Beef Tenderness Survey. *J. Anim. Sci.* 69:3274-3283.
- Oduanya, S. O. and A. O. Okubanjo. 1983. A research note: Shear force values for steaks from the semitendinosus muscle of pre-rigor leg-twisted beef carcasses. *J. Food Sci.* 48:1577-1578.
- SAS. 1994. SAS User's Guide: Statistics. SAS Inst. Inc., Cary, NC.
- Smith, G. C., T. G. Jambers, Z. L. Carpenter, T. R. Dutson, R. L. Hostetler, and W. M. Oliver. 1979. Increasing the tenderness of forage-fed beef. *J. Anim. Sci.* 49:1207-1218.

- Sonaiya, E. B. and J. R. Stouffer. 1982. A Research Note: Mechanical tensioning of electrically stimulated carcasses for improved tenderness. *J. Food Sci.* 47:1010-1015.
- Wang, H., J. R. Claus, and N. G. Marriott. 1994. Selected skeletal alterations to improve tenderness of beef round muscles. *J. Muscle Foods.* 5:137-147.
- Wang, H., J. R. Claus, and N. G. Marriott. 1996. Prerigor treatment and endpoint temperature effects on U. S. choice beef tenderness. *J. Muscle Foods.* 7:45-54.

Tables

Table 1. Means, standard deviations, and minimum and maximum values for live weight and carcass traits

Trait	Mean	SD	Minimum	Maximum
Live weight, lb	1234	30	1189	1281
Hot carcass wt, lb	762	21	727	798
Adjusted fat thickness, in	0.48	0.14	0.25	0.90
Longissimus muscle area, in ²	12.5	1.2	10.7	15.3
Actual kidney, pelvic, and heart fat, %	3.5	0.7	2.4	5.1
USDA yield grade	3.3	0.7	2.0	4.9
Overall maturity ^a	154	11	130	180
Marbling score ^b	413	57	330	570

^a100 = A⁰⁰; 200 = B⁰⁰; etc.

^b300 = Slight⁰⁰; 400 = Small⁰⁰; etc.

Table 2. Least squares means for initial carcass length drop and total 24-h carcass length drop of control and treated sides

Trait	Treatment ^a						P
	A	B	C	D	E	F	
Initial carcass length drop, in	0.00 ^b	1.68 ^{cd}	1.25 ^c	1.60 ^{cd}	1.94 ^d	2.93 ^e	< 0.001
Total 24-h carcass length drop, in	1.70 ^b	2.78 ^c	2.85 ^c	2.38 ^{bc}	3.11 ^c	4.18 ^d	< 0.001

^aA = Control; B = Saw pelvis at the sirloin-round junction; C = Separate the pelvic-femur joint; D = Saw femur at the mid-point; E = Combination of B and C; F = Combination of B and D.

^{b,c,d,e}Means within a row lacking a common superscript letter differ ($P < 0.05$).

Table 3. Least squares means for sarcomere length (μm) of muscles from control and treated sides

Muscle	Treatment ^a						P
	A	B	C	D	E	F	
Round							
Semimembranosus (top round)	1.82 ^b	2.00 ^{ed}	1.91 ^{cd}	1.88 ^{cb}	2.04 ^e	1.96 ^{cde}	< 0.001
Adductor (top round)	1.88 ^b	2.02 ^{cd}	2.13 ^d	2.03 ^{cd}	2.02 ^{cd}	1.93 ^{bc}	0.005
Biceps Femoris (bottom round)	1.86	1.92	1.92	1.90	1.89	1.86	0.429
Semitendonosis (eye of round)	2.19 ^b	2.46 ^{cd}	2.19 ^b	2.39 ^c	2.45 ^c	2.54 ^d	< 0.001
Knuckle							
Vastus Lateralis (sirloin tip)	1.99	1.99	2.00	2.01	2.12	2.12	0.107
Rectus Femoris (sirloin tip)	2.26	2.46	2.40	2.28	2.37	2.49	0.192
Sirloin							
Gluteus Medius (top sirloin)	1.79 ^{bc}	1.96 ^d	1.93 ^{cd}	1.87 ^{bcd}	2.11 ^e	1.76 ^b	< 0.001
Psoas Major (tenderloin)	3.52 ^b	2.15 ^d	2.31 ^d	3.22 ^c	2.29 ^d	2.09 ^d	< 0.001

^aA = Control; B = Saw pelvis at the sirloin-round junction; C = Separate the pelvic-femur joint; D = Saw femur at the mid-point; E = Combination of B and C; F = Combination of B and D.

^{b,c,d,e}Means within a row lacking a common superscript letter differ ($P < 0.05$).

Table 4. Least squares means for shear force (lb) of cooked steaks from control and treated sides

Muscle	Treatment ^a						P
	A	B	C	D	E	F	
Round							
Semimembranosis (top round)	9.80	9.21	9.91	10.18	9.14	10.15	0.386
Adductor (top round)	9.27	9.34	9.03	9.51	8.92	9.89	0.291
Biceps Femoris (bottom round)	11.81	11.04	12.05	11.52	11.30	11.88	0.768
Semitendonosis (eye of round)	8.29 ^b	9.23 ^{cd}	8.48 ^{bc}	9.34 ^d	8.65 ^{bcd}	8.90 ^{bcd}	0.053
Knuckle							
Vastus Lateralis (sirloin tip)	10.57	11.10	10.71	10.04	10.40	9.98	0.394
Rectus Femoris (sirloin tip)	9.43 ^b	9.07 ^{bc}	7.33 ^{de}	7.62 ^{de}	8.15 ^{cd}	7.06 ^e	< 0.001
Sirloin							
Gluteus Medius (top sirloin)	8.76	7.95	7.86	7.92	8.39	8.72	0.252
Biceps Femoris (top sirloin cap)	7.02	7.17	6.73	7.11	6.91	6.75	0.878
Psoas Major (tenderloin)	6.82 ^b	7.40 ^b	7.81 ^{bc}	7.68 ^b	6.62 ^b	9.05 ^c	0.013

^aA = Control; B = Saw pelvis at the sirloin-round junction; C = Separate the pelvic-femur joint; D = Saw femur at the mid-point; E = Combination of B and C; F = Combination of B and D.

^{b,c,d,e}Means within a row lacking a common superscript letter differ ($P < 0.05$).

Table 5. Correlation coefficients between sarcomere length and shear force for different muscles

Muscle	r	P
Round		
Semimembranosis (top round)	0.11	0.418
Adductor (top round)	-0.28	0.028
Biceps Femoris (bottom round)	0.02	0.884
Semitendonosis (eye of round)	0.26	0.046
Knuckle		
Vastus Lateralis (sirloin tip)	-0.26	0.044
Rectus Femoris (sirloin tip)	-0.31	0.015
Sirloin		
Gluteus Medius (top sirloin)	-0.19	0.149
Psoas Major (tenderloin)	-0.36	0.053